



Linking the non-visual effects of light exposure with occupational health

Price, Luke L. A.; Udovii, Ljiljana; Behrens, Thomas; van Drongelen, Alwin; Garde, Anne Helene; Hogenelst, Koen; Jensen, Marie Aarrebo; Khazova, Marina; Nowak, Kamila; Rabstein, Sylvia; Romanus, Erik; Wolska, Agnieszka

Published in:
International Journal of Epidemiology

DOI:
[10.1093/ije/dyz131](https://doi.org/10.1093/ije/dyz131)

Publication date:
2019

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/)

Citation for published version (APA):
Price, L. L. A., Udovii, L., Behrens, T., van Drongelen, A., Garde, A. H., Hogenelst, K., ... Wolska, A. (2019). Linking the non-visual effects of light exposure with occupational health. *International Journal of Epidemiology*, 48(5), 1393-1397. <https://doi.org/10.1093/ije/dyz131>



Opinion

Linking the non-visual effects of light exposure with occupational health

Luke L A Price,^{1*} Ljiljana Udovičić,² Thomas Behrens,³
 Alwin van Drongelen,⁴ Anne Helene Garde,^{5,6} Koen Hogenelst,⁴
 Marie Aarrebo Jensen,^{5,6} Marina Khazova,¹ Kamila Nowak,⁷
 Sylvia Rabstein,³ Erik Romanus,² and Agnieszka Wolska⁷

¹Radiation Dosimetry Department, Public Health England (PHE), Centre for Radiation, Chemical and Environmental Hazards, Didcot, UK, ²Physical Agents Unit, Federal Institute for Occupational Safety and Health (BAuA), Dortmund, Germany, ³Center of Epidemiology, Institute for Prevention and Occupational Medicine of the German Social Accident Insurance (IPA), Institute of the Ruhr-Universität Bochum, Bochum, Germany, ⁴Healthy Living Unit, The Netherlands Organisation for Applied Scientific Research (TNO), The Hague, The Netherlands, ⁵Department of Psychosocial Work Environment, National Research Centre for the Working Environment (NFA), Copenhagen, Denmark, ⁶Department of Public Health, University of Copenhagen, Copenhagen, Denmark and ⁷Central Institute for Labour Protection - National Research Institute (CIOP-PIB), Warszawa, Poland

*Corresponding author. Radiation Dosimetry Department, Public Health England (PHE), Centre for Radiation, Chemical and Environmental Hazards, Didcot, OX11 0RQ, UK. E-mail: Luke.Price@phe.gov.uk

Editorial decision 31 May 2019; Accepted 10 June 2019

Introduction

In May 2018, the Federal Institute for Occupational Safety and Health (BAuA, Germany) hosted ‘Light and Health at Work’, a workshop concerning occupational health issues relating to non-visual effects of light of both indoor and shift workers. The agenda reflected a common interest in translational research linking laboratory findings with occupational and public health outcomes, and resulted in the founding of the European scientific network NoVEL (standing for Non-Visual Effects of Light). This article sets out the network participants’ shared goals to improve the scientific evidence about light’s non-visual effects, circadian disruption and well-being, using light exposure interventions with high quality assessment of light.

The main work conditions that impair exposure profiles that support healthy circadian regulation are daytime indoor work that reduces light exposures and night-shift work that increases light-at-night (LAN).

Daytime indoor work

Daylight reduces differences in circadian rhythm phase,¹ but indoor workers spend on average 3.5 h less outdoors on workdays.² This negative influence on circadian regulation may induce social jetlag³ and, over time, have an adverse impact on health.^{4–6} Potentially, this could be counteracted by increased indoor light levels or short-wavelength enriched light.^{7–12} However, other reviews of light interventions highlight that there is still limited evidence of the effects of daylight exposure or of increased daytime light levels at work, and studies may not have been sufficient in size, duration, follow-up or range of outcome measures to elucidate the full health impacts.^{13,14} This situation is particularly noticeable for workplace interventions and long-term health outcomes.

Night-shift work

Night work has been associated with cardiovascular disease (CVD), diabetes, impaired cognition, mental disorders

and cancers, including breast cancer.^{15–23} Several mechanisms including LAN, circadian disruption and insufficient sleep²⁴ have been put forward to link night work to adverse health effects. Light exposure interventions to mitigate the adverse effects of shift work^{25,26} have produced mixed results. Tailored interventions based on individual characteristics, such as chronotype and age, can lead to higher participation rates and better results.²⁷

To date, many studies lack a sufficiently rigorous exposure metrology and a basis in chronobiology to test these possibilities.²⁸ Chronobiological disruption from exposure to light can be masked in field studies that do not measure and allow for variations in individual exposures and/or the timing of the individual's internal clock.²⁹ There are similarities between the apparent effects of night-shift and early morning work and insufficient sleep,¹⁸ but properly testing the possible causal mechanisms presents a substantial challenge. A co-ordinated multidisciplinary approach that meets the complex demands of the field is overdue.

The sections below illustrate important specializations that can and should be considered as part of a systematic investigation into light and health at work.

Non-visual effects of light

Light is perceived through five photopigments: melanopsin, rhodopsin and three cone opsins. Intrinsically-photosensitive retinal ganglion cells (ipRGCs) respond to light directly through melanopsin, and indirectly due to signals originating from absorption of light by the other four pigments.^{30–35} Thus ipRGCs combine the photic signals in the non-visual system, signals which can either entrain or disrupt the circadian timing of genetic expression in the suprachiasmatic nuclei (SCN), which in turn regulate numerous peripheral circadian rhythms, to be found in almost all cell types. Downstream effects may depend on stimulation of just melanopsin or of any combination of the five photopigments, so light weighted for all five photopigment sensitivities should be measured (see Figure 1) and reported, to unravel the contribution of each photoreceptor type in determining non-visual responses in the brain and body.³⁵

Light exposure measurement

When measuring light exposures in occupational health studies, it is necessary to consider light level, spectrum, timing and duration of exposure and the exposure history of the participant,³⁶ preferably captured with personal dosimeters (or actigraphs) as well as *in situ* lighting measurements. Until recently, almost all field studies and many laboratory studies reported only sample measurements of illuminance and correlated colour temperature (CCT)^{7,37,38} or compared a




Figure 1. Non-visual effects of light on circadian regulation and long-term health. Top: the non-visual system comprises five different photoreceptor-based spectral sensitivities. Bottom: ipRGCs combine photic information from these five channels to drive entrainment of the central clock in the SCN. Clock signals provide circadian information to entrain downstream responses, in anticipation of daily exogenous and endogenous demands, which also act as biomarkers of subjective physiological timing. Important biomarker rhythms include plasma melatonin, cortisol and core body temperature. Short term responses, including sleep, appetite, mood and cognitive performance are highly non-linear: work and sleep schedules and social factors can also influence or disrupt these rhythms.

particular type of lamp or CCT versus another,^{7,11} or indirect versus direct lighting,³⁹ without determining actual light exposures. Relying on metrics based on human brightness perception and colour vision (i.e. photometry and colorimetry) for describing experimental light conditions is now strongly discouraged,⁴⁰ because they do not reflect non-visual effects of light.

Spectroradiometry would provide the ideal information, but spectrometers are bulky, especially compared with actigraphs incorporating light sensors. At present, the optical performance of such wearables is highly variable, and none of them effectively measures more than two of the five spectral channels recommended for research on non-visual effects of light,^{41–45} demanding careful selection and prior characterization of light dosimeters. Advances in technology may improve this situation in the future.

Shift work and light exposure metrics

Shift work characteristics including timing of shifts, shift rotation and on-off rostering all have implications for LAN and daytime light exposures that may cause regular

circadian disruption. It has been recommended that the duration, timing and intensity of light exposure at work plus number of years worked, shift pattern and rotation should be included in the description of shift work in epidemiological studies.²⁴ However, the characteristics of light exposure including LAN that may predict different health aspects are still unclear.⁴⁶ It has also been suggested that studies include data on exposures both at work and outside working hours.²⁴ The NoVEL network participants reflect an increasing number of study groups that are conducting field research in which daylight levels and light exposure information are collected continuously over 24 h each day.

To date, only a limited number of metrics directly based on the features of the non-visual system have been proposed which offer insights for circadian lighting design. The most cited metrics are 'Circadian Stimulus – CS',^{47–49} and melanopic irradiance, with or without temporal smoothing.^{35,50,51} These metrics are not fully developed, and determining the best approach will require further dedicated research data, in both the field and the laboratory. Future exposure metrics could differ with regard to different shift types or shift rotation patterns. For example, workplace light exposure data of daytime office work may include more information on sunlight exposure, including ultraviolet radiation (UVR) for vitamin D synthesis, and be concerned less with LAN.

Translational field studies

Controlled laboratory studies enable monitoring of the participants, to support them in achieving higher levels of compliance. Whereas they can simulate broad-brush aspects of work, findings obtained in laboratory settings may not always be readily generalizable, as it is difficult to introduce the same level of control over conditions in real work environments, especially for shift work.

Although field studies lack the same level of control of the environment, they provide—in contrast to laboratory work—valuable information on improving real life working conditions. Participants in field studies are exposed to individual variables such as personal habits (e.g. the quantity of sleep and commuting). They follow everyday routines (e.g. housekeeping, preparing food, taking care of children and pets). Group level variables may include socioeconomic links to light pollution (living in the city), cultural aspects (e.g. as reflected in architecture), season, latitude etc. Such uncontrollable exposures and complex behaviours are part of everyone's lives, as well as being unavoidable aspects of working out of phase with the light cycle. This inherent unpredictability should always be taken into account when estimating the effectiveness of interventions in real-life settings.

Circadian biomarkers

Measurement of circadian rhythms requires the use of biomarkers. Salivary cortisol provides a good indication of plasma levels,⁵² whereas a recent study questioned whether the same is true of salivary melatonin,⁵³ possibly due to additional melatonin produced in the salivary glands. Alternatively, field researchers looking for more reliable data on nocturnal melatonin amplitude might consider analysis of urine samples.⁵⁴

Participants in field studies of workers are primarily healthy adults, for whom the variations in biomarkers can be relatively small.⁵⁵ Hence it is crucial to use precise and sensitive methods under analytical control, based on biomarkers that are stable during handling and shipping.⁵⁶

Cognitive performance testing and ecological momentary assessment (EMA)

To capture feelings, thoughts and behaviour as they evolve during a real-life intervention trial, ecological momentary assessment (EMA; or experience sampling methodology) can be used.⁵⁷ EMA has several advantages to using questionnaires before and after a trial, including greater ecological validity with minimal retrospective bias. The frequency of assessments provides higher sensitivity to intervention effects and data concerning patterns in daily life.⁵⁸

EMA is well suited to assess the non-visual effects of light on mental health, such as mood disorders.⁵⁹ The effects of exposure conditions and disruption of the circadian rhythm on cognition processes, such as processing speed, alertness, attentional control, inhibition, planning and, short-term and working memory, should also be considered.^{21,23,60–62} Associations linking both social interaction and mood to changing light exposure have been shown using EMA coupled with a wearable light dosimeter.⁶³ EMA methodology may be integrated within interventions, often referred to as EMA/I, where feedback or advice is presented based on an individual's responses to the EMA questions.⁶⁴

Summary

The impact of light exposures on the health and well-being of shift workers and indoor daytime workers has not been explored in sufficient detail. Advice to promote positive health outcomes for these groups should be based on evidence taken from, or validated by, well-designed field studies into the effects of light exposure. These effects are not just driven by artificial lighting in the workplace, but may depend on exposures to daylight and the influence working hours have on 24-h exposures and workers' lifestyles.

To obtain the best foundation for future health advice on optimal light conditions at work and light exposures on working days, there is a need to develop better field studies on circadian rhythms and other non-visual effects of light, combining diverse techniques and areas of expertise, including:

- experimental designs to unravel causality for long-term health outcomes, including detailed descriptions of shift characteristics;
- accurate personal light exposure measurements of the five recommended spectrally-weighted channels, to reveal the aspects of light ultimately affecting health;
- robust validated biomarkers to measure diurnal rhythms, and high-quality cognitive performance techniques, including interviews concerning chronometry and organizational aspects of the workplace; and
- EMA to monitor changes in behaviour and well-being on an ultradian scale, together with input from participants and wider stakeholders, which inform the deployment of individualized and appropriately timed interventions.

To effectively tackle the occupational and public health challenges of light exposure, particularly in shift work, we call for a greater use of tailored multidisciplinary prospective approaches with sufficient follow-up to address long-term consequences of lifestyle-like interventional changes.

Conflict of interest: None declared.

References

1. Wright KP Jr, McHill AW, Birks BR, Griffin BR, Rusterholz T, Chinoy ED. Entrainment of the human circadian clock to the natural light-dark cycle. *Curr Biol* 2013;**23**:1554–58.
2. Baczyńska KA, Khazova M, O'Hagan JB. Sun exposure of indoor workers in the UK – survey on the time spent outdoors. *Photochem Photobiol Sci* 2019;**18**:120–28.
3. Roenneberg T, Allebrandt KV, Meroow M, Vetter C. Social jet-lag and obesity. *Curr Biol* 2012;**22**:939–43.
4. Antunes LC, Levandovski R, Dantas G, Caumo W, Hidalgo MP. Obesity and shift work: chronobiological aspects. *Nutr Res Rev* 2010;**23**:155–68.
5. Kivimaki M, Batty GD, Hublin C. Shift work as a risk factor for future type 2 diabetes: evidence, mechanisms, implications, and future research directions. *PLoS Med* 2011;**8**:e1001138.
6. Morris CJ, Yang JN, Garcia JI, Myers S, Bozzi I, Wang W. Endogenous circadian system and circadian misalignment impact glucose tolerance via separate mechanisms in humans. *Proc Natl Acad Sci U S A* 2015;**112**:E2225–34.
7. Viola AU, James LM, Schlangen LJ, Dijk DJ. Blue-enriched white light in the workplace improves self-reported alertness, performance and sleep quality. *Scand J Work Environ Health* 2008;**34**:297–306.
8. Vetter C, Juda M, Lang D, Wojtysiak A, Roenneberg T. Blue-enriched office light competes with natural light as a zeitgeber. *Scand J Work Environ Health* 2011;**37**:437–45.
9. Figueiro MG, Steverson B, Heerwagen J, Kampschroer K, Hunter CM, Gonzales K. The impact of daytime light exposures on sleep and mood in office workers. *Sleep Health* 2017;**3**:204–15.
10. Hubalek S, Brink M, Schierz C. Office workers' daily exposure to light and its influence on sleep quality and mood. *Light Res Technol* 2010;**42**:33–50.
11. Mills PR, Tomkins SC, Schlangen LJ. The effect of high correlated colour temperature office lighting on employee wellbeing and work performance. *J Circadian Rhythms* 2007;**5**:2.
12. Canazei M, Pohl W, Bliem HR, Weiss EM. Acute effects of different light spectra on simulated night-shift work without circadian alignment. *Chronobiol Int* 2017;**34**:303–17.
13. Sack RL, Auckley D, Auger RR *et al*. Circadian rhythm sleep disorders: part I, basic principles, shift work and jet lag disorders. An American Academy of Sleep Medicine review. *Sleep* 2007;**30**:1460–83.
14. Pachito DV, Eckeli AL, Desouky AS *et al*. Workplace lighting for improving alertness and mood in daytime workers. *Cochrane Database Syst Rev* 2018;**3**:CD012243.
15. Cordina-Duverger E, Menegaux F, Popa A *et al*. Night shift work and breast cancer: a pooled analysis of population-based case-control studies with complete work history. *Eur J Epidemiol* 2018;**33**:369–79.
16. Vyas MV, Garg AX, Iansavichus AV *et al*. Shift work and vascular events: systematic review and meta-analysis. *BMJ* 2012;**345**:e4800.
17. Behrens T, Rabstein S, Wichert K *et al*. Shift work and the incidence of prostate cancer: a 10-year follow-up of a German population-based cohort study. *Scand J Work Environ Health* 2017;**43**:560–68.
18. Kecklund G, Axelsson J. Health consequences of shift work and insufficient sleep. *BMJ* 2016;**355**:i5210.
19. Ijaz S, Verbeek J, Seidler A *et al*. Night-shift work and breast cancer - a systematic review and meta-analysis. *Scand J Work Environ Health* 2013;**39**:431–47.
20. Gan Y, Yang C, Tong X *et al*. Shift work and diabetes mellitus: a meta-analysis of observational studies. *Occup Environ Med* 2015;**72**:72–78.
21. Rouch I, Wild P, Ansiau D, Marquie JC. Shiftwork experience, age and cognitive performance. *Ergonomics* 2005;**48**:1282–93.
22. Marquie JC, Tucker P, Folkard S, Gentil C, Ansiau D. Chronic effects of shift work on cognition: findings from the VISAT longitudinal study. *Occup Environ Med* 2015;**72**:258–64.
23. Titova OE, Lindberg E, Elmståhl S, Lind L, Schiöth HB, Benedict C. Association between shift work history and performance on the trail making test in middle-aged and elderly humans: the EpiHealth study. *Neurobiol Aging* 2016;**45**:23–29.
24. Stevens RG, Hansen J, Costa G *et al*. Considerations of circadian impact for defining 'shift work' in cancer studies: IARC Working Group Report. *Occup Environ Med* 2011;**68**:154–62.
25. Neil-Sztramko SE, Pahwa M, Demers PA, Gotay CC. Health-related interventions among night shift workers: a critical review of the literature. *Scand J Work Environ Health* 2014;**40**:543–56.
26. Pallesen S, Bjorvatn B, Mageroy N, Saksvik IB, Waage S, Moen BE. Measures to counteract the negative effects of night work. *Scand J Work Environ Health* 2010;**36**:109–20.
27. van Drongelen A, Boot CR, Hlobil H, Twisk JW, Smid T, van der Beek AJ. Evaluation of an mHealth intervention aiming to

- improve health-related behavior and sleep and reduce fatigue among airline pilots. *Scand J Work Environ Health* 2014;**40**:557–68.
28. Erren TC, Morfeld P, Stork J *et al*. Shift work, chronodisruption and cancer? The IARC 2007 challenge for research and prevention and 10 theses from the Cologne Colloquium 2008. *Scand J Work Environ Health* 2009;**35**:74–79.
 29. Saksvik IB, Bjorvatn B, Hetland H, Sandal GM, Pallesen S. Individual differences in tolerance to shift work - a systematic review. *Sleep Med Rev* 2011;**15**:221–35.
 30. Melyan Z, Tarttelin EE, Bellingham J, Lucas RJ, Hankins MW. Addition of human melanopsin renders mammalian cells photoreceptive. *Nature* 2005;**433**:741–45.
 31. Qiu X, Kumbalasing T, Carlson SM *et al*. Induction of photosensitivity by heterologous expression of melanopsin. *Nature* 2005;**433**:745–49.
 32. Drouyer E, Rieux C, Hut RA, Cooper HM. Responses of suprachiasmatic nucleus neurons to light and dark adaptation: relative contributions of melanopsin and rod-cone inputs. *J Neurosci* 2007;**27**:9623–31.
 33. Guler AD, Ecker JL, Lall GS *et al*. Melanopsin cells are the principal conduits for rod-cone input to non-image-forming vision. *Nature* 2008;**453**:102–05.
 34. Lall GS, Revell VL, Momiji H *et al*. Distinct contributions of rod, cone, and melanopsin photoreceptors to encoding irradiance. *Neuron* 2010;**66**:417–28.
 35. Lucas RJ, Peirson SN, Berson DM *et al*. Measuring and using light in the melanopsin age. *Trends Neurosci* 2014;**37**:1–9.
 36. International Commission on Illumination (CIE). *CIE 218: 2016 Research Roadmap for Healthful Interior Lighting Applications*. Vienna: CIE, 2016.
 37. Shamsul BMT, Sia CC, Ng YG, Karmegan K. Effects of light's colour temperatures on visual comfort level, task performances, and alertness among students. *Am J Public Health Res* 2013;**1**:159–65.
 38. Lavoie S, Paquet J, Selmaoui B, Rufiange M, Dumont M. Vigilance levels during and after bright light exposure in the first half of the night. *Chronobiol Int* 2003;**20**:1019–38.
 39. Fostervold KI, Nersveen J. Proportions of direct and indirect indoor lighting—the effect on health, well-being and cognitive performance of office workers. *Light Res Technol* 2008;**40**:175–200.
 40. International Commission on Illumination (CIE). *CIE S 026/E: 2018 System for Metrology of Optical Radiation for ipRGC-influenced Responses to Light*. Vienna: CIE, 2018.
 41. Price LLA, Khazova M, O'Hagan JB. Performance assessment of commercial circadian personal exposure devices. *Light Res Technol* 2012;**44**:17–26.
 42. Figueiro MG, Hamner R, Bierman A, Rea MS. Comparisons of three practical field devices used to measure personal light exposures and activity levels. *Light Res Technol* 2013;**45**:421–34.
 43. Markvart J, Hansen ÅM, Christoffersen J. Comparison and correction of the light sensor output from 48 wearable light exposure devices by using a side-by-side field calibration method. *LEUKOS* 2015;**11**:155–71.
 44. Price LL, Lyachev A, Khazova M. Optical performance characterization of light-logging actigraphy dosimeters. *J Opt Soc Am A* 2017;**34**:545–57.
 45. Price LLA, Lyachev A. Modification of a personal dosimetry device for logging melanopic irradiance. *Light Res Technol* 2017;**49**:922–27.
 46. Rabstein S, Behrens T, Bruning T. Light exposure and melatonin among rotating shift nurses - letter. *Cancer Epidemiol Biomarkers Prev* 2012;**21**:557.
 47. Rea MS, Figueiro MG, Bierman A, Bullough JD. Circadian light. *J Circadian Rhythms* 2010;**8**:2.
 48. Rea M, Figueiro M. Light as a circadian stimulus for architectural lighting. *Light Res Technol* 2018;**50**:497–510.
 49. Figueiro MG, Nagare R, Price L. Non-visual effects of light: How to use light to promote circadian entrainment and elicit alertness. *Light Res Technol* 2018;**50**:38–62.
 50. Al Enezi J, Revell V, Brown T, Wynne J, Schlangen L, Lucas R. A “melanopic” spectral efficiency function predicts the sensitivity of melanopsin photoreceptors to polychromatic lights. *J Biol Rhythms* 2011;**26**:314–23.
 51. Price L. On the role of exponential smoothing in circadian dosimetry. *Photochem Photobiol* 2014;**90**:1184–92.
 52. El-Farhan N, Rees DA, Evans C. Measuring cortisol in serum, urine and saliva - are our assays good enough? *Ann Clin Biochem* 2017;**54**:308–22.
 53. van Faassen M, Bischoff R, Kema IP. Relationship between plasma and salivary melatonin and cortisol investigated by LC-MS/MS. *Clin Chem Lab Med* 2017;**55**:1340–48.
 54. Leung M, Tranmer J, Hung E, Korsiak J, Day AG, Aronson KJ. Shift work, chronotype, and melatonin patterns among female hospital employees on day and night shifts. *Cancer Epidemiol Biomarkers Prev* 2016;**25**:830–38.
 55. Hansen AM, Høgh A, Persson R. Frequency of bullying at work, physiological response, and mental health. *J Psychosom Res* 2011;**70**:19–27.
 56. Kamel A, Prakash C. High performance liquid chromatography/atmospheric pressure ionization/tandem mass spectrometry (HPLC/API/MS/MS) in drug metabolism and toxicology. *Curr Drug Metab* 2006;**7**:837–52.
 57. Shiffman S, Stone AA, Hufford MR. Ecological momentary assessment. *Annu Rev Clin Psychol* 2008;**4**:1–32.
 58. Smyth JM, Stone AA. Ecological momentary assessment research in behavioral medicine. *J Happiness Stud* 2003;**4**:35–52.
 59. aan het Rot M, Hogenelst K, Schoevers RA. Mood disorders in everyday life: a systematic review of experience sampling and ecological momentary assessment studies. *Clin Psychol Rev* 2012;**32**:510–23.
 60. Bjorvatn B, Stangenes K, Øyane N *et al*. Subjective and objective measures of adaptation and readaptation to night work on an oil rig in the North Sea. *Sleep* 2006;**29**:821–29.
 61. Ozdemir PG, Selvi Y, Ozkol H *et al*. The influence of shift work on cognitive functions and oxidative stress. *Psychiatry Res* 2013;**210**:1219–25.
 62. Kazemi R, Haidarimoghadam R, Motamedzadeh M, Golmohamadi R, Soltanian A, Zoghhipaydar MR. Effects of shift work on cognitive performance, sleep quality, and sleepiness among petrochemical control room operators. *J Circadian Rhythms* 2016;**14**:1.
 63. aan het Rot M, Moskowitz DS, Young SN. Exposure to bright light is associated with positive social interaction and good mood over short time periods: a naturalistic study in mildly seasonal people. *J Psychiatr Res* 2008;**42**:311–19.
 64. Heron KE, Smyth JM. Ecological momentary interventions: incorporating mobile technology into psychosocial and health behaviour treatments. *Br J Health Psychol* 2010;**15**(Pt 1):1–39.