

ABSTRACT BOOK



8th Congress of European Microbiologists

In collaboration with  sfam
society for
applied microbiology



7-11 July 2019 | Glasgow, Scotland | www.fems2019.org



TABLE OF CONTENT

Oral presentations (O001 – O302).....	1
Poster Presentations (PM001 – PW498).....	301
Monday (PM001 – PM408).....	301
Tuesday (PT001 – PT412).....	714
Wednesday (PW001- PW498).....	1131

PW408 Presence of non-typhoidal Salmonella adapted to diverse stresses in raw chicken meat, Portugal

Sofia Ribeiro¹, Andreia Rebelo^{1;2}, Joana Vanessa Cordeiro Melro Mourão¹, Carla Novais¹, Luisa Maria Vieira Peixe¹, Patricia Antunes^{1;3}

¹UCIBIO/REQUIMTE, Microbiology Laboratory, Faculty of Pharmacy, University of Porto, Portugal

²School of Health, Polytechnic Institute of Porto, Portugal

³Faculty of Nutrition and Food Science, University of Porto, Porto, Portugal

Background: *Salmonella* causes frequently foodborne infections mainly associated with eggs/poultry products. A decreasing trend in human salmonellosis, particularly of *S. Enteritidis*, has been observed in EU due to successful control programs at the avian production level. However, the effectiveness of control practices on the elimination of *Salmonella*, and particularly of EU targeted serotypes, in poultry has been scarcely explored

Objectives: Here, we investigated the presence of *Salmonella*, and their clinically-relevant serotypes, among fresh chicken-meat samples at final processing level.

Methods: Pooled chicken-meat samples (n=53; each sample=25g of neck skin from 10 carcasses; 29 producers) were collected in 2018 (spring/summer). Samples were processed following ISO-6579-1:2017 cultural standard and a molecular approach (PCR-*invA* gene) applied directly in the pre-enrichment/enrichments broths. Search of EU targeted serotypes (*Enteritidis*/*Typhimurium*/4,5,12:i:-) and their antibiotic/metal resistance markers were performed by PCR. Ability to survive/grow at acidic pH (2-6.5) was evaluated by broth-microdilution.

Results: *Salmonella* was detected in 4%-two samples of fresh chicken-meat from different poultry-farms/both seasons (cultural+molecular approaches). The isolates belonged to a non-H₂S-producing serotype *S.4,5,12:i:-* (n=6 isolates/spring sample), with the typical antibiotic/metal resistance genotypic features of clones currently circulating in Europe, or *S. Enteritidis* (n=3 isolates/summer sample). Isolates grew at minimal pH=4-4.5 and survived until pH=3.5-4. A low occurrence of *Salmonella* in chicken carcasses was observed, although with detection of two serotypes of public health significance with ability to grow under diverse stresses. Our results alert for the need to evaluate current biosafety measures to prevent the spread of these pathogens in the poultry production through the consumer.