

Influence of cutting height on seasonal composition of *Moringa oleifera* in the rainforest zone of Nigeria

AA Fadiyimu^A, JA Alokun^B, AN Fajemisin^B and RD Aladesanwa^C

^A Department of Animal Production, Federal College of Agriculture, Akure, Nigeria

^B Department of Animal Production and Health

^C Department of Crop and Pest Management, Federal University of Technology, Akure, Nigeria

Contact email: yemifadiyimu@yahoo.com

Keywords: *Moringa oleifera*, rainforest zone, seasonal composition.

Introduction

Moringa oleifera, a native of sub-Himalayan regions of northwestern India, is now indigenous to many countries in Africa. Matured leaves and young branches of *Moringa* are potentially useful livestock fodder, for ruminants (Akinbamijo *et al.* 2004) and non-ruminants (Ly *et al.* 2001). However, in Nigeria it has been grossly underexploited as it is restricted to the arid northern zone where it is used mainly as live fence and as vegetable salad. The study was initiated because not enough information on the nutritive value of *Moringa* foliage and its change with season, harvesting or grazing had been accumulated in Nigeria especially in the rainforest zone.

Materials and Methods

The experiment was carried out in Akure (7°16'N, 5°12'E) in the tropical rainforest zone of Nigeria. The zone is characterized by two distinct seasons: wet or rainy (April to November) and dry season (December to March). The experiment was laid out as a randomized complete block design into which was fitted a 2×3 (cutting season versus cutting height) factorial arrangement. The cutting heights were 50, 100 and 150 cm. Seedlings of *Moringa oleifera* were transplanted on field plots at a spacing of 40 × 40 cm in August during the peak of the rainy season. Each treatment was replicated on twelve 4 m² plots. After twelve weeks of establishment (just before flowering stage), the plant foliage were cut back to 50, 100 or 150 cm above the ground level and thereafter they were not pruned to these respective heights until the next flowering stage i.e. every twelve weeks. Fodder yield for the dry season treatment was recorded from November to March and that for the wet season from April to October.

Samples of leaves and tender stems of *Moringa* from each treatment were collected separately over a period of one year and divided into dry and wet season samplings. They were dried in a forced air oven at 60°C for 48 hours, ground to pass through 1mm sieve and analyzed for dry matter, crude protein, ether extract and ash (AOAC 1995), Neutral Detergent Fiber (NDF), Acid Detergent Fiber (ADF) and Acid Detergent Lignin (ADL) according to the procedures of Van Soest *et al.* (1991). Na and K were determined by flame photometry

while P was determined by the vanado-molybdate method. Ca and Fe were determined after wet digestion with a mixture of nitric acid, sulphuric acid and perchloric acid using atomic absorption spectrophotometer (AOAC 1995). The data were subjected to analysis of variance using the general linear model (GLM) procedure of MINITAB (2000). Means were separated by Least Significant Difference (LSD).

Results

The significantly higher CP content of *M. oleifera* in the wet season than that in the dry season confirms earlier report by Jung and Vogel (1992) that tropical forage plants are highly palatable and nutritious during the rainy season but as the dry season approaches the nutritive value (particularly CP content) decreases (Table 1). However, the extent of this reduction seems to be worse for herbaceous forages than for shrubs and trees as the latter still retain enough nutrients to meet both the maintenance and production requirements of grazing ruminants during the dry months (Dicko and Sikena 1992).

NDF and ADF contents of *M. oleifera* at 150 cm height of pruning were significantly higher than at lower cutting heights probably due to greater residual carbohydrate reserves at the higher cutting heights (Stur *et al.* 1998). The lower Ca and Na contents and higher P content during the wet season are consistent with earlier findings (Kallah *et al.* 1999).

Conclusion

From the result of this experiment, a cutting height of 100 to 150 cm is recommended when *M. oleifera* is cultivated for fodder production.

References

- Akinbamijo OO, Adediran SA, Nouala S, Saecker J (2004) *Moringa* fodder in ruminant nutrition in the Gambia. International Trypanotolerance Centre, Banjul, the Gambia. www.moringanews.org/documents/Fodder.doc
- AOAC (1995) Official methods of analysis of the Association of Official Analytical Chemists. 16th ed. Arlington, Virginia, USA.
- Dicko MS, Sikena LK (1992) Fodder trees and shrubs in range and farming systems in dry tropical Africa. In 'Legume

Table 1. Effect of cutting height, season and their interaction on nutrient compositions (%) of *M. oleifera*.

Treatments	DM	CP	ASH	EE	NDF	ADF	ADL	Ca	P	Na	K	Fe	
Height (cm)													
50	20.75	26.98	8.28	3.96	23.76ab	19.03b	4.98	3.06	0.12	0.26	2.55	0.14	
100	21.29	27.18	8.44	4.36	22.67b	18.42b	4.29	2.68	0.11	0.29	2.83	0.12	
150	21.84	27.05	8.79	4.53	25.90a	22.94a	4.98	2.47	0.12	0.29	2.96	0.15	
Season													
Wet	21.02	30.14a	8.92	4.07	23.81	20.36	3.88b	2.30b	0.14a	0.26b	2.76	0.13	
Dry	21.57	24.00b	8.09	4.50	24.41	19.90	5.53a	3.17a	0.09b	0.30a	2.80	0.14	
Height × Season													
50	Wet	19.96	29.56	8.66	3.64	23.21	17.67	4.17	2.46	0.15	0.24	2.63	0.14
	Dry	21.55	24.40	7.90	4.29	24.31	20.40	5.80	3.66	0.10	0.29	2.47	0.14
100	Wet	21.43	30.22	8.86	4.27	22.75	20.08	3.38	2.40	0.14	0.28	2.79	0.13
	Dry	21.15	24.14	8.02	4.45	22.59	16.76	5.20	2.97	0.09	0.29	2.87	0.11
150	Wet	21.66	30.65	9.24	4.31	25.48	23.33	4.17	2.04	0.14	0.27	2.85	0.13
	Dry	22.02	23.44	8.35	4.76	26.32	22.55	5.59	2.90	0.09	0.32	3.07	0.17
Overall Mean		21.29	27.07	8.50	4.29	24.11	20.13	4.72	2.74	0.12	0.28	2.78	0.14
Statistical significance													
Height	NS	NS	NS	NS	*	*	NS	NS	NS	NS	NS	NS	
Season	NS	***	NS	NS	NS	NS	*	**	***	*	NS	NS	
Height × Season	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	

Means within the same column with different letters are significantly different; NS = Not Significant.

- trees and other fodder trees as protein sources for livestock'. (Eds A Speedy, P Pugliese) FAO Animal Production and Health Paper 102. (FAO Rome, Italy)
- Jung HJ, Vogel KP (1992) Lignification of switchgrass and big bluestem plant parts during maturation and its effect on fibre degradability. *Journal of Science of Food and Agriculture* **59**, 169.
- Kallah MS, Muhammad IR, Baba M, Lawal R (1999) The effect of maturity on the composition of hay and silage made from Columbus grass (*Sorghum almum*) grown in Nigeria. *Tropical Grassland*, **33**, 46-50.
- Ly J, Pok Samkol, Preston PR (2001) Nutritional evaluation of tropical leaves for pigs: Pepsin/pancreatin digestibility of thirteen plant species. *Livestock Research for Rural Development* **13**(5) <http://www.lrrd.org/lrrd13/5/ly135.htm>
- MINITAB (2000) Minitab Statistical Software, Release 10.2. Minitab Inc., State College, PA, USA.
- Stur WW, Shelton HM, Gutteridge RC (1998) Defoliation management of forage tree legumes. In 'Forage tree legumes in tropical agriculture'. (Eds RC Gutteridge, HM Shelton) (Tropical Grassland Society of Australia Inc) <http://www.fao.org/ag/AGP/AGPC/doc/Publicat/Gutt-shel/x5556e0h.htm>
- Van Soest PJ, Robertson JB, Lewis BA (1991) Methods for dietary fibre, neutral detergent fibre and non-starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* **74**, 3583-3597.