



Garbage in the diet of carnivores in an agricultural area

Łukasz Jankowiak¹, Anna W. Malecha², Agata J. Krawczyk³

¹Department of Vertebrate Zoology and Anthropology, Institute for Research on Biodiversity, University of Szczecin, Wąska 13, PL-71-415 Szczecin, Poland, Corresponding author, E-mail: jankowiakl@gmail.com

²Institute of Zoology, Poznań University of Life Sciences, Wojska Polskiego 71C, 60-625 Poznań, Poland

³Department of Systematic Zoology, Institute of Environmental Biology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland

ABSTRACT

Human food waste is considered to be richer in carbohydrates, lipids and proteins than most natural food supplies; however, it is very well digested in scats. So, as an indication of this kind of food in the diet, we have used each indigestible, anthropogenic origin element found in faeces (e.g., glass, plastic, rubber, etc.). There are few studies discussing the importance of garbage in the diet of mammalian predators living in farmland; definitely, most focus on this issue in urban areas. We studied the contribution of garbage in the diet of raccoon dog (*Nyctereutes procyonoides*), red fox (*Vulpes vulpes*), marten (*Martes* sp.), polecat (*Mustela putorius*), stoat (*Mustela erminea*), American mink (*Neovison vison*) and Eurasian otter (*Lutra lutra*) in the agricultural areas of western Poland in 2006-2010. In addition, we examined the spatial changes in the diet of red fox and polecat. The largest contribution of garbage was found in scats of raccoon dog (8.8%), red fox (4.8%) and marten (4.3%). The diet of polecat, stoat and Eurasian otter contained 2.5%, 1.7% and 0.2% garbage items respectively. The most frequent item was plastic. Our analysis showed that garbage consumption by red fox and polecat was greater closer to human settlements. The results reveal a continuous gradient in the garbage consumption that corresponds with the degree of synanthropization of particular species.

KEYWORDS

carnivores, diet, garbage, generalist species, synanthropization, human settlements

 © 2016 Ł. Jankowiak et al.

This is an open access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivs license

INTRODUCTION

Human food waste is considered to be an important supplement to natural food supplies for some mammals and may positively affect urban populations (Contesse et al. 2004; Gehrt 2004). Some species can easily change their behaviour and learn how to find and use human food waste and, as a consequence, become partially independent from fluctuations in natural food resources (Goldyn et al. 2003; Ditchkoff et al. 2006). On the other hand, they may be forced to use human food waste due to a high density of their population (Contesse et al. 2004), an increase in urban areas (Pickett et al. 2011) and the lack of natural prey (Martina & Gallarati 1997). It is believed that the most common human food wastes of many wild animals are fruits, vegetables and scavenged meat (Harris 1981; Contesse et al. 2004; Oro et al. 2013; Tryjanowski et al. 2015).

In most studies of the carnivore diet, garbage are omitted and usually not included in the analysis and discussion

(e.g., Lindström 1994; Tryjanowski 1997; Lanszki et al. 1999; Lanszki 2003), and mostly discussed in surveys of species living in urban areas (Harris 1981; Saunders et al. 1993; Contesse et al. 2004) and rarely those in farmland areas (Cavallini & Volpi 1996; Goldyn et al. 2003). Even rarer are studies discussing the importance of garbage in communities of carnivorous species. In fact, only Prigioni et al. (2008) roughly discussed the occurrence of garbage in diets of four carnivore species. There is also little information about the types of garbage found in scats.

Most species whose diet is supplemented with anthropogenic food are highly opportunistic and use garbage dumps as a source of food (Martina & Gallarati 1997). The best known such mammal is red fox, which has a partly urbanized population (*Vulpes vulpes*; Harris 1981; Goldyn et al. 2003). The surveys revealed that raccoon dog (*Nyctereutes procyonoides*) could consume anthropogenic food (Kauhala et al. 1998); however, there is no obvious confirmation of preference to hu-

man settlements, at least in the European population (Drygala et al. 2008). Literature of this species indicates the occurrence of raccoon dogs in the cities, however it was found in Japan (Saeki 2001). Mustelids are also typical consumers of garbage (Fedriani et al. 2001), especially stone marten (*Martes foina*) (Broekhuizen 1999; Goszczyński et al. 2007). The European polecat (*Mustela putorius*) is considered to be a generalist feeder with an almost exclusively carnivorous diet, and has had a recent tendency to occur in human settlements (Baghli et al. 2002). Stoat (*Mustela erminea*) is regarded as a specialist predator feeding mostly on rodents, but also as a semi-generalist – feeding occasionally on other food resources (Korpimäki et al. 1991; Martinoli et al. 2001). American mink (*Neovison vison*) is characterized as an opportunistic predator which establishes territories along rivers or other water bodies (Sidorovich 2000). Contrastingly, Eurasian otter (*Lutra lutra*) is described as a fish-eater, specialist in Central Europe (Lanszki & Sallai 2006; Krawczyk et al. 2016). Stoat and riparian species like Eurasian otter, American mink are thought to prefer human settlements less (Carss 1995). However, when food resources are highly available, otters could use fish farms areas. Home ranges for red fox are established as 20-30 km² (Goszczyński 2002), for female polecat as 0.1 – over 2 km², and for male polecat as 0.2 – 12 km² (home range changes in seasons; Brzeziński & Romanowski, 1997).

The objectives of our study were to show the different types of garbage in the diet of carnivores, and the inter-specific variability in the amount of garbage remains in the carnivores' diet, along with a variation within the single species according to a distance to human settlements. In accordance with the above facts about opportunistic species, we predict that the specimens with home ranges closer to human settlements should often use the anthropogenic food like garbage dump, and it should be manifested by increased occurrence of garbage items in scats.

1. MATERIALS AND METHODS

The study was carried out in the Wielkopolska region, near the town of Odolanów (51°34'N, 17°40'E) in western Poland. The study area (220 km²) is characterized by extensive farmland with a mosaic of arable fields, meadows, wasteland and scrub. There are also many natural and artificial streams. Forests, mainly coniferous, occupy only a small percentage of the study area. Our scope of research was the administrative part of the Odolanów, Sulmierzyce and Przygodzice district, which are characterized as urban-rural with mean density as 77,9 humans/km². One garbage dump is localized in the study area. A large part of garbage produced by households were kitchen waste – 17% at rural and 26% at urban areas, which gave 231 tons of kitchen waste in Odolanów district in 2006 (GUS; Szyszkowski 2008).

Scats of surveyed species were collected from 2006 to 2010 and analyzed following the standard procedures. The scats were identified according to shape, size and place where

the faeces were found (for details see Romanowski 1998). They were collected and placed into plastic bags and stored for later analysis. Later on, faeces were soaked in water, crumbled and examined for macroscopic remains identified using a light microscope (for details see Jędrzejewska & Jędrzejewski 1998). The contribution of garbage was expressed as the percentage of scats in which they occurred. We separated scats in a warm season (April-November) and a cold season (December-March) to undertake seasonal comparisons. The details of collected sample size are shown in Table 1.

Because human food waste is very well digested in scats, we have used each indigestible, anthropogenic origin element found in faeces (e.g., glass, plastic, rubber) as an indication of this kind of food in the diet. To test the relationship between distance to settlements and garbage presence in scats, these were tested using a generalized linear model with a binomial distribution and logit-link function. We also computed the odds ratio to show how the probability of garbage items' presence changed per unit change in relation to the distance to nearest buildings from the place where scat was found. We did not do this analysis for Eurasian otter, stoat and American mink, stone marten because of the small sample size of garbage in their diet.

The significance was accepted at $p < 0.05$. The whole statistical analysis were done by STATISTICA 8.0v software (StatSoft Inc. 2007)

2. RESULTS

The examination of scats revealed the occurrence of garbage of 12 different types in the carnivores' diet (Table 1). Plastic occurred in the scats of five of seven species, and for four species, it was the most frequent garbage. Glass was the most often type of garbage found in the diet of Eurasian otter. The analysis of stoat faeces revealed the presence of only one type of garbage – wool, which was found only once.

For red fox and polecat, we found that the probability of finding the garbage item in scat decreased if the distance to settlements increased (Table 2, Figure 1). The odds ratio shows that with every 100 m increase in the distance to settlements, the chance to find a garbage item decreased by 15% for red fox and by 19% for polecat.

3. DISCUSSION

Our survey suggests continuous gradient in the garbage consumption. From the most synanthropized species – red fox, which had the most numerous garbage in the diet, to American mink, which did not consumed garbage at all (Gołdyn et al. 2003; Zabala et al. 2005; Goszczyński et al. 2007). On the other hand, the Eurasian otter, American mink and stoat did not show signs of synanthropization. The analysis showed that red fox and polecat, whose scats were found closer to human settlements, probably used garbage as an easily accessible food source. The main source of garbage for these animals

Table 1. Types of garbage in diets of Carnivora species in the study area.

Garbage item	raccoon dog (<i>Nyctereutes procyonoides</i>)		red fox (<i>Vulpes vulpes</i>)		marten (<i>Martes</i> sp.)		polecat (<i>Mustela putorius</i>)		stoat (<i>Mustela erminea</i>)	Eurasian otter (<i>Lutra lutra</i>)	American mink (<i>Neovison vison</i>)
	N	%	N	%	N	%	N	%	N	N	N
Plastic	6	60	15	33.3	4	50.0	10	32.3		1	
Aluminium foil					1	12.5	7	22.6		1	
Styrofoam			13	29.0			2	6.5			
Paper	1	10	6	13.3	1	12.5	4	12.9			
Buckshot							1	3.2			
Lime					1	12.5	2	6.4			
Paint							2	6.4			
Wool	1	10	1	2.2			2	6.5	1		
Hemp twine							1	3.2			
Glass	2	20	4	8.9						2	
Rubber			1	2.2							
Sponge			1	2.2	1	12.5					
Unidentified			4	8.9							
Total number of garbage		10		45		8		31	1	4	0
Total garbage occurrence		8.8		4.8		4.3		2.5	1.7	0.2	0
Total number of scats		215		936		161		1068	59	2269	115

Table 2. The results of generalised linear models showing the relationship between the presence of garbage item in the scats of two species and the distance to buildings from the place where the scat was found.

Model	Term	Coefficient	Standard Error	Wald Chi-square	p-value	Odds ratio
red fox (<i>Vulpes vulpes</i>)	intercept	-1.9364	0.3692			
	distance to buildings	-0.0015	0.0005	8.9135	<0.01	0.9985
polecat (<i>Mustela putorius</i>)	intercept	-2.6452	0.2852			
	distance to buildings	-0.0019	0.0005	13.9582	<0.001	0.9981




Figure 1. The negative relationships between probability of finding the garbage item in the scats of red fox *Vulpes vulpes* (A) and polecat *Mustela putorius* (B) with the distance to nearest buildings from place where the scat was found.

could be small garbage dumps and containers found in farmyards or gardens, or even some rubbish near roads (Ditchkoff et al. 2006). The most frequent garbage items found in the diet of surveyed species was plastic. Probably during the foraging of rubbish dumps, animals tear plastic bags to get the remains of human food and accidentally eat it. Human garbage often constitutes a food source which is more calorific and richer in carbohydrates, lipids and proteins than most natural food items (Peirce and Van Daele 2006). They are available all year round and independent of the environmental conditions (Panek and Bresiński 2002; Gołdyn et al. 2003). Peirce and Van Daele (2006) have shown that eating garbage may provide earlier maturation, higher growth rates, and gain some calorific benefit. However, foraging close to human settlements may be risky, because many of the carnivorous species are regarded as pests, and are persecuted because of the possibility of being poisoned (Birks 1998; Dip et al. 2003). There is also a possible risk of damage to or blockage of the digestive track (Sadove & Morreale 1990). Staying in the human neighbourhood may also increase road mortality (Ditchkoff et al. 2006). It should be noted that signs of poultry consumption found as feather remains in scats could also be an evidence of synanthropization. A paper from the study area shows that the occurrence of poultry remains in scats increases in territories of red fox closer to human settlements (Jankowiak et al. 2008).

We assessed the relative contribution of garbage to carnivores' diet by quantifying the proportion of scats with non-digestible garbage remains (e.g., rubber, plastic and glass). However, the relative contribution of digestible material may be underestimated because human food waste can be completely digested and its remains cannot be found (Litviatis 2000). Nevertheless, it is this kind of material that truly forms a part of the diet (i.e., not the non-digestible items, such as rubber or plastic that do not contribute nutrients nor energy to whom ingested it, thus by definition do not form part of the diet) and will contribute to carnivores' fitness in the form of human-mediated subsidizing. We assumed according to Cavallini and Volpi (1996)

survey that there is a relationship between digestible and non-digestible material.

The input of nutrients from human food waste can positively benefit the population density (Fedriani et al. 2001; Bino 2010; Oro et al. 2013; Tryjanowski et al. 2015). On the other hand, feeding on garbage could be a result of high population densities, as due stronger competition, the carnivores are forced to forage closer to human settlements (Contesse et al. 2004). However, the ability to adapt to this unusual habitat, gaining some benefits and avoiding risks arising from the human proximity, could be important factors determining the development of these populations. The subject of this study covers an important ecological issue, due to the invasion of natural wastes by human settlements and activities.

References

- Baghli, A., Engel, E & Verhagen, R. (2002) Feeding habits and trophic niche overlap of two sympatric Mustelidae, the polecat *Mustela putorius* and the beech marten *Martes foina*. *Zeitschrift für Jagdwissenschaft*, 48, 217–225.
- Bino, G., Dolev, A., Yosha, D., Guter, A., King, R., Saltz, D. & Kark, S. (2010) Abrupt spatial and numerical responses of overabundant foxes to a reduction in anthropogenic resources. *Journal of Applied Ecology*, 47, 1262–1271.
- Birks, J.D.S. (1998) Secondary rodenticide poisoning risk arising from winter farmland use by the European polecat *Mustela putorius*. *Biological Conservation*, 85, 233–240.
- Broekhuizen, S. (1999) *Martes foina*. In: A. Mitchell-Jones, G. Amori, W. Bogdanowicz, B. Krystufek, P.J.H. Reijnders, F. Spitzenberger, M. Stue, J.B.M. Thissen, V. Vohralik & J.T. Zima (Eds.) *The atlas of European mammals* (pp. 342–343). London: Poyser AD.
- Brzeziński, M. & Romanowski, J. (1997) *Tchórz*. Warszawa: Wydawnictwo Świat. [in Polish]

- Carss, D.N. (1995) Foraging behaviour and feeding ecology of the otter *Lutra lutra*: a selective review. *Hystrix, the Italian Journal of Mammalogy*, 7, 1–2.
- Cavallini, P. & Volpi, T. (1996) Variation in the diet of the red fox in a Mediterranean area. *Revue d'Ecologie (Terre et Vie)*, 51, 173–189.
- Contesse, P., Hegglin, D., Gloor, S., Bontadina, F. & Deplazes P. (2004) The diet of urban foxes (*Vulpes vulpes*) and the availability of anthropogenic food in the city of Zurich, Switzerland. *Mammalian Biology*, 69, 81–95.
- Dip, R., Hegglin, D., Deplazes, P., Dafflon, O., Koch, H. & Naegeli, H. (2003) Age- and sex-dependent distribution of persistent organochlorine pollutants in urban foxes. *Environmental Health Perspectives*, 111, 1608–1612.
- Ditchkoff, S., Saalfeld, S. & Gibson, C. (2006) Animal behavior in urban ecosystems: Modifications due to human-induced stress. *Urban Ecosystems*, 9, 5–12.
- Drygala, F., Stier, N., Zoller, H., Boegelsack, K., Mix, H.M. & Roth, M. (2008) Habitat use of the raccoon dog *Nyctereutes procyonoides* in north-eastern Germany. *Zeitschrift für Säugetierkunde*, 73, 371–378.
- Fedriani, J.M., Fuller, T.K. & Sauvajot, R.M. (2001) Does availability of anthropogenic food enhance densities of omnivorous mammals? An example with coyotes in southern California. *Ecography*, 24, 325–331.
- Gehrt, S.D. (2004) Ecology and management of striped skunks, raccoons, and coyotes in urban landscapes. In: N. Fascione, A. Delach & M.A. Smith (Eds.). *People and predators: from conflict to coexistence*. Washington: Island Press.
- Gołdyn, B., Hromada, M., Surmacki, A. & Tryjanowski P. (2003) Habitat use and diet of the red fox *Vulpes vulpes* in an agricultural landscape in Poland. *Zeitschrift für Jagdwissenschaft*, 49, 191–200.
- Goszczyński, J. (2002) Home ranges in red fox: territoriality diminishes with increasing area. *Acta Theriologica*, 47, 103–114.
- Goszczyński, J., Pośluszny, M., Pilot, M. & Gralak, B. (2007) Patterns of winter locomotion and foraging in two sympatric marten species: *Martes martes* and *Martes foina*. *Canadian Journal of Zoology*, 85, 239–249.
- GUS Central Statistical Office (2012) Available from <http://www.stat.gov.pl> [23 March 2012].
- Harris, S. (1981) The food of suburban foxes (*Vulpes vulpes*), with special reference to London. *Mammalian Review*, 11, 151–168.
- Jankowiak, Ł., Antczak, M., & Tryjanowski, P. (2008) Habitat use, food and the importance of poultry in the diet of the red fox *Vulpes vulpes* in extensive farmland in Poland. *World Applied Science Journal*, 4, 886–890.
- Jędrzejewska, B. & Jędrzejewski, W. (1998) Predation in vertebrate communities. The Białowieża Primeval Forest as a case study. Berlin: Springer Ecological Studies. Springer-Verlag.
- Kauhala, K., Laukkanen, P. & von Rége, I. (1998) Summer food composition and food niche overlap of the raccoon dog, red fox and badger in Finland. *Ecography*, 21, 457–463.
- Korpimäki, E., Norrdahl, K., & Rinta Jaskari, T. (1991) Responses of stoats and least weasel to fluctuating food abundances: is the low phase of the vole cycle due to mustelid predation? *Oecologia*, 88, 552–561.
- Krawczyk, A.J., Bogdziewicz, M., Majkowska, K. & Glazaczow, A. (2016) Diet composition of the Eurasian otter *Lutra lutra* in different freshwater habitats of temperate Europe: A review and meta-analysis. *Mammal Review*, 46, 106–113.
- Lanszki J., Körmendi, S., Hancz, C. & Zalewski, A. (1999) Feeding habits and trophic niche overlap in a Carnivora community of Hungary. *Acta Theriologica*, 44, 429–442.
- Lanszki, J. (2003) Feeding habits of stone martens in a Hungarian village and its surroundings. *Folia Zoologica*, 52, 367–377.
- Lanszki, J. & Sallai, Z. (2006) Comparison of the feeding habits of Eurasian otters on a fast flowing river and its backwater habitats. *Mammalian Biology*, 71, 336–346.
- Lindström, E.R. (1994) Large prey for small cubs - on crucial resources of a boreal red fox population. *Ecography*, 17, 17–22.
- Litvaitis, J.A. (2000) Investigating food habits of terrestrial vertebrates. In: L. Boitani & T.K. Fuller (Eds.). *Research techniques in animal ecology: controversies and consequences* (pp. 165–190). New York: Columbia University Press.
- Martina, A. & Gallarati, M. (1997) Use of garbage dump by some mammal species in the Majella massif (Abruzzo, Italy). *Hystrix*, 9, 23–29.
- Martinoli, A., Preatoni, D.G., Chiarenzi, B., Wauters, L.A. & Tosi, G. (2001) Diet of stoats (*Mustela erminea*) in an Alpine habitat: The importance of fruit consumption in summer. *Acta Oecologica*, 22, 45–53.
- Oro, D., Genovart, M., Tavecchia, G., Fowler, M.S. & Martínez-Abraín, A. (2013) Ecological and evolutionary implications of food subsidies from humans. *Ecology Letters*, 16, 1501–1514.
- Panek, M. & Bresiński, W. (2002) Red fox *Vulpes vulpes* density and habitat use in a rural area of western Poland in the end of 1990s, compared with the turn of 1970s. *Acta Theriologica*, 47, 433–442.
- Peirce, K.N. & Van Daele, L.J. (2006) Use of a garbage dump by brown bears in Dillingham, Alaska. *Ursus* 17, 165–177.
- Pickett, S.T.A., Cadenasso, M.L., Grove, J.M., Boone, C.G., Irwin, E., Groffman, P.M., et al. (2011) Urban ecological systems: Scientific foundations and a decade of progress. *Journal of Environmental Management*, 92, 331–362.
- Prigioni, C., Balestrieri, A., Remonti, L. & Cavada, L. (2008) Differential use of food and habitat by sympatric carnivores in the eastern Italian Alps. *Italian Journal of Zoology*, 75, 173–184.
- Romanowski, J. 1998. Śladami zwierząt. PWRiL [in Polish].
- Sadove, S.S. & Morreale, S.J. (1990) Marine mammal and sea turtle encounters with marine debris in the New York Bight and the Northeast Atlantic. In: R.S. Shomura & M.L. Godfrey (Eds.). *Proceedings of the Second International Conference on Marine Debris*, 2-7 April 1989, Honolulu, Hawaii.
- Saeki, M. (2001) Ecology and conservation of the raccoon dog (*Nyctereutes procyonoides*) in Japan. Oxford University Press.
- Saunders, G., White, P.C.L., Harris, S. & Rayner, J.M.V. (1993) Urban foxes (*Vulpes vulpes*): food acquisition, time and energy budgeting of a generalised predator. *Symposia of the Zoological Society of London*, 65, 215–234.

- Sidorovich, V.E. (2000) Seasonal variation in the feeding habits of riparian mustelids in river valleys of NE Belarus. *Acta Theriologica*, 45, 233–242.
- StatSoft Inc. (2007) STATISTICA (data analysis software system), version 8.0. www.statsoft.com.
- Szyszkowski, P. (2008) Plan of waste management for Wielkopolska region in 2008-2011 with prospects for 2012-2019. Actualization. Poznań: ARCADIS Profil.
- Tryjanowski, P. (1997) Food of the Stone marten (*Martes foina*) in Nietoperek Bat Reserve. *Zeitschrift für Säugetierkunde* 62, 318–320.
- Tryjanowski, P., Skórka, P., Sparks, T. H., Biaduń, W., Brauze, T., Hetmański, T., et al. (2015). Urban and rural habitats differ in number and type of bird feeders and in bird species consuming supplementary food. *Environmental Science and Pollution Research*, 22, 15097–15103.
- Zabala, J., Zuberogoitia, I. & Martínez-Climent, J.A. (2005) Site and landscape features ruling the habitat use and occupancy of the polecat (*Mustela putorius*) in a low density area, a multiscale approach. *European Journal of Wildlife Research*, 51, 157–162.