



## The DNA polymerases of *Drosophila melanogaster*

Steven J. Marygold, Helen Attrill, Elena Speretta, Kate Warner, Michele Magrane, Maria Berloco, Sue Cotterill, Mitch McVey, Yikang Rong & Masamitsu Yamaguchi

To cite this article: Steven J. Marygold, Helen Attrill, Elena Speretta, Kate Warner, Michele Magrane, Maria Berloco, Sue Cotterill, Mitch McVey, Yikang Rong & Masamitsu Yamaguchi (2020): The DNA polymerases of *Drosophila melanogaster*, Fly, DOI: [10.1080/19336934.2019.1710076](https://doi.org/10.1080/19336934.2019.1710076)

To link to this article: <https://doi.org/10.1080/19336934.2019.1710076>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



[View supplementary material](#)



Published online: 14 Jan 2020.



[Submit your article to this journal](#)



[View related articles](#)



[View Crossmark data](#)

## The DNA polymerases of *Drosophila melanogaster*

Steven J. Marygold <sup>a</sup>, Helen Attrill <sup>a</sup>, Elena Speretta <sup>b</sup>, Kate Warner <sup>b</sup>, Michele Magrane <sup>b</sup>, Maria Berloco <sup>c</sup>, Sue Cotterill <sup>d</sup>, Mitch McVey <sup>e</sup>, Yikang Rong <sup>f</sup>, and Masamitsu Yamaguchi<sup>g</sup>

<sup>a</sup>FlyBase, Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge, UK; <sup>b</sup>UniProt, European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Cambridgeshire, UK; <sup>c</sup>Dipartimento di Biologia, Università degli Studi di Bari "Aldo Moro", Bari, Italy; <sup>d</sup>Department Basic Medical Sciences, St Georges University London, London, UK; <sup>e</sup>Department of Biology, Tufts University, Medford, MA, USA; <sup>f</sup>School of Life Sciences, Sun Yat-sen University, Guangzhou, China; <sup>g</sup>Department of Applied Biology and Advanced Insect Research Promotion Center, Kyoto Institute of Technology, Kyoto, Japan

### ABSTRACT

DNA synthesis during replication or repair is a fundamental cellular process that is catalyzed by a set of evolutionary conserved polymerases. Despite a large body of research, the DNA polymerases of *Drosophila melanogaster* have not yet been systematically reviewed, leading to inconsistencies in their nomenclature, shortcomings in their functional (Gene Ontology, GO) annotations and an under-appreciation of the extent of their characterization. Here, we describe the complete set of DNA polymerases in *D. melanogaster*, applying nomenclature already in widespread use in other species, and improving their functional annotation. A total of 19 genes encode the proteins comprising three replicative polymerases (alpha-primase, delta, epsilon), five translesion/repair polymerases (zeta, eta, iota, Rev1, theta) and the mitochondrial polymerase (gamma). We also provide an overview of the biochemical and genetic characterization of these factors in *D. melanogaster*. This work, together with the incorporation of the improved nomenclature and GO annotation into key biological databases, including FlyBase and UniProtKB, will greatly facilitate access to information about these important proteins.

### ARTICLE HISTORY

Received 25 October 2019  
Revised 19 December 2019  
Accepted 23 December 2019  
Published online 14 January 2020

### KEYWORDS



*Drosophila melanogaster*;  
DNA polymerase; DNA  
synthesis; DNA replication;  
translesion synthesis


## Introduction

Multiple DNA polymerases are required for DNA synthesis in eukaryotic cells [1]. Their functions have been characterized in many different model systems and found to be highly conserved through evolution. The alpha-primase, delta and epsilon polymerases exist as multi-subunit complexes and are responsible for the bulk of nuclear DNA replication during S phase. These 'replicative polymerases' belong to the B family of polymerases and are characterized by relatively high processivity of 5'-3' DNA synthesis and high fidelity. The alpha-primase complex is responsible for synthesis of short RNA primers and initial DNA synthesis. On the lagging strand, it repeatedly synthesizes short RNA-DNA primers that are then extended by polymerase delta to generate Okazaki fragments. After initiating synthesis on the leading strand, alpha-primase is replaced by polymerase epsilon to perform the bulk of leading strand

DNA elongation, though polymerase delta may be used as the leading strand polymerase in some circumstances and in certain species. Polymerase epsilon is further distinguished from delta by showing little dependence on the proliferating cell nuclear antigen (PCNA) protein for its activity. Both delta and epsilon possess a 3'-5' exonuclease ('proof-reading') activity that ensures replication fidelity. (For a recent review on replicative polymerases, see [2].)

Other specialized polymerases are required for DNA synthesis during nuclear DNA repair and/or translesion synthesis (TLS). TLS allows replication past residual DNA lesions that would otherwise block the progress of the replication machinery. On encountering a lesion, replicative DNA polymerases are switched for specific TLS polymerases (e.g. zeta, eta, iota, Rev1, theta) that then either bypass or repair lesions before the normal machinery re-associates and replication continues

**CONTACT** Steven J. Marygold  [sjm41@cam.ac.uk](mailto:sjm41@cam.ac.uk)  FlyBase, Department of Physiology, Development and Neuroscience, University of Cambridge, Cambridge CB2 3DY, UK

 Supplemental data for this article can be accessed [here](#)

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

downstream of the damage. Owing to their specialized functions, TLS polymerases exhibit lower fidelity and processivity than replicative polymerases. TLS is mutagenic when the incorrect base is inserted opposite the DNA lesion. (For a recent review on TLS polymerases, see [3].)

TLS polymerases exhibit different preferences for types of lesions and differ in their nucleotide selection and mutagenic potential. For example, polymerase zeta (another member of the B family) is involved in the response to a large variety of DNA damage, showing high fidelity for the bypass of some lesions but low fidelity for others [4]. Polymerase eta (Y family) functions primarily to bypass cyclobutane pyrimidine dimers, which are the predominant lesions resulting from ultraviolet (UV) radiation, and can also bypass 7,8-dihydro-8-oxyguanine lesions and abasic sites [5]. In contrast to other DNA polymerases, the metazoan-specific polymerase iota (Y family) is able to use non-canonical Hoogsteen interactions for nucleotide base pairing, allowing it to incorporate nucleotides opposite various lesions in the DNA template that impair Watson-Crick interactions [6]. Rev1 (Y family) is unique in that its DNA synthesis activity is restricted to the incorporation of one or two molecules of dCTP regardless of the nature of the template nucleotide – in this sense, it is better classified as a deoxycytidyl transferase than a polymerase *per se*. Furthermore, Rev1 also has non-catalytic functions in DNA synthesis, serving as a scaffolding protein for the assembly of multiprotein TLS complexes [3]. DNA polymerase theta (belonging to the A family) is another polymerase restricted to metazoa and is unique in possessing a helicase domain associated with DNA-dependent ATPase activity [7,8]. Theta has an indispensable role in repair synthesis of double-strand breaks via an alternative non-homologous end-joining pathway and also functions in repair/bypass of interstrand crosslinks.

In contrast to the panel of polymerases functioning in the synthesis of nuclear DNA, DNA polymerase gamma (A family) is solely responsible for the replication and repair of mitochondrial DNA (mtDNA), at least in non-mammalian species (reviewed in [9]). It is highly accurate and processive, containing both polymerase and 3'-5' exonuclease activities, as well as a 5'-deoxyribose phosphate lyase activity that functions in base excision repair.

Its polymerase activity is stimulated by association with the mitochondrial single-stranded DNA binding (mtSSB). A primase for mtDNA replication has not been identified – instead, transcription is thought to provide the priming event for replication. In mammals, a second polymerase, PrimPol, is also required for repair of mtDNA damage, acting to reinitiate stalled replication forks in mitochondria [10].

While much of the work characterizing the identity and function of eukaryotic DNA polymerases has been conducted in yeast and mammalian cell culture, there is also a long and rich history of DNA polymerase research in the fruit fly, *Drosophila melanogaster*, which continues through to this day. DNA polymerase activity was first detected in fly embryos in the 1970s [11,12], which led to the purification and characterization of several replicative polymerases and their component subunits in the 1980s and 1990s. Most of the genes encoding these proteins were then cloned by individual labs in the 1990s, prior to the publication of the complete *D. melanogaster* genome sequence in 2000 [13]. Within the past two decades, all of the TLS polymerases of *D. melanogaster* have also been functionally characterized, albeit to different extents. Many fly polymerase genes have also been studied genetically, taking advantage of the relative ease and power of this approach in this model organism. These methods have been extended in recent years to create fly models of human diseases associated with polymerase dysfunction.

Despite this large body of work, the DNA polymerases of *D. melanogaster* have not yet been systematically cataloged or reviewed. This has led to inconsistent and out-dated nomenclature for this gene set and several deficiencies in their functional annotations within biological databases, which in turn has led to an under-appreciation of the extent of their characterization. Here, we seek to rectify these points. We identify all *D. melanogaster* DNA polymerases using a combination of bioinformatic and literature-based searches. In so doing, we have substantially revised and improved the Gene Ontology (GO) terms used to annotate these genes/proteins in biological databases [14]. We also propose an updated, unified nomenclature for these factors, based on that already in use in the wider

field. Finally, we provide an overview of the history and key findings of the research characterizing the composition and function of DNA polymerases in flies.

## Defining the set of *D. melanogaster* DNA polymerases

We took a combinatorial approach to ensure we identified the full set of genes known or predicted to encode DNA polymerases in *D. melanogaster*. We searched the FlyBase [15], UniProtKB [16], QuickGO [17] and KEGG [18] databases for *D. melanogaster* entries annotated with relevant GO and/or Enzyme Commission terms. We also searched for additional literature references using FlyBase and PubMed, and identified *D. melanogaster* orthologs of the established set of human and yeast (*Saccharomyces cerevisiae*) DNA polymerases using the DIOPT and HCOP ortholog prediction tools [19,20]. The final collection of verified *D. melanogaster* polymerases/polymerase subunits is presented in Table 1.

A key part of this survey was to review and improve the GO annotations associated with the fly DNA polymerases, adding new or more specific annotations where appropriate and removing any erroneous annotations. In particular, we ensured that catalytic activity GO terms were applied only to catalytic, and not regulatory/accessory, subunits of DNA polymerase complexes. Overall, we added 140 new annotations (of which 135 are based on experimental data) and removed 10 erroneous annotations. The full set of DNA polymerase-relevant GO annotations for this set of genes/proteins following our review is provided in Supplementary Table 1.

Table 1 also shows our proposal for an updated and systematic nomenclature for the *D. melanogaster* DNA polymerase genes/proteins. This is based on the approved nomenclature used for human and vertebrate polymerases and includes the use of English rather than the traditional Greek characters, which can be difficult to interpret/process in reading the regular literature and in database representation. Previous symbols and other notable synonyms are

**Table 1.** *D. melanogaster* genes encoding (subunits of) DNA polymerases.

| Family                          | Pol     | Gene Symbol              | Notable synonym(s)               | FlyBase gene ID | UniProt protein accession | <i>H. sapiens</i> ortholog (identity) <sup>a</sup> | <i>S. cerevisiae</i> ortholog (identity) <sup>a</sup> |
|---------------------------------|---------|--------------------------|----------------------------------|-----------------|---------------------------|--|---|
| <b>Replicative polymerases</b>  |         |                          |                                  |                 |                           |  |   |
| B                               | ALPHA   | <b>PolA1</b>             | <i>DNApol-α180</i>               | FBgn0259113     | P26019                    | POLA1 (40%)  | POL1 (32%)  |
|                                 |         | <b>PolA2</b>             | <i>DNApol-α73</i>                | FBgn0005696     | Q9VB62                    | POLA2 (30%)  | POL12 (23%)   |
|                                 |         | <b>Prim1</b>             | <i>DNApol-α50</i>                | FBgn0011762     | Q24317                    | PRIM1 (42%)  | PRI1 (36%)  |
|                                 |         | <b>Prim2</b>             | <i>DNApol-α60</i>                | FBgn0259676     | Q9VPH2                    | PRIM2 (33%)  | PRI2 (30%)  |
|                                 |         | <b>PolD1</b>             | <i>DNApol-δ</i>                  | FBgn0263600     | P54358                    | POLD1 (58%)  | POL3 (51%)  |
| B                               | DELTA   | <b>PolD2<sup>b</sup></b> | <i>Pol31</i>                     | FBgn0027903     | Q9W088                    | POLD2 (43%)  | POL31 (30%)   |
|                                 |         | <b>PolD3<sup>b</sup></b> | <i>Pol32, rd</i>                 | FBgn0283467     | Q9Y118                    | POLD3 (22%)  | POL32 (n/a) <sup>c</sup>                              |
|                                 |         | <b>PolE1</b>             | <i>DNApol-ε255</i>               | FBgn0264326     | Q9VCN1                    | POLE (55%)   | POL2 (39%)  |
| B                               | EPSILON | <b>PolE2</b>             | <i>DNApol-ε58</i>                | FBgn0035644     | Q9VRQ7                    | POLE2 (42%)  | DPB2 (25%)  |
|                                 |         | <b>PolE3</b>             | <i>Chrac-14</i>                  | FBgn0043002     | Q9V444                    | POLE3 (47%)  | DPB4 (25%)  |
|                                 |         | <b>PolE4</b>             | <i>Mes4</i>                      | FBgn0034726     | Q9W256                    | POLE4 (38%)  | DPB3 (28%)  |
|                                 |         | <b>PolZ1</b>             | <i>mus205, rev3, DNApol-ζ</i>    | FBgn0002891     | Q9GSR1                    | REV3L (27%)  | REV3 (25%)  |
| B                               | ZETA    | <b>PolZ2</b>             | <i>rev7</i>                      | FBgn0037345     | Q9VNE1                    | MAD2L2 (29%)                                       | REV7 (18%)  |
|                                 |         | <b>PolD2<sup>b</sup></b> | <i>Pol31</i>                     | FBgn0027903     | Q9W088                    | POLD2 (43%)  | POL31 (30%)   |
|                                 |         | <b>PolD3<sup>b</sup></b> | <i>Pol32, rd</i>                 | FBgn0283467     | Q9Y118                    | POLD3 (22%)  | POL32 (n/a) <sup>c</sup>                              |
|                                 |         | <b>PolH</b>              | <i>DNApol-η</i>                  | FBgn0037141     | Q9VNX1                    | POLH (29%)   | RAD30 (23%)   |
| Y                               | ETA     | <b>PolH</b>              | <i>drad30A</i>                   |                 |                           |  |   |
| Y                               | IOTA    | <b>PolI</b>              | <i>DNApol-ι, drad30B</i>         | FBgn0037554     | Q9VHV1                    | POLI (31%)   | -   |
| Y                               | REV1    | <b>Rev1</b>              | -                                | FBgn0035150     | Q9W0P2                    | REV1 (30%)   | REV1 (24%)  |
| A                               | THETA   | <b>PolQ</b>              | <i>mus308, DNApol-θ</i>          | FBgn0002905     | O18475                    | POLQ (29%)   | -   |
| <b>Mitochondrial polymerase</b> |         |                          |                                  |                 |                           |  |   |
| A                               | GAMMA   | <b>PolG1</b>             | <i>tam, pol γ-α, DNApol-γ125</i> | FBgn0004406     | Q27607                    | POLG (44%)   | MIP1 (33%)  |
|                                 |         | <b>PolG2</b>             | <i>pol γ-β, DNApol-γ35</i>       | FBgn0004407     | Q9VJV8                    | POLG2 (24%)  | -   |

a. Percentage amino acid identity between the *D. melanogaster* protein and the human/yeast ortholog according to DIOPT [19].

b. PolD2 and PolD3 are part of the polymerase delta and zeta complexes.

c. PolD3/Pol32 subunits are known to show extreme divergence at the primary sequence level [21], but FBgn0283467 has been identified as *D. melanogaster* PolD3/Pol32 [59].

listed in the table. (A full listing of previous and proposed nomenclature is given in Supplementary Table 2.) These revisions bring the *D. melanogaster* nomenclature into line with that used in the wider field and will make it easier for researchers from diverse backgrounds to access the fly data.

It is notable that several human/mammalian DNA polymerases lack a clear ortholog in *D. melanogaster* [19,20], namely beta (POLB), delta subunit 4 (POLD4), kappa (POLK), lambda (POLL), mu (POLM), nu (POLN), PRIMPOL and terminal deoxynucleotidyltransferase (TDT/DNTT). Conversely, *D. melanogaster* possess the iota and theta DNA polymerases that are not found in yeast.

## Research on *D. melanogaster* DNA polymerases

Below, we present an overview and historical perspective of the key research characterizing the composition and function of the DNA polymerases in flies, focusing on their roles in DNA synthesis (summarized in Table 2). This is not intended to be a comprehensive review of all literature concerning these factors, or how DNA polymerases function within the wider contexts of replication control and DNA repair. Readers

**Table 2.** Characterization status of *D. melanogaster* DNA polymerases.

|       | Biochemistry     | Cloning | Expression | Genetics      |
|-------|------------------|---------|------------|---------------|
| PolA1 | [28–30,33–40]    | [41,42] | [24,41–45] | [47]          |
| PolA2 | [25,33–36]       | [48]    | [44,45,49] | [47]          |
| Prim1 | [28–36,50,51]    | [51]    | -          | [47]          |
| Prim2 | [28–30,33–36,50] | [52]    | -          | [53]          |
| PolD1 | [37,56–58]       | [60]    | [58]       | [47,58]       |
| PolD2 | [58]             | n/a     | [58]       | [58]          |
| PolD3 | [58]             | n/a     | [58]       | [58,59,63,64] |
| PolE1 | [68,69]          | [44,70] | [44,70]    | [47,71,72]    |
| PolE2 | [69]             | n/a     | -          | [73]          |
| PolE3 | -                | n/a     | -          | -             |
| PolE4 | -                | n/a     | -          | -             |
| PolZ1 | [74,75]          | [76]    | -          | [59,76]       |
| PolZ2 | [74,75]          | n/a     | -          | -             |
| PolH  | [80]             | [80]    | -          | [59,81]       |
| Poll  | [80]             | [80]    | -          | -             |
| Rev1  | [74,82]          | n/a     | -          | [59]          |
| PolQ  | [86–88]          | [83]    | [87]       | [88–91]       |
| PolG1 | [26,94–106]      | [105]   | [118]      | [107–113]     |
| PolG2 | [26,94–104]      | [114]   | [118]      | [108,117]     |

The references shown report biochemical data, cloning, transcript/protein expression or genetic data (i.e. phenotypic analysis using classical mutations or transgenes) relevant to the role of the gene/protein in DNA synthesis. (Cloning status is given as 'n/a' where cloning was not reported prior to the release of the *D. melanogaster* genome sequence.)

interested in those topics are referred to excellent recent reviews [22,23].

## Replicative polymerases

**Alpha ( $\alpha$ )-primase.** *Alpha-primase initiates synthesis on the leading strand and repeatedly synthesizes short primers on the lagging strand during nuclear DNA replication. The canonical complex comprises two primase subunits and two polymerase subunits, with each subcomplex containing one catalytic and one accessory protein (reviewed in [27]).*

The *D. melanogaster* alpha-primase was first purified and characterized from embryos in a series of studies by Lehman and colleagues in the early 1980s [28–35]. This and other work [36,37] established the tetrameric subunit structure of the complex: a 182 kDa catalytic polymerase subunit (PolA1), a non-catalytic 73 kDa subunit (PolA2), and two smaller subunits of 50kDa and 60kDa (Prim1 and Prim2, respectively) associated with primase activity.

Additional research on the isolated PolA1 subunit further characterized its properties and revealed it possesses a cryptic 3'-5' exonuclease that functions as a proofreading activity to increase replication fidelity [38–40]. The gene encoding PolA1 was cloned in 1991–1992 [41,42]. Those and other studies [43–45] determined that levels of the protein are relatively high in unfertilized eggs and early embryos owing to maternal loading, with its subcellular location being mainly nuclear during non-mitotic phases of the cell cycle. PolA1 interacts physically and genetically with the PrSet7 methyltransferase, suggesting that this interaction and the regulation of histone monomethylation play a role in DNA replication [46]. Consistent with a key role in DNA synthesis, global RNAi-mediated knockdown of PolA1 is lethal, whereas eye-specific RNAi results in a small eye [47].

The gene encoding the 73 kDa PolA2 subunit was cloned in 1992 [48] and is expressed similarly to PolA1 [44,45,49]. Its molecular function in flies remains undefined, but is assumed to play an essential role in the initiation of DNA replication based on the characterization of orthologs [27]. Similar to PolA1, knockdown of PolA2 results in lethality or a small eye phenotype [47].

The primase subcomplex was isolated and shown capable of synthesizing primers in the



absence of the other two subunits [50]. The gene encoding Prim1 was cloned in 1994 and this subunit alone was demonstrated to have primase activity [51]. The protein is detected throughout embryogenesis but is only evident in adult females (ovaries) at later development stages, suggesting maternal expression. The Prim2-encoding gene was cloned in 1999 and its expression found to be similar to Prim1 [52]. A specific function has not been ascribed to Prim2 in *D. melanogaster*, though it may stabilize Prim1 and/or act to translocate Prim1 to the nucleus based on analysis of Prim2 orthologs [27]. Knockdown of Prim1 expression causes lethality [47]. Similarly, strong *Prim2* mutations are lethal, whereas hypomorphic *Prim2* mutants show cell cycle perturbations [53].

**Delta ( $\delta$ ).** DNA polymerase delta (PolD) primarily acts to extend DNA on the lagging strand, though it can also serve as the leading strand polymerase. The canonical eukaryotic complex is composed of three to four subunits, one of which possesses catalytic activity (reviewed in [54,55]).

The *D. melanogaster* PolD holoenzyme was purified from embryos and characterized to be a highly processive, PCNA-dependent DNA polymerase with a 3'-5' proof-reading activity [37,56,57]. It was initially identified as a heterodimer [57], but a more recent study has shown it to be a heterotrimer [58]. (Vertebrate PolD has an additional subunit, but this appears to be lacking in invertebrates.) *D. melanogaster* PolD3 (aka Pol32) facilitates nuclear localization of the PolD complex and increases PolD processivity [58,59], while PolD2 (aka Pol31) acts as a structural component, bridging the other two subunits [58].

Despite being cloned in 1995 [60], there has been relatively little genetic characterization of the *PolD1* gene – several recessive lethal mutations have been isolated [61,62] and RNAi-mediated knockdown of *PolD1* causes lethality [47]. PolD2 is similarly uncharacterized from a genetic perspective. However, genetic approaches have been used to show that PolD3 is essential for DNA replication in early embryogenesis [58,63]. Interestingly, PolD3 is also required for the repair of double-strand breaks by homologous recombination involving extensive DNA synthesis [59] and for break-induced replication during repair of single broken strands [64],

though it's unclear whether PolD functions within the polymerase delta or zeta complex in these roles (see below). Genetic interactions between PolD3 and the other two delta subunits have been reported, supporting functional associations *in vivo* [58].

**Epsilon ( $\epsilon$ ).** Polymerase epsilon (PolE) is the major leading strand polymerase. The canonical complex is composed of one catalytic subunit and three non-catalytic subunits (reviewed in [65,66]). Recent work in mammals has shown that the two smaller PolE subunits form a stable sub-complex that acts as a histone H3-H4 chaperone to facilitate nucleosome assembly during DNA replication [67].

The *D. melanogaster* catalytic subunit (PolE1) was purified from embryos and characterized to be a highly processive DNA polymerase with a 3'-5' exonuclease activity [68,69]. The gene was cloned in 2000 [70] and its protein levels found to be highest in unfertilized eggs and early embryos, similar to PolA subunits [44,70]. Consistent with its molecular function, *PolE1* mutation or knock-down results in inhibition of DNA synthesis, under-sized tissues and early lethality [47,71,72]. Knock-down of *PolE1* expression also causes defective endoreplication in larval salivary glands [71,72]. Remarkably, the mitotic replication defects caused by PolE1 knock-down could be rescued by expression of the non-catalytic C-terminal domain of PolE1, whereas the endoreplication defects could not [71]. This suggests that the polymerase/exonuclease activities of PolE1 can be compensated for by other replicative polymerases (most likely PolD) during regular DNA replication in mitotic cycles, whereas these catalytic domains play a specific and critical role during endoreplication.

A 58 kDa subunit of DNAPol epsilon (PolE2) was also purified and shown to directly associate with PolE1 [69]. Its specific function is unknown in flies, but it likely plays an essential structural role from studies of its orthologs [65,66]. A null mutation in the *PolE2* gene has been shown to be pupal lethal and cause defects in S phase progression in mitotic and endoreplicative cell cycles, similar to mutations in *PolE1* [73].

*D. melanogaster* PolE3 (aka Chrac-14) and PolE4 (aka Mes4) have not been studied in the

context of DNA replication. Nonetheless, they are likely to be part of the *D. melanogaster* PolE holoenzyme based on studies of their orthologs and may perform similar roles to their mammalian counterparts in facilitating nucleosome assembly during replication [67].

### Translesion/DNA repair polymerases

**Zeta ( $\zeta$ ).** DNA polymerase zeta (PolZ) performs TLS in response to a variety of DNA damage. Studies in yeast and humans have shown that it exists as a heterotetramer, in which a single subunit has catalytic activity and two of the accessory subunits are shared with PolD (reviewed in [4]).

*D. melanogaster* PolZ1 was purified and characterized as a high processivity polymerase and, while lacking any 3'-5' exonuclease function, it shows high fidelity for DNA synthesis [74]. The same study demonstrated that PolZ2 directly interacts with PolZ1 without influencing its DNA synthetic activity [74]. Interestingly, the PolZ complex may be involved in the repair of abasic sites in flies – this function is performed by DNA polymerase beta in mammals, but flies (and many other organisms) do not have a POLB ortholog [75].

The gene encoding the catalytic subunit, *PolZ1* (aka *mus205* or *rev3*), was cloned in 2001 [76]. This, and several previous genetic studies [77–79], showed that mutants were hypersensitive to alkylating agents and UV, suggesting PolZ1 is involved in lesion bypass/TLS. Further genetic characterization has demonstrated that PolZ1 also functions in the repair of double-strand breaks [59]. In contrast, there have been no genetic studies on the PolZ2 (aka *Rev7*) subunit to date.

While it has not been experimentally shown that the *D. melanogaster* PolZ complex includes the PolD2 and PolD3 subunits of polymerase delta, this seems likely based on studies in other species [4]. In this regard, it is possible that PolD3 acts to promote the nuclear localization of the PolZ complex in flies, as described for its function within PolD [58].

**Eta ( $\eta$ ).** The gene encoding *D. melanogaster* polymerase eta (PolH) was cloned in 2001 [80]. The same study demonstrated PolH has DNA polymerase activity and can bypass UV-induced lesions: *cis-syn*-cyclobutane pyrimidine dimers are

bypassed error-free, whereas ([4–6])-photoproducts are bypassed in a highly error-prone manner [80]. Consistent with that study, genetic approaches have shown that PolH mutants are extremely sensitive to UV radiation [59,81]. PolH may also function in the repair synthesis of double-strand breaks [59]. Notably, the fly PolH mutants may provide *in vivo* models of a variant form of xeroderma pigmentosum, a disease characterized by increased incidence of UV-induced skin cancers caused by mutations in human PolH [81].

**Iota ( $\iota$ ).** Similar to PolH, the gene encoding *D. melanogaster* polymerase iota (PolI) was cloned in 2001 and the purified protein shown to have DNA polymerase activity [80]. It too can bypass *cis-syn*-cyclobutane pyrimidine dimers UV photoproducts in an error-free manner, though it is not able to bypass ([4–6])-photoproducts at all [80]. PolI function has not been studied using genetic approaches to date, and it remains an open question whether it plays roles outside of being a 'backup' to PolH.

**Rev1.** A scaffolding/co-ordinating function for *D. melanogaster* Rev1 is supported by studies demonstrating physical interactions between it and other TLS polymerases including PolH, PolI, PolZ1 and PolZ2 [74,82]. Genetic investigations lend further support to this view, showing that loss of Rev1 results in high sensitivity to ionizing radiation and affects the degree of homologous recombination repair synthesis [59]. One idea arising from these studies is that Rev1 coordinates the initial recruitment of other translesion polymerases and, in so doing, prevents replicative polymerases from acting during early repair synthesis [59]. There are no biochemical data supporting a deoxycytidyl transferase/DNA polymerase activity of Rev1 in flies, though this is assumed to be present from characterization of its orthologs.

**Theta ( $\theta$ ).** PolQ is the best studied TLS polymerase in *D. melanogaster*. The gene encoding PolQ was cloned in 1996 [83] and shown to correspond to the well-studied *mutagen-sensitive 308* (*mus308*) locus, mutations in which conferred strong sensitivity to DNA cross-linking agents [84,85]. The purified

PolQ protein was demonstrated to have both polymerase and, uniquely amongst DNA polymerases, DNA-dependent ATPase activity [86–88]. It is expressed throughout all stages of development [87].

The ATPase activity of PolQ is associated with a N-terminal helicase-like domain, which is separated from the C-terminal polymerase domain by an unstructured central region [83]. Both domains are required for resistance to cross-linking agents *in vivo* and purified PolQ can bypass structures representative of unhooked crosslinks *in vitro* [88]. These data suggest that the helicase activity may stimulate DNA unwinding and strand displacement to facilitate synthesis by the polymerase domain.

A series of genetic studies by the McVey laboratory have delineated a second, crucial function for PolQ in repair of double-strand breaks via a process termed microhomology-mediated end-joining (MMEJ) [88–91]. MMEJ occurs when short, complementary DNA sequences located at broken DNA ends anneal and serve as primers for fill-in synthesis. While only the polymerase activity of PolQ is required for MMEJ, loss of the helicase domain affects the spectrum of repair junctions that are recovered [88]. PolQ is also important for repair of double-strand breaks arising from replication fork collapse during endoreplication in eggshell-producing follicle cells [91].

### Mitochondrial polymerase

**Gamma ( $\gamma$ ).** DNA polymerase gamma (PolG) is responsible for the replication and repair of mtDNA (reviewed in [9]). The subunit composition of the holoenzyme varies across species, ranging from a single catalytic subunit in yeast and nematodes to a heterodimer (one accessory subunit) in insects, and a heterotrimer (including a dimeric accessory subunit) in vertebrates [92].

PolG is the best characterized *D. melanogaster* polymerase, having been comprehensively analyzed by the Kaguni laboratory over several years [93]. The complex was first purified from embryos in 1986 and found to comprise two polypeptides of 125 and 35 kDa [94]. Several subsequent studies defined the holoenzyme as a high fidelity polymerase associated with a 3'-5' exonuclease (proof-reading) activity, which requires the mitochondrial single stranded binding protein (mtSSB) to stimulate its processivity [95–104].

The gene encoding the large subunit, PolG1, was cloned in 1996 [105]. This and other studies established that PolG1 harbors both the polymerase and exonuclease activities, as well as a mitochondrial targeting sequence [102,105,106]. Mutations in this gene were first identified via a screen for altered larval responses to light (resulting in the gene being named 'tamas' – Sanskrit for 'dark inertia') – indeed, these were the first reported mutations in any PolG gene [107]. Mutants die during late larval stages, exhibiting impaired mtDNA replication and decreased mtDNA content [107–109]. Ubiquitous RNAi-mediated knock-down of PolG1 depletes mtDNA, decreases mitochondrial respiratory activity and results in lethality, while neuronal-specific knock-down causes progressive behavioral deficits [110,111]. Other studies have utilized sophisticated genetic techniques to engineer flies to express either polymerase- or exonuclease-deficient PolG1 – the former resulted in mtDNA depletion, whereas the latter led to accumulation of mutations/deletions of mitochondrial DNA [112,113].

The gene encoding the smaller accessory subunit, PolG2, was cloned in 1997 [114]. It is located in the genome within 9 kilobases of the *PolG1* gene and the encoded protein contains a mitochondrial targeting sequence [102,114]. Biochemical analyses have demonstrated a critical role for PolG2 in the catalytic efficiency of the PolG1 subunit [102], while structural modeling suggests it is also involved in primer recognition and processive DNA elongation [115]. These findings are consistent with work on the mammalian ortholog [116] and define the PolG2 subunit as a processivity factor for the PolG1 polymerase. Null mutations in *PolG2* have been generated and result in lethality during pupal stages, with mutant cells exhibiting loss of mtDNA and reduced proliferation [108,117].

Despite their proximity in the genome, the *PolG1* and *PolG2* genes exhibit differential expression: *PolG1* is expressed highly in very early embryos but is then largely undetectable until adult stages, whereas *PolG2* expression increases during embryonic stages, remains present at lower levels during larval and pupal development, and is then expressed highly again in adults [118]. One reason for this difference is that transcription of *PolG2*, but not



*PolG1*, is in part governed by a DNA replication-related element (DRE) in its promoter [118]. Interestingly, DREs have also been identified in the promoters of the *PolA1*, *PolA2*, *PolE4* and *PCNA* genes [41,49,119,120] and thus may be widely used to control the expression of DNA replication-related factors.

Mutations in human *POLG* and *POLG2* are associated with several diseases including mtDNA depletion syndromes, such as Alpers syndrome, and mtDNA deletion disorders, such as progressive external ophthalmoplegia [121]. Significantly, several studies in the last decade have manipulated the orthologous *D. melanogaster* genes to provide important insights into the etiology of, and possible therapies for, these diseases [108–113,122]. For example, two studies generated flies expressing exonuclease- or polymerase-deficient versions of PolG1 in order to investigate how defects in these two different activities of the enzyme may contribute to pathophysiology [112,113]. Two other investigations studied the effects of mutations or knock-down of *D. melanogaster* *PolG1* or *PolG2* specifically in the nervous system to probe how mitochondrial dysfunction may lead to neuropathies and neurodegenerative disorders in humans [108,110]. As a final example, Siibak et al. engineered the endogenous *PolG1* gene in flies to have mutations equivalent to those found in patients to probe their molecular and physiological consequences, finding that both mutations cause mtDNA depletion [109].

## Perspective

The DNA polymerases of *D. melanogaster* have been researched for almost 50 years. In this brief report, we have gathered together the fruits of these investigations and presented the full set of fly DNA polymerases alongside the key references characterizing them. In so doing, we have proposed a revised nomenclature that is in line with wider usage in the field and greatly improved the quality and coverage of the functional (GO) annotations associated with these factors. These improvements in annotation and their integration into core biological databases, including FlyBase ([www.flybase.org](http://www.flybase.org)) and UniProtKB ([www.uniprot.org](http://www.uniprot.org)), will greatly enhance access to, and the impact of, *D. melanogaster* DNA polymerase research. Furthermore, all UniProtKB records for

these proteins have been manually annotated/reviewed and a dedicated ‘Gene Group’ page for the DNA polymerases has been added to FlyBase (<http://flybase.org/reports/FBgg0001200>).

Much of what we know about eukaryotic polymerases has been conducted in yeast and mammalian cell culture. Flies provide a way to investigate polymerase function in the context of an intact multi-cellular organism with unrivaled possibilities for sophisticated genetic analyses to inform biological function. Indeed, the genetic approach, often combined with biochemical assays, has been key to the several landmark discoveries about DNA polymerases that were made through research using *D. melanogaster*. These include the first characterization of mutations in PolG subunits [107,117], which ultimately led to a genetic dissection of the roles of the exonuclease and polymerase domains of PolG1 [112,113]; the function of PolQ within the MMEJ repair pathway [88–90]; and the role of PolD3 in facilitating the nuclear import of PolD1 [58]. Several of the mechanisms elucidated by these studies provided precedent for other organisms, particularly so in the case of PolQ. For example, studies of PolQ in *C. elegans*, mice, and human cells have clearly shown that both its domains are important for alternative end joining/MMEJ processes [123–125], as originally described in flies. The role of *D. melanogaster* PolQ in the repair of double-strand breaks arising from replication fork collapse [91] has also been found to be conserved [126–128]. Early studies on the fly PolQ identified its role in interstrand crosslink repair – a similar role for mammalian PolQ has been debated [7], but a recent study has demonstrated that human POLQ is required for repair of mitomycin C-induced damage, which introduces interstrand crosslinks [127]. These examples emphasize the relevance of work on *D. melanogaster* DNA polymerases to other eukaryotic animals, including humans.

Nonetheless, it’s equally clear that flies must do a few things differently, at least compared to the well-studied yeast and mammalian polymerase systems. For instance, flies have two additional polymerases (iota and theta) compared to yeast, but lack several of the TLS polymerases present in mammals (Table 1). Presumably, the

'missing' polymerases are either not required for DNA synthesis in yeast/fly cells or their functions are performed by one or more of the polymerases that are present [75]. The sub-unit composition of the delta and gamma polymerases also differs between these three species (Table 1), leading to functional differences at least in the latter case [92]. As a final example, several fly cell types exhibit endoreplication, whereby cells undergo multiple S phases without entering mitosis or cytokinesis, resulting in giant cells with a polyploid nucleus [129]. This variant cell cycle is common in other arthropods, nematodes and plants, but is rarely found in mammals and is absent from yeast. While regulation of endoreplication occurs primarily at the level of core components of the cell cycle machinery, it's apparent that some fly DNA polymerases, including PolE and PolQ have also developed endocycle-specific regulatory mechanisms [71–73,91].

We anticipate that future research using *D. melanogaster* will continue to provide pioneering insights into DNA polymerase biology that are applicable across a wide range of species. Pursuing and developing additional fly models of the several diseases associated with lesions in human DNA polymerases [121,130] is likely to be a particularly useful research direction. We hope the work presented in this review will facilitate these new studies and aid the integration of the ensuing knowledge.

## Acknowledgments

We thank Brian Calvi for critical reading of the manuscript, Sandra Orchard for overseeing UniProt curation, and three anonymous reviewers for insightful comments.

## Disclosure statement










No potential conflict of interest was reported by the authors.

## Funding

S.M. is funded by National Human Genome Research Institute at the US National Institutes of Health [U41HG000739] to Norbert Perrimon (PI), Nicholas Brown (co-PI); H.A. is funded by the UK Medical Research Council [MR/N030117/1]; E.S., K.W. and M.M. are funded by the

National Eye Institute (NEI), National Human Genome Research Institute (NHGRI), National Heart, Lung, and Blood Institute (NHLBI), National Institute of Allergy and Infectious Diseases (NIAID), National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), National Institute of General Medical Sciences (NIGMS), and National Institute of Mental Health (NIMH) of the National Institutes of Health under Award Number U24HG007822 (the content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health) and by European Molecular Biology Laboratory (EMBL) core funds.

## ORCID

Steven J. Marygold  <http://orcid.org/0000-0003-2759-266X>  
 Helen Attrill  <http://orcid.org/0000-0003-3212-6364>  
 Elena Speretta  <http://orcid.org/0000-0003-1506-7438>  
 Kate Warner  <http://orcid.org/0000-0001-8705-181X>  
 Michele Magrane  <http://orcid.org/0000-0003-3544-996X>  
 Maria Berloco  <http://orcid.org/0000-0003-3826-8427>  
 Sue Cotterill  <http://orcid.org/0000-0001-9913-0259>  
 Mitch McVey  <http://orcid.org/0000-0001-8883-0188>  
 Yikang Rong  <http://orcid.org/0000-0002-9787-9669>

## References

- [1] Jain R, Aggarwal AK, Rechkoblit O. Eukaryotic DNA polymerases. *Curr Opin Struct Biol.* 2018;53:77–87.
- [2] Burgers PMJ, Kunkel TA. Eukaryotic DNA replication fork. *Annu Rev Biochem.* 2017;86:417–438.
- [3] Yang W, Gao Y. Translesion and repair DNA polymerases: diverse structure and mechanism. *Annu Rev Biochem.* 2018;87:239–261.
- [4] Makarova AV, Burgers PM. Eukaryotic DNA polymerase zeta. *DNA Repair (Amst).* 2015;29:47–55.
- [5] Yang W. An overview of Y-Family DNA polymerases and a case study of human DNA polymerase eta. *Biochemistry.* 2014;53:2793–2803.
- [6] Makarova AV, Kulbachinskiy AV. Structure of human DNA polymerase iota and the mechanism of DNA synthesis. *Biochemistry (Mosc).* 2012;77:547–561.
- [7] Black SJ, Kashkina E, Kent T, et al. DNA polymerase theta: a unique multifunctional end-joining machine. *Genes (Basel).* 2016;7:E67.
- [8] Wood RD, Double S. DNA polymerase theta (POLQ), double-strand break repair, and cancer. *DNA Repair (Amst).* 2016;44:22–32.
- [9] Graziewicz MA, Longley MJ, Copeland WC. DNA polymerase gamma in mitochondrial DNA replication and repair. *Chem Rev.* 2006;106:383–405.
- [10] Torregrosa-Munumer R, Forslund JME, Goffart S, et al. PrimPol is required for replication reinitiation after mtDNA damage. *Proc Natl Acad Sci U S A.* 2017;114:11398–11403.

- [11] Margulies L, Chargaff E. Survey of DNA polymerase activity during the early development of *Drosophila melanogaster*. *Proc Natl Acad Sci U S A*. 1973;70:2946–2950.
- [12] Brakel CL, Blumenthal AB. Multiple forms of *Drosophila* embryo DNA polymerase: evidence for proteolytic conversion. *Biochemistry*. 1977;16:3137–3143.
- [13] Adams MD, Celniker SE, Holt RA, *et al*. The genome sequence of *Drosophila melanogaster*. *Science*. 2000;287:2185–2195.
- [14] The Gene Ontology, C. The gene ontology resource: 20 years and still GOing strong. *Nucleic Acids Res*. 2019;47:D330–D338.
- [15] Thurmond J, Goodman JL, Strelets VB, *et al*. FlyBase 2.0: the next generation. *Nucleic Acids Res*. 2019;47:D759–D765.
- [16] UniProt C. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res*. 2019;47:D506–D515.
- [17] Binns D, Dimmer E, Huntley R, *et al*. QuickGO: a web-based tool for gene ontology searching. *Bioinformatics*. 2009;25:3045–3046.
- [18] Kanehisa M, Sato Y, Furumichi M, *et al*. New approach for understanding genome variations in KEGG. *Nucleic Acids Res*. 2019;47:D590–D595.
- [19] Hu Y, Flockhart I, Vinayagam A, *et al*. An integrative approach to ortholog prediction for disease-focused and other functional studies. *BMC Bioinformatics*. 2011;12:357.
- [20] Braschi B, Denny P, Gray K, *et al*. Genenames.org: the HGNC and VGNC resources in 2019. *Nucleic Acids Res*. 2019;47:D786–D792.
- [21] Johansson E, Garg P, Burgers PM. The Pol32 subunit of DNA polymerase delta contains separable domains for processive replication and proliferating cell nuclear antigen (PCNA) binding. *J Biol Chem*. 2004;279:1907–1915.
- [22] Hua BL, Orr-Weaver TL. DNA replication control during *Drosophila* development: insights into the onset of S phase, replication initiation, and fork progression. *Genetics*. 2017;207:29–47.
- [23] Sekelsky J. DNA repair in *Drosophila*: mutagens, models, and missing genes. *Genetics*. 2017;205:471–490.
- [24] Yamaguchi M, Hayashi Y, Hirose F, *et al*. Distinct roles of E2F recognition sites as positive or negative elements in regulation of the DNA polymerase alpha 180 kDa catalytic subunit gene promoter during *Drosophila* development. *Nucleic Acids Res*. 1997;25:3847–3854.
- [25] Kuroda K, Ueda R. Phosphorylation and dephosphorylation of the B subunit of DNA polymerase alpha-primase complex in the early embryogenesis of *Drosophila*. *Biochem Biophys Res Commun*. 1999;254:372–377.
- [26] Fan L, Kaguni LS. Multiple regions of subunit interaction in *Drosophila* mitochondrial DNA polymerase: three functional domains in the accessory subunit. *Biochemistry*. 2001;40:4780–4791.
- [27] Muzi-Falconi M, Giannattasio M, Foiani M, *et al*. The DNA polymerase alpha-primase complex: multiple functions and interactions. *ScientificWorldJournal*. 2003;3:21–33.
- [28] Banks GR, Boezi JA, Lehman IR. A high molecular weight DNA polymerase from *Drosophila melanogaster* embryos. Purification, structure, and partial characterization. *J Biol Chem*. 1979;254:9886–9892.
- [29] Villani G, Sauer B, Lehman IR. DNA polymerase alpha from *Drosophila melanogaster* embryos. Subunit structure. *J Biol Chem*. 1980;255:9479–9483.
- [30] Villani G, Fay PJ, Bambara RA, *et al*. Elongation of RNA-primed DNA templates by DNA polymerase alpha from *Drosophila melanogaster* embryos. *J Biol Chem*. 1981;256:8202–8207.
- [31] Conaway RC, Lehman IR. Synthesis by the DNA primase of *Drosophila melanogaster* of a primer with a unique chain length. *Proc Natl Acad Sci U S A*. 1982;79:4585–4588.
- [32] Conaway RC, Lehman IR. A DNA primase activity associated with DNA polymerase alpha from *Drosophila melanogaster* embryos. *Proc Natl Acad Sci U S A*. 1982;79:2523–2527.
- [33] Kaguni LS, Rossignol JM, Conaway RC, *et al*. Association of DNA primase with the beta/gamma subunits of DNA polymerase alpha from *Drosophila melanogaster* embryos. *J Biol Chem*. 1983;258:9037–9039.
- [34] Kaguni LS, Rossignol JM, Conaway RC, *et al*. Isolation of an intact DNA polymerase-primase from embryos of *Drosophila melanogaster*. *Proc Natl Acad Sci U S A*. 1983;80:2221–2225.
- [35] Kaguni LS, DiFrancesco RA, Lehman IR. The DNA polymerase-primase from *Drosophila melanogaster* embryos. Rate and fidelity of polymerization on single-stranded DNA templates. *J Biol Chem*. 1984;259:9314–9319.
- [36] Kuroda K, Kagiya-Takahashi R, Shinomiya T. Immunoaffinity purification and properties of *Drosophila melanogaster* DNA polymerase alpha-primase complex. *J Biochem*. 1990;108:926–933.
- [37] Peck VM, Gerner EW, Cress AE. Delta-type DNA polymerase characterized from *Drosophila melanogaster* embryos. *Nucleic Acids Res*. 1992;20:5779–5784.
- [38] Cotterill S, Chui G, Lehman IR. DNA polymerase-primase from embryos of *Drosophila melanogaster*. The DNA polymerase subunit. *J Biol Chem*. 1987;262:16100–16104.
- [39] Cotterill SM, Reyland ME, Loeb LA, *et al*. A cryptic proofreading 3'–5' exonuclease associated with the polymerase subunit of the DNA polymerase-primase from *Drosophila melanogaster*. *Proc Natl Acad Sci U S A*. 1987;84:5635–5639.
- [40] Reyland ME, Lehman IR, Loeb LA. Specificity of proofreading by the 3'–5' exonuclease of the DNA polymerase-primase of *Drosophila melanogaster*. *J Biol Chem*. 1988;263:6518–6524.
- [41] Hirose F, Yamaguchi M, Nishida Y, *et al*. Structure and expression during development of *Drosophila*

- melanogaster gene for DNA polymerase alpha. *Nucleic Acids Res.* **1991**;19:4991–4998.
- [42] Melov S, Vaughan H, Cotterill S. Molecular characterisation of the gene for the 180 kDa subunit of the DNA polymerase-primase of *Drosophila melanogaster*. *J Cell Sci.* **1992**;102(Pt 4):847–856.
- [43] Yamaguchi M, Kuroda K, Hirose F, et al. Distribution of DNA polymerase alpha during nuclear division cycles in *Drosophila melanogaster* embryo. *Cell Struct Funct.* **1992**;17:105–112.
- [44] Yamaguchi M, Hirose F, Nishimoto Y, et al. Expression patterns of DNA replication enzymes and the regulatory factor DREF during *Drosophila* development analyzed with specific antibodies. *Biol Cell.* **1995**;85:147–155.
- [45] Kuroda K, Ueda R. A 130 kDa polypeptide immunologically related to the 180 kDa catalytic subunit of DNA polymerase alpha-primase complex is detected in early embryos of *Drosophila*. *J Biochem.* **1995**;117:809–818.
- [46] Sahashi R, Crevel G, Pasko J, et al. DNA polymerase  $\alpha$  interacts with PrSet7 and mediates H4K20 monomethylation in *Drosophila*. *J Cell Sci.* **2014**;127:3066–3078.
- [47] Kohzaki H, Asano M, Murakami Y. DNA replication machinery is required for development in *Drosophila*. *Front Biosci (Landmark Ed).* **2018**;23:493–505.
- [48] Cotterill S, Lehman IR, McLachlan P. Cloning of the gene for the 73 kD subunit of the DNA polymerase alpha primase of *Drosophila melanogaster*. *Nucleic Acids Res.* **1992**;20:4325–4330.
- [49] Takahashi Y, Yamaguchi M, Hirose F, et al. DNA replication-related elements cooperate to enhance promoter activity of the *drosophila* DNA polymerase alpha 73-kDa subunit gene. *J Biol Chem.* **1996**;271:14541–14547.
- [50] Cotterill S, Chui G, Lehman IR. DNA polymerase-primase from embryos of *Drosophila melanogaster*. DNA primase subunits. *J Biol Chem.* **1987**;262:16105–16108.
- [51] Bakkenist CJ, Cotterill S. The 50-kDa primase subunit of *Drosophila melanogaster* DNA polymerase alpha. Molecular characterization of the gene and functional analysis of the overexpressed protein. *J Biol Chem.* **1994**;269:26759–26766.
- [52] Huikeshoven H, Cotterill S. Cloning and characterisation of the gene for the large subunit of the DNA primase from *Drosophila melanogaster*. *Biochim Biophys Acta.* **1999**;1445:359–362.
- [53] Chen X, Li Q, Fischer JA. Genetic analysis of the *Drosophila* DNAPrim gene. The function of the 60-kD primase subunit of DNA polymerase opposes the fat facets signaling pathway in the developing eye. *Genetics.* **2000**;156:1787–1795.
- [54] Prindle MJ, Loeb LA. DNA polymerase delta in DNA replication and genome maintenance. *Environ Mol Mutagen.* **2012**;53:666–682.
- [55] Tahirov TH. Structure and function of eukaryotic DNA polymerase delta. *Subcell Biochem.* **2012**;62:217–236.
- [56] Chiang CS, Mitsis PG, Lehman IR. DNA polymerase delta from embryos of *Drosophila melanogaster*. *Proc Natl Acad Sci U S A.* **1993**;90:9105–9109.
- [57] Aoyagi N, Matsuoka S, Furunobu A, et al. *Drosophila* DNA polymerase delta. Purification and characterization. *J Biol Chem.* **1994**;269:6045–6050.
- [58] Ji J, Tang X, Hu W, et al. The processivity factor Pol32 mediates nuclear localization of DNA polymerase delta and prevents chromosomal fragile site formation in *Drosophila* development. *PLoS Genet.* **2019**;15:e1008169.
- [59] Kane DP, Shusterman M, Rong Y, et al. Competition between replicative and translesion polymerases during homologous recombination repair in *Drosophila*. *PLoS Genet.* **2012**;8:e1002659.
- [60] Chiang CS, Lehman IR. Isolation and sequence determination of the cDNA encoding DNA polymerase delta from *Drosophila melanogaster*. *Gene.* **1995**;166:237–242.
- [61] Brizuela BJ, Elfring L, Ballard J, et al. Genetic analysis of the brahma gene of *Drosophila melanogaster* and polytene chromosome subdivisions 72AB. *Genetics.* **1994**;137:803–813.
- [62] Cooper MT, Kennison JA. Molecular genetic analyses of polytene chromosome region 72A–D in *Drosophila melanogaster* reveal a gene desert in 72D. *PLoS One.* **2011**;6:e23509.
- [63] Tritto P, Palumbo V, Micale L, et al. Loss of Pol32 in *Drosophila melanogaster* causes chromosome instability and suppresses variegation. *PLoS One.* **2015**;10:e0120859.
- [64] Bhandari J, Karg T, Golic KG. Homolog-dependent repair following dicentric chromosome breakage in *Drosophila melanogaster*. *Genetics.* **2019**;212:615–630.
- [65] Pospiech H, Syvaaja JE. DNA polymerase epsilon - more than a polymerase. *ScientificWorldJournal.* **2003**;3:87–104.
- [66] Hogg M, Johansson E. DNA polymerase epsilon. *Subcell Biochem.* **2012**;62:237–257.
- [67] Bellelli R, Belan O, Pye VE, et al. POLE3-POLE4 is a Histone H3-H4 chaperone that maintains chromatin integrity during DNA replication. *Mol Cell.* **2018**;72(112–126):e115.
- [68] Aoyagi N, Oshige M, Hirose F, et al. DNA polymerase epsilon from *Drosophila melanogaster*. *Biochem Biophys Res Commun.* **1997**;230:297–301.
- [69] Oshige M, Takeuchi R, Ruike T, et al. Subunit protein-affinity isolation of *Drosophila* DNA polymerase epsilon catalytic subunit. *Protein Expr Purif.* **2004**;35:248–256.
- [70] Oshige M, Yoshida H, Hirose F, et al. Molecular cloning and expression during development of the *Drosophila* gene for the catalytic subunit of DNA polymerase epsilon. *Gene.* **2000**;256:93–100.
- [71] Suyari O, Kawai M, Ida H, et al. Differential requirement for the N-terminal catalytic domain of the DNA



- polymerase epsilon p255 subunit in the mitotic cell cycle and the endocycle. *Gene*. 2012;495:104–114.
- [72] Verma A, Sengupta S, Lakhota SC. DNAPol-epsilon gene is indispensable for the survival and growth of *Drosophila melanogaster*. *Genesis*. 2012;50:86–101.
- [73] Sahashi R, Matsuda R, Suyari O, et al. Functional analysis of *Drosophila* DNA polymerase epsilon p58 subunit. *Am J Cancer Res*. 2013;3:478–489.
- [74] Takeuchi R, Oshige M, Uchida M, et al. Purification of *Drosophila* DNA polymerase zeta by REV1 protein-affinity chromatography. *Biochem J*. 2004;382:535–543.
- [75] Takeuchi R, Ruike T, Nakamura R, et al. *Drosophila* DNA polymerase zeta interacts with recombination repair protein 1, the *Drosophila* homologue of human abasic endonuclease 1. *J Biol Chem*. 2006;281:11577–11585.
- [76] Eeken JC, Romeijn RJ, de Jong AW, et al. Isolation and genetic characterisation of the *Drosophila* homologue of (SCE)REV3, encoding the catalytic subunit of DNA polymerase zeta. *Mutat Res*. 2001;485:237–253.
- [77] Boyd JB, Harris PV. Mutants partially defective in excision repair at five autosomal loci in *Drosophila melanogaster*. *Chromosoma*. 1981;82:249–257.
- [78] Brown TC, Boyd JB. Postreplication repair-defective mutants of *Drosophila melanogaster* fall into two classes. *Mol Gen Genet*. 1981;183:356–362.
- [79] Boyd JB, Shaw KE. Postreplication repair defects in mutants of *Drosophila melanogaster*. *Mol Gen Genet*. 1982;186:289–294.
- [80] Ishikawa T, Uematsu N, Mizukoshi T, et al. Mutagenic and nonmutagenic bypass of DNA lesions by *Drosophila* DNA polymerases dpoleta and dpoliota. *J Biol Chem*. 2001;276:15155–15163.
- [81] Wallace HA, Merkle JA, Yu MC, et al. TRIP/NOPO E3 ubiquitin ligase promotes ubiquitylation of DNA polymerase eta. *Development*. 2014;141:1332–1341.
- [82] Kosarek JN, Woodruff RV, Rivera-Begeman A, et al. Comparative analysis of in vivo interactions between Rev1 protein and other Y-family DNA polymerases in animals and yeasts. *DNA Repair (Amst)*. 2008;7:439–451.
- [83] Harris PV, Mazina OM, Leonhardt EA, et al. Molecular cloning of *Drosophila* mus308, a gene involved in DNA cross-link repair with homology to prokaryotic DNA polymerase I genes. *Mol Cell Biol*. 1996;16:5764–5771.
- [84] Boyd JB, Sakaguchi K, Harris PV. mus308 mutants of *Drosophila* exhibit hypersensitivity to DNA cross-linking agents and are defective in a deoxyribonuclease. *Genetics*. 1990;125:813–819.
- [85] Leonhardt EA, Henderson DS, Rinehart JE, et al. Characterization of the mus308 gene in *Drosophila melanogaster*. *Genetics*. 1993;133:87–96.
- [86] Oshige M, Aoyagi N, Harris PV, et al. A new DNA polymerase species from *Drosophila melanogaster*: a probable mus308 gene product. *Mutat Res*. 1999;433:183–192.
- [87] Pang M, McConnell M, Fisher PA. The *Drosophila* mus 308 gene product, implicated in tolerance of DNA interstrand crosslinks, is a nuclear protein found in both ovaries and embryos. *DNA Repair (Amst)*. 2005;4:971–982.
- [88] Beagan K, Armstrong RL, Witsell A, et al. *Drosophila* DNA polymerase theta utilizes both helicase-like and polymerase domains during microhomology-mediated end joining and interstrand crosslink repair. *PLoS Genet*. 2017;13:e1006813.
- [89] Chan SH, Yu AM, McVey M. Dual roles for DNA polymerase theta in alternative end-joining repair of double-strand breaks in *Drosophila*. *PLoS Genet*. 2010;6:e1001005.
- [90] Yu AM, McVey M. Synthesis-dependent microhomology-mediated end joining accounts for multiple types of repair junctions. *Nucleic Acids Res*. 2010;38:5706–5717.
- [91] Alexander JL, Beagan K, Orr-Weaver TL, et al. Multiple mechanisms contribute to double-strand break repair at rereplication forks in *Drosophila* follicle cells. *Proc Natl Acad Sci U S A*. 2016;113:13809–13814.
- [92] Ciesielski GL, Oliveira MT, Kaguni LS. Animal Mitochondrial DNA Replication. *Enzymes*. 2016;39:255–292.
- [93] Kaguni LS. DNA polymerase gamma, the mitochondrial replicase. *Annu Rev Biochem*. 2004;73:293–320.
- [94] Wernette CM, Kaguni LS. A mitochondrial DNA polymerase from embryos of *Drosophila melanogaster*. Purification, subunit structure, and partial characterization. *J Biol Chem*. 1986;261:14764–14770.
- [95] Wernette CM, Conway MC, Kaguni LS. Mitochondrial DNA polymerase from *Drosophila melanogaster* embryos: kinetics, processivity, and fidelity of DNA polymerization. *Biochemistry*. 1988;27:6046–6054.
- [96] Kaguni LS, Olson MW. Mismatch-specific 3'→5' exonuclease associated with the mitochondrial DNA polymerase from *Drosophila* embryos. *Proc Natl Acad Sci U S A*. 1989;86:6469–6473.
- [97] Olson MW, Kaguni LS. 3'→5' exonuclease in *Drosophila* mitochondrial DNA polymerase. Substrate specificity and functional coordination of nucleotide polymerization and mispair hydrolysis. *J Biol Chem*. 1992;267:23136–23142.
- [98] Williams AJ, Wernette CM, Kaguni LS. Processivity of mitochondrial DNA polymerase from *Drosophila* embryos. Effects of reaction conditions and enzyme purity. *J Biol Chem*. 1993;268:24855–24862.
- [99] Olson MW, Wang Y, Elder RH, et al. Subunit structure of mitochondrial DNA polymerase from *Drosophila* embryos. Physical and immunological studies. *J Biol Chem*. 1995;270:28932–28937.
- [100] Williams AJ, Kaguni LS. Stimulation of *Drosophila* mitochondrial DNA polymerase by single-stranded DNA-binding protein. *J Biol Chem*. 1995;270:860–865.
- [101] Farr CL, Wang Y, Kaguni LS. Functional interactions of mitochondrial DNA polymerase and single-stranded DNA-binding protein. Template-primer DNA binding and initiation and elongation of DNA strand synthesis. *J Biol Chem*. 1999;274:14779–14785.

- [102] Wang Y, Kaguni LS. Baculovirus expression reconstitutes *Drosophila* mitochondrial DNA polymerase. *J Biol Chem.* **1999**;274:28972–28977.
- [103] Farr CL, Matsushima Y, Lagina AT 3rd, et al. Physiological and biochemical defects in functional interactions of mitochondrial DNA polymerase and DNA-binding mutants of single-stranded DNA-binding protein. *J Biol Chem.* **2004**;279:17047–17053.
- [104] Ciesielski GL, Bermek O, Rosado-Ruiz FA, et al. Mitochondrial single-stranded DNA-binding proteins stimulate the activity of DNA polymerase gamma by organization of the template DNA. *J Biol Chem.* **2015**;290:28697–28707.
- [105] Lewis DL, Farr CL, Wang Y, et al. Catalytic subunit of mitochondrial DNA polymerase from *Drosophila* embryos. Cloning, bacterial overexpression, and biochemical characterization. *J Biol Chem.* **1996**;271:23389–23394.
- [106] Luo N, Kaguni LS. Mutations in the spacer region of *Drosophila* mitochondrial DNA polymerase affect DNA binding, processivity, and the balance between Pol and Exo function. *J Biol Chem.* **2005**;280:2491–2497.
- [107] Iyengar B, Roote J, Campos AR. The *tamas* gene, identified as a mutation that disrupts larval behavior in *Drosophila melanogaster*, codes for the mitochondrial DNA polymerase catalytic subunit (DNAPol-gamma125). *Genetics.* **1999**;153:1809–1824.
- [108] Baqri RM, Turner BA, Rheuben MB, et al. Disruption of mitochondrial DNA replication in *Drosophila* increases mitochondrial fast axonal transport in vivo. *PLoS One.* **2009**;4:e7874.
- [109] Siibak T, Clemente P, Bratic A, et al. A multi-systemic mitochondrial disorder due to a dominant p.Y955H disease variant in DNA polymerase gamma. *Hum Mol Genet.* **2017**;26:2515–2525.
- [110] Humphrey DM, Parsons RB, Ludlow ZN, et al. Alternative oxidase rescues mitochondria-mediated dopaminergic cell loss in *Drosophila*. *Hum Mol Genet.* **2012**;21:2698–2712.
- [111] Rodrigues APC, Camargo AF, Andjelkovic A, et al. Developmental arrest in *Drosophila melanogaster* caused by mitochondrial DNA replication defects cannot be rescued by the alternative oxidase. *Sci Rep.* **2018**;8:10882.
- [112] Bratic A, Kauppila TE, Macao B, et al. Complementation between polymerase- and exonuclease-deficient mitochondrial DNA polymerase mutants in genomically engineered flies. *Nat Commun.* **2015**;6:8808.
- [113] Samstag CL, Hoekstra JG, Huang CH, et al. Deleterious mitochondrial DNA point mutations are overrepresented in *Drosophila* expressing a proofreading-defective DNA polymerase gamma. *PLoS Genet.* **2018**;14:e1007805.
- [114] Wang Y, Farr CL, Kaguni LS. Accessory subunit of mitochondrial DNA polymerase from *Drosophila* embryos. Cloning, molecular analysis, and association in the native enzyme. *J Biol Chem.* **1997**;272:13640–13646.
- [115] Fan L, Sanschagrin PC, Kaguni LS, et al. The accessory subunit of mtDNA polymerase shares structural homology with aminoacyl-tRNA synthetases: implications for a dual role as a primer recognition factor and processivity clamp. *Proc Natl Acad Sci U S A.* **1999**;96:9527–9532.
- [116] Lee YS, Lee S, Demeler B, et al. Each monomer of the dimeric accessory protein for human mitochondrial DNA polymerase has a distinct role in conferring processivity. *J Biol Chem.* **2010**;285:1490–1499.
- [117] Iyengar B, Luo N, Farr CL, et al. The accessory subunit of DNA polymerase gamma is essential for mitochondrial DNA maintenance and development in *Drosophila melanogaster*. *Proc Natl Acad Sci U S A.* **2002**;99:4483–4488.
- [118] Lefai E, Fernandez-Moreno MA, Alahari A, et al. Differential regulation of the catalytic and accessory subunit genes of *Drosophila* mitochondrial DNA polymerase. *J Biol Chem.* **2000**;275:33123–33133.
- [119] Suyari O, Ida H, Yoshioka Y, et al. Identification of the *Drosophila* *Mes4* gene as a novel target of the transcription factor DREF. *Exp Cell Res.* **2009**;315:1403–1414.
- [120] Yamaguchi M, Nishida Y, Moriuchi T, et al. *Drosophila* proliferating cell nuclear antigen (cyclin) gene: structure, expression during development, and specific binding of homeodomain proteins to its 5'-flanking region. *Mol Cell Biol.* **1990**;10:872–879.
- [121] Stumpf JD, Copeland WC. Mitochondrial DNA replication and disease: insights from DNA polymerase gamma mutations. *Cell Mol Life Sci.* **2011**;68:219–233.
- [122] Yu Z, O'Farrell PH, Yakubovich N, et al. The Mitochondrial DNA Polymerase Promotes Elimination of Paternal Mitochondrial Genomes. *Curr Biol.* **2017**;27:1033–1039.
- [123] Wyatt DW, Feng W, Conlin MP, et al. Essential roles for polymerase theta-mediated end joining in the repair of chromosome breaks. *Mol Cell.* **2016**;63:662–673.
- [124] Mateos-Gomez PA, Kent T, Deng SK, et al. The helicase domain of Poltheta counteracts RPA to promote alt-NHEJ. *Nat Struct Mol Biol.* **2017**;24:1116–1123.
- [125] Schimmel J, Kool H, van Schendel R, et al. Mutational signatures of non-homologous and polymerase theta-mediated end-joining in embryonic stem cells. *Embo J.* **2017**;36:3634–3649.
- [126] Roerink SF, van Schendel R, Tijsterman M. Polymerase theta-mediated end joining of replication-associated DNA breaks in *C. elegans*. *Genome Res.* **2014**;24:954–962.
- [127] Feng W, Simpson DA, Carvajal-Garcia J, et al. Genetic determinants of cellular addiction to DNA polymerase theta. *Nat Commun.* **2019**;10:4286.
- [128] Wang Z, Song Y, Li S, et al. DNA polymerase theta (POLQ) is important for repair of DNA double-strand breaks caused by fork collapse. *J Biol Chem.* **2019**;294:3909–3919.
- [129] Zielke N, Edgar BA, DePamphilis ML. Endoreplication. *Cold Spring Harb Perspect Biol.* **2013**;5:a012948.
- [130] Beuning PJ. DNA polymerases: from molecular mechanisms to human disease, a special issue. *Chem Res Toxicol.* **2017**;30:1921.