

Nutritional adequacy of meals from an independent catering facility versus chain-restaurants for young adults

1

2 **Charoula Konstantia Nikolaou¹, Catherine Ruth Hankey², Michael Ernest John Lean²**

3

4 **Affiliations:** ¹SSH/JURI - Institut pour la recherche interdisciplinaire en sciences juridiques (JUR-I),
5 Catholic University of Louvain, Belgium, ²Human Nutrition Section, College of Medical, Veterinary
6 and Life Sciences, University of Glasgow

7

8 **Correspondence Author**

9 Charoula-Konstantia Nikolaou

10 Universite Catholique de Louvain

11 Collège Thomas More

12 Place Montesquieu 2

13 Louvain-La-Neuve

14 1348

15 E-mail: charoulanikolaou@yahoo.co.uk

16

17

18 **Word Count = 2,481 (word limit=3,000)**19 **Running Title:** Nutritional Quality of independent and chain restaurants20 **Key Words:** Public Health, Catering, nutrition, obesity21 **Conflict of interest:** The authors declare there is no conflict of interest.22 **Funding:** The first author received a scholarship from the Greek State Scholarship Foundation and
23 the European Union

24

25

26 **Abstract (word count= 200, word limit=250)**

27

28 **Background/Objectives:** Eating out-of-home has been associated with the increasing prevalence of
29 obesity. While some chain-restaurants provide nutritional information for their products, smaller
30 independent catering facilities may not provide such information. The aim of this study was to
31 assess the nutritional adequacy of meals provided to young adults at an independent catering facility
32 and compare them with meals provided by chain-restaurants.

33 **Subjects/Methods:** Meals were analysed in 2014 in the UK in relation of nutrient provision to
34 targets for macro- and micro-nutrients. One-way ANOVA was performed to compare menus
35 between the restaurants included in the analyses.

36 **Results:** 2,056 meal-combinations were analysed, 210 from the student accommodation and 1,846
37 from five largest national chain-restaurants. Mean (SD) nutritional content was; student
38 accommodation: 1193(269)kcal, fat 52.0(22)g, saturated fat 24.5(14.5)g, protein 42.4(28.5)g,
39 carbohydrate 117.0(30)g, chain-restaurants: 922(160)kcal, fat 40.0(9.7)g, saturated fat 14.5(5.8)g,
40 protein 31.2(6.5)g, carbohydrate 104.2(16.6)g. Meals from the student accommodation presented
41 significantly more calories than the meals in all five chain-restaurants ($p=0.0015$).

42 **Conclusions:** Meal-provision in the student accommodation was in excess of energy-requirements
43 and higher than the meals offered in chain-restaurants. Regulating or setting nutritional standards
44 for all places that provide food is essential as current food provision may favour unwanted weight-
45 gain and diet-related diseases.

46 **Background/Objectives**

47 Food eaten out of home has been suggested as a contributing factor to the obesity epidemic (Jeffery
48 et al. 2006), and portion sizes have increased steadily over the past 20 years (Benson, 2009). Chain-
49 restaurants have been particularly criticised both for the nutritional quality of the food they provide
50 and the way they advertise their products, especially to children. Chain-restaurants, however,
51 account only for approximately 25% of the total sales of food eaten outside home in the UK, while
52 small independent catering facilities account for the majority (75%) (DEFRA,2011). More
53 importantly, catering facilities in institutions such as universities, hospitals, and schools that usually
54 provide more than one meal/day to service users are subject to minimal nutritional control posing a
55 greater risk for the development of obesity and diet-related diseases. Only since April 2015,
56 compulsory food standards were introduced in hospitals in England but still food quality varies
57 greatly between hospitals and areas. Attention has been drawn to the nutritional content of chain-
58 restaurant food, but no studies exist on nutritional provision in settings such as student
59 accommodation, which provide food on a repeated daily basis. Catered accommodation takes the
60 pressure off young adults from preparing and cooking their meals, while cost is also included in the
61 accommodation price but this convenience comes with uncertain health implications. Food eaten
62 outside home is often higher in calories and of poorer nutritionally quality (Prentice & Jebb, 2003)
63 and people rarely can estimate the calorie content or the nutritional value of the food on offer
64 (Chandon & Wansink, 2007). This is a particular public health concern because young adulthood;
65 once a period of optimal health is now a period where rapid weight gain occurs (Nikolaou et
66 al.2015, Crombie et al.2009) and poor eating habits can be established (Demory-Luce et al 2004).
67 Students also can often be food insecure and relying solely on the food provided by the university
68 due to the cost of studying and the cost of food (Hughes, 2012). Poor diet is not only a major
69 contributing factor for chronic diseases, but can also affect the academic performance of students

70 (Florence et al. 2008). An awareness of the energy contents of meals may be valuable for
71 interventions to prevent unwanted weight gain, diet-related diseases, and nudge caterers to review
72 and modify the nutritional content of the food on offer

73 The present study analysed the nutritional content of the menu offered to university students living
74 in catered accommodation and 1) compare it to current dietary recommendations and 2) with
75 popular meals provided by the five largest chain restaurants in the UK.

76 **Subjects and Methods**

77 This study was exempt from IRB approval.

78 **Location of catering facilities included in the study**

79 1) The catered accommodation, located in an urban university, provides accommodation for
80 approximately 120 students, mostly 1st year. Breakfast and evening meals are included in the
81 accommodation price. The hall is located a 40-minute walk or 10-15 minute bus journey
82 away the main University's campus. The closest retail grocery outlets were a 20 minute-
83 walk away. Very limited cooking facilities are available on-site, so most students rely on
84 meals provided in the hall.

85 2) The top five chain restaurants holding the largest share in the UK market were included in
86 this study. Those were, according to the turnover in millions 1) McDonalds (£1,872m) 2) JD
87 Wetherspoons (£ 1,038m) Greggs (£708m), 4) KFC (£614m) and 5) Costa coffee (£610m)
88 (Statistica, 2013) Outlets of these five chain-restaurants were within the same postcode as
89 the catered accommodation (Figure 1).

90 **Menus**

91 1) The menu comprised a five-week cycle, developed by the catering staff employed by the
92 University, without any nutritional guidance. The main focus of caterers was on low food-

93 costs and wastage. All meals were prepared and cooked on site by in-house catering staff
94 using standardised recipes. For meat and fish products, standard portions were purchased
95 (110g per portion of meat and 170g per portion for fish) and other cooked food (eg rice,
96 soup, potatoes) was served using standard utensils allowing the estimation of the calorie
97 content of meals. Recipes, including ingredient lists, for all dishes served in the hall, were
98 made available to the researcher for analysis by catering staff. Stock rooms were checked
99 and all commercially prepared ingredients used for cooking or preparations of the meals
100 were recorded.

101 2) All of the chain-restaurants had nutritional information of the products on sale available on-
102 line and at the point of sale. Nutritional information available on-line included number of
103 calories, amount of fat, saturated fat and salt. Three of these restaurants (McDonalds,
104 Greggs, KFC) have made voluntarily pledges to reduce the calories consumed and started
105 posting calorie information at the point of sale from 1 Jan 2013, under the English Health
106 Department Responsibility Deal (DoH, 2014). The two remaining (JD Wetherspoons and
107 Costa coffee) are partners in the Responsibility Deal since 2011.

108

109 **Meal combinations**

110 1) Evening meals comprised three courses. From a choice of three different main course
111 dishes, students could only choose one, as well as a starter, dessert and a piece of fruit, or
112 two pieces of fruit, or a piece of fruit and a pot of yogurt (125ml). The side dishes (rice,
113 chips, and vegetables) were served using the same utensils to ensure portion-size
114 consistency. In view of the numerous possible meal combinations, for this study it was
115 assumed that each service user would choose the three-course meal. The meal-options for
116 analysis were created by two formulae: meal-option 1 = starter + main course (from a choice

117 of 3) + side dishes + dessert; meal-option 2 = starter + main course (from a choice of 3) +
118 side dishes + item of fruit.

119 2) Meal-combinations were created from the items listed using main course food items and a
120 dessert, and a side dish. A side dish was used in order to be directly comparable with the
121 meals provided at the hall.

122

123 **Data Analyses/Statistics**

124 All possible meal-combinations provided in the hall were compiled using an Access database
125 (Microsoft Office Access, 2013). Nutritional composition of macronutrients and micronutrients of
126 meal choices of the evening meal were determined using nutrient analysis software (WinDiets,
127 Robert Gordon University, 2010). When commercial ingredients used in the preparation of meals
128 were not listed, then that item was added to the database using manufacturer's nutrient values.

129 All possible meal-combinations provided in the five chain-restaurants were compiled in an Excel
130 database using the nutritional information provided on the websites of the companies.

131 After data had been checked for normalcy, mean values and distributions of macronutrients and
132 micronutrients were calculated and related to the UK Dietary Reference Values (DRVs) for
133 macronutrients and Reference Nutrient Intake (RNIs) for micronutrients (DoH, 1991).

134 One-way analysis of variance (ANOVA) was used to test for differences between the catered
135 accommodation menu and the chain-restaurants' menus for calories, macronutrients, and sodium,
136 using SPSS 21 (SPSS, Chicago). Post hoc comparisons between pairs were also conducted. In order
137 to keep the probability of type 1 error below 0.05, our main comparison pair was between the
138 means of the catered accommodation menu and the means of all chain restaurants' menus.

139 **Results**

140 **Accommodation Meals**

141 A total of 210 combinations for evening meal choices were created from the five-week menu cycle,
142 105 with a dessert (meal-option 1) and 105 replacing the dessert with two pieces of fruit (meal-
143 option 2). Mean values for macronutrients and micronutrient contents are shown on **Table 1**. The
144 mean (SD) nutritional contents of meal-option 1 were 1193(269)kcal, fat 52.0(22)g, saturated fat
145 24.5(14.5)g, protein 42.4(28.5)g, carbohydrate 117.0(30)g, vitamin C 74(78)mg, iron 14(9)g,
146 calcium 217(211)mg. Proportional energy contents were carbohydrate 39%, protein 14%, fat 39%,
147 sat fat 18%. Mean nutritional contents of meal-option 2 were; 896(215)kcal, fat 29.0(17)g,
148 saturated fat 9.9(8)g, protein 35.0(22)g, carbohydrate 125.0(24)g, vitamin C 90(75)mg, iron 6.3(2)g,
149 calcium 206(189)mg. Proportional energy contents for meal-option 2 were carbohydrate 56%,
150 protein 16%, fat 29%, sat fat 10%.

151

152 **Nutritional adequacy of meal-options.**

153 In general, almost all meal-options (86%-100%) exceeded the 30% recommendations for energy
154 and for macronutrients. While most meal-options were adequate for most micronutrients, there
155 were marked short-falls for some, especially iodine and calcium.

156

157 **Meal-option 1**

158 *Young men:* Expressed as % of Guideline Daily Amount (GDA) the mean values of macronutrients
159 for this option were 48% of energy, with 55% of fat, 80% of saturated fat, 45% of protein and 98%
160 of carbohydrate. Almost all meals exceeded the 30% recommendation for energy (n=105, 100%),
161 for fat (n=103, 98%), for saturated fat (n=88, 84%), for protein (n=95, 90%), and for carbohydrate
162 (n=105, 100%). The 30% recommendation was only met by 33% (n=35) of the meals for iodine,

163 13% (n=15) for calcium. A significant proportion of meals exceeded the 100% recommendation for
164 fat (n=10, 9%) and for saturated fat (n=20, 18%).

165 *Young women:* Expressed as %GDA, this option provided means of 60% of energy, 74% of fat,
166 120% of saturated fat, 56% of protein, and 130% of carbohydrate of the GDA. All meals exceeded
167 the 30% recommendation for energy (n=105, 100%), for fat (n=105, 100%), for saturated fat
168 (n=105, 100%), and for carbohydrate (n=105, 100%). The 30% recommendation was only met by
169 10% (n=9) of the meals for iodine, and 5% (n=5) for calcium. A significant proportion of meals
170 exceeded the 100% recommendation for fat (n=15, 14%) and for saturated fat (n=75, 71%).

171

172 **Meal-option 2**

173 *Young men:* Expressed as %GDA, this option provided means 36% of energy, 30% of fat, 33% of
174 saturated fat, 37% of protein, and 104% carbohydrate. A high proportion of meals exceeded the
175 30% recommendation for energy (n=90, 86%), for fat (n=88, 84%), for saturated fat (n=40, 38%),
176 for protein (n=90, 86%), and for carbohydrate (n=105, 100%). The 30% recommendation was only
177 met by 18% (n=20) of meals for iodine, 20% (n=18) for calcium. Only a small proportion of meals
178 exceeded the 100% recommendation for saturated fat (n=5, 5%).

179 *Young women:* Expressed as %GDA, this option provided means of 45% of energy, 41% of fat,
180 50% of saturated fat, 47% of protein, and 139% of carbohydrate. Most meals exceeded the 30%
181 recommendation for energy (n=100, 95%), for fat (n=96, 91%), for saturated fat (n=55, 52%), for
182 protein (n=103, 98%), and for carbohydrate (n=105, 100%). The 30% recommendation was only
183 met by 18% (n=20) of meals for iodine, 20% (n=18) for calcium. Only a small proportion of meals
184 exceeded the 100% recommendation for saturated fat (n=15, 14%).

185

186

187 **Chain-restaurants Menus**

188 In total 1,846 meal-combinations were created for the five chain-restaurants; 1) McDonalds=799, 2)
189 JD Wetherspoons=496, 3) Greggs=180, 4) KFC=143, 5) Costa coffee=228. Mean values, standard
190 deviation, and the range for calories, fat, saturated fat, protein, carbohydrate and sodium are shown
191 in **Table 2** (JD Wetherspoons provides only the calorie content of meals and Greggs provides does
192 not provide information on saturated fat).

193

194 **Comparison between independent catering facility and chain-restaurants menus**

195 Catered accommodation and chain restaurants menus were compared for calorie and macronutrient
196 content. Meal-option 1 was significantly higher in calories than all the meals offered in all of the
197 five-chain restaurants ($p=0.015$), fat ($p= 0.02$) and saturated fat ($p<0.001$). Meal option-2 was not
198 different than the menus in the five chain restaurants.

199

200 **Discussion**

201 The current study aimed to assess the suitability of the nutritional composition of meals offered to
202 young adults provided in a small independent catering facility within a UK University, compared
203 with guidelines and with menus offered from chain-restaurants. The hall provided meals in excess
204 of the 30% of requirements and so did the chain restaurants. Guidance from the former UK Food
205 Standards Agency (FSA), now Department of Health, recommends that an evening meal should
206 provide 30% of Guideline Daily Amount (GDA) of energy requirements (FSA, 2011), i.e. 600 kcal
207 for women and 750 kcal for men. The meals offered to young adults in this hall greatly exceeded

208 this 30% energy recommendation, two-fold for energy and up to three-fold for certain
209 macronutrients. For many micronutrients, most meal-options were ample, and indeed exceeded
210 100% of daily requirements, but low contents were seen for some essential nutrients like iodine and
211 calcium and vitamin D. Young adults often follow chaotic lifestyles and may commonly skip
212 meals, or need extra energy to sustain sporting activities, in which case providing extra energy at
213 evening meals might be desirable. However, since few young adults now engage in regular sporting
214 activity and many are prone to rapid unwanted weight gain (Nikolaou et al.2015), routinely
215 providing energy well above average requirements is unwise, at least without some warning. A
216 youth choosing the highest calorie three-course option would exceed these recommended amounts
217 for the evening meal by 1,630 kcal for women and 1,480 kcal for men. Making such high energy
218 choices every day is improbable, but would lead to weight gains of about 6 kg per month, assuming
219 that each kg weight gain requires a surplus intake of 7,000 kcal (Garrow, 1974). Assuming that
220 young adults had average requirements and over time chose a range of meal-options which provided
221 the average energy content (1193 kcal, about 400kcal above requirement), this excess from evening
222 meals alone would still lead to weight gain would still be about 1.6 kg/ month.

223 When the meals provided in the accommodation were compared with the menus offered in chain
224 restaurants, they proved to be 8%-35% higher in calorie content. Since, large chain-restaurants
225 accounts for only a small proportion of the food eaten outside home, targeting only those, as in New
226 York (Bernell, 2010), or under the English DoH 'Responsibility Deal' (DoH, 2014) to provide
227 calorie information will have only a minor impact on the obesity crisis. A catered student
228 accommodation is an interesting and important example of a smaller independent catering outlets,
229 as it provides meals to young adults on a daily basis, which may have cumulative effects on health.
230 It is likely that the patterns of nutrient contents will be similar in other commercial outlets.
231 Consumers need better provision, in terms of menu and recipe design, at least to allow nutritionally

232 balanced meals to be chosen by those who want them, and to warn them when energy or nutrient
233 contents are undesirable for long-term health.

234

235 Without nutritional information, it would be difficult for consumers to guess how the energy or
236 nutrient contents relate to their needs or to recommended values for a meal. Even trained health
237 professionals, such as dietitians, struggle to estimate nutritional contents of foods accurately
238 (Backstard et al. 1997) or to identify when a menu is meeting the requirements for specific
239 macronutrients. Commercial ready meals are currently nutritionally chaotic (Celnik et al.2012). It
240 has been suggested that to improve health meals should be designed to provide 30% of
241 macronutrients and micronutrients, as a 'default position'. This can easily be achieved by modifying
242 recipe-ingredients, as shown by modifying traditional pizza recipes to match nutritional guidelines
243 (Combet et al.2014).

244

245 There are several limitations in this study with the most important being the inability to measure the
246 nutrient content of meals on offer with direct calorimetry. However, all meals are cooked on site
247 with fresh ingredients and the principal researcher worked closely with the catering staff in order to
248 get details of each recipe used. Also, quantities of ingredients used for preparing each recipe were
249 confirmed by examining the ingredients orders and the stock room. Students residing in the hall
250 were also observed during meal times hence the meal-options created and analysed for this study
251 were realistic (Nikolaou et al. 2015).

252

253 Nutritional analyses of meal-options offered to young adults in an independent catering facility
254 revealed excessive energy, carbohydrate, fat and saturated fat, and variable micronutrient contents
255 compared to current recommendations. While desirable in certain cases, excess energy content is
256 hard to identify and may promote unwanted weight gain and consumption of nutritionally

257 unbalanced diets. The nutritional profile of those menus was worse when compared with menus
258 offered from chain-restaurants.

259

260

261

262

263 **References**

- 264 Backstrand, JR. Wootan, MG. Young, LR. Hurley, J. (1997) Fat Chance: A Survey of
 265 Dietitians' Knowledge of the Calories and Fat in Restaurant Meals.. Washington, DC:
 266 Center for the Science in the Public Interest.
- 267 Benson,C.(2009) 'Increasing portion size in Britain, *Society, biology and human affairs*,
 268 74(2):4-20.
- 269 Bernell, B. (2010). History and Impact of the New York City Menu Labeling Law, *The. Food &*
 270 *Drug LJ*, 65, 839.
- 271 Celnik, D, Gillespie, L. and Lean, MEJ. (2012) Time-scarcity ready-meals, ill-health and the
 272 obesity epidemic. *Trend in Food Science & Technology*, 4-11.
- 273 Chandon, P., & Wansink, B. (2007). The biasing health halos of fast-food restaurant health
 274 claims: lower calorie estimates and higher side-dish consumption intentions. *Journal of*
 275 *Consumer Research*, 34(3), 301-314.
- 276 Crombie, AP. Ilich, JZ. Dutton, GR. Panton, LB. Abood, DA. (2009) The freshman weight gain
 277 phenomenon revisited. *Nutr Rev*, 67(2):83-94.
- 278 Combet, E. Jarlot, A. Aidoo, K. Lean, MEJ. (2014) Development of a nutritionally balanced
 279 pizza as a functional meal designed to meet published dietary guidelines. *Pub Health*
 280 *Nut*; 17(11):2577-86.
- 281 Demory-Luce, D. Morales, M. Nicklas, T. Baranowski, T. Zakeri, I. Berenson, G. (2004)
 282 Changes in food group consumption patterns from childhood to young adulthood: the
 283 Bogalusa Heart Study. *Journal of the American Dietetic Association*, 104(11), pp.1684-
 284 1691.
- 285 Department for Environment, Food and Rural Affairs. Family Food.
 286 https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/193804/familyfood-2011report.pdf. Accessed on 23/08/2014
- 287
 288 Department of Health. <https://responsibilitydeal.dh.gov.uk/tag/calorie-labelling/> Accessed on
 289 25/09/2014.
- 290 Department of Health. Report on Health and Social Subjects No 41. (1991) Dietary Reference
 291 Values for Food Energy and Nutrients for the United Kingdom. London: HMSO.
- 292 Florence, M. Asbridge, M. Veugelers, P. (2008) Diet quality and academic performance. *J Sch*
 293 *Health*, 78(4):209-215.

- 294 Food Standards Agency. FSA nutrient and food based guidelines for UK institutions.
295 <http://www.food.gov.uk/multimedia/pdfs/nutguideuk.pdf>.
- 296 Garrow, JS. (1974) Methods for measuring energy expenditure: In energy balance and obesity
297 in man.. Amsterdam. North-Holland Publishing Company.
- 298 Hughes, R. (2012) Food insecurity amongst university students [online]. *Nutridate*, 23 (2): 2-4.
299 Availability:<<http://search.informit.com.au/documentSummary;dn=733756016916007;res=IELHEA>> ISSN: 1835-5080. [cited 11 Nov 16].
- 300
- 301 Jeffery, RW. Baxter, J. McGuire, M. Linde, J. (2006) Are fast food restaurants an environmental
302 risk factor for obesity *International Journal of Behavioral Nutrition and Physical*
303 *Activity*, 3:2.
- 304 Leading ten chain restaurants in the United Kingdom (UK) in 2013, by sales revenue (in million
305 GBP). The statistics Portal. 2013. [https://www.statista.com/statistics/307866/top-ten-](https://www.statista.com/statistics/307866/top-ten-chain-restaurants-in-the-united-kingdom-by-revenue/)
306 [chain-restaurants-in-the-united-kingdom-by-revenue/](https://www.statista.com/statistics/307866/top-ten-chain-restaurants-in-the-united-kingdom-by-revenue/) (accessed on 14/11/2016).
- 307 Nikolaou, CK. Hankey, CR. Lean, ME. (2015) Weight changes in young adults: a mixed-
308 methods study. *International journal of obesity*, Mar 1;39(3):508-13.
- 309 Nikolaou, C. K., Hankey, C. R., & Lean, M. E. J. (2015). Effects of calorie labelling on macro-
310 and micro-nutrients in main-meal choices made by young adults. *European journal of*
311 *clinical nutrition*.
- 312 Prentice, AM. Jebb, SA. (2003) Fast foods, energy density and obesity: a possible mechanistic
313 link. *Obes Rev*, 4:187-194.
- 314
- 315
- 316

318 **Table 1: Nutritional compositions of evening meal-options from the UK for macronutrients and**
 319 **micronutrients, analysed in 2014**

Nutrient	Meal-option 1		Meal-option 2		DRV/RNI	
	Mean	Range	Mean	Range	Females	Males
Energy (kcal)	1193	858-1774	896	594-1431	2000	2500
Fat (g)	52	17.6-105.5	29	5.6-78.5	70	95
Saturated Fat (g)	24.5	4.7-60.3	9.9	0.8-34.9	20	30
Protein (g)	42.4	15.8-129.3	35	15-119	75	94
Carbohydrate (g)	117	86-188	125	83-166	90	120
Vitamin A (ug)	1092	28-3390	1167	202-3255	600	700
Thiamin (mg)	24	0.12-433	24	0.2-433	0.8	1.1
Riboflavin (mg)	0.81	0.09-3.58	0.87	0.2-3.7	1.1	1.3
Niacin (mg)	25	2.4-105	25	2-105	14	18
Vitamin B6 (mg)	2.9	0.39-15	3.3	0.9-15.5	1.2	1.5
Vitamin B12 (ug)	1.5	0-5	1.2	0-5	1.5	1.5
Folate (ug)	105	30-330	124	54-359	200	200
Vitamin C (mg)	74	4.5-237	90	23.2-256	40	40
Vitamin D (ug)	4.5	0-13	5.3	1-13.5	200	200
Calcium (mg)	217	37-819	206	47-849	800	1000
Magnesium (mg)	102	38-228	180	31-114	300	300
Sodium (mg)	1295	141-2972	1281	49-2987	6000	6000
Potassium (mg)	1514	622-3419	604	234-1373	3.5	3.5
Iron (g)	14	2.5-59	6.3	3.8-12.2	14.8	8.7
Zn (mg)	19	0.6-149	16	2-150	7	9.5
Mn (mg)	3.4	0.3-28.5	3.1	0.7-29	1.4	1.4
Se (ug)	26	1.1-163.4	28	3-165	60	70
I (ug)	210	8-2678	145	0-2678	150	150

320 DRV=Dietary Reference Value, RNI=Reference Nutrient Intake


321 **Table 2:** Mean nutritional content of meals provided to the five biggest chain-restaurants in the UK for energy and key macronutrients, analysed in
 322 2014
 323







	Calories (kcal)		Fat (g)		Saturated Fat (g)		Carbohydrate (g)		Protein (g)		Sodium (mg)	
	Mean (SD)	Range	Mean (SD)	Range	Mean (SD)	Range	Mean (SD)	Range	Mean (SD)	Range	Mean (SD)	Range
McDonalds	820 (289)	292-1240	26.4(16.9)	8.0-67.0	14.5 (9.0)	2.0-27.0	88.9 (29.9)	40.0-137	32.2(12.8)	11.0-49.0	945 (315)	433-1575
JD Wetherspoons	1,101 (267)	688-1,456	-	-	-	-	-	-	-	-	-	-
Greggs	885 (181)	630-1,170	45.2 (12.1)	27.0-59.0	-	-	107.3 (13.6)	88.5-116.5	26.7 (4.4)	22.0-31.5	1,221 (276)	788-1,536
KFC	773 (61)	430-1,100	39.8 (9.9)	19-54	11.7 (3.8)	5.5-47	80.9 (15.9)	48-122	24.5 (6.3)	17-37	1,604.3 (363.4)	880-2,300
Costa Coffee	855 (149)	648-1,132	35 (9.1)	18.5-49.0	12 (6.6)	3.0-24.5	105.7 (22.6)	79.8-142.8	26.6 (5.4)	19.9-36	947 (313)	501-1,410
Meal 1	1,193 (268)	858-1,774	52 (22)	17.6-105.5	24.5 (14.4)	4.7-60.3	117 (30)	86-188	42.4 (28.5)	15.8-129.3	1,295 (793)	141-2,972
Meal 2	896 (215)	594-1,431	29 (17)	5.6-78.5	9.9 (8)	0.8-34.9	125 (24)	83-166	35 (22)	15-119	1,281 (753)	49-2,987

324 ***JD Wetherspoons only provided information on the calorie content of food items on the menu**

325 ***Greggs did not provide the saturated fat content of food items on sal3**

Figure 1: Location of the catering outlets compared in this study in Scotland, UK



-  Costa coffee
-  Greggs
-  KFC
-  McDonalds
-  Student accommodation
-  Wetherspoons

All outlets are within the same postcode, G20.