



This is a repository copy of *Science requirement document (SRD) for the European Solar Telescope (EST) (2nd edition, December 2019)*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/154969/>

Version: Submitted Version

Article:

Schlichenmaier, R., Rubio, L.R.B., Collados, M. et al. (23 more authors) (Submitted: 2019) Science requirement document (SRD) for the European Solar Telescope (EST) (2nd edition, December 2019). arXiv. (Submitted)

© 2019 The Author(s). For reuse permissions, please contact the Author(s).

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Science Requirement Document (SRD)
for the
European Solar Telescope (EST)

2nd edition, December 2019

Written by: Schlichenmaier, R.¹; Bellot Rubio, L.R.²; Collados, M.^{3,4}; Erdelyi, R.^{5,6}; Feller, A.⁷; Fletcher, L.^{8,13}; Jurčák, J.⁹; Khomenko, E.³; Leenaarts, J.¹⁰; Matthews, S.¹¹; Belluzzi, L.^{12,1}; Carlsson, M.^{13,14}; Dalmasse, K.¹⁵; Danilovic, S.¹⁰; Gömöry, P.¹⁶; Kuckein, C.¹⁷; Manso Sainz, R.⁷; Martínez González, M.³; Mathioudakis, M.¹⁸; Ortiz, A.^{13,14}; Riethmüller, T.L.⁷; Rouppe van der Voort, L.^{13,14}; Simoes, P.J.A.¹⁹; Trujillo Bueno, J.^{3,20}; Utz, D.^{2,21}; Zuccarello, F.²²

(Author affiliations given on page 3.)

Discussed and approved by the EST Science Advisory Group (SAG)



“This project has received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No 739 500”

EST Science Advisory Group (SAG) members

1	Luis	Bellot Rubio	AA, Spain	Chair of Sect. II/1
2	Luca	Belluzzi	IRSOL, Switzerland	
3	Mats	Carlsson	UiO, Norway	
4	Sanja	Danilovic	SU, Sweden	
5	Robertus	Erdelyi	HSPF, Hungary	Chair of Sect. II/2
6	Alex	Feller	MPS, Germany	Chair of Sect. II/8
7	Lyndsay	Fletcher	University of Glasgow, UK	Chair of Sect. II/6
8	Peter	Gömöry	AISAS, Slovakia	
9	Jan	Jurčák	CAS, Czech Republic	Chair of Sect. II/4
10	Elena	Khomenko	IAC, Spain	Chair of Sect. II/7
11	Christoph	Kuckein	AIP, Germany	
12	Jorrit	Leenaarts	SU, Sweden	Chair of Sect. II/3
13	Arturo	López Ariste	CNRS, France	
14	María Jesús	Martínez González	IAC, Spain	
15	Mihalis	Mathioudakis	QUB, UK	
16	Sarah	Matthews	UCL, UK	Chair of Sects II/5 & II/9
17	Ada	Ortiz	UiO, Norway	
18	Rolf	Schlichenmaier	KIS, Germany	Chair of EST SAG
19	Javier	Trujillo Bueno	IAC, Spain	
20	Dominik	Utz	IGAM, Austria	
21	Luc	Roupe van der Voort	UiO, Norway	
22	Francesca	Zuccarello	Univ. of Catania & INAF, Italy	

Author affiliations:

- ¹: Leibniz-Institut für Sonnenphysik (KIS), Schöneckstr. 6, 79104 Freiburg, Germany
- ²: Instituto de Astrofísica de Andalucía (IAA), Glorieta de la Astronomía s/n, Granada, Spain
- ³: Instituto de Astrofísica de Canarias (IAC), Vía Láctea, 38205 La Laguna, Tenerife, Spain
- ⁴: Departamento de Astrofísica, Universidad de La Laguna, 38205 La Laguna, Tenerife, Spain
- ⁵: SP2RC, University of Sheffield, Sheffield, UK
- ⁶: Dept of Astronomy, Eötvös University, Budapest, Hungary
- ⁷: Max-Planck-Institut für Sonnensystemforschung (MPS), Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany
- ⁸: SUPA School of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, UK
- ⁹: Astronomical Institute of the Czech Academy of Sciences, Fričova 298, 25165 Ondřejov, Czech Republic
- ¹⁰: Institute for Solar Physics, Department of Astronomy, Stockholm University, Albanova University Center, 10691 Stockholm, Sweden
- ¹¹: UCL Mullard Space Science Laboratory, Holmbury St Mary, Dorking RH5 6NT, UK
- ¹²: Istituto Ricerche Solari Locarno (IRSOL), Via Patocchi 57, CH-6605 Locarno Monti, Switzerland
- ¹³: Rosseland Centre for Solar Physics, University of Oslo, PO Box 1029, Blindern 0315, Oslo, Norway
- ¹⁴: Institute of Theoretical Astrophysics, University of Oslo, PO Box 1029, Blindern 0315, Oslo, Norway
- ¹⁵: IRAP, Université de Toulouse, CNRS, CNES, UPS, F-31028 Toulouse, France
- ¹⁶: Astronomical Institute of the Slovak Academy of Sciences, 05960 Tatranská Lomnica, Slovakia
- ¹⁷: Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany
- ¹⁸: Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, Belfast, BT7 1NN, Northern Ireland, U.K.
- ¹⁹: Centro de Rádio Astronomia e Astrofísica Mackenzie, Escola de Engenharia, Universidade Presbiteriana Mackenzie, São Paulo, Brazil
- ²⁰: Consejo Superior de Investigaciones Científicas, Spain
- ²¹: IGAM/Institute of Physics, Karl-Franzens University Graz, Austria
- ²²: Dipartimento di Fisica e Astronomia "Ettore Majorana", University of Catania, Via S. Sofia 78, 95123 Catania, Italy

Contents

I	Introduction	7
II	Top-level science goals	8
1	Structure and evolution of magnetic flux	8
1.1	Formation and disappearance of kG flux concentrations in the solar photosphere	8
1.2	Internal structure of small-scale flux concentrations	9
1.3	Magnetic bright points	10
1.4	Small-scale flux emergence in the quiet Sun	10
1.5	Magnetic flux cancellations in the quiet Sun	11
1.6	Quiet Sun internetwork fields	12
1.7	Polar magnetic fields	13
1.8	Network dynamics	13
2	Wave coupling throughout solar atmosphere	15
2.1	MHD waves in localized quiet Sun structures	16
2.2	Magnetic twist and torsion	17
2.3	Wave propagation in active regions	19
2.4	Wave propagation in the quiet Sun	20
3	Chromospheric dynamics, magnetism, and heating	22
3.1	Magnetic structure at supergranular scales	22
3.2	Spicules and jets	23
3.3	Structure of small-scale chromospheric jets	24
3.4	Wave propagation, mode conversion and wave damping	24
3.5	Flux emergence and reconnection events	24
3.6	Observational determination of electric currents	25
3.7	Temperature structure of the solar chromosphere	26
3.8	Magnetic field measurements using Ca II H&K	26
3.9	Summary of requested instrument capabilities	26
4	Large scale magnetic structures: sunspots, prominences and filaments	27
4.1	Stability of the umbra	28
4.2	Umbral dots	28
4.3	Structure of cool sunspot umbrae	28
4.4	Umbral flashes as a probe of fine structure in the umbra chromosphere	29
4.5	Penumbral and umbral micro-jets	29
4.6	Light bridges	29
4.7	Evolution of an individual penumbral filament	30
4.8	Formation and decay of sunspot penumbrae	30
4.9	Relation between moat flows and sunspot decay	31
4.10	Fine structure of prominences and filaments	31
4.11	Are quiescent and active region filaments the same phenomenon?	32
4.12	Magnetic field and dynamics of tornado prominences	32

5	Coronal Science	33
5.1	Sunspot light-bridges/light walls	33
5.2	Light Walls	33
5.3	Origins of the solar wind	33
5.4	Probing pre-flare triggers	34
5.5	Macrospicules/spicules and Transition Regions Quakes (TRQs)	34
5.6	Ellerman bombs	34
6	Solar Flares and Eruptive Events	35
6.1	Thermal structure of flare chromosphere and photosphere: line observations	35
6.2	Thermal structure of flare chromosphere and photosphere: continuum observations	36
6.3	Velocity structure of the flaring atmosphere	36
6.4	Diagnostics for non-thermal particles	37
6.5	Oscillatory phenomena in flares	37
6.6	Sunquakes	37
6.7	Large-scale structure and evolution of the magnetic field	38
6.8	Filaments in flaring active regions	39
6.9	Coronal Mass Ejections	39
6.10	Structure and evolution of the magnetic field at small scales	39
7	Coupling in partially ionized solar plasma	39
7.1	Dynamics of partially ionized prominence plasma	40
7.2	Influence of partial ionization on spicules	41
7.3	Detection of partial ionization effects in the photosphere	42
7.4	Multi-fluid physics of chromospheric waves, shocks and swirls	42
7.5	Flares and energetic events	42
8	Scattering physics and Hanle-Zeeman diagnostics	43
8.1	Investigation of the small-scale magnetism of the quiet solar photosphere	44
8.2	Investigation of the magnetism of the solar chromosphere	47
8.3	Deepening our understanding of the physics of scattering polarization	50
9	Solar science exploration between 350 and 400 nm	51
9.1	White-light emission from flares - Continuum diagnostics in vicinity of Balmer jump	51
9.2	Coronal forbidden lines in the visible	51
9.3	Ca II H&K spectroscopy and spectropolarimetry	51
9.4	Multi-line spectropolarimetry	52
10	Tables for Observing Programmes	53
III	Discussion and Requirements	120
11	What are the particular strengths of the EST design?	120
11.1	EST preliminary design study 2008 - 2011	120
11.2	Design developments 2011 – 2019	120

12 Discussion	121
12.1 Discussion on photon flux	121
12.2 Discussion on field-of-view (FOV)	122
12.3 Discussion on pointing requirements	122
12.4 Discussion on wavelength range	122
12.5 Discussion on instrumentation and light distribution	123
13 Scientific Requirements	123
13.1 Telescope specifications	123
13.2 Requirements for instrumentation and light distribution:	124
IV Appendix, References, & Abbreviations	126
A Signal-to-noise ratio S/N	126
A.1 Optimum spatio-temporal resolution	126
A.2 Number of photons at diffraction limit	126
B References	126
C List of Abbreviations	138

Part I

Introduction

The European Solar Telescope (EST)¹ is a research infrastructure for solar physics. It is planned to be an on-axis solar telescope with an aperture of 4 m and equipped with an innovative suite of spectro-polarimetric and imaging post-focus instrumentation. The EST project was initiated and is driven by EAST², the European Association for Solar Telescopes. EAST was founded in 2006 as an association of 14 European countries. Today, as of December 2019, EAST consists of 26 European research institutes from 18 European countries.

The Preliminary Design Phase of EST was accomplished between 2008 and 2011. During this phase, in 2010, the first version of the EST Science Requirement Document (SRD) was published. After EST became a project on the ESFRI³ roadmap 2016, the preparatory phase started. This phase is partially supported by EU funding through the PRE-EST H2020 project⁴. The goal of the preparatory phase is to accomplish a final design for the telescope and the legal governance structure of EST. A major milestone on this path is to revisit and update the Science Requirement Document (SRD).

The EST Science Advisory Group (SAG) has been constituted by EAST and the Board of the PRE-EST⁴ EU project in November 2017 and has been charged with the task of providing with a final statement on the science requirements for EST. Based on the conceptual design, the SRD update takes into account recent technical and scientific developments, to ensure that EST provides significant advancement beyond the current state-of-the-art.

The present update of the EST SRD has been developed and discussed during a series of EST SAG meetings:

- 1st telecon meeting on Nov 5th, 2017
- 2nd meeting in Freiburg, Nov 24, 2017
- 3rd telecon meeting, Dec 15, 2017
- 4th telecon meeting, March 26, 2018
- 5th meeting in Belfast, April 16 & 17, 2018
- 6th meeting in Naxos, June 16, 2018
- 7th telecon meeting, January 14, 2019
- 8th telecon meeting, October 11, 2019
- 9th telecon meeting, October 22, 2019
- 10th telecon meeting, December 3, 2019

The SRD develops the top-level science objectives of EST into individual science cases. Identifying critical science requirements is one of its main goals. Those requirements will define the capabilities of EST and the post-focus instrument suite. The technical requirements for the final design of EST will be derived from the SRD.

The science cases presented in Part II (Sects. 1 to 8) are not intended to cover all the science questions to be addressed with EST, but rather to provide a precise overview of the capabilities that will make of EST a competitive state-of-the-art telescope to push the boundaries of our knowledge over the next few decades. The science cases contain detailed observing programmes specifying the type of observations needed to solve specific science problems. An effort is being made to define the parameters of the required observations as accurately as possible, taking into account both present capabilities and technological developments expected in the near future. The tables of the observing programmes corresponding to the science cases are compiled in Sect. 10. The EST science cases represent challenging observations that put strong constraints on the telescope and its instrument suite. Ultimately, they will be translated into Technical Requirement Document (TRD) leading to the final EST design to be implemented during the construction phase.

The unique design advantages of the EST concept is presented in Section 11. The effect of the science cases on the EST design are discussed in Section 12 and summarized in Section 13.

¹EST web link: <http://www.est-east.eu>

²EAST web link: <http://www.est-east.eu/est/index.php/people/>

³ESFRI: European Strategy Forum on Research Infrastructures. Web link to ESFRI roadmap in March 2016: <http://www.esfri.eu/roadmap-2016>

⁴PRE-EST: The EST Preparatory Phase has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement Nr. 739 500.

Part II

Top-level science goals

1 Structure and evolution of magnetic flux

Authors: Luis Bellot Rubio, Dominik Utz, Sanja Danilovic

Magneto-convection is an ubiquitous process in the solar surface. The interaction of magnetic fields and granular convection leads to the formation of magnetic features and a broad range of flows and waves on very small spatial and temporal scales. This makes the photosphere highly dynamic and the place where heating of the upper solar atmosphere could be initiated via small-scale flux emergence, flux cancellation, braiding of magnetic field lines, or generation and upward propagation of waves.

These processes need to be studied at their intrinsic spatial and temporal scales, of order 100 km and 10 s, respectively. For the most part, small-scale magnetic fields in the photosphere and chromosphere are weak, producing very small polarisation signals. To detect them, high polarimetric sensitivity is needed. The three requirements have been difficult to meet with existing telescopes and, as a result, temporal resolution has usually been sacrificed. Thus, our knowledge of the temporal evolution of quiet sun magnetic fields is very limited. With its large aperture primary mirror and dedicated suite of instruments, EST will for the first time provide the necessary sensitivity and spatial resolution to study the structure, dynamics, and evolution of the weak fields pervading the quiet solar surface, which according to recent estimates harbor orders of magnitude more magnetic flux than active regions and sunspots.

1.1 Formation and disappearance of kG flux concentrations in the solar photosphere

Outside of active regions, the solar magnetic field is organized in small-scale (~ 100 km) magnetic flux tubes and flux sheets (Howard & Stenflo 1972; Stenflo 1973; Stenflo et al. 1984; Berger et al. 2004). These features are observed in intergranular lanes and form the photospheric network, located at the borders of supergranular cells. Small-scale flux tubes are believed to play an important role in the irradiance variations of the Sun (Riethmüller et al. 2014) and the heating of the chromosphere and corona through a variety of dynamical processes, including the channeling of waves, shock fronts, and solar tornadoes (see Wedemeyer-Böhm et al. 2012).

The magnetic field of network flux tubes is as strong as 1500 G (e.g., Bellot Rubio et al. 2000b; Utz et al. 2013), i.e., well above the equipartition value of a few hundred G. Therefore, in addition to the kinematic concentration of flux by horizontal convective motions, another mechanism capable of enhancing the field up to kG strengths is required. In the late 1970s, it was theoretically proposed that the field is amplified by a thermal instability known as convective collapse (Parker 1978; Spruit 1979). This process leads to rapid downflows in the tube's interior, causing a strong evacuation of the magnetic element and a concentration of the field. The spatial and temporal scales associated with the formation of intense flux tubes are on the order of 100 km and 30 s, according to numerical simulations by Danilovic et al. (2010a).

Unfortunately, convective collapse has proven very difficult to detect. Only a limited number of single case studies and statistical analyses have been presented in the literature (e.g., Bellot Rubio et al. 2001; Nagata et al. 2008; Fischer et al. 2009; Narayan 2011). Thus, we still lack a clear, convincing observational picture of the formation of kG flux concentrations on the solar surface. One of the main problems is that the intensification events recorded so far do not produce stable kG features. In most cases, rapid upflows develop while the field is amplified, leading to the disappearance of the magnetic features (Bellot Rubio et al. 2001; Socas-Navarro & Manso Sainz 2005). Some recent studies suggest that the newly created elements undergo oscillations in field strength and size (Requerey et al. 2014; Utz et al. 2014). This may partly explain the observed lack of stability: the features would seem to disappear if the polarisation signals go below the detection threshold during specific phases of the oscillation. Better polarimetric sensitivity than currently available is needed to verify this conjecture.

Interestingly, the rapid photospheric downflows generated by convective collapse should produce even faster flows in the chromosphere, due to its lower density. This provides another means to detect convective collapse events. However, only Fischer et al. (2009) have studied the response of the chromosphere to convective collapse. Using Dopplergrams in the $\text{Mg I } b_2$ line taken by the Hinode NFI, they concluded that a significant fraction of the field amplification events observed in the photosphere show rapid downflows near the temperature minimum region. Clearly, a full description of the formation process of kG features calls for simultaneous spectropolarimetric measurements in the photosphere, temperature minimum region, and chromosphere, at very high cadence (~ 10 s). The required multi-line observations are currently out of reach for existing telescopes, due to the lack of proper instrumentation (integral field spectropolarimeters or a suite of imaging polarimeters working in parallel).

The evolution and physical processes that lead to the disappearance of kG flux tubes are not well understood either. On large scales, network regions do not change significantly in the course of hours/days, but on the smallest scales the

flux evolves rapidly (e.g., Gošić et al. 2016). Small-scale magnetic elements change shape, move around pushed by granular convection, and interact with other flux concentrations, merging or cancelling with them (Requerey et al. 2014). During their lifetime, they may brighten considerably and be detected as magnetic bright points (MBPs). It is possible to study these features using broad-band imaging. However, the brightness enhancement is transitory and MBPs usually disappear much earlier than the magnetic field itself. We still do not know why. To understand these processes, we need ultra-high resolution imaging and spectropolarimetry, in order to determine the full vector magnetic field. Unfortunately, spectropolarimetric measurements are scarce, due to the very small spatial and temporal scales involved. The evolution and eventual disappearance of flux tubes would be a natural consequence of their interaction with convective motions if they are a surface phenomenon, as indicated by recent numerical simulations. On the other hand, the persistence of some network features suggests that they are rooted below the solar surface. In that case, other mechanisms might be responsible for the disappearance of the flux.

OP 1.1.1 Formation and evolution of intense flux tubes in the solar atmosphere

See table on page 54.

1.2 Internal structure of small-scale flux concentrations

Our understanding of small-scale magnetic flux concentrations heavily relies upon magneto-convection simulations (e.g., Steiner et al. 1998; Vögler et al. 2005; Stein & Nordlund 2006; Yelles Chaouche et al. 2009; Rempel 2014). Conversely, an accurate observational characterisation of their internal structure may help validate and improve current computer models. Particularly interesting are the interfaces between magnetic and field-free regions. One example of such interfaces are the canopies of flux tubes expanding with height in the solar atmosphere. These are regions where one can expect electric currents, Joule dissipation, and a variety of dynamical flows.

Magnetic canopies and the flows associated with them were soon recognized as essential ingredients for explaining the asymmetries of the polarisation profiles that emerge from flux concentrations at a resolution of 1". However, a direct observational detection of such interfaces was not possible at the time. Only with Hinode and SUNRISE have we started to resolve the internal structure of magnetic elements, including their canopies.

Rezaei et al. (2007) presented evidence of canopies in magnetic elements from the spatial variation of the Stokes V area asymmetry of the Fe I 630 nm lines recorded by Hinode. The observed variations appear to agree well with those predicted by radiative magneto-hydrodynamic simulations. Martínez González et al. (2012) characterised the canopy of a large network flux concentration from an analysis of Fe I 525.02 measurements taken by SUNRISE. They too found a spatial distribution of Stokes V area asymmetries that agrees well with the results of MHD models. In particular, they located the height of the canopy as a function of radial distance from the center of the flux patch, finding an expanding structure, and determined the jumps in field strength and velocity across the discontinuity. More recently, Buehler et al. (2015) derived the 3D structure of plage flux concentrations using a spatially coupled inversion of the Fe I lines recorded by Hinode. For the first time, these authors were able to map the height variation of the vector magnetic field and flow field across the magnetic element. They verified the existence of a ring of strong downflows surrounding the flux concentration in deep photospheric layers and found hints of opposite magnetic polarities associated with the downflows. These features go well beyond the simple picture of magnetic flux tubes.

The next step is to determine the temporal evolution of flows and fields across magnetic elements, relating the observed changes to interactions with the surrounding granular convection (Requerey et al. 2014, 2015). This demands time series with cadences of less than 1 minute that cannot be obtained with existing telescopes due to insufficient photon flux.

It is also important to examine the differences between magnetic canopies in internetwork, network and plage regions through statistical analyses. How high are magnetic canopies located in the quietest and more active regions? What fraction of the surface is covered by strongly inclined fields in the chromosphere? What are their average lifetimes? To answer these questions, a good height coverage of the chromosphere and high polarimetric sensitivity are required.

Magnetic interfaces do not exist only as canopies in flux concentrations, but also as true discontinuities in other solar structures, e.g., magnetic field lines emerging in the photosphere or horizontal chromospheric fields (Schaffenberger et al. 2006). Hinode observations show a non-negligible fraction of Stokes V profiles with extreme asymmetries. Such asymmetries can only be produced by sharp magnetic field discontinuities in structures that do not seem to fit the picture of classical flux tubes (Sainz Dalda et al. 2012). The observed profiles change in the course of minutes, indicating a rapid evolution of the field and/or associated flows. EST, with its ultra-high resolution capabilities, will contribute to the characterisation of those fields and their temporal evolution in the photosphere and the chromosphere.

OP 1.2.1 Internal structure and evolution of magnetic elements

See table on page 55.

1.3 Magnetic bright points

Interestingly, the cross-section of small-scale kG flux tubes and flux sheets can be identified as bright points when observed in Fraunhofer's G band region of the solar spectrum (e.g., Steiner et al. 2001; Bodnárová et al. 2013). These features are called G-band bright points or more generally magnetic bright points (MBPs). MBPs are important because they represent the photospheric part of flux tubes reaching up to the upper atmosphere. Their buffeting and violent displacement by the surrounding granulation (Muller et al. 1989) create magnetohydrodynamic waves (e.g., Mathioudakis et al. 2013) traveling up into the upper atmosphere (see Sect. 2). Moreover, they have different total and spectral irradiances compared with their surroundings. Thus, changes in the appearance of MBPs over the solar cycle have consequences for the total solar irradiance variation and therefore for the climate on Earth.

Although MBPs are fundamental building blocks of the solar magnetism (Zwaan 1987), many basic questions are still unanswered. For example, it is not known if these flux elements can be created with arbitrarily small sizes. All studies trying to reveal the size distribution of MBPs so far had the problem that MBPs can be found all the way down to the diffraction limit of the telescope (e.g., Abramenko et al. 2010). Thus we need larger telescopes to detect the lower size limit of these features. Besides, even the shape of the size distribution itself is under discussion. Clearly, the physics creating an unlimited exponential distribution toward the smallest scales would be different from the physics creating a lognormal or Gaussian distribution of sizes.

Similarly, there have been problems in measuring a fundamental property of MBPs - their magnetic field strength. Due to their small size and highly dynamic evolution, the inferred field strength distributions contain a lot of noise and uncertainty (see, for example, Beck et al. 2007; Utz et al. 2013). Therefore, it is impossible to say if MBPs always appear with field strengths of roughly 1.3 kG as predicted by theory (Spruit 1979) and supported by MHD simulations (Riethmüller et al. 2014), or if they indeed possess a significant weak field component as deduced from real observations. To answer this question we need higher spatial and temporal resolution spectro-polarimetric data with increased sensitivity.

Another fundamental property of MBPs is their movement on the solar surface. We do not know if they follow random and erratic paths or more deterministic, regular trajectories. This is important in at least two ways. Firstly, as mentioned before, the driving of MBPs will create MHD waves which travel into the upper atmosphere and deposit energy there, thus contributing to chromospheric/coronal heating (e.g., Hasan & Ulmschneider 2004). Clearly, the kind of motion experienced by MBPs will strongly influence the type and amplitude of the resulting MHD waves (Muller et al. 1994). Secondly, the way MBPs move tells us about the diffusion of magnetic fields (Utz et al. 2010; Abramenko et al. 2011), hence by studying them we gain knowledge also on how large-scale structures such as sunspots disappear.

Besides of these basic properties we need to study the evolution of MBPs in more detail (e.g., Utz et al. 2014). While we believe they are created via the convective collapse process (see Sect. 1.1), we do not know much about their dissolution or about the processes they experience between creation and dissolution. This is due to their very short lifetimes (of the order of 2.5 min). With the cadences delivered by existing spectroscopic instruments, generally in the order of 30 s (Abramenko et al. 2010; Utz et al. 2010; Liu et al. 2018), we only have about 5 measurements of the properties of MBPs from creation to dissolution, which prohibits detailed studies of the involved physical processes. With EST it will be possible to increase the spatial resolution and, more importantly, the temporal cadence. This will enable us to study not only the basic properties of MBPs but also their whole evolution from birth to death, teaching us in that way fundamental plasma physics.

OP 1.3.1 Magnetic bright points

See table on page 56.

OP 1.3.2 Dynamic parameters and evolution of MBPs

See table on page 57.

1.4 Small-scale flux emergence in the quiet Sun

In the last two decades, high-resolution observations by ground-based and space-borne telescopes have led to the discovery of different modes of magnetic flux emergence in the quiet Sun. These events occur at the spatial scale of granular convection and seem to be very frequent.

Centeno et al. (2007) and Ishikawa et al. (2008) described the appearance of low-lying magnetic loops in granules. The loops are characterised by footpoints located in intergranular lanes and horizontal fields in between, above granular cells. It has been suggested that they emerge from subsurface layers, carried by the upflows of granular convection. After emergence, the distance between the footpoints increases rapidly, and the linear polarisation signals indicating horizontal fields disappear in only a few minutes. About 25% of the loops seem to rise into the chromosphere, producing measurable polarisation signals and brightenings on their way up (Martínez González & Bellot Rubio 2009). The loops have a range

of sizes and are observed to emerge down to the smallest spatial scales of ~ 0.2 achievable nowadays (Martínez González et al. 2012).

Orozco Suárez et al. (2008) reported on the emergence of apparently vertical fields in quiet Sun granules. The typical lifetimes of those events are of order 10-20 minutes. The flux concentrations appear at the center of granules and grow with time, associated with upward motions. They fade gradually until they become undetectable below the noise level. Sometimes, the magnetic flux is observed to disappear with the hosting granule. Throughout the process, no opposite polarities are detected in the field of view. Such events may represent the emergence of fields from subsurface layers. An alternative explanation is that the fields do not come from subsurface layers, but are present in the photosphere all the time until they show up above the noise level by the action of some unknown mechanism.

The events described by Orozco Suárez et al. (2008) may just be individual examples of the unipolar flux appearances observed in the solar internetwork by Hinode and SUNRISE (e.g., Lamb et al. 2010; Gošić et al. 2016; Smitha et al. 2017). Unipolar appearances are very common in the quiet Sun and may contribute a significant fraction of the total internetwork flux, but their origin is a mystery.

An observational characterisation of the magnetic properties (strengths, inclinations) and evolution of both small-scale loops and unipolar flux patches in the quiet Sun still needs to be performed. To that end, multi-line spectropolarimetry at high cadence is required, given the short lifetimes of the features.

Furthermore, the total amount of flux carried to the solar surface by emerging loops and unipolar features has not been evaluated properly so far. Due to the limited sensitivity of the measurements, a significant fraction of the flux emerging on the surface may have gone unnoticed, although estimates from Hinode and SUNRISE suggest that the amount of flux carried by internetwork flux concentrations is much larger than that of active regions (Thornton & Parnell 2011; Gošić et al. 2016; Smitha et al. 2017). To solve this problem, long time sequences of multi-line spectropolarimetric observations covering a full supergranule are needed, at the highest sensitivity possible.

Finally, it is important to clarify the role of emerging internetwork loops and unipolar patches in heating the solar atmosphere. These features are known to interact with pre-existing fields, merging and cancelling with them. Cancellations of opposite-polarity flux patches are likely to result in magnetic reconnection and energy release at chromospheric heights. This process needs to be quantified in the solar internetwork.

OP 1.4.1 Emergence and evolution of magnetic fields in granular convection

See table on page 58.

OP 1.4.2 Properties of magnetic fields emerging in the quiet Sun

See table on page 59.

1.5 Magnetic flux cancellations in the quiet Sun

Flux cancellation occurs when magnetic features of opposite polarity come into close contact and disappear partially or totally as a consequence of the interaction. In that way, magnetic flux is removed from the solar surface.

Cancellation events are very common, particularly among the small flux features of the quiet Sun internetwork. However, we still do not know if they are the result of the submergence of Ω -loops or the rise of U-loops, with or without previous reconnection of unrelated magnetic flux systems. All these mechanisms may be a source of chromospheric/coronal heating and transient events, due to the release of magnetic energy. To distinguish among the different possibilities, simultaneous observations of the photosphere and chromosphere need to be carried out at the highest angular resolution possible. By analyzing the photospheric and chromospheric flow fields at the cancellation sites, and the timing difference between the events occurring in the different layers, it will be possible to draw definite conclusions about the mechanisms responsible for the removal of magnetic flux from the solar surface.

If magnetic reconnection is the main driver behind cancellations, then the goal is to determine how the magnetic topology of the cancelling features changes, the height at which the reconnection takes place, and whether current MHD simulations are able to quantitatively reproduce observations as we go down to extremely small spatial scales. We also need to determine how much these processes contribute to the overall heating of the solar atmosphere.

Recent observations show that current instruments are unable to resolve these processes and reliably detect pre- and post-reconnection loops (Iida et al. 2010; Kubo et al. 2010; Kubo et al. 2014). Furthermore, sampling of the layers where the magnetic energy gets deposited is difficult because of the rapid temperature and density increase which can result in dramatic changes of the formation heights of the observables (Danilovic et al. 2017).

This observational program requires the highest spatial resolution possible combined with high spectropolarimetric sensitivity so that the magnetic field vector can be constrained reliably. Additionally, simultaneous observations in multiple

spectral lines is mandatory in order to retrieve the 3D configuration of these events. This is especially true for determining the location and length of the current sheet and tracing the path of the outflowing material.

OP 1.5.1 Magnetic field topology, dynamics and energy release at flux cancellation sites

See table on page 60.

1.6 Quiet Sun internetwork fields

A substantial (perhaps dominant) fraction of the total magnetic flux of the solar photosphere is stored in internetwork regions, i.e., the interior of supergranular cells (Gošić et al. 2014). Internetwork fields produce very small polarisation signals either because they are intrinsically weak, or because the magnetic filling factor is small, or both. Thus, their observational characterisation represents a significant challenge. Prior to Hinode, our understanding of these fields was mainly based on long integration (1 min) snapshots with spatial resolutions not better than 1'' (Sánchez Almeida & Lites 2000; Khomenko et al. 2003; Lites & Socas-Navarro 2004; López Ariste et al. 2007; Martínez González et al. 2008)

Measurements with the Hinode spectropolarimeter at a resolution of 0.3 revealed that internetwork fields are weak and highly inclined. The field strength distribution peaks at hG values, monotonically decreasing toward stronger fields. The field inclination distribution shows a maximum at 90 degrees, which corresponds to purely horizontal fields. However, the exact shape of the distribution is still under debate, with some authors favoring a quasi-isotropic distribution (Asensio Ramos 2009; Asensio Ramos & Martínez González 2014) and others pointing out that the distribution corresponds to very inclined but not isotropic fields (Orozco Suárez & Bellot Rubio 2012; Danilovic et al. 2016). Furthermore, the magnetic filling factor (the fractional area of the resolution element covered by fields) amounts to 0.2-0.3, even at the resolution of Hinode. Thus, there is still room for subpixel structuring of the field on very small scales. To investigate this possibility, a resolution of 0.1 or better is required.

While circular polarisation signals are seen in almost 100% of the internetwork surface area (Lites et al. 2008), linear polarisation is much more difficult to detect due to the weakness of the field. This represents a serious drawback for the accurate determination of the magnetic properties of the internetwork, in particular the field inclination. Linear signals exist, but they are buried in the noise (Bellot Rubio & Orozco Suárez 2012; Martínez González et al. 2016). To improve the field inclination estimates and resolve the controversy about the shape of the distribution, it is necessary to detect those signals and invert them together with Stokes V. This requires much more sensitive observations than are feasible with any existing instrument, by at least one order of magnitude.

The height variation of internetwork fields is an important aspect of the quiet Sun magnetism. It has been addressed only recently by Danilovic et al. (2016), using Hinode measurements that do not show linear polarisation signals except in a tiny fraction of the field of view. Here too the analysis would benefit from increased sensitivity leading to the detection of linear polarisation all over the field of view. Another way to improve the accuracy of the results is to observe as many spectral lines as possible simultaneously, in order to extract information on the different atmospheric layers in a more direct way and not through extrapolation.

Another aspect that needs to be studied is the temporal evolution of internetwork fields. The Stokes profiles emerging from internetwork flux patches are often asymmetric and show multiple lobes, indicating a very dynamic nature. It is therefore important to resolve their evolution on short time scales. The large photon collecting power of EST will make it possible to obtain the time series required for that.

Observing the temporal evolution of internetwork fields is also important to understand their nature. A fundamental question is whether internetwork fields are the result of a global or a local dynamo. Recent magneto-convection simulations of local dynamo action seem to explain the predominance of inclined fields deduced from Hinode observations (Schüssler & Vögler 2008; Danilovic et al. 2010b; Rempel 2014). However, other mechanisms can also produce inclined fields in the quiet Sun internetwork, such as the emergence of horizontal fields into the solar surface (Steiner et al. 2008). One way to distinguish between scenarios is to compare the short-term evolution of internetwork fields with existing MHD simulations, which will be possible for the first time with EST.

Lastly, it is necessary to study the interaction of internetwork fields with other internetwork and network fields. These processes release energy and can be expected to contribute to chromospheric heating in the quiet Sun. First indications that this is the case have been gathered by Gošić et al. (2018), but the observations lack the sensitivity required to detect the weakest internetwork fields, and therefore a large fraction of the quiet Sun surface remains unexplored.

OP 1.6.1 Physical properties of internetwork magnetic fields

See table on page 60.

OP 1.6.2 Short-term evolution of internetwork fields

See table on page 61.

1.7 Polar magnetic fields

The magnetic fields in the poles of the Sun are believed to be an essential ingredient of the solar dynamo. As the activity cycle proceeds, flux from mid-latitude decaying active regions is transported to the poles by supergranular diffusion and meridional circulation. This flux accumulates in the polar caps and eventually reverses the magnetic polarity of the poles. Among other reasons, the polar caps are important because the fast solar wind emanates from them and because they host a variety of magnetic structures such as faculae (one of the main sources of solar irradiance variations), polar plumes (observed in the EUV), and small-scale X-ray polar jets (Shimojo et al. 2007; Cirtain et al. 2007).

Far from the direct influence of the lower-latitude activity belt, the polar region is also the obvious place to investigate whether or not small-scale photospheric flux concentrations are the result of local dynamo action. Observations with Hinode suggest that the only difference between the magnetism of the quiet Sun at the disk center and the polar region is the existence of strong, relatively large unipolar flux patches, with the small-scale fields showing nearly the same distribution of parameters (Tsuneta et al. 2008; Ito et al. 2010). This seems to favor a turbulent dynamo. However, most of the solar surface in those regions remains unexplored as a consequence of the strong projection effects, which reduce both the amplitude of the polarisation signals and the effective spatial resolution of the observations. EST, with its enormous light collecting power, will give access to very weak polarisation signals that current instruments cannot detect. Polar field studies will also benefit from the high spatial resolution capabilities of EST, which permit a significant decrease in the solar surface area sampled by the resolution element.

Of particular interest are the properties of polar faculae in the lower atmospheric layers and how they appear and disappear. The short-term evolution of polar faculae is essentially unknown. However, the results of Kaithakkal et al. (2015) based on Hinode measurements suggest that these features are formed by advection of flux by supergranular flows rather than by flux emergence. Eventually polar faculae fade and disappear on the spot, with or without previous fragmentation. This mode of disappearance is not well understood and requires further investigation. One possibility is that the magnetic patches fragment in elements too small to be detected. The fragments may later cancel out with opposite polarity elements that reach the polar region around the maximum of the solar cycle. EST will confirm or refute this scenario.

Other scientific questions that need to be addressed include how the flux from the old cycle is replaced in the polar caps (either by reconnection, field submergence, or field expulsion), whether or not the properties of strong flux concentrations are the same in the polar regions and the equatorial belt, and how the magnetic field drives the various phenomena observed at high latitudes, in particular X-ray jets. These problems should be investigated at their relevant spatial scales (100 km).

Further progress in our understanding of polar fields will be allowed by magnetic stereoscopy from coordinated observations of EST and Solar Orbiter. Magnetic stereoscopy is the only reliable way to solve the 180° azimuth ambiguity of Zeeman measurements, which is particularly problematic near the poles as it renders the orientation and even the polarity of the field unknown (e.g., Ito et al. 2010). For meaningful comparisons of the same structure from the two vantage points, the high spatial resolution of EST is essential, as other ground-based telescopes will not be able to isolate the small surface area contained in the Solar Orbiter resolution element. Thus, coordination of EST and Solar Orbiter will likely result in new ground-breaking science.

OP 1.7.1 Structure of polar faculae

See table on page 62.

OP 1.7.2 Properties, distribution and evolution of polar magnetic fields

See table on page 63.

1.8 Network dynamics

The magnetic network is a well-known large-scale feature of the solar photosphere that outlines the borders of supergranular cells (e.g., Orozco Suárez et al. 2012). It contains a significant fraction of the total magnetic flux of the quiet Sun, and is relatively stable on supergranular time scales (~ days). The magnetic network also leaves clear imprints on the chromosphere, where it can be easily identified in Ca II H filtergrams as a bright network structure indicating excess temperatures but also shock fronts and waves caused by the dynamics of the underlying photospheric network (e.g., Hasan & van Ballegoijen 2008).

Open questions concerning the magnetic network include its obvious unipolar character on large scales (compared to the smaller bipolar internetwork fields populating the interior of supergranules), its flux balance (Schrijver et al. 1997), and its evolution over the solar cycle (Thibault et al. 2014).

Moreover, strong network elements have lifetimes of several minutes to hours only (e.g., de Wijn et al. 2005), while the network itself persists for days. Thus, while appearing stable, the network is fully dynamic: strong magnetic elements are continually formed, reshuffled, and dissolved there. During those processes, magnetic flux can be lost due to submergence,

reconnection, cancelation of opposite polarity elements, and various other mechanisms (Schrijver et al. 1997). As a result, the network flux needs to be replenished by new flux elements appearing on the solar surface. The source of such elements is still unknown. Recent papers suggest that quite a large fraction can be provided by internetwork magnetic features, which are swept to the network via supergranular flows (see Gošić et al. 2014). A contradicting older hypothesis would state that most of the network flux is actually coming from decaying ephemeral regions.

For a more thorough understanding of the flux balance of the magnetic network and its dynamics, it is necessary to lower the detection threshold of internetwork magnetic features in terms of both field strength and size, to capture the more abundant lower end of the distributions. Only then the contribution of internetwork fields to the maintenance of the network will be fully understood. On the other hand, a better characterisation of the interaction between internetwork and network fields is necessary to understand how internetwork fields get converted into network fields and what this means for the dynamics of the upper atmosphere.

OP 1.8.1 Network evolution and dynamics

See table on page 64.

2 Wave coupling throughout solar atmosphere

Authors: Robertus Erdélyi, Elena Khomenko, Mihalis Mathioudakis

The Sun generates a wealth of waves, mostly acoustic in their generated character, by turbulent convection beneath the photosphere (Lighthill 1952; Stein 1967; Goldreich & Kumar 1990; Goldreich et al. 1994; Bi & Li 1998; Musielak et al. 1994; Nordlund & Stein 2001; Stein & Nordlund 2001). The energy of these waves in the photosphere is believed to be sufficiently large to account for the heating of the upper layers (Ulmschneider 1989; Musielak et al. 1994; Fawzy et al. 2002; Fossum & Carlsson 2006; Banerjee et al. 2007; Erdélyi & Ballai 2007; Taroyan & Erdélyi 2009; Jess et al. 2009; Arregui 2015). For this to manifest, however, during their propagation from the (sub)photosphere to higher, magnetically dominated regions of the solar atmosphere, the waves may need to transform their acoustically dominated sub-photospheric nature to waves that account for the presence of the atmospheric magnetic fields. Once they reach the mid to upper atmosphere, they often dissipate there, at least in part, and release their non-thermal energy in the form of heat. Studying the wave coupling through the solar atmosphere consists of studying how the waves propagate and evolve, interact, transform and dissipate during their passage through the photosphere, chromosphere, transition region and corona of the Sun (Erdélyi 2006). Such research includes studying the waves in the presence of atmospheric stratification, the wave interactions with the often dynamic magnetic structures present everywhere over the surface of the Sun. Wave interaction with these structured and stratified magnetic fields results in the generation of a large variety of wave modes. At the same time structuring, whether magnetic or thermodynamic, also serves as guide for the progression of wave energy and momentum, and connect all the layers up to the corona and interplanetary space. The common line for all observational studies of the wave coupling is the need of simultaneous data at several layers of the solar atmosphere, which EST will routinely provide in unprecedented spatial, temporal and spectral resolution.

Traditionally, many wave studies focus on a single layer of the solar atmosphere (e.g., Bellot Rubio et al. 2000a; De Pontieu et al. 2003; Tomczyk et al. 2007; Erdélyi & Taroyan 2008; Vecchio et al. 2009; Reardon et al. 2009; Fujimura & Tsuneta 2009; Cauzzi et al. 2009; Martínez González et al. 2011; Rajaguru et al. 2012). However, there is increasing evidence that observing a single layer does not give us sufficient information to understand plasma heating or to carry out plasma diagnostics by means of solar magneto-seismology (SMS) in the required details. Nowadays, the state-of-the-art in wave studies consists of using multi-instrument multi-layer data combined simultaneously to follow the wave propagation over several layers. Examples of recent observations of such kind can be found elsewhere (De Pontieu et al. 2005; Taroyan et al. 2005; Jefferies et al. 2006; McIntosh & Jefferies 2006; Centeno et al. 2006; De Pontieu et al. 2007c; Felipe et al. 2010b; Kontogiannis et al. 2010; Reznikova et al. 2012; Jess et al. 2013; Yuan et al. 2014; Freij et al. 2014; Krishna Prasad et al. 2017). For recent review of multi-wavelength studies of waves in the chromosphere see Jess et al. (2015).

Observational studies are frequently accompanied by realistic or semi-realistic simulations which have been specifically designed to study wave dynamics (Erdélyi et al. 2007a; Malins & Erdélyi 2007; Nutto et al. 2012; Kuridze et al. 2009; Vigeesh et al. 2009; Felipe et al. 2010b; Kato et al. 2011; Fedun et al. 2011; Vigeesh et al. 2012; Mumford et al. 2015; Mumford & Erdélyi 2015; Shelyag et al. 2016; Rijs et al. 2016). Some of the simulations include heights up to the corona (Pascoe et al. 2011; Heggland et al. 2011; Santamaria et al. 2016; Krishna Prasad et al. 2015). Yet another promising direction focuses on data-driven simulations (De Pontieu et al. 2007a; Carlsson & Stein 1997; Felipe et al. 2011; Krishna Prasad et al. 2015; De Pontieu et al. 2017; Snow et al. 2018).

The wave studies can be sub-divided into two large categories: (1) when the waves are studied in order to measure the energy being carried and dissipated to heat the atmosphere, (2) when the observed wave properties are used to infer the properties of the waveguides (i.e. solar magneto-seismology, see reviews in Gizon & Birch 2005; Erdélyi et al. 2007b; Cally 2007). In each of these two cases, a particular manifestation of the wave phenomena depends on the magnetic structure they interact with: e.g., sunspots, pores, a range of magnetic elements in the quiet Sun, prominences or filaments.

Regarding sunspot waves, our current understanding is summarized in Moradi et al. (2010); Khomenko & Collados (2015). The global picture of the wave propagation in these structures is addressed in details successfully and it is understood that waves observed in these structures are different manifestations of the same phenomena. While there is still no agreement whether sunspot waves are internally or externally excited, there is an agreement that the umbral chromosphere contains slow magneto-acoustic waves, propagating along the field lines and forming shocks (Centeno et al. 2006). In the penumbra, a mixture of fast and slow waves are observed, the periods of the slow waves following the rule of the ramp effect, affecting the cutoff frequency for inclined fields (Jess et al. 2013; Yuan et al. 2014). Clear detections of Alfvén waves are still elusive (Tomczyk et al. 2007; Erdélyi & Fedun 2007; Van Doorselaere et al. 2008; Jess et al. 2009). Recent studies using DST, ROSA and HMI/AIA data by Krishna Prasad et al. (2015) demonstrate how the particular wave fronts can be traced propagating from the photospheric source to the corona. These kind of observations are, however, sparse. It remains to determine how sunspot waves interact with fine structure of the magnetic field in the umbra (umbral dots) and penumbra, how they propagate through the transition region, and their energy content and dissipation mechanisms in the chromosphere and corona. For that, one would need high resolution and high polarimetric precision data to finally measure elusive oscillations of the magnetic field. The question is even more challenging to be

answered for smaller magnetic structures, e.g. from pores to tiny inter-granular magnetic flux concentrations.

The coupling between the different solar layers by means of waves, propagating in quiet Sun structures has received a considerable attention. High-frequency acoustic halos surrounding active regions have been recently attempted to be explained in terms of the wave mode conversion physics (Khomenko & Collados 2009; Rijs et al. 2016). Waves with photospheric periodicities are systematically observed overall in the mid to upper atmosphere, even in the solar corona. Nevertheless, it is still an open question how the p -mode energy would propagate efficiently up there because of a set of obstacles that waves encounter on their way up, such as: cut-off layer, wave speed gradients, transition region, multi-component plasma, etc. (De Pontieu et al. 2004, 2005; McIntosh & Jefferies 2006; Centeno et al. 2009). Vortex motions (magnetic tornadoes), observed connecting all layers are another potential source of coupling and energy transport to the upper atmosphere (Wedemeyer-Böhm et al. 2012; Shelyag et al. 2013; Iijima & Yokoyama 2017; Yang et al. 2018; Liu et al. 2018).

Studying the processes related to coupling observationally requires resolving and measuring the magnetic field strength and topology at smallest scales. These kind of acquisitions are already at the limit of the current instrumentation, especially when it comes to measuring the magnetic field in the chromosphere and higher up. Photospheric flows may alter the magnetic field topology not only in the photosphere but also in higher layers of the solar atmosphere. Therefore, the chromospheric magnetic fields may become non-potential and electric currents are generated. The currents can then be dissipated through various dissipative processes, e.g., ambipolar diffusion neutral-related mechanisms, yielding important highly localized energy deposits and allowing the dissipation of even magnetic non-compressible waves in an efficient manner. Dissipation of the Alfvén waves seems to be extremely difficult to test observationally with the existing instrumentation, because these waves have velocity and magnetic field perturbations perpendicular to the magnetic field lines (i.e. frequently perpendicular to the line of sight) within constant magnetic surfaces. Each such surface may have its own frequency-dependent transversal magnetic oscillation. It is a real observational challenge to determine unambiguously these constant magnetic surfaces (Erdélyi & Fedun 2007; Jess et al. 2009). The wave physics must also be explored in the range of very high frequencies. It is not yet established how the high-frequency and small-wavelength waves contribute to the energetic connectivity between the various layers of the solar atmosphere (Fossum & Carlsson 2006; Fleck et al. 2010). Simulations and observations have been controversial in this respect, mainly because observational detections are extremely hard and challenging to be made.

Wave diagnostics based on polarimetry or filter imaging have their advantages and shortcomings, and the information is often masked by radiative transfer effects and limited spatial and temporal resolution. The combination of different techniques with the high-resolution instrumental capabilities of EST will provide the necessary tool to clarify the connectivity by means of waves in the Sun.

2.1 MHD waves in localized quiet Sun structures

Energy propagation is intimately linked to the magnetic field strength and topology. Magnetic fields in the quiet Sun are swept to the boundaries of intergranular lanes, and to the boundaries of larger network cells, forming kG magnetic field concentrations. These flux tube-like structures expand with height due to the pressure fall-off, producing a carpet of strongly inclined fields at chromospheric heights. The plasma- β changes with height in the solar atmosphere, from pressure dominated regions in the photosphere (including sunspots) to the magnetically dominated regions in the chromosphere and above. Measuring the strength and topology of the fields is crucial for studies of the wave propagation. For recent review on measurements of photospheric and chromospheric magnetic fields see Lagg et al. (2017).

MHD wave propagation in MBPs Magnetic Bright Points (MBPs) are small-scale kG field magnetic flux concentrations found in the dark intergranular lanes (Keys et al. 2013, 2014; Liu et al. 2018). Formed by the process of convective collapse, these highly dynamic features are ubiquitous across the solar surface. They exhibit transverse motions of a few km sec⁻¹ caused by granular buffeting and can lead to the excitation of MHD waves. Due to their strong localised field, MBPs can act as wave guides and transfer energy into the upper parts of the solar atmosphere (de Wijn et al. 2009; Mathioudakis et al. 2013). Their ubiquitous nature means that they can be a significant contributor to the non-thermal energy budget of the solar atmosphere. Jess et al. (2012) and Stangalini et al. (2013) showed that MBPs display upward propagating waves and that longitudinal MBP oscillations in the photosphere can drive H-alpha spicule oscillations in the chromosphere. EST observations will allow us to observe variations in the Stokes profiles in the presence of MHD waves at the location of MBPs in the presence of MHD waves. We will be able to *determine the phase relations* between the Stokes profiles, Doppler shifts and intensity perturbations of MHD waves at these locations. Therby we can determine uniquely the energy associated with these wave modes and their contribution to localised heating.

OP 2.1.1 Sausage and kink oscillations in MBPs

A range of magneto-hydrodynamic waves have now been observed in pores and sunspots (including MHD sausage and kink modes). Whether similar waves could be present in smaller concentrations of strong magnetic field, such as MBPs

is, however, currently unknown. The abundant presence of such waves, if they are observed to propagate through the solar atmosphere, could contribute significant amounts of energy into the solar corona where it can be dissipated. Using extremely high-resolution imaging data from EST, it will be possible to identify whether area perturbations (consistent with the presence of sausage mode oscillations) can be identified in MBPs.

See table on page 65.

OP 2.1.2 Magneto-acoustic waves in spicules on disk

RBEs and RREs are the disk counterparts of the H-alpha spicules (Kuridze et al. 2012, 2013, 2015a). It is known that spicules are excellent waveguides for a wide range of MHD waves whether slow or fast, whether sausage or kink. It is even claimed that torsional Alfvén waves are detected in spicules at limb. However, limb observations are hard, it requires a detection of higher photon count than at disk. Therefore, the hunt for the various MHD waves present in concentrations of strong magnetic field, such as RBEs/RREs is, more feasible. Similar to MBPs, pores or sunspots, the propagation of such waves, could carry significant amounts of energy flux contributing into the non-thermal energy of the upper solar atmosphere where it can be dissipated. Using the combination of extremely high-resolution spectro-polarimetric imaging data from EST, it will be possible to identify whether these perturbations (consistent with the presence of waves in spicules) can be identified and their roles assessed for energising the local plasma in RBEs/RREs.

See table on page 65.

OP 2.1.3 Magneto-acoustic waves in spicules at limb

A key challenge with spicule observations off-limb require to disentangle the superposition of structures along the line of sight. One may employ Fe I 630.2, H α , Ca II and He I 1083.0 to measure magnetic field across the lower solar atmosphere. Point the telescope at the limb or close to limb harbouring the spicules. Sit and stare may be used for 2D spectroscopy. Full-Stokes polarimetry of Fe I 630.2, Ca II 854.2 and He I 1083.0 is suggested to measure the magnetic field in the chromosphere, spectroscopy in H α to track the spicules to larger heights, spectroscopy in Ca II K to have the highest spatial resolution if possible. Context imaging using FP or BBF is desirable. Coordination with space missions is highly desirable to trace spicule evolution in the (E)UV to greater (e.g. transition region or low coronal) heights and temperatures. Science questions can be addressed such as: How do the fine structure of spicules manifest and evolve; What type of wave motions do spicules exhibit (MA longitudinal, kink, sausage, torsional AW, etc.); What is the energy and momentum flux and power spectrum of the waves? What is the magnetic field structure in spicules?

See table on page 66.

OP 2.1.4 Torsional Alfvén wave (TAW) propagation in spicules

Here, the main focus is to find conclusive evidence for the most elusive waves in solar magnetic structures, the Alfvén wave. Given the likely axially symmetric structure of spicules, the most expected form of these magnetic waves would be the torsional Alfvén waves (TAWs).

See table on page 67.

2.2 Magnetic twist and torsion

Given their size, pores (or sunspots) are ideal magnetic structures for studying MHD waves in spite of their complexity. In 1942, Hannes Alfvén theoretically predicted the existence of a very special type of MHD wave in which the sole restoring force is provided by the tension of the magnetic field lines. In 1970 he won the Nobel prize in Physics for this contribution to the field and for the numerous applications in plasma physics that derive from it. Indeed, Alfvén waves are expected to play a key role in heating and energy dissipation within both laboratory and space plasmas including the solar atmosphere. Despite the many laboratory proofs, and several claims in the astrophysical literature focusing on the upper layers of the Sun's atmosphere, Alfvén waves have not been *directly* observed in the lower solar atmosphere, where torsional oscillations of magnetic flux iso-contours can now be measured directly. Thus, Alfvén waves represent the most elusive and physically intriguing class of MHD waves.

The aim is to find not only the $m = 0$ torsional Alfvén wave but also its higher harmonic, the $m = \pm 1$. Discovering the existence of the latter could be carried out by identifying (a)symmetric torsional motions e.g. in the lobes of a pore (or sunspot) separated by e.g. its light bridge. Directly proving the existence of these torsional Alfvén modes would be a significant step to observationally prove the completeness of eigenmodes of the MHD Hilbert operator.

OP 2.2.1 Magnetic torsion and torsional oscillations of pores or sunspots

See table on page 67.

OP 2.2.2 Observation of magnetic vortices and tornadoes

2.2.2.1: Swirl penetration into the chromosphere

Intensity (density) swirls have been widely observed from the photosphere up to the corona in both quiet-Sun region and coronal holes (e.g., Wedemeyer-Böhm et al. 2012). It is then natural to infer the presence of *magnetic* swirls and tornadoes in the solar atmosphere, having considered the magnetic nature of the solar plasma. However, high-spatial and temporal observations of magnetic swirls and tornadoes are still urgently needed to answer the following scientific questions: 1) Are intensity and magnetic swirls coupled with each other in space and time? 2) How are they coupled? 3) What is the role of MHD waves (especially Alfvén waves) in the propagation of intensity and magnetic swirls from the photosphere into the chromosphere? Recently, Liu et al. (2018) developed an automated swirl detection algorithm (ASDA) aiming to automatically detect vortex motions at different layers of the solar atmosphere. This algorithm will also be adapted to magnetic field observations to perform automated detection of magnetic swirls.

2.2.2.2: Swirls in very high-cadence observations

Having considered the very short lifetime of previously detected swirls ($> 70\%$ less than 6 sec), it is further proposed to investigate the coupling of intensity and magnetic swirls with very high-cadence observations. The penetration of swirls from the photosphere into the chromosphere and whether tornadoes are formed from swirls will also be further studied into high details using ultra-high cadence observations.

2.2.2.3: Observation of swirls under 2-fluid condition

At a time scale less than 1 sec, the single-fluid MHD approximation will be broken down and the two-fluid condition will apply. We propose to further investigate photospheric intensity swirls under the two-fluid condition. Extremely high cadence (0.1 sec) will be needed.

See table on page 68.

OP 2.2.3 Formation of magnetic swirls in intergranular lanes

It has been found by their statistical study (Liu et al. 2018), that, more than 70% of swirls are located in intergranular lanes. The small-scale photospheric magnetic swirls are widely hypothesised to form as a natural consequence to granular flows (e.g., Wang et al. 1995; Attie et al. 2009) in the quiet Sun and could play a key role in the supply of energy to the upper solar atmosphere either, for example, through the build up of magnetic energy in twisted field lines in the corona or through the channeling of MHD waves (e.g., Velli & Liewer 1999; Shelyag et al. 2011). However, the exact mechanism of how these magnetic swirls are generated in the intergranular lanes remain unclear. High-resolution observations of the velocity, intensity and magnetic field inhomogeneity in the intergranular lanes are needed to perform both case and statistical studies on the role that MHD instabilities resulting from granular flows may play in generating magnetic and intensity swirls, as suggested by Liu et al. (2019).

See table on page 69.

OP 2.2.4 Vortex flows in the lower solar atmosphere

Due to their small-scale in the solar photosphere, not much insight into vortex flows, especially on small scales such as isolated flux tubes (about 200 km diameter), has been obtained in this layer from direct observations so far (Bonet et al. 2008; Liu et al. 2018). While the observational evidence is quite scarce, in simulations, however, vortex flows seem to be abundant both in hydrodynamical simulations (see, e.g., Calvo et al. 2016) as well as in MHD simulations (Shelyag et al. 2011). Thus, vortex features and the associated swirling vortex motions may play an important role in the creation and amplification of magnetic fields as well as in the transportation of energy into the higher layers via the generation of MHD waves (Fedun et al. 2011). Moreover, it seems as if magnetic fields and vortex motions are to a great extent coupled phenomena.

For tracking down the vortex motions directly in observations extremely highly resolved images with a sufficient high cadence are necessary, as can be seen, e.g., in Liu et al. (2018), who studied MBPs, proxies for isolated strong small-scale magnetic flux tubes, and have shown a case of an apparently turning bright point. However, due to the still limited observational capabilities (1m diameter telescope; ~ 70 km diffraction limit; feature size ~ 200 km) they were not able to reinforce this single case study measurement by statistical data. Clearly, a 4 m class telescope would be necessary to directly track the rotation of a feature via its morphologic changes.

On the other hand, indirect methods exist and are applied for instance in the case of Alfvén wave detection (Jess et al. 2009) where rotational motions of small-scale features can be detected as spectro-polarimetric variations in line widths. To conclude, for successful revelations of the true nature of vortex flows and their interaction with the solar magnetic field a 4 m class solar telescope with imaging and spectro-polarimetric capabilities in the photosphere and chromosphere is needed.

See table on page 69.

2.3 Wave propagation in active regions

Sunspots are structures with strong magnetic fields in active regions which are thought to be “portals” through which energy of e.g. acoustic p -modes can penetrate into the upper atmosphere via a range of wave coupling processes.

One of such processes is mode conversion. Theoretical mode conversion models suggest it is a two-step process. In the first place, acoustic p -modes convert to fast and slow magneto-acoustic waves at heights where the plasma- β is equal to unity. The slow mode (acoustic in nature) would then continue along the magnetic field lines to the upper chromosphere. The fast (more magnetic in nature) mode would refract and reflect due to the gradients in the Alfvén speed. Around heights of this reflection the Alfvén mode itself would be generated through the second mode transformation. All these processes are strongly dependent on the wave parameters (e.g. frequency) and magnetic field inclination and azimuth. This theoretical process could provide a way of transferring the energy to the chromosphere efficiently via the generation of Alfvén waves at heights close to the transition region, amplifying their possibility to escape this barrier without reflection. Observational confirmation of this process is still missing.

Among other wave-coupling theoretical mechanisms acting in sunspots (and pores) are the resonant coupling of p -modes and the formation of photosphere-chromosphere cavity.

OP 2.3.1 Excitation mechanisms of sunspot waves

Several theories exist regarding the mechanisms of excitation of sunspot waves, and regarding the relation between wave phenomena observed in different layers. Using multi-instrument and multi-layer sunspot observations, Zhao et al. (2016) has recently shown that helioseismic p -mode waves are able to channel up from the photosphere through the chromosphere and transition region into the corona, and that the magneto-acoustic waves observed in different atmospheric layers are the same waves originating from the photosphere but exhibiting differently under distinct physical conditions. Do this observation suggest that waves are locally excited in sunspots by the weak convection? How do these waves continue traveling up to the corona, and excite oscillations in coronal loops?

Long-time series of data are needed at moderate resolution, bi-dimensional Doppler velocity data at several height. Magnetic field information is required to determine the wave propagation speeds and directions. The observed area should be large to cover the sunspot and the nearby quiet Sun.

See table on page 70.

OP 2.3.2 Alfvén waves in sunspots

This observing program aims at finding observational evidence for Alfvén waves in sunspots. It aims answering the following questions: at what height are the Alfvén waves generated, depending on magnetic field strength and topology; how large is the energy carried by Alfvén waves up to the transition region and corona; are the Alfvén waves able to penetrate into the corona without significant reflections; what is the power spectrum of the velocity component transverse to the magnetic field.

This observing campaign would require low-noise polarimetric data to measure oscillations of the magnetic field and velocity, magnetic field topology, and their mutual relation. Data should be taken at different locations across the disc.

See table on page 71.

OP 2.3.3 Magnetic field oscillations in sunspots

Detecting oscillations of magnetic field has been illusive (Bellot Rubio et al. 2000a; Fujimura & Tsuneta 2009). High-precision polarimetry should make such detection possible. As discussed in the literature, detected oscillations in the magnetic field are not always due to “real” oscillations in the magnetic field vector, but can be rather due to opacity effects, shifting the formation heights of spectral line up and down (Khomenko et al. 2003). In chromospheric lines, such as Ca II 854, the effects of shifts of line formation heights on the passage of shocks are able to produce detection of “false” magnetic field oscillations with amplitudes as large as 50 – 100 G, while intrinsic oscillations do not exceed few G in amplitude (Felipe et al. 2014). In order to separate real oscillations from these effects, one has to have information about magnetic field gradient, and also compare data from different spectral lines, providing complimentary information.

Magnetic field oscillations, together with oscillations in other quantities, and the phase information of propagating waves at several layers would be required to identify the wave modes observed in sunspots, depending on the region (umbra, penumbra) and height. Several lines with different sensitivity to temperature need to be observed simultaneously. The resolution can be moderate, with the primarily goal to have low-noise signal.

See table on page 71.

OP 2.3.4 Sunspot penumbral waves in the photosphere and above

Running penumbral waves (RPW) are disturbances, detected mainly in chromospheric lines, that appear to be emitted from the umbra and expand concentrically with constant velocity around 10 – 15 km/s (Zirin & Stein 1972; Giovanelli 1972). Reznikova et al. (2012); Jess et al. (2013); Kobanov et al. (2013) have investigated the role of the magnetic field topology in the propagation characteristics of umbral and running penumbral waves at chromospheric, transition region and coronal layers. They found an increase of the oscillatory period of brightness oscillations as a function of distance from the umbral center. Reznikova et al. (2012) and Jess et al. (2013) concluded that running penumbral wave phenomena are the chromospheric signature of upwardly propagating magneto-acoustic waves generated in the photosphere. Recently, Löhner-Böttcher & Bello González (2015) have found signatures of RPWs that are found at photospheric layers, with some evidence for being there a relation between field inclination and observed wave propagation velocities. Possible connections between RPW and umbral flushes has to be explored further, following the waves through several layers of the atmosphere simultaneously. It is not known well what happens to the penumbral waves at layers above the chromosphere, how they penetrate through the transition region to the corona and if the inclination facilitate this process.

The purpose of this particular observing campaign is to track the wavefronts of RPWs from photospheric heights where they presumably are generated, across the penumbra in the chromosphere and higher up. It also aims at finding observational evidences for simultaneous up- and downward wave propagation in the penumbral chromosphere, suggestive for possible reflections from the transition region. The campaign would reveal how RPWs are generated and if they can efficiently penetrate to the corona, depending on their frequency and field inclination.

Fine structure of the penumbra and its relation to waves The photospheric penumbra is very inhomogeneous. The question to be answered here is how does this inhomogeneity affect wave propagation? Why do running penumbral waves apparently disappear at the boundary of the visible penumbra? The observational setup of this campaign is the same as OP2.3.4 above.

MHD wave propagation in pores Pores are intermediate-scale kG field magnetic flux concentrations found around sunspots and around/in active regions. They exhibit a range of perturbations, from sausage, transverse kink or Alfvén modes. Due to their strong and compact magnetic field, pores can act as wave guides and transfer energy into the upper parts of the solar atmosphere (de Wijn et al. 2009; Mathioudakis et al. 2013; Freij et al. 2014, 2016). In spite of their less ubiquity when compared to e.g. MBPs, they may still have an important contributor to the non-thermal energy budget of the solar atmosphere. EST observations will allow us to track variations in their area, intensity and Stokes profiles pointing towards the presence of MHD waves in pores. Phase relations between the Stokes profiles, Doppler shifts and intensity perturbations of MHD waves at these locations will enable the unique identification of wave modes in pores, therefore, allowing us to determine the associated energy flux with these wave modes with their contribution to localised heating.

See table on page 72.

OP 2.3.5 Sausage and kink oscillations in pores

Here, the most important aspect is to have a combined intensity (area) and LoS component magnetic field observations at high cadence.

See table on page 72.

OP 2.3.6 (Torsional) Alfvén waves in pores

Here, the aim is to find observational evidence for (torsional) Alfvén waves in pores, i.e., at what height are the (T)AWs generated, depending on magnetic field strength and topology; how large is the energy carried out by (T)AWs up to the transition region and corona; are (T)AWs able to penetrate into the corona without significant reflections; what is the power spectrum of the velocity component transverse to the magnetic field.

This observing campaign would require low-noise polarimetric data to measure oscillations of the magnetic field and velocity, magnetic field topology, and their mutual relation. Data should be taken at different locations across the disc.

See table on page 73.

2.4 Wave propagation in the quiet Sun

Relations between waves and magnetic topology have been hard to determine for waves in the quiet Sun, due to more complex and weaker fields. Unlike in sunspots, magnetic field in the quiet regions is generally weaker, less organized, and, as a consequence, it is more challenging to measure the field components.

OP 2.4.1 High-frequency wave propagation and dissipation from the photosphere to the chromosphere

The granular buffeting and vortical motions detected at the top of the solar convection zone can lead to the generation of MHD waves either directly or as a result of reconnection in the magnetic canopy region. The photospheric driver of MHD waves is typically of too low a frequency for significant dissipation on its transit of the chromosphere. However, if sufficient wave reflection occurs at the transition region it may be possible for a turbulent cascade to be set up in the chromosphere. Theoretical predictions suggest that dissipation mechanisms, such as resonant absorption, are very efficient at high frequencies (Verth et al. 2010). Observations obtained with the ROSA high cadence imager, suggest that there is a frequency-dependent velocity power of transverse waves. Morton et al. (2014) show that the velocity power of transverse waves in the chromosphere as a function of frequency has increased power at higher frequencies. In certain small-scale solar structures higher frequency waves transport more energy through the chromosphere.

Knowledge of the energy budget in the high-frequency domain is therefore crucial for atmospheric heating. EST observations will allow the simultaneous study of several photospheric and chromospheric spectral lines. This will provide the spectropolarimetric precision required to study the photospheric motions and their interaction with the magnetic field, and the process of energy transfer via MHD wave excitation and propagation.

See table on page 74.

OP 2.4.2 Time-dependent behaviour of chromospheric jets

The chromosphere is constantly perturbed by short lived (1 -10 minutes), jet-like extrusions that reach heights of 2000 - 10000 km above the photosphere. These thin jets are formed in the vicinity of photospheric magnetic field concentrations. They can be a consequence of shock waves channeled along the magnetic field lines. It is not clear how different periodicities are created, and what is the role of the high-frequency waves likely excited in the photosphere. For this reason, this observing campaign will measure the dominant periods of waves depending on the strength and topology of the fields, concentrating on the longitudinally propagating waves. It will then be determined whether the shock behaviour is always present in such jets. Local energy dissipation and transfer to the thermal energy should be possible to trace via localized small-scale brightening in chromospheric lines.

See table on page 75.

OP 2.4.3 Network and plage oscillations

There seems to be still no agreement on the explanation of the presence of ubiquitous long-period waves in the chromosphere above network bright points and plages. Different observations give evidences both for inclined and vertical propagation (both upward and downward), at least for the network elements, while theoretical models seem to point toward the inclined propagation as a dominant mechanism. Inclined propagation receives more observational support especially for the strong field plage regions where it is expected that the slow acoustic waves propagate field-aligned in the low- β regime. This observing campaign addresses the propagation in weaker structures, accompanied by observations of more network and inter-network elements. The requirements are similar to the program “Excitation mechanisms of sunspot waves”.

See table on page 75.

Alfvén wave detection depending on topology The aims and the instrumental setup of this campaign are similar to “Alfvén waves in sunspots or pores”, except that the target areas are quiet areas of the Sun: network, internetwork and plage.

Transverse waves observed at the limb Theoretical models for acoustic halos and shadows indicate that both may have a similar origin, tracing mode transformation process, and fast mode refraction and reflection at the inclined field of the magnetic canopy. It remains to clarify the relation between the acoustic halos and glories, present on the same locations. Some theoretical models for halos suggest that the power enhancement in horizontal velocity component should be significantly stronger than in the vertical one. This campaign will study halos and shadows at locations off the disc centre to shed more light on their nature. Models of transformation to Alfvén waves suggest that they will be reinforced at the periphery of active regions. Conforming this by observations is an aim of this campaign.

Acoustic shadows and glories across the disc center The observed distributions of photospheric and chromospheric power of long- and short-period waves seems to be distinct over network and plage/facular regions. In the network, long-period 5 min waves seem to be transmitted in the close proximity of magnetic elements, while short-period 3 min waves are “shadowed” due to the interaction of magneto-acoustic waves with the more horizontal fields of the canopies. In the stronger-field plage regions, short-period (3 min, $\nu = 5 - 7$ mHz) halos dominate both in the photosphere and in the chromosphere, and the propagation of long-period waves is enhanced for inclined fields, but vertical propagation has

also been reported. The reason for this different behaviour may be the variation of the height of the magnetic canopy $\beta = 1$ layer, being lower for stronger plage fields, thus modifying the spectrum of waves reaching the heights sampled by photospheric and chromospheric observations.

This campaign will perform a high-resolution study, comparing regions with different magnetic fluxes, both at the disc centre and closer to the limb to get information about horizontal velocities. Simultaneous measurements of the magnetic field vector are also required.

3 Chromospheric dynamics, magnetism, and heating

Authors: Jorrit Leenaarts, Mats Carlsson, Peter Gömöry, Christoph Kuckein, Ada Ortiz Carbonell

The chromosphere is the interface between the photosphere and the corona. In the chromosphere the dynamics change from gas-pressure driving to magnetic-force driving, radiation transport changes from optically thick to optically thin, the gas state changes from neutral to ionised and from local thermodynamic equilibrium (LTE) to non-equilibrium conditions, and the MHD approximation is not sufficient to fully describe its physics. The magnetic field is the key quantity in the physics of the chromosphere. Understanding its physics thus requires determination of the magnetic field at all locations in the chromosphere. Major open questions are: How large is the non-thermal energy input into the chromosphere and how and where is it dissipated? How does the chromosphere regulate the mass and energy flow into the corona? How does magnetic flux rise through the chromosphere? It is also an astrophysical laboratory where MHD and plasma physics can be studied at scales inaccessible on Earth, and serves as a test case for understanding chromospheres of other stars.

The chromosphere is relatively difficult to study compared to the photosphere: Evolution timescales are shorter, changes are observed down to a timescale of one second. Chromospheric lines are often broad and deep, and thus have a relatively small photon flux. The magnetic field in the chromosphere is volume filling, and the resulting flux densities and thus polarisation signals are weaker than in the photosphere. The fundamental spatial scales are expected to be as small or even smaller than the resolution of even a 4-meter telescope. Because the chromosphere is highly stratified, one cannot sample the entire chromosphere with a single spectral line. Simultaneous and co-spatial observations in multiple spectral lines are required to sample all heights.

Observations of the chromosphere will thus always require a trade-off. High signal-to-noise is desired in order to measure weak polarisation signals, but simultaneously the highest resolution and short integration times are needed to resolve the relevant spatial and temporal scales. Signal-to-noise is therefore crucial for observing the chromosphere. The overall requirements on EST for studying the chromosphere are thus:

- High photon efficiency of the telescope-instrument-camera system. Experience with the Swedish 1-m Solar Telescope indicates that an image derotation system is not essential. Such a system should not be added if it would add additional mirrors or lenses.
- Polarimetric accuracy of at least 10^{-4} .
- A spatial PSF that minimizes straylight. Straylight decreases the contrast at small spatial frequencies, and thus requires higher signal-to-noise to reach diffraction-limited resolution.
- A light distribution system that allows simultaneous observations in at least three and preferably four or five spectral lines.
- The required spectral resolution of each instrument, in particular for FPs, should be considered carefully. A too high spectral resolution lowers the SNR but might not lead to an increased science output (see for an example in Ca II 854.2 the tests by de la Cruz Rodríguez et al. 2012).

3.1 Magnetic structure at supergranular scales

The magnetic structure in the photosphere at supergranular scales is relatively well understood (e.g., Gošić et al. 2014). The magnetic field in the chromosphere is much less well constrained owing to the weaker polarization signal compared to photospheric lines. This is especially true outside active regions.

OP 3.1.1 Magnetic field structure in the quiet Sun chromosphere

See table on page 77.

This OP aims to answer the following questions: How do network field and internetwork field look and connect? How does the magnetic carpet look in the chromosphere? At what heights is the magnetic canopy located? How does the height of the $\beta = 1$ surface vary? To do so a full supergranule should be scanned with a slit spectrograph, covering many

lines from the photosphere to the chromosphere, simultaneously and co-spatially to allow for multi-line inversions. Such data can be inverted using multi-line non-LTE inversion codes (de la Cruz Rodríguez et al. 2016) to get full 3D magnetic and atmospheric structure. It can also be used for field extrapolations and construction of magnetic connectivity maps. The aim is to measure weak fields, so spatial resolution is of secondary concern, signal to noise and spectropolarimetric accuracy are more important. The Na I lines are needed to get sensitivity in the upper photosphere and low chromosphere, in the sensitivity gap between Fe I 630.1/630.2 and Ca II 854.2.

OP 3.1.2 Fibrillar structure of the chromosphere

See table on page 77.

One of the most intriguing features of the solar chromosphere is its fibrillar topology in almost all chromospheric structures. This behaviour, beautifully manifested in high resolution $H\alpha$ images, clearly originates from the presence of smallest-scale magnetic elements. Measurements at the diffraction limit of the Swedish 1-m Solar Telescope in the Ca II K line show that these fibrillar structures cover a large fraction of the solar chromosphere: they extend from active regions far into the quiet Sun, where they become more diffuse and at some point too faint for detection (Pietarila et al. 2009). There is an indication that such quiet Sun fibrils can effectively suppress the dominant 3-minute oscillatory power in the chromosphere. Fibrils carry transverse waves that can propagate with Alfvénic speeds (Morton et al. 2012). A thorough investigation of these fibrillar structures requires spectropolarimetric measurements at the highest attainable spatial resolution since the average width of fibrils is comparable to or smaller than the diffraction limit of a 1-m class telescope. This program focusses on their spatial structure at the highest resolution and dynamics on timescales of ~ 30 s or longer.

3.2 Spicules and jets

The chromosphere produces a plethora of jet-like phenomena. The most ubiquitous are Type I and type II spicules (De Pontieu et al. 2007b). Type I spicules are slow-mode magneto-acoustic shocks, and are rather well modelled and understood (De Pontieu et al. 2007a). Type II spicules are faster, and not so well understood. Martínez-Sykora et al. (2017) proposed a model for Type II spicule acceleration by magnetic tension forces. A crucial aspect in this model is the effect of ion-neutral interactions, which allows the magnetic field to diffuse from the photosphere into the chromosphere and amplifies the magnetic tension. Ion-neutral interactions also drive heating in the spicules. It is highly interesting to study Type II spicules to test the model predictions. Spicules are narrow (< 200 km) and not very long (up to ~ 10 arcsec) and their acceleration sites are compact (< 1 arcsec). They evolve fast (< 2 min) and flows reach high speeds (100 km s^{-1}). They are thus best studied with 2D imaging spectropolarimetry. Imaging spectropolarimeters have the additional advantage that the S/N cadence trade-off can be made after the observation is made. Fabry-Perot instruments will likely be too slow to capture the evolution owing to the need for spectral scanning, while slit spectrographs are fast but require slow spatial scanning if imaging information is required.

OP 3.2.1 Type II spicule acceleration on disk

See table on page 77.

The aim is to observe the edge of a network or plage region that harbors Type II spicules, with sit and stare using 2D spectroscopy. Full-Stokes polarimetry in Fe I 630.1/630.2, Na I 589 and Ca II 854.2 are required to get the magnetic field in the photosphere and low chromosphere. Spectroscopy in $H\alpha$ is used to follow the spicule to larger heights, spectroscopy in Ca II K should ideally be used to obtain images of the acceleration site at the highest spatial resolution.

OP 3.2.2 Type II spicule evolution on disk

See table on page 78.

Similar to 3.2.1, but now a focus on the further evolution after launching. Simulations indicate that the spicule axis and the magnetic field vector need not be parallel. He I 1083.0 observations are needed to measure magnetic field in the upper chromosphere.

OP 3.2.3 Type II spicule evolution off-limb

See table on page 78.

This program is similar to 3.2.2, but here the aim is to observe at the limb. Off-limb observations are required to have less superposition of structures along the line of sight. Context imaging using a Fabry-Perot or broad-band imager is desirable. Coordination with space missions is highly desirable to trace spicule evolution at UV wavelengths to higher layers and temperatures. Example scientific questions that can be addressed with this OP are: how does spicule fine structure look and evolve? What kind of wave motions do spicules exhibit? What is the energy flux and power spectrum of the waves? What is the magnetic field structure in spicules?

OP 3.2.4 Magnetic field of spicules

See table on page 78.

Spicules are intrinsically magnetic structures. Measuring the magnetic field has been difficult owing to the small width, fast evolution and relatively weak fields in spicules. This OP focusses on measuring the magnetic field vector in many spicules using spectropolarimetric observations in the He I 587.6 nm (e.g. López Ariste & Casini 2005) and 1083 nm lines (e.g. Centeno et al. 2010). Both lines form in the triplet system of He I, and observing both lines simultaneously constrains the magnetic field and thermodynamic state of spicules better than single-line observations (e.g. Casini et al. 2009). The large spectral separation of the lines requires atmospheric refraction compensation in the spectrograph to get strictly co-spatial and co-temporal spectra.

3.3 Structure of small-scale chromospheric jets

OP 3.3.1 Small-scale chromospheric jets

See table on page 79.

Some numerical simulations indicate an occurrence of bidirectional or counterstreaming plasma flows within chromospheric jets, which have not yet been observationally confirmed. Fast plasma flows within jets may trigger increased turbulence and instabilities of the Kelvin-Helmholtz type at interface of the jet and the ambient plasma. However, the internal structure, dynamics and possible instabilities within the jets are beyond the resolution limit of current 1-m class telescopes. This OP aims to study the internal structure and dynamics of jets using imaging only.

3.4 Wave propagation, mode conversion and wave damping

The chromosphere is pervaded by compressive (magnetic-acoustic) waves, as well as transverse and torsional Alfvénic waves (e.g., Jess et al. 2015). There are indications for sausage-modes. A large fraction of these waves are excited in the photosphere by granular buffeting, and through motions of photospheric magnetic elements. Mode conversion occurs at the $\beta = 1$ surface, wave damping occurs, as well as wave reflection at the $\beta = 1$ surface and the transition region. Acoustic waves have dominant periods of around 3 minutes and speeds of order 10-30 km s⁻¹, while Alfvénic waves have been observed with periods down to 30 s, but waves with shorter periods are also present. The Alfvén speed in the chromosphere can be much higher (up to a few hundred km s⁻¹) than the sound speed. Observations of Alfvénic waves thus require much higher cadence than magneto-acoustic waves to resolve periods and track propagation.

OP 3.4.1 Alfvén waves

See table on page 79.

Transverse and torsional waves that propagate at Alfvénic speeds are observed in spicules and fibrils. This OP aims to derive phase speeds, periods, amplitudes, damping lengths and damping timescales. A large FOV is required to track propagation. High-cadence imaging in CaII K, H α is required to catch short period oscillations and oscillations with high phase speeds, but this comes at the price of sampling the lines at only a few wavelength positions. Spectropolarimetry at lower cadence in Fe I 630.2, CaII 854.2 is required to constrain the magnetic field.

OP 3.4.2 Acoustic wave interaction with chromospheric field structure

See table on page 79.

The aim of this OP is to track how acoustic waves propagate upward and reflect, deflect, and/or mode-convert when interacting with the chromospheric magnetic field canopy, and determine the wave energy flux carried by the waves. Waves propagate both vertically and horizontally, so a decent FOV is required. Knowledge of the photospheric and chromospheric magnetic field is required, so spectropolarimetry in Fe I 630.2 and CaII 854.2 nm is needed. Wave amplitudes are expected to be of order 5–20 km s⁻¹.

3.5 Flux emergence and reconnection events

OP 3.5.1 Highly variable phenomena in the chromosphere

See table on page 80.

SST observations have shown numerous examples of small-scale, highly variable phenomena with associated plane-of-the-sky velocities well above 100 km s⁻¹ (Lin et al. 2012; Scullion et al. 2015). These events appear to be related to re-configurations of the magnetic field, possibly as a result of reconnection. Rapid intensity variations can also be observed at sites of magnetic flux emergence. To resolve these phenomena, both high spatial resolution and high temporal

resolution are needed. Fabry-Perots have large FOV but require spectral scanning and thus allow only a few wavelengths to fulfil the cadence requirement.

OP 3.5.2 Highly variable phenomena in the chromosphere

See table on page 80.

As observing program 5, but now with 2D spectroscopy to get full spectral information. Horizontally propagating disturbances cannot be tracked over a large distance.

OP 3.5.3 Reconnection at different heights

See table on page 80.

Reconnection of magnetic field lines is thought to be a way of releasing large amounts of magnetic energy into the solar atmosphere. That released magnetic energy can be transformed into heating of the chromosphere and/or kinetic energy of the accelerated plasma that may be expelled from the reconnection site. Ellerman bombs have been shown to be the signal of reconnection at upper-photospheric levels (e.g. Vissers et al. 2015; Danilovic 2017) UV bursts (previously called IRIS bombs Peter et al. 2014) appear to be the signature of reconnection in the upper chromosphere and transition region (Hansteen et al. 2017).

Vissers et al. (2015) also describe flaring active region fibrils that appear very bright in UV lines, but also visible in H I 656.3, and likely also in Ca II 393.4. They are likely signs of reconnection too, but possibly in a different geometry than Ellerman bombs. Many unanswered questions remain: What are the fundamental differences / similarities between Ellerman Bombs, UV bursts, microflares, and flaring active-region fibrils? Do Ellerman bombs trigger reconnection higher up? How does reconnection happen as a function of height? Because reconnection is an inherently magnetic process, this OP requires high polarimetric sensitivity to constrain the magnetic field.

Such observations require a high cadence and a large spectral range because of the high speeds of reconnection jets and the very wide line profiles observed in such reconnection events. Co-observing with space based UV instruments is essential to track the evolution of reconnection events at higher temperatures.

3.6 Observational determination of electric currents

Dissipation of electric currents are one of the candidates proposed to explain the heating of the upper solar atmosphere. Measuring not only the magnitude but also the direction of the electric currents is important to quantify the contribution of Ohmic dissipation to the overall heating. However, this is a very difficult measurement to make, and only a few attempts have been done (Solanki et al. 2003; Socas-Navarro 2005) The electric current is determined by taking the curl of the magnetic field. The magnetic field must be determined by inversion of spectropolarimetric measurements. The curl operation amplifies any noise or errors made in the determination of the magnetic field.

The measurements are best done by simultaneously observing a number of photospheric and chromospheric lines. Because the atmosphere and magnetic fields evolve, there is a trade-off between spatial resolution, time resolution, S/N and FOV. Various different observations with different trade-offs must be made.

OP 3.6.1 Electric currents and heating in active regions

See table on page 81.

The magnetic field in the chromosphere of active regions is relatively strong (typically ~100-500 G) and a lower S/N is sufficient, This allows for the use of Fabry-Pérot instruments with an acceptable time resolution.

OP 3.6.2 Electric currents and heating outside active regions

See table on page 81.

Spectro-polarimetric scans of a quiet region (network and internetwork). Field strengths are lower than in active regions, so a higher S/N is required. In order to perform inversions (quasi-) simultaneous spectra are required. This cannot be done with narrowband filtergraphs, so a slit spectrograph is required.

OP 3.6.3 Electric currents and heating outside active regions at small spatial scales

See table on page 81.

Similar to OP 3.6.2. This program focusses on small scale structure and fast time evolution using integral field units.

3.7 Temperature structure of the solar chromosphere

The temperature of the solar chromosphere is time-varying and highly inhomogeneous. Using off-limb observations and the center-to-limb variation in the solar 4.7- μm rovibrational bands of carbon monoxide. Ayres & Rabin (1996) found evidence for a surprisingly cool atmospheric component at chromospheric heights, as tracked by the presence of CO molecules. Observations and inversions of optical and IR lines indicate temperatures down to 4000 K, but these methods are insensitive to lower temperatures because of the non-LTE scattering formation of the diagnostic spectral lines. Theoretical studies indicate that a non-magnetic solar chromosphere should cool down to temperatures below 1500 K (Leenaarts et al. 2011). Ion-neutral interactions can dissipate sufficient energy in cool bubbles to avoid cooling to such low temperatures (Martínez-Sykora et al. 2012, 2017) The location and temperature of such cool clouds is thus still unclear. Many of the previous observing programs can be used to infer temperatures using inversions (e.g. de la Cruz Rodríguez et al. 2012).

There are more options to study temperatures and the next OPs use different techniques to study cool phases of the chromosphere.

OP 3.7.1 Measurement of CO clouds

See table on page 82.

Long-slit spectroscopy in the 4700 nm CO bands has been a proven tool to study cool chromospheric gas. The large diameter of EST allows such observations at previously unattainable resolution. The polarisation in the CO lines offers a unique window on the magnetic field in cool gas.

OP 3.7.2 Temperature bifurcation diagnostics by scattering polarization in Ca II K

See table on page 82.

Scattering polarization in CaII K spectra taken at the solar limb has been used by Holzreuter et al. (2006) to find further evidence of a temperature bifurcation in the chromosphere, independently from techniques based on CO spectroscopy or inversions. Observed polarization signals are of order 0.5%-1%. Slit spectroscopy can catch such signals in 30 s, allowing much better time resolution than the 30 min of the data in Holzreuter et al. (2006).

3.8 Magnetic field measurements using Ca II H&K

Martinez Pillet et al. (1990) measured circular polarization in the Ca II H&K lines for quiet sun, plage, a sunspot umbra and penumbra and a flare. A Stokes V signal of order 5–15% was measured in all but the quiet Sun observations. Polarization measurements in these lines could thus in principle be an interesting tool to infer magnetic fields in the formation height range between Ca II 854.2 nm and He I 1083 nm. Spectropolarimetric data can be obtained with slit spectrographs, Fabry-Perot instruments and/or integral field units. All previous OPs that include the line could thus in principle be extended with polarimetry. However, as the line has a very low photon flux, it takes long integration times to get a high SNR. As an example program we include an observing program with a slit spectrograph aimed at measuring the magnetic field structure in plage.

OP 3.8.1 Magnetic field determination in plage including Ca II H&K

See table on page 82.

The aim of this OP is to determine the 3D spatial variation of the magnetic field in plage using multiple lines, including Ca II H&K.

3.9 Summary of requested instrument capabilities

Almost all observing programs require (near-) simultaneous multi-line observations. Popular lines are photospheric Fe I lines (557.6, 630.1/630.2) to get magnetic fields in the photosphere, Ca II 854.2 to get magnetic field in the middle chromosphere and He I 1083 for upper chromospheric magnetic fields. Imaging spectroscopy (but not spectropolarimetry) in the Ca II H and/or K line is often requested too, as is imaging in H α 656.3. Other lines that are requested are He I 587.6, Na I 589.0/589.6, Si I 1082.7, and the CO band at 4700 nm.

The FOV requested for imaging spectropolarimetry is typically 30'' \times 30'', except where entire supergranules or active regions are observed, in which case a larger FOV of typically 60'' \times 60'' is requested. OPs focussing on events evolving on timescales of seconds typically request a small 10'' \times 10'' FOV.

Ideally the light distribution system is thus split into at least four domains: < 500 nm, 500-700 nm, 700-900 nm, and > 900 nm channels, allowing four lines, from the photosphere to upper chromosphere to be observed simultaneously using FPs or IFUs.

Popular instrument requests and capability per channel (we do not include the infrequent requests for > 1100 nm) can be divided as

	IMSP		FP		SP	
	requested	polarimetry	requested	polarimetry	requested	polarimetry
< 500 nm	yes	no	yes	no	yes	yes
500-700 nm	yes	yes	yes	yes	yes	yes
700-900 nm	yes	yes	yes	yes	yes	yes
> 900 nm	yes	yes	yes	yes	yes	yes

4 Large scale magnetic structures: sunspots, prominences and filaments

Authors: Jan Jurčák, Marian Martínez González, Luc Rouppe van der Voort

Sunspots and filaments (also known as prominences when observed off-limb) are large-scale magnetic structures that cover a significant fraction of the solar disk. EST is designed to excel at resolving the smallest spatial scales which inevitably implies that the field of view is limited as compared to the extent of the largest sunspots and filaments. However, fundamental physical processes that define these structures occur at spatial scales that are currently beyond the reach of present-day instrumentation. The capabilities of EST will be crucial for the full characterisation of the smallest building blocks of these structures so that fundamental questions about the nature of sunspots and filaments can finally be addressed.

Sunspots are large-scale concentrations of strong magnetic fields and the primary manifestation of solar activity. The umbra is the dark central area of a sunspot and harbors the strongest and most vertical magnetic field. It is surrounded by the filamentary penumbra where the magnetic field is more inclined and in part nearly horizontal. The strong magnetic field in sunspots inhibits the normal convective overturning motion that results in the granulation pattern found all over the solar surface outside active regions. However, recent advances in observations and numerical simulations have arrived at the view that fine-scale structures in sunspots, such as umbral dots, light bridges and penumbral filaments, result from a form of convective motions that is heavily affected by the strong magnetic field.

For example, spectropolarimetric observations of sunspot penumbrae (van Noort 2012; Tiwari et al. 2013), observations of orphan penumbrae (Kuckein et al. 2012; Jurčák et al. 2014b; Zuccarello et al. 2014), and advanced MHD simulations of sunspots (e.g., Rempel 2011) provide a coherent picture of sunspot penumbrae resulting from convection in an inclined magnetic field. The penumbral filaments contain horizontal fields and carry a plasma flow, historically known as the Evershed flow. These regions with horizontal magnetic field are interlaced with more vertical and typically stronger field (so-called “spines”). The penumbral topology of rapidly varying magnetic field inclination is often referred to as the “uncombed” magnetic field (Solanki & Montavon 1993).

Filaments consist essentially of chromospheric material found at great altitude as magnetically levitating clouds embedded in the hot corona (Leroy 1989; Tandberg-Hanssen 1995; Trujillo Bueno et al. 2002). When observed off-limb, filaments are traditionally referred to as prominences. Broadly speaking, filaments can be segregated in active region (AR) and quiescent (QS) structures. Both lie above the polarity inversion line, an observationally defined line that separates the two dominant opposite polarity magnetic fields.

QS filaments are very long structures suspended at more than 10 Mm above the surface that often live for weeks or even months. AR filaments are formed in active regions, often in recurrent flaring areas, and are shorter in length and life time as compared to QS filaments. They are difficult to be seen as prominences at the limb because they lie lower in the atmosphere and are hidden behind other features along the line of sight. Though having different parameters, their nature is generally believed to be the same. In particular, the magnetic field plays a fundamental role in the formation, support, and eruption of filaments.

The diagnostic of magnetic fields in such structures is mainly based on spectro-polarimetry in the He I multiplet at 1083.0 nm and in the He I D₃ line at 587.6 nm. The reason is that these lines are basically absent in the photospheric spectrum and its mere detection provides a fingerprint on the presence of dense chromospheric structures.

The radiation produced in these lines (either in absorption or in emission) is linearly polarised due to scattering processes. When atoms are illuminated by the anisotropic radiation field from the underlying photosphere, it induces atomic-level polarisation (population imbalances and coherences between magnetic sublevels). This linear polarisation is modified via partial decoherence in the presence of a magnetic field (Hanle effect), and circular polarisation is typically generated by Zeeman splitting (e.g., Leroy 1989; Trujillo Bueno et al. 2002). Hence, to constrain the magnetic field vector in filaments

we need to observe the full Stokes vector. These signals are typically weak, and we need a signal to noise ratio much better than 2000 to properly infer the magnetic configuration.

4.1 Stability of the umbra

Recent studies show that umbrae are stable if the vertical component of the magnetic field (B_{ver}) is stronger than a certain invariant value (1867 G as estimated from Hinode spectropolarimetric data, Jurčák et al. 2018). If B_{ver} is lower, these originally dark umbral areas are prone to be transformed into penumbrae, light bridges, or granulation depending on the magnetic field inclination and location within the umbra (Jurčák et al. 2015, 2017). The theoretical study by Wentzel (1992), and the advanced MHD simulations by Rempel (2011), suggest a mechanism to turn predominantly vertical field lines into horizontal ones: from the convectively unstable sub-photosphere, hot buoyant plasma pushes upwards along the umbral field lines; when cooling, the mass load exerted by the plasma will bend and incline field lines where the vertical field has reduced strength; only in areas with strong vertical magnetic fields is the magnetic tension as a restoring force strong enough to prevent the field lines from bending over and becoming horizontal. This scenario which explains the fundamental property of sunspot structure has yet to be confirmed by observations. This requires observations with high temporal and spatial resolution to measure the interplay between the convective motions and the magnetic field at the boundaries between umbra and penumbra, granulation, and light bridges both in the solar photosphere and chromosphere.

OP 4.1.1 Stability of the umbra - interplay between the convection and magnetic forces

See table on page 83.

4.2 Umbral dots

Umbral dots are very fine-scale structures in sunspot umbrae and pores that have sub-structure that is barely resolved with present-day instrumentation. Umbral dots were successfully reproduced in advanced MHD simulations (Schüssler & Vögler 2006), where they correspond to the top of magneto-convective cells in the strong and vertical magnetic field of the umbra. Current spectropolarimetric observations of highest spatial resolution are in agreement with the modeled structures (see, e.g., Ortiz et al. 2010), but the observations still do not fully spatially resolve umbral dots. Spectropolarimetric observations of better spatial resolution achieved simultaneously in a number of spectral lines probing different heights in the atmosphere would further clarify the nature of umbral dots and their impact on the sunspot chromosphere.

OP 4.2.1 Multi-wavelength analysis of umbral dots

See table on page 84.

4.3 Structure of cool sunspot umbrae

The presence of molecular lines near the 630 nm Fe I lines and the IR Fe I lines at 1.56 μm hinder the reliable characterization of the magnetic and the thermal structure of cool sunspot umbrae. The Ti I multiplet at 2.2 μm , sensitive to properties of cool plasma, is free from molecular blends and therefore represents an ideal tool for the diagnostics of the coolest regions on the solar surface. Additionally, this multiplet is extremely Zeeman sensitive allowing for a reliable determination of the magnetic structure of cool umbrae. Alternatively, cool sunspot umbrae can be studied using observations in the visible and 1.6 μm lines, but this requires to account for the molecular lines in the inversion codes that are used to interpret the Stokes profiles.

Such observations would allow us to investigate the possible presence of granular-scale patterns in sunspot umbra that were not yet detected, but their signatures are present in the MHD simulations of sunspot umbrae (Schüssler & Vögler 2006; Vitas 2011).

OP 4.3.1 Probing the structure of cool sunspot umbrae

See table on page 85.

4.4 Umbral flashes as a probe of fine structure in the umbra chromosphere

Umbral flashes are sudden brightenings in the Ca II H & K and IR triplet lines observed in the chromosphere above the sunspot umbra. They were first reported by Beckers & Tallant (1969) and are commonly thought to be formed by magnetoacoustic waves that originate in the photosphere and steepen into shocks as they propagate to chromospheric heights (Bard & Carlsson 2010). The increased brightness provided by the umbral flashes has revealed fine structure of both cool and hot material intermixed on very small spatial scales (Socas-Navarro et al. 2009; de la Cruz Rodríguez et al. 2013). These fine structures could be created by the shocks or become apparent when they are illuminated by the umbral flashes. The horizontal orientation and general stability of these small-scale fibrillar structures have questioned the general belief that the magnetic field topology in the umbra chromosphere is uniformly vertical (Henriques et al. 2015). High-resolution imaging in Ca II K or H combined with full-Stokes polarimetry in other spectral lines will allow us to gain insight into the inhomogeneous nature of the chromosphere above the sunspot umbra. The lack of intensity in the umbra and the strong photospheric field make this project particularly challenging and well suited for the polarimetric capabilities of EST.

OP 4.4.1 Umbral flashes

See table on page 85.

4.5 Penumbral and umbral micro-jets

In recent years, various chromospheric jet-like phenomena have been discovered in the sunspot umbra and penumbra. There are impulsive penumbral and umbral microjets (Katsukawa et al. 2007; Bharti et al. 2013), and slower, longer-lived, and more ubiquitous sunspot dynamic fibrils (in the umbra also referred to as umbral spikes, Rouppe van der Voort & de la Cruz Rodríguez 2013; Yurchyshyn et al. 2014).

The latter are a shorter version of the well-known dynamic fibrils found in plage regions and are well understood (De Pontieu et al. 2007a). It is still unclear how the impulsive micro-jets are related or affected by the dynamic fibrils and large-scale sunspot waves (umbral flashes and running penumbral waves). It is likely that they are driven by magnetic reconnection. Recent studies of IRIS data suggest that some chromospheric microjets may be driven from above by reconnection in the transition region rather than from below (Samanta et al. 2017). With the unprecedented spatial resolution of EST, we will be able to better characterise the dynamic properties of sunspot jets. A recent study with the Swedish 1-meter Solar Telescope (Drews & Rouppe van der Voort 2017) indicated that not all microjets were spatially resolved (the SST diffraction limit at Ca II 8542 is 130 km) so the high spatial resolution of EST is needed to establish the possible existence of a ubiquitous population of narrow microjets that can presently not be detected. Combined with high-sensitivity spectro-polarimetry, we will be able to address the formation mechanism of microjets. The strong field environment of sunspots, which gives strong signals in spectro-polarimetric diagnostics, makes sunspot microjets an ideal target to study the details of the fundamental process of magnetic reconnection.

OP 4.5.1 Penumbral and umbral micro-jets

See table on page 86.

4.6 Light bridges

Light bridges are long bright lanes that cross the dark umbrae of sunspots. Light bridges display a large variety of morphology and fine structure: some harbor cells that resemble the granulation cells of the normal quiet Sun, some are intrusions of penumbral filaments, while others appear as the alignment of bright grains and seem to be associated with the isolated umbral dots found in the umbra (see, e.g., Rimmele 2008; Rouppe van der Voort et al. 2010; Lagg et al. 2014; Schlichenmaier et al. 2016).

It is now firmly established that light bridges are regions that harbor convective flows of relatively weakly magnetized plasma in the strongly magnetized surroundings of the umbra. It will be of great interest to utilize the capabilities of EST to characterize in detail the conditions in the different types of light bridges to explain their morphology and fine structure. With these light bridge observations, together with the detailed observations of umbral dots and penumbral filaments as described in other observing programs, EST will provide a major advancement in the understanding of magneto-convection in the strong field regime.

It has been noted that light bridges often show enhanced chromospheric activity, with ($H\alpha$) surges and chromospheric jets reported in a number of analyses (see, e.g., Roy 1973; Shimizu et al. 2009; Louis et al. 2014). There have been strong indications that magnetic reconnection is driving these surges (Robustini et al. 2016, 2018). Tian et al. (2018) differentiate between two types of jets above light bridges: a short type that is virtually continuously present and are caused by upward leakage of magnetoacoustic waves from the photosphere, similar as for dynamic fibrils (De Pontieu et al. 2007a), and more

occasionally occurring long and fast surges that are caused by intermittent reconnection. Sometimes, the periodically re-occurring jets above light bridges are referred to as light walls (Yang et al. 2017; Hou et al. 2017). The comprehensive study of the chromospheric and transition-region properties of light bridges by Rezaei (2018) confirms that they are complex multi-temperature structures associated with enhanced energy deposition, but demonstrates the need for further multi-height investigation of the magnetic and dynamic structure through the photosphere and chromosphere, in order to identify heating mechanisms and the height at which energy is deposited.

OP 4.6.1 Structure and dynamics of light-bridges

See table on page 87.

4.7 Evolution of an individual penumbral filament

There is general consensus about the structure of individual penumbral filaments, both from an observational and a theoretical point of view. Penumbral filaments are convective cells heavily influenced by the strong and inclined magnetic field of the penumbra (Tiwari et al. 2013; Rempel 2011). However, there still remain questions about the details of filament formation and decay processes, their influence on the larger-scale magnetic field topology, and the dynamical evolution of the plasma flow field and the magnetic field within the filaments. This is closely related to the OP 4.1.1 that focus on a different aspect of the interplay between the convection and the magnetic field.

OP 4.7.1 Evolution of an individual penumbral filament

See table on page 88.

4.8 Formation and decay of sunspot penumbrae

While individual penumbral filaments have a typical life time of less than an hour, the penumbra as a whole is generally a rather stable structure that remains present around a sunspot for days or even weeks. The formation and decay of the sunspot penumbra, on the other hand, is a process that only takes a few hours. For example, Schlichenmaier et al. (2010) analysed the formation of a penumbra around a protospot that took only 4.5 hours. They concluded that this protospot developed a penumbra while accumulating magnetic flux from the active region flux emergence site. Rezaei et al. (2012) found signatures of a canopy at the photospheric level around the protospot before the penumbra appearance at the solar surface and this is in agreement with analyses of Shimizu et al. (2012) and Romano et al. (2013), who found indications of a penumbral halo at the chromospheric level prior to the formation of a penumbra in the photosphere. Jurčák et al. (2014a) found an increase in magnetic field inclination at the protospot boundary prior to the onset of penumbra formation. Romano et al. (2013), Romano et al. (2014a), and Murabito et al. (2016) put these findings into the context of magnetic flux at chromospheric height bending down to the photosphere leading to penumbra formation at the solar surface. They also found signatures of (counter Evershed) inflowing material similar to those reported by Schlichenmaier et al. (2011), and they discuss on the siphon nature of those in the context of falling flux tubes.

From the modeling point of view, several studies help to put into context the observational findings: Simon & Weiss (1970), Jahn & Schmidt (1994), and Hurlburt & Rucklidge (2000) found that with increasing magnetic flux, the inclination of the field becomes increasingly more horizontal. Numerical simulations by MacTaggart et al. (2016) show that magnetic canopies naturally form during the emergence of a twisted flux tube. According to Rempel (2012), these canopies have sufficiently inclined magnetic field to form extended penumbrae. Wentzel (1992) proposed a penumbral model fed by flux tubes fallen onto the photosphere owing to the upwelling of a mass flow in the inner footpoint (within the umbra) of field lines closing up (submerging) in the surroundings of a sunspot.

A decrease in the size of the penumbra marks the beginning of the end for mature spots. The penumbra eventually disappears and only the umbra remains visible in white-light images. It has been suggested that this process contributes to the removal of flux from active regions (see Martínez Pillet 2002, and references therein). In that case, it would be an essential ingredient of the solar activity cycle. Penumbral decay may also be a source of localized chromospheric and coronal heating (Wang et al. 2004; Deng et al. 2005).

Despite its far-reaching implications, the disappearance of the penumbra is not well understood. Bellot Rubio et al. (2008), Watanabe et al. (2014), and Verma et al. (2018) analysed the disappearance of penumbra and explained it by the rise of photospheric magnetic field lines into the chromosphere, i.e., an inverse mechanism to the predicted formation mechanism of the penumbra. Similar changes of the magnetic field orientation were also described in case of flare-induced rapid penumbral decay (Wang et al. 2004; Deng et al. 2005).

For the moment, this idea remains speculative. To identify the exact mechanism responsible for the disappearance of the penumbra, EST should perform spectropolarimetric observations of mature spots throughout their decay phase. This

requires high and stable data quality on long time scales. Furthermore, it is important to measure magnetic field orientation both in photosphere and in chromosphere with high spatial resolution.

OP 4.8.1 Capturing the formation and decay of penumbrae

See table on page 89.

4.9 Relation between moat flows and sunspot decay

Mature sunspots are usually surrounded by a so-called moat region that harbors a conspicuous flow pattern with radially outward moving magnetic features (MMFs, Sheeley 1972; Hagenaar & Shine 2005). The sizes of sunspot moats are typically close to supergranular cell sizes and the moat flow velocities reach several hundred meters per second. The origin of sunspot moats is not well understood. Observations (e.g., Vargas Domínguez et al. 2008) show that the appearance of moat flows is closely related to the presence of a penumbra. However, this view has been challenged by Löhner-Böttcher & Schlichenmaier (2013) as well as by Rempel (2015) who find that the presence of the moat flow does not depend on the existence of an Evershed flow and/or penumbra around the umbra.

It has also been suggested that the MMFs are extensions of penumbral filaments which turn back below the visible solar surface at the sunspot magnetopauses and reappear as MMFs in the moats. Observations by Sainz Dalda & Bellot Rubio (2008) indeed support this so-called sea-serpent model (Schlichenmaier 2002). As argued by Martínez Pillet (2002), MMFs, while probably part of the decay process, have an origin not related to the decay process itself. However, there are also studies relating the observed decay rates of the sunspot magnetic flux to the flux carried by MMFs (Hagenaar & Shine 2005; Kubo et al. 2008).

Detailed observations of the relationship between sunspots and their surrounding quiet regions are required to solve the puzzling appearance of sunspot moats. Particularly, high spatial and temporal resolution observations of the dynamics of sunspot penumbrae and MMFs, including full magnetic field vector and flow velocity measurements, as will be provided by the EST, shall lead to substantial progress in the understanding of large-scale dynamic processes in and around sunspots and how these processes influence sunspot decay. The observational setup is comparable to the study of penumbra formation and decay, but need to capture a larger FOV.

OP 4.9.1 Observations of moat flow properties and its impact on sunspot decay

See table on page 90.

4.10 Fine structure of prominences and filaments

A very important issue, with respect to expected capabilities of the EST, is the fine structure of the magnetic field in filaments (and prominences but we'll use the word filament since both are the same phenomenon observed with different geometries). Direct imaging shows that filaments appear to be highly dynamic at different spatio-temporal scales, tend to form very fine filamentary structures, and show counterintuitive behaviors arising from the competition between the vertical gravitational stratification and magnetic forces operating in any other direction. However, such fine structure has never been reported in magnetic field measurements. In fact, the magnetic fields inferred with the best spatial resolution achievable at present (0.5'') show a very smooth field (Casini et al. 2003; Orozco Suárez et al. 2014; Martínez González et al. 2015). Understanding if the fields follow the intensity structures or not is of vital importance to understand how these structures are sustained against gravity and insulated from the million degree corona. In particular, various theoretical models predict magnetic dips in which the filament plasma is condensed, but such dips have not been observationally identified so far. Confirming or denying this will support the works that claim that magnetic fields in filaments are horizontal to the surface and unrelated to what we observe in intensity (e.g., Orozco Suárez et al. 2014; Levens et al. 2016) or aligned with the plasma structures (e.g., Zirker et al. 1998; Martínez González et al. 2015). Spectro-polarimetric data with much higher spatial resolution and better polarimetric accuracy than achieved today are needed for this.

OP 4.10.1 Determining the evolution of the magnetic and dynamic fine structure of prominences and filaments

See table on page 91.

OP 4.10.2 Determining the magnetic and dynamic fine structure of prominences and filaments

See table on page 91.

4.11 Are quiescent and active region filaments the same phenomenon?

Active region filaments as well as quiescent ones are seen as dark absorption features in the core of many chromospheric spectral lines. Their properties seem to differ mainly in the life time and the heights they can reach. However, they are assumed to have the same origin, the only difference being the region where they appear, that somehow has an impact in their lifetimes and heights of formation.

At present, the few existing measurements of the full magnetic field vector of QS filaments show that a controversy exists concerning the magnetic properties of both structures. Spectropolarimetric observations of quiescent filaments have revealed magnetic fields strengths of the order of a tens of Gauss (Merenda et al. 2006; Orozco Suárez et al. 2014; Martínez González et al. 2015), while AR filaments have been found with magnetic fields from 100G up to 700G (Sasso et al. 2007, 2011; Xu et al. 2012b; Kuckein et al. 2012). However, recently, the strongest fields have been put in doubt by Díaz Baso et al. (2016). In light of these results it is clear that, from the observational point of view, the precise topology of magnetic fields in filaments is still a matter of debate. The unprecedented spatial resolution and polarimetric accuracy expected of the EST will help overcoming the observational challenges that restrain the study of these structures in detail. In particular, the main aim with the EST observations will be to unveil if both QS and AR filaments harbor weak fields or not. In other words, EST observations will help unveil if QS and AR chromospheric structures share the same origin and are the same magnetic phenomenon seen in different perspectives.

Recently, it has been shown from empirical data of AR filaments that these show complex Stokes profiles, revealing that gradients of the physical properties (including the magnetic field) happen along the line of sight (Díaz Baso et al. 2018, submitted). However, the analysis of such signals using more realistic models including gradients or many atmospheric components has not been successful because of the low signal to noise ratio. Improving the polarimetric accuracy by collecting more photons will definitely help a proper inference of the gradients of the magnetic and thermodynamical properties in AR filaments.

OP 4.11.1 Comparison of the magnetic field properties in QS prominences and AR filaments

See table on page 92.

4.12 Magnetic field and dynamics of tornado prominences

The name of solar tornadoes was first introduced by Pettit (1932) to label a specific kind of solar prominences that appeared like "vertical spirals or tightly twisted ropes". Recently, the high spatial resolution and the time cadence and continuity of the coronal data provided by the Atmospheric Imaging Assembly (AIA) onboard the Solar Dynamics Observatory (SDO) has reawakened the interest on the study of solar tornadoes. Solar tornadoes are nowadays associated to vertical funnel-shaped dark structures in the coronal Fe XII line at 17.1 nm. These structures are found to be hosted in the legs of some quiescent prominences (e.g., Su et al. 2012; Wedemeyer et al. 2013), and are directly linked to the evolution and final fate of the prominence.

The magnetic and dynamic properties of solar tornadoes are still a matter of debate and a hot topic in the recent literature. Concerning the magnetic configuration, Martínez González et al. (2016, 2015) argue that they harbour vertical, helical fields that connect the main body of the prominence with the underlying surface. These findings are in agreement with previous claims of vertical fields in filament barbs (or prominence legs; Zirker et al. 1998). On the contrary, Schmieder et al. (2015) and Levens et al. (2016) infer magnetic fields that are almost parallel to the surface, in agreement with the magnetic field measured in a photospheric barb endpoint (López Ariste et al. 2006). A major advancement in observational capabilities such as EST will provide is required to settle the debate.

Concerning the dynamics of tornado prominences, a debate exists on the real or apparent rotation of the structure. Su et al. (2012) reported on the oscillatory pattern in the plane-of-the-sky motions of solar tornadoes as seen in SDO/AIA 17.1 nm, claiming that these structures are rotating with periods of about 50 min. Subsequent works (Orozco Suárez et al. 2012; Wedemeyer et al. 2013; Su et al. 2014, Levens et al. 2015) confirm the rotation scenario measuring opposite sign Doppler shifts at both sides of prominence legs. Panasenco et al. (2014) propose that the rotation of solar tornadoes as seen in SDO/AIA 17.1 nm is only apparent and could be explained by oscillations. In particular, Luna et al. (2016) propose that this apparent rotational motion can be caused by large amplitude oscillations along the magnetic field. The work of Mghbrishvili et al. (2015) explore both the rotation and the wave scenario and find both compatible with SDO/AIA 17.1 nm data of solar tornadoes. They also propose that, most likely, both rotation and waves are at work, identifying the oscillation as a magnetohydrodynamic kink mode. Consistent with this view, Martínez González et al. (2016) argue that (1) if rotation exists, it is intermittent, lasting no more than one hour, and (2) the observed velocity pattern is also consistent with an oscillatory velocity pattern.

The true nature of tornadoes and their relationship to prominences requires an understanding of their magnetic structure and dynamical properties. In particular for the dynamics, the EST spatial resolution and photon collector capabilities will be the key to understand if tornadoes are really rotating structures or if the rotation is just a visual illusion due to waves.

OP 4.12.1 Magnetism and dynamics of tornado prominences

See table on page 92.

5 Coronal Science

Authors: Sarah Matthews, Robertus Erdelyi, Mihalis Mathioudakis

EST is not optimised to carry out coronal science as one of its primary objectives, but it is well suited to providing complementary observations of the underlying photosphere and chromosphere that will significantly advance our understanding of coronal physics in a number of areas. In order to achieve this, co-ordination with space-based facilities will be required, and on the timescale of EST first-light the new space facilities that are envisaged to be operating are Solar Orbiter and Solar C EUVST. Co-ordination with Solar Orbiter will require detailed planning due to the complexity of its orbit. EST will be a critical resource for Solar C EUVST which, as a single instrument spacecraft, will provide high spatial, spectral and temporal resolution spectroscopic measurements of the transition region and corona, but have no on-board magnetic field capability.

5.1 Sunspot light-bridges/light walls

Sunspot light-bridges are one topic where EST combined with space observations can provide new insights into the magnetic and dynamic structure through the photosphere and chromosphere, and up into the overlying transitions region, allowing us to identify heating mechanisms and the height at which energy is deposited. Further details on a proposed observing programme to determine the variation of the magnetic field and dynamics of light-bridges can be found in Section 4 of this document.

5.2 Light Walls

One particular aspect of activity in light-bridges are surges, which have been found to form light walls that may penetrate deeper into the low corona, and also present a possible application for the SMS techniques, assuming asymmetric waveguides. Whilst their true nature is uncertain, the main features of light walls are the observed groups of surges, brightenings, and other magnetic structures in the chromosphere rooted in sunspot light bridges. They have been demonstrated to guide MHD waves driven by nearby disturbances (Yan et al. 2015) and provide an excellent testbed for the recently developed theory of asymmetric magnetic slabs ((Allcock & Erdélyi 2017); (Zsámberger et al. 2018)) allowing the actual geometry of light walls to be determined. Coupled with observations from space by Solar Orbiter and Solar C EUVST it may also be possible to determine where the plasma surrounding light walls has a turning point from high-beta to low-beta.

5.3 Origins of the solar wind

Hinode EIS has demonstrated that active regions upflows are common, but combined studies with in situ observations have showed clearly that not all upflows will become outflows measurable in the solar wind, and debates continue about where in the atmosphere the upflows originate. Solar Orbiter SPICE/SWA and Solar C EUVST will provide multi-vantage points measurements of coronal outflows, from the chromosphere to the corona in the case of Solar C EUVST, but need corresponding chromospheric magnetic field measurements in order to determine the processes that produce them, e.g. low-altitude reconnection, presence of small-scale open field at active region boundaries.

OP 5.3.1 Determining which coronal upflows become outflows

See table on page 94.

This observing plan aims to explore the role of small-scale photospheric and chromospheric dynamics in driving the solar wind at the edges of active regions and coronal holes where persistent upflows are observed in coronal lines. It will measure the magnetic field strength and direction from the photosphere to the upper chromosphere using spectropolarimetry, and the plasma dynamics using 2D spectroscopy to determine intensity, line widths, and line of sight velocities. Measurements should be coordinated with space platforms to provide the link to coronal spectroscopy, and ideally with in-situ measurements of plasma composition.

5.4 Probing pre-flare triggers

Spectroscopy of the transition region and corona indicates significant dynamics can be present prior to the onset of flares and eruptions (e.g. (Woods et al. 2017) and references therein), which are likely signatures of flux emergence and/or cancellation, tether cutting reconnection, and flux rope formation. Imaging spectroscopy indicates that these dynamics typically occur on small-scales and can subsequently lead to large-scale destabilization and explosive energy release. While these higher atmospheric signatures will be well-probed with Solar Orbiter EUV, SPICE and STIX, or with Solar C EUVST, the small scale chromospheric dynamics and magnetic field evolution that are critical to answering questions about how the flare/eruption is triggered, and when and where in the atmosphere flux ropes are formed, requires high resolution observations of the chromospheric magnetic field and associated dynamics.

OP 5.4.1 Measuring the photospheric and chromospheric dynamics before flares

See table on page 95.

This observing plan aims to explore the role of small-scale photospheric and chromospheric dynamics in the period before flare/CME onset and its relationship to TR and coronal dynamics. It will measure the magnetic field strength and direction from the photosphere to the upper chromosphere using spectropolarimetry, and the plasma dynamics using 2D spectroscopy to determine intensity, line widths, and line of sight velocities. Measurements should be coordinated with space platforms to provide the link to coronal spectroscopy.

5.5 Macrospicules/spicules and Transition Regions Quakes (TRQs)

Spicules and macrospicules are most likely generated in the lower solar atmosphere. Their impact on the solar corona is fundamentally important, as e.g. a mere 1% of spicule material may be sufficient to supply the mass needed for solar wind. Therefore, the diagnostics from their footpoints in the photosphere/low chromosphere up to the corona is an important question. Again, Joint EST and space observations may be able to identify and track MHD waves in these structures, enabling us to construct the magnetic skeleton of these jet structures. Details of observing programmes related to spicules and macro-spicules can be found in Section 2.

TRQS ((Scullion et al. 2011)) are energetic waves that have been detected in the transition region, and evolve in a similar manner to waves on 2D elastic waveguides. The question of what drives these TRQs remains open, with one possible explanation being that they are driven by (macro) spicules. Here, combining EST with space observations of the TR and corona will provide new insights into the effects of spicules on the TR and low corona, allowing us to identify the low coronal signatures of TRQs. To achieve this, a large sample of TRQs will be constructed, using IRIS/Solar Orbiter/Solar C EUVST in imaging mode, observed jointly with EST. The discovery of a link between these waves and RBEs identified in CRISP H α data is potentially very far reaching. Work by (Henriques et al. 2016) has already found convincing evidence of links between RBEs and coronal transient events. However, they did not study whether their coronal features corresponded to coherent waves (i.e., TRQs manifesting as propagating wave fronts in IRIS 140 nm data). This study on TRQs is vital for expanding our knowledge of the coupling between the lower solar atmosphere and the TR/low corona, and in particular searching for alternative drivers of TRQs (e.g. shocks) using chromospheric and NIR line profiles.

OP 5.5.1 Rising spicule signatures

See table on page 96.

5.6 Ellerman bombs

Ellerman bombs (EBs; (Ellerman 1917)) have been widely studied within ARs over the past century ((Nelson et al. 2015)) due to the hypothesis that they are driven by photospheric magnetic reconnection. Recent research highlighting similar events in the quiet Sun (QSEBs; (Roupe van der Voort et al. 2016)) suggests that magnetic reconnection can happen almost continuously throughout the photosphere. Using H α datasets (multi-positions and cadences 5 s) sampled at varying latitudes at the solar limb (where identification of QSEBs is less ambiguous) could lead to fundamental improvements in our understanding of QSEBs through analysis of their signatures in both EST and IRIS. EST will investigate the spatial coverage of QSEBs with respect to latitude to understand whether these events (and potentially small-scale magnetic reconnection in general) occur over the whole solar disk, including the polar regions, or whether they are confined to active latitudes (like sunspots). Further, obtaining a large sample of QSEBs will allow us to discover any links to jet-like events (e.g., spicules, macro-spicules) that penetrate the upper solar atmosphere using EST/Solar Orbiter/Solar C EUVST data. Section 2 outlines sample observing programmes for Ellerman bombs in more detail.

OP 5.6.1 Ellerman bombs in the lower solar atmosphere

See table on page 96.

6 Solar Flares and Eruptive Events

Authors: Lyndsay Fletcher, Francesca Zuccarello, Christoph Kuckein, Kévin Dalmasse, Paulo Simões

In a solar flare the reconfiguration of the current-carrying coronal magnetic field, in response to the development of a coronal magnetic instability, results in the liberation of stored magnetic energy. This is converted to thermal energy, kinetic energy of accelerated particles, and ultimately radiated energy, and mass motion if the flare is accompanied by a coronal mass ejection (CME). To understand the 3-D magnetic configuration and electrical current distribution before, during and after a solar eruptive event requires measurement of the photospheric and chromospheric vector magnetic field. As most of the radiated energy in a flare originates in the chromosphere, a concerted study of this region is necessary also to understand flare energetics and to diagnose flare energy transport.

The energy input from a solar flare has a profound impact on the thermal structure of the chromosphere, and may also influence the photosphere. Different mechanisms of energy input (particle beams, MHD waves, thermal conduction) and dissipation will lead to different temperature, density and velocity structures. This gives, in principle, some power to discriminate between models (Kerr et al. 2016). Spectroscopic observations in multiple lines, and continua, through the solar chromosphere, and at the highest possible spatial and temporal resolution are needed to track the evolution of the flaring chromosphere as it responds to intense energy input. It is already well-known from lightcurves in radio, X-rays and $H\alpha$ that while timescales of 10s of seconds are characteristic of major bursts of energy release in a flare (Grigis & Benz 2005), they also exhibit variability on shorter timescales (Radziszewski et al. 2007). Spatial structures on sub-arcsecond scales have also been detected in flare chromospheric ribbons and photospheric sources (Isobe et al. 2007; Sharykin & Kosovichev 2014). The high spatial and temporal resolution offered by EST is therefore required to probe the elementary energy release, transport and dissipation processes in a flare.

Particularly in larger events, an eruption of CME plasma away from the Sun, driven by unbalanced Lorentz and gravitational forces, accompanies the flare. Analysis of coronal magnetic configurations prone to eruption are currently based on photospheric magnetograms, giving non-force-free boundary conditions for force-free field extrapolations. Measurements of the chromospheric field, which is closer to force-free, will significantly advance our ability to identify the conditions for eruptions, when used in state-of-the-art extrapolations and in data-driven simulations. Also of interest are changes that take place in the photospheric and chromospheric magnetic field in response to the rapid field reconfiguration happening above.

Electric currents are fundamental to flares and CMEs. At large scales, the magnetic field component associated with the current appears to store the free magnetic energy for a flare (Kontogiannis et al. 2017). At very small scales, they provide localised regions for dissipating and releasing the free energy through magnetic reconnection or Joule dissipation (Sharykin & Kosovichev 2014). By providing vector magnetic field measurements at different heights from the photosphere to the chromosphere, EST will for the first time enable the detailed analysis of the 3D properties of electric currents in the lower solar atmosphere, as well as the study of their evolution during solar flares and CMEs.

Observations and simulations strongly suggest that one particular current-carrying structure is often at the heart of a flare and CME - a highly sheared or twisted flux rope. In many cases this supports material in the form of a prominence, which can be analysed spectroscopically to reveal aspects of the distribution of mass and magnetic field. This prominence material can constitute the inner part, or bright core, of a coronal mass ejection (CME), that can have important consequences for Space Weather conditions. Therefore the ability to map out the prominence field prior to the eruption will improve boundary conditions for the evolving field distribution of a CME in space. Similar phenomena are now seen to happen in simulations and observations of much smaller jet-like events as well.

These different aspects of flares and eruptive events demonstrates a need, which will be met by EST, for chromospheric radiation and magnetic diagnostics, at high spatial and temporal resolution, but also with the ability to observe or construct large (active-region scale) fields-of-view on a longer timescale pre-eruption.

6.1 Thermal structure of flare chromosphere and photosphere: line observations

The nominal association between the formation temperature of a particular ion, and height in the atmosphere, as well as the difference in optical depth in line core and wings, provides a means to determine the atmospheric properties as a function of height (e.g. Kuridze et al. 2017), and potentially also time. In a converging chromospheric magnetic field, the physical size of sources emitted by different atoms and ions, and at different wavelengths is also expected to vary with height (Xu et al. 2012a). To address the evolution of the flaring chromosphere requires that we observe in spectral lines formed at different chromospheric heights, with as high a cadence as possible. In the upper chromosphere flows can be strong so a broad spectral window is necessary, while as large as possible a FOV optimises the chance of catching a flare. Only Stokes I is required. These high-cadence programmes should be sequenced with programmes to map the vector field, to identify associations with e.g. electric current density, Lorentz forces, and other 3-D magnetic properties.

OP 6.1.1 Thermal structure of flare: line observations

See table on page 97.

6.2 Thermal structure of flare chromosphere and photosphere: continuum observations

‘White light’ emission is often detectable in highly energetic flare events, but is also a feature of some much smaller flares (Jess et al. 2008) The significance of this broad-band optical emission is that it embodies a very large fraction of the radiated energy (Milligan et al. 2014). The most likely source for flare-enhanced WL emission is recombination emission in the hydrogen Balmer and Paschen continuum (at UV and EUV wavelengths enhancement of the Lyman continuum of H and He is clearly seen). This indicates a central role for ionisation and recombination in the thermodynamics of the flaring chromosphere; the change of state from a neutral to an ionised plasma species is an extremely efficient ‘thermostat’ - until ionisation is complete. The ionisation of both H and He will have significant roles to play in flare thermodynamics, and can be examined by following the continuum emission at different parts of the optical spectrum. Broad-band continuum observations, even at low spectral resolution, will be capable of diagnosing this basic physical process. One way to distinguish between chromospheric and photospheric origins of flare white light emission is to look at differences in the continuum intensity around the H recombination edges. Though the Balmer jump has been detected before (Hiei 1982) it did not resolve this question. The Paschen jump (near 815nm) has been observed only once in flare (Neidig & Wiborg 1984) but should provide a robust observational test for the white-light emission mechanism. The relatively simple measurements proposed here should suffice to finally settle the white-light flare mystery and allow us to use future continuum data for diagnostics of the flaring atmosphere.

However, observations in the optical are always going to be challenging, particularly in small flares, because of the small contrast compared to the photosphere (though this is alleviated if the flare ribbon extends into a sunspot). According to modeling (Simões et al. 2017) extension into the infrared beyond $2\mu\text{m}$ - provides higher contrast than in optical flares, and is synchronised in time with flare energy input to within 0.2s. The primary signature of flare energy - the HXRs that are generated by flare-accelerated electrons, cannot in the foreseeable future be imaged with the spatial and temporal resolution achievable with the EST so observations in the infrared will provide a unique window into flare temporal and spatial evolution.

These high-cadence programmes should be sequenced with programmes to map the vector field, to identify associations with e.g. electric current density, Lorentz forces, and other 3-D magnetic properties.

OP 6.2.1 Thermal structure of flare: continuum observations

See table on page 98.

OP 6.2.2 Thermal structure of flare: detection of the H Paschen Jump

See table on page 99.

6.3 Velocity structure of the flaring atmosphere

If the energy that is input to the lower solar atmosphere cannot be radiated or conducted rapidly away then the plasma will expand. Strongly confined by the magnetic field, the flows will be predominantly along the field. The emission and absorption by upwards and downwards moving material leads to complex line profiles which must be interpreted with care; for example absorption in upflowing material can lead to an apparently red-shifted profile (Kuridze et al. 2015b; Capparelli et al. 2017). Features in $H\alpha$ formed in the upper chromosphere indicate flows reaching tens of km s^{-1} ; The possibility also exists of using multi-line observations, together with overall understanding of spectral line formation gleaned from models, to map out the bulk flow structure of the flare-excited chromosphere.

The presence of plasma turbulence in the flare chromosphere is also to be expected, in association with intense heating, shocks and rapid distortion of the field as it responds to the coronal reconfiguration. Spectral lines emitted in turbulent regions are expected to be broadened either by magnetohydrodynamic “macroturbulence”, or because of out-of-equilibrium distributions of emitting ions, having non-Maxwellian velocity distributions, offering the possibility of examining the development of non-equilibrium velocity distributions evolving on rapid timescales. These high-cadence programmes should be sequenced with programmes to map the vector field, to identify associations with e.g. electric current density, Lorentz forces, and other 3-D magnetic properties.

OP 6.3.1 Velocity structure of the flaring atmosphere

See table on page 100.

6.4 Diagnostics for non-thermal particles

Chromospheric hard X-ray radiation is non-thermal bremsstrahlung, diagnosing flare-accelerated electrons. The dominant model of energy transport places the source of these electrons in the corona, with the electrons arriving in a beam directed along the magnetic field. However HXR spectra interpreted as having two components - a directly-viewed component and an albedo component - are not consistent with an anisotropic distribution. While Coulomb collisions or wave-particle interactions may effectively isotropise the electron distribution, a possible - and long-sought- diagnostic of anisotropic particle distributions is the collisional linear impact polarization generated when appropriate angular momentum sub-states are preferentially excited according to the angular distribution of the exciter particles (Hénoux & Chambe 1990). Claims of observations of linear polarization in lines such as H α remain contentious (e.g. Bianda et al. 2005), as non-simultaneous measurements of Stokes parameters can easily lead to spurious apparent polarization signatures (as, in certain lines, can opacity effects (Judge et al. 2015)). The search for a flare-associated linear polarization signature, as well as measurement of its direction with respect to the solar radial direction (which can also distinguish between electron-generated and proton-generated emission) must therefore be a priority.

OP 6.4.1 Diagnostics for non-thermal particles

See table on page 101.

6.5 Oscillatory phenomena in flares

Recent observations (Nakariakov & Melnikov 2009; Van Doorsselaere et al. 2016) provide indications of quasi-periodic emission during flares, observed in all the electromagnetic spectrum, from radio waves, through H α , WL, EUV, SXR and HXR wavelengths. Among these phenomena, the so called quasi-periodic pulsations (QPPs), which owe their name to well-observed modulations in amplitude and period, can be characterised by periods ranging from fraction of seconds to minutes. However, it is also important to stress that statistical studies have shown that the distribution of the observed periodicities might be affected by the time resolution of the instruments used for their detection (McLaughlin et al. 2018). It is not clear whether QPPs with different periods and modulations are different manifestations of the same physical process or whether the different periods indicate different physical mechanisms. Other questions concern whether QPPs detected in different phases of flares are due to different mechanisms or whether QPPs detected in thermal and non-thermal emission are different.

According to some authors, QPPs characterized by periods ranging from few seconds to several minutes can be attributed to MHD waves; QPPs with periods of fraction of seconds to fast sausage modes, while for longer periods other mechanisms, like slow magnetoacoustic and fast kink mode have been invoked (Foullon et al. 2005; Van Doorsselaere et al. 2011; Su et al. 2012)

Another mechanism that has been taken into account is based on the so-called oscillatory reconnection, occurring in distinct bursts that results in periodic emission when particles are accelerated and the plasma is heated. In this context, there are some studies (e.g. Murray et al. 2009; McLaughlin et al. 2018) based on the assumption that oscillatory reconnection can be caused by magnetic flux emergence in a pre-existing field, indicating that there should be a relationship between the initial submerged magnetic field strength and the observed QPPs periods: longer periods should be related to stronger magnetic fields.

OP 6.5.1 QPPs in different phases of flares

See table on page 102.

OP 6.5.2 QPPs and oscillatory reconnection

See table on page 103.

6.6 Sunquakes

Sunquakes are phenomena characterised by the propagation of acoustic waves that refract deep in the convection zone and appear in photospheric images as ripples, centered around sites hosting M- or X-class flares, and often associated with CMEs (Donea 2011). They have energy between 10^{27} and 10^{29} erg, originate within an area of the order of 10 Mm^2 and show traveling speeds of the order of tens of km s^{-1} . Observations sometimes indicate a co-spatiality of the ripple center of the sunquake with hard X-ray, white light, gamma-ray emission, as well as with sites showing abrupt and permanent changes in the photospheric line-of-sight magnetic field.

The proposed mechanisms causing the sunquakes can be divided into two categories: those based on impulsive heating and those related to the action of the Lorentz force. In the first category the acoustic wave is generated by impulsive heating

that might be related to: i) thick-target heating of the chromosphere by energetic electrons (Kosovichev 2006); ii) heating of the photosphere due to backwarming or deeply penetrating protons (Zharkova & Zharkov 2007); iii) wave heating of the photosphere and chromosphere (Russell & Fletcher 2013). In the second category, the driver is the Lorentz force acting on the photospheric plasma (Fisher et al. 2012). This arises from rapid changes in the photospheric magnetic field observed in many major flares, caused by restructuring in the coronal field. It is often observed during flares associated with sunquakes.

Moreover, another issue concerning sunquakes is related to the fact that the propagation of the acoustic wave in the solar interior seems to be triggered by a process of energy release occurring in the corona, and in order to drive an acoustic disturbance in the solar interior, the energy must propagate through nine pressure scale heights. Therefore, it would also be important to investigate how this energy transfer can take place.

OP 6.6.1 Sunquakes initiated by impulsive heating during flares

See table on page 104.

OP 6.6.2 Sunquakes initiated by changes in the Lorentz force during flares

See table on page 105.

6.7 Large-scale structure and evolution of the magnetic field

It is now established that the energy released during flares (in the form of thermal, kinetic and radiative energy) is previously stored in stressed, non-potential magnetic field configurations until magnetic reconnection takes place. Before the flare occurrence, there is a slow phase of energy storage, which can be due to emergence of new magnetic flux from the subphotospheric layers or to shearing motions that can contribute to a continuous increase of the magnetic energy through build-up of electric currents, on a timescale much longer than the Alfvén timescale in the solar atmosphere. This eventually leads to force-free field configurations, with the coronal magnetic field (at least in ARs) appearing to evolve through a succession of quasi-force-free states

Only in a few cases do the WL images show changes in the sunspot structure after the flare occurrence, generally limited to major (WL) flares. However, it is difficult to single out whether these changes depend on the magnetic field global topology, on the magnetic energy released during the event, on the layer where the magnetic reconnection takes place, or it is mainly related to the spatial resolution of the instruments used.

During the last decade, there has been increasing evidence of rapid, irreversible enhancements of both the longitudinal and horizontal components of the photospheric magnetic field, as well as of magnetic shear at some sections of the magnetic neutral line. In this respect, it would be important to verify whether the reconnection that causes the coronal field reconfiguration leads to similar abrupt, non-reversing changes also in the chromospheric field, in concomitance with the passage of the flare ribbons.

OP 6.7.1 Changes in the magnetic field configuration during/after flares

See table on page 106.

Spectropolarimetric data taken by SDO/HMI used to study the properties of the large-scale vector field, and in particular the evolution of the magnetic helicity flux, indicate that differences in the accumulation of the magnetic helicity flux in the corona can be attributed to the overall magnetic configuration and to the location of flux ropes in active regions. However, differing scenarios exist. Romano et al. (2014b) find that energy injection provided by the shearing motions, pointed out by high values of the shear and the dip angles along the main polarity inversion line, is more significant than the energy injected by magnetic flux emergence. However Liu & Schuck (2012) report that the energy flux associated with magnetic flux emergence contributes to 60% of the total energy, with only 40% arising from shearing. New observations are important to determine the relative importance of these. With appropriate lines chosen throughout the photosphere and chromosphere, it will be possible to assess the helicity flux and accumulation at different heights, to understand how magnetic helicity is transported through the atmosphere.

OP 6.7.2 Determine magnetic helicity accumulation, dip and shear angle in flaring active regions

See table on page 107.

6.8 Filaments in flaring active regions

Filaments are large scale objects embedded in the corona and chromosphere of the Sun. They are located on top of polarity inversion lines (PILs) which separate the positive and negative polarities of the photospheric magnetic field. Especially active region filaments are often associated with very narrow PIL channels which often produce flares. As a consequence, the filament can get ejected into space. However, it is still not understood why some filaments are ejected while others remain stable after the flare. The magnetic environment in such a scenario is extremely difficult to interpret. Sasso et al. (2014) found up to five different magnetic components associated with the flaring filament in the chromosphere. The authors used ground-based spectropolarimetric observations in the He I 10830 spectral range, which is often used to infer the magnetic field in the chromosphere. The derived LOS velocities of the flaring filament were supersonic in both directions (toward and against the solar surface).

OP 6.8.1 Filaments in flaring active regions

See table on page 108.

6.9 Coronal Mass Ejections

During a coronal mass ejection an amount of magnetised plasma of the order of $10^{14} - 10^{16}$ g is expelled from the Sun at velocities ranging between $10^2 - 10^3$ km s⁻¹. The energy involved in these phenomena is of the order of $10^{28} - 10^{32}$ erg.

There are still some difficulties in determining the relationship between flares and CMEs, due to the use of different instruments needed to observe phenomena on the disk and in the outer corona. This circumstance can have different implications, like for instance the fact that there can be a loss of information during the time when the plasma is travelling within the region covered by the occulting disk of the coronagraph. Also, it is not always straightforward to spatially associate a CME with a flare, i.e., to determine the exact location on the solar disk where the eruptive phenomenon started. Another issue arises from the fact that the statistical analyses show that in some cases the flare occurs before the CME, while in others the opposite is true, posing some problems on the initiation models.

OP 6.9.1 CME's sources and temporal relation with flares

See table on page 109.

6.10 Structure and evolution of the magnetic field at small scales

As a complement to the large-scale observation of the magnetic field, high temporal resolution observations of small regions are also required, for at least two purposes. Firstly, simultaneous sampling of magnetically sensitive photospheric and chromospheric spectral lines at high cadence would enable us to study how the magnetic field changes, albeit only in limited areas, during the most dynamic phase of the flare, when the timescale of evolution is seconds. Such observations will also allow a search for signatures of flare-associated reconnection not only in the corona, but also in the lower atmosphere (including component reconnection). Secondly, there are small-scale analogues to the flare process in the form of jets, which have similar magnetic topologies but a much smaller scale, at typically 40'' - 60'', and can involve the ejection of a mini-filament (Raouafi et al. 2016; Moore et al. 2015; Sterling et al. 2016; Wyper et al. 2017). This suggests that these smaller and more readily-observed events share some of the magnetic and MHD properties of larger flare and CME events, and may also have similar chromospheric radiation and non-thermal particle populations. Selected OPs from above can be run with smaller FOVs at higher cadence, and the possibility of capturing an entire event simultaneously.

7 Coupling in partially ionized solar plasma

Authors: Elena Khomenko, Robertus Erdelyi, Mihalis Mathioudakis, Sarah Matthews, Lyndsay Fletcher

The solar photosphere and chromosphere are only partially ionized. The ionization fraction is below 10^{-3} in the photosphere, increasing to about 0.5 in the chromosphere (with 1 being completely ionized plasma). The importance of this fact has not been considered to its full extent in the past in the solar physics community. Only relatively recently, when powerful computing techniques and codes have become accessible, we have started to be in a position to simulate complex partial ionization effects and understand their profound consequences. The influence of partial ionization of the solar plasma on its dynamics has been considered in analytical and numerical models by e.g. Arber et al. (2007); Soler et al. (2010); Zaqarashvili et al. (2012); Leake et al. (2012); Khomenko et al. (2014a); Martínez-Sykora et al. (2012, 2016); Shelyag et al. (2016).

The great majority of the studies of photospheric, chromospheric and coronal plasma dynamics uses MHD as the main tool for quite successfully understanding the complex structure and dynamical processes of these solar atmospheric layers. Nevertheless, the MHD approach overlooks a number of dissipative and dispersive non-ideal mechanisms associated with ion-neutral interactions in weakly ionized and weakly collisional solar plasmas, especially important under chromospheric conditions. In recent few years it has been repeatedly demonstrated that processes related to the non-ideal plasma behaviour due to neutrals may be the key to solve the problem of chromospheric heating, dynamics and fine structure (e.g., Martínez-Sykora et al. 2012; Khomenko & Collados 2012; Martínez-Sykora et al. 2016; Shelyag et al. 2016).

A suitable alternative to the MHD approach is a multi-fluid approach where all the plasma species are considered as separate fluids interacting by collisions. A multi-fluid treatment is essential for the weakly collisionally coupled chromosphere because the relevant processes for energy transport and conversion happen at spatial and temporal scales similar to ion-neutral collisional scales (Ballester et al. 2018). For example, the use of the multi-fluid formalism allows to write an equation for the drift velocity between ions and neutrals (Bittencourt 1986). The existence of these drifts between species is a direct consequence of the partial ionization, and reflects that the coupling between the fluids is not strong enough to ensure a behaviour as a single fluid.

The processes related to partial ionization happen at short spatial and temporal scales. Nevertheless, these scales are not as small as typical plasma scales of the fully ionized plasmas, because they are related to ion-neutral, and not ion-electron collisional scales. The values of these scales depend on the details of the process and the values of the physical parameters. Estimates, and simulations, made for the case of the solar atmosphere in e.g. Khomenko & Collados (2012); Martínez-Sykora et al. (2012); Khomenko et al. (2014a) suggest that temporal resolution of a fraction of a second is necessary to start resolving these effects.

Observations of partial ionization effects lag significantly behind the theoretical developments. Thanks to the powerful combination of instruments, allowing simultaneous observations in several spectral lines, and to the high photon efficiency, the EST will be the only telescope in the world able to push for direct observations of partially ionized plasma processes, allowing our community for the first time to observationally test the limits of classical magneto-hydrodynamics.

7.1 Dynamics of partially ionized prominence plasma

One way of detecting partial ionization effects is the direct measurements of the dynamics of the different components of the plasma. Simultaneous measurements using spectral lines of elements with different degrees of ionization, or different atomic mass, may provide indications of different dynamics of plasma components. The major obstacle for interpreting such measurement is the opacity of the solar plasma. When measured over the solar disc, spectral lines of different elements can form over different height ranges, and therefore provide information of dynamics of different volumes of the plasma. Measurements off-limb may help palliating this problem. The material of solar prominences can frequently be considered optically thin. In addition, the physical conditions in prominences are expected to give rise to significant partial ionization effects, with a considerable amount of neutral atoms. Therefore, different dynamical events in prominence plasmas can be used to test the existence of deviations from classical magneto-hydrodynamics.

Processes related to destabilisation of prominences are not well understood. Studying wave dissipation, instabilities and reconnection in weakly ionised prominence plasmas will allow confirming or rejecting the role of neutrals in destabilisation of prominences.

OP 7.1.1 Waves in prominences observed in neutral and ionized spectral lines

See table on page 110.

Velocities typically measured in solar prominences reveal the existence of important mass flows and waves with different periodicities, as well as instabilities. Instabilities may also lead to mass motions that can have a different impact on neutrals and ions (Soler et al. 2012; Khomenko et al. 2014b). Several recent publications reported simultaneous measurements of prominence dynamics in ionized and neutral spectral lines (Khomenko et al. 2016; Anan et al. 2017). While Khomenko et al. (2016) show a detection of small differences in the ionized Ca II 854.2 nm and in the neutral He I 1083.0 nm velocities in the observed prominence, Anan et al. (2017) concluded that similar differences exist also between the velocities of atoms of the same species emitted by different spectral lines (HI 397 nm, HI 434 nm, Ca II 397 nm, and Ca II 854).

With this campaign the study of ion-neutral effects in prominences will be expanded to the comparison of velocities measured in spectral lines of several neutral and ionized spectral lines strictly simultaneously. These measurements will be accompanied by measurements of the magnetic field vector in order to find possible relations between drift velocities and the magnetic field strength and orientation.

OP 7.1.2 Prominence-corona transition region (PCTR) dynamics

See table on page 111.

The prominence-corona interface is a thin layer separating prominence material from the corona, where the transition from coronal to prominence values in all thermodynamic parameters happen. This region is subject to perturbations, instabilities and turbulence flows. Numerical simulations of Rayleigh-Taylor instability (RTI) at PCTR from Khomenko et al. (2014b) suggest that the drift velocity between ions and neutrals acquire maximum values at this layer. Another conclusion from similar studies (Soler et al. 2012) is that, thanks to the presence of the destabilising effect by neutral atoms, Rayleigh-Taylor instability at PCTR can develop at the smallest scales possible despite the stabilising effect of the magnetic field. The purpose of this campaign is to push the capabilities of EST to observe the dynamics of PCTR at the highest spatial and temporal resolutions possible, using simultaneous measurements in neutral and ionized spectral lines.

OP 7.1.3 Draining of neutral material from prominences and filaments

See table on page 111.

Theoretical work by Gilbert et al. (2002) suggest that cross-field diffusion of neutral material in filaments/ prominences is an important mechanism of their mass loss. As a consequence of incomplete collisional coupling of prominence plasma, neutral helium and hydrogen drains out of prominences. Since the atomic mass of helium and hydrogen is different, the elements have different diffusion speeds, and it takes different times for them to cross the prominence structure. Gilbert et al. (2007) found observational confirmation of this effect, by measuring the relative He/H abundance in filaments across the disc. They found a relative helium deficit in the upper parts of the prominence compared to a relative helium surplus in the lower regions. This was attributed to be a consequence of the large loss timescale for neutral helium compared to neutral hydrogen.

In this campaign we will observe filaments and prominences at different positions of the solar disc with the purpose to study the relative abundance of He, H and Ca II . The emphasis will be on high spatial resolution and large fields of view covering all of the structure. The time resolution is less important since the drainage is a slow process. Magnetic field will be measured simultaneously to find the relation between the relative abundances of He, H and Ca II and the magnetic field structure.

7.2 Influence of partial ionization on spicules

Spicules observed off-limb, as well as their on-disc counterparts are good candidates to show non-MHD effects. Spicules are made of partially ionized chromospheric material. These thin and elongated structures show dynamics on rather short time scales. Theoretical works and numerical simulations suggest that partial ionization effects play an important role in the formation of spicules (De Pontieu & Haerendel 1998; Martínez-Sykora et al. 2017).

OP 7.2.1 Dynamics of spicules observed in neutral and ionized spectral lines

See table on page 111.

This observing program will measure velocities of ionized and neutral species associated with spicules at the highest cadence possible in order to verify how similar they are. Simultaneously, the magnetic field vector in spicules needs to be measured.

OP 7.2.2 Alignment between the magnetic field in fibrils and disc counterparts of spicules

See table on page 112.

One is often making the assumption that linear features seen in observations in spectral lines formed in the chromosphere are aligned with the magnetic field vector. This is expected in the case of ideal MHD where the magnetic field is frozen in and particles can only move along the magnetic field. When ion-neutral effects are taken into account, this is no longer necessarily true. Numerical simulations show that the magnetic field is often not well aligned with chromospheric features (Martínez-Sykora et al. 2016). There are also observations that indicate occasional misalignment between the magnetic field and chromospheric features (de la Cruz Rodríguez & Socas-Navarro 2011). An observing program to address the alignment between the magnetic field and linear chromospheric features needs imaging in a chromospheric line and full vector magnetic field measurements in the chromosphere. High signal to noise is crucial for the accuracy of the magnetic field determination while temporal cadence and field-of-view is less critical.

7.3 Detection of partial ionization effects in the photosphere

OP 7.3.1 Neutral and ionized material Evershed flow in sunspots

See table on page 113.

This campaign aims at detecting ion-neutral effects in the photosphere. Strong magnetic field of sunspots, through the increase of the ion gyro frequency, might lead to stronger decoupling effects. These effects, if detectable, should be present on very small spatial and temporal scales. Recently, Khomenko et al. (2015) have measured the amplitudes of the Evershed flow using pairs of carefully selected Fe I and Fe II spectral lines. They compared azimuthally averaged amplitudes of the Evershed flow extracted from neutral and ion lines and found measurable differences in the radial component of the flow. All five pairs of lines show the same tendency, with a few hundred m/s larger amplitude of the flow measured from Fe I lines compared to Fe II lines. This tendency is preserved at all photospheric heights and radial distances in the penumbra. The origin of this effect is not entirely clear. Spatially and temporally resolved measurements of the Evershed flow, together with measurements of the magnetic field are necessary to conclude about the presence of ion-neutral effects in the photosphere of sunspots.

7.4 Multi-fluid physics of chromospheric waves, shocks and swirls

Waves in the solar atmosphere are usually classified as slow and fast magneto-acoustic waves and Alfvén waves, assuming the complexity of the waveguide allows this simplification. The propagation of magneto-acoustic waves, driven by the combination of total pressure and magnetic tension, is guided by the magnetic field, though other physical quantities may also influence wave propagation. For example, in the upper solar atmosphere the magnetic field may be considered as locally uniform, however, density may vary across the waveguide. Alfvén waves are driven by the magnetic tension and carry energy along the magnetic field. The low ionization fraction of the solar photosphere and chromosphere, and the strong change of the ionization fraction with height strongly affects the propagation of magneto-acoustic waves and Alfvén waves. The motion of the neutrals in the partially ionized atmosphere will not follow the motion of the magnetic flux, questioning the use of atmospheric structures to constrain magnetic field extrapolation methods (Martínez-Sykora et al. 2016). In a partially ionized plasma (even without structuring or stratification) the wave characteristics are different from those of their counterparts in a plasma that is described by the MHD approximation. Such characteristics may be the wave modes themselves, their frequencies or their spatial behavior (e.g. amplitudes, wave eigenfunctions). Theory has already predicted these changes at various degrees (Soler et al. 2013; Mather et al. 2018). The high temporal cadence and spatial resolution of EST measurements will be used to pick up these deviations in the wave characteristics caused by the presence of partial ionization. In general, it is necessary to accurately measure the wave frequencies in individual waveguides, as well as their eigenfunctions (i.e. spatial wavelength), and amplitudes. Once measured, these quantities can be used a diagnostic even for the degree of partial ionization using methods of solar magneto-seismology (Ruderman & Erdélyi 2009).

OP 7.4.1 Observation of magnetic swirls under 2-fluid condition

See table on page 113.

We propose to further investigate photospheric intensity swirls under the two-fluid condition. Statistical results in Liu et al. (2018) have shown that most (> 70%) swirls detected have lifetime less than 6 sec. Extremely high cadence (0.1 sec) will be needed. Spatial resolution plays vital influence in the number and parameters of swirls detected. The swirls are usually located in intergranular lanes. The 2D field of view of IFU is essential in the observation of these features.

7.5 Flares and energetic events

OP 7.5.1 Observations of reconnection and plasmoids in partially ionized plasma

See table on page 114.

It is widely accepted that magnetic reconnection plays a critical role in flares and CMEs, in particular for abruptly and efficiently releasing magnetic energy (and converting it into kinetic and thermal energy), but also as a possible driver of these events. Most of our (incomplete) understanding of magnetic reconnection is based on observations, theory and simulations of reconnection in the fully ionized corona, but it is becoming clear that reconnection in the partially ionized chromosphere is also important, and simulations indicate that partial ionization effects lead to the development of fast reconnection as the result of the onset of the tearing mode instability without the need for anomalous resistivity (e.g., Leake et al. 2012; Singh et al. 2015; Ni et al. 2015). Singh et al. (2015) in particular predict that the reconnection will proceed through different stages based on the level of ionization present, and that in very strong chromospheric field conditions kinetic scales can be reached. While simulations seem to show good agreement with the properties of small-scale chromospheric jets, little work has so far been done to investigate the role and evolution of chromospheric reconnection in solar flares. Observations by Innes et al. (2015) and Rouppe van der Voort et al. (2017) indicate that

both imaging and spectroscopy can be used to infer the presence of plasmoids produced by the onset of the tearing mode instability, but accompanying magnetic field measurements are critical to help confirm the magnetic nature of the islands.

OP 7.5.2 Wave damping by ion-neutral friction as a possible cause of flare chromosphere heating, wight-light flares, and sunquakes

See table on page 115.

According to the thick-target model, the heating of the low atmospheric layers and the consequent chromospheric evaporation during flares is caused by particles that are accelerated at the reconnection site and then travel towards the lower atmosphere. However, there are still some open questions concerning this mechanism, like for instance the large number of particles that should be involved, when compared to coronal densities.

In recent years, taking into account that Alfvén waves produced during reconnection can deliver concentrated Poynting flux to the chromosphere, the role of MHD waves during flares has been investigated (Russell & Fletcher 2013; Reep & Russell 2016). In particular, simulations of Alfvénic waves propagating from the corona to the chromosphere have shown that these waves can be damped due to ion-neutral friction and that the energy lost by these waves can heat the chromosphere and contribute to the evaporation of the chromospheric plasma. The damping of the Alfvén waves in the chromosphere strongly depends on the wave frequency and for periods of 1 s or fractions of s, an amount of energy in the range 37 % - 100 % that enters the chromosphere can be damped by ion-neutral friction. Moreover, if coronal waves are trapped in closed coronal structures, the total transmission results to be higher, due to multiple incidences.

These simulations also indicate that the propagation of Alfvén waves during flares could be important to explain WL flares (they in fact provide a direct mechanism for heating the temperature minimum region, where the bulk of white light flare emission is formed) and sunquakes (waves with periods of several seconds can enter the chromosphere and pass undamped into the solar interior).

Because the atmospheric response is nearly identical, the heating due to Alfvén waves can appear extremely similar to that caused by electron beams. Therefore up to now it has not been possible to distinguish observationally these two mechanisms.

8 Scattering physics and Hanle-Zeeman diagnostics

Authors: Luca Belluzzi, Alex Feller, Javier Trujillo Bueno, Rafael Manso Sainz

The great diagnostic potential of scattering polarization and the Hanle effect is today widely recognized. During the last decades, a series of novel diagnostic methods based on these physical mechanisms have been developed and successfully applied, especially for the investigation of the magnetism of the solar atmosphere in domains that are not accessible through the standard Zeeman-effect techniques (Trujillo Bueno et al. 2006, 2017; Casini & Landi Degl’Innocenti 2008; Trujillo Bueno 2009, 2010; Casini et al. 2017).

The birth and rapid growth of this new research field has certainly been possible thanks to the recent technological advances, which allowed the development of new polarimeters with increasingly high sensitivities (presently up to 10^{-5} in the degree of polarization). On the other hand, it is clear that in order to fully exploit the potentialities of these new diagnostic methods, a deep and solid understanding of the physics of scattering polarization is necessary. Unfortunately, the theoretical modeling of scattering polarization has turned out to be a very complicated task. A solid description of matter-radiation interaction within the framework of Quantum Electrodynamics is actually necessary, a series of atomic physics aspects need to be taken into account, and efficient numerical techniques for the solution of the radiative transfer (RT) problem in non-local thermodynamic equilibrium (NLTE) conditions, in the presence of polarization phenomena (NLTE problem of the 2nd kind), need to be applied.

The clearest manifestation of scattering polarization is certainly the so-called Second Solar Spectrum, namely the linearly polarized spectrum of the solar radiation coming from quiet regions close to the limb (Stenflo & Keller 1997; Gandorfer 2000, 2002, 2005). This spectrum is rich of signals and spectral details of double scientific interest: on the one hand, they encode a wealth of information about the physical conditions present in the solar atmosphere. On the other hand, the physical origin of several of these signals remains unclear, and they thus represent precious observational signatures (often not reproducible in laboratory plasmas) of physical phenomena that still need to be completely understood. The Second Solar Spectrum is in fact a precious window also for improving our understanding of the physics of scattering polarization, and since many years it represents a key test-bench for the theories of the generation and transfer of polarized radiation. The observation of many details of the Second Solar Spectrum requires rather high polarimetric sensitivities, which today can generally be achieved only by completely sacrificing the spatial and temporal resolution. With a 4 m aperture telescope like EST, it will finally be possible to observe the spectral details of the Second Solar Spectrum with an unprecedented spatial and temporal resolution, thus further increasing the potentialities of this spectrum as a diagnostic tool, and possibly revealing new aspects of the physics of scattering.

In this chapter, we present a series of observing programs (OPs) focused on a selection of spectral lines that produce scattering polarization signals of particular interest, either because of their diagnostic potential for the investigation of the properties of the solar atmosphere, or because their physical origin is still unclear. The OPs are specifically designed so as to exploit the specific advantages of the EST, in particular the combination of high polarimetric sensitivity and high spatio-temporal resolution. The OPs are arranged on the basis of their main scientific goal as follows:

1. *Investigation of the small-scale magnetism of the quiet solar photosphere via the Hanle effect in atomic and molecular lines.*
 - OP 8.1.1: Spatial fluctuations of scattering polarization in Sr I 4607 Å
 - OP 8.1.2: Spatial fluctuations of scattering polarization in C₂ molecular lines around 5140 Å
 - OP 8.1.3: Spatial fluctuations of scattering polarization in the Ti I multiplet around 4530 Å
 - OP 8.1.4: Simultaneous observations in Sr I 4607 Å and C₂ around 5140 Å
2. *Investigation of the magnetism of the chromosphere via the combined action of Hanle, Zeeman, and magneto-optical effects in strong resonance lines.*
 - OP 8.2.1: Ca I 4227 Å resonance line
 - OP 8.2.2: Ca II K and H (3934, 3968 Å) resonance lines
 - OP 8.2.3: Ca II IR triplet (8498, 8542, 8662 Å)
3. *OPs of interest for deepening our understanding of the physics of scattering polarization*
 - OP 8.3.1: Na I D₂ and D₁ (5890, 5896 Å) lines

Some of the above-mentioned scientific goals have already been addressed in previous sections of this document. The reason for gathering in a separate section all the OPs focused on scattering polarization signals is that they involve common dedicated observational and instrumental requirements. These include:

- increased polarimetric sensitivity (generally higher than what is needed for Zeeman measurements);
- interest to perform observations closer to the limb (or even off-limb), where the use of adaptive optics (AO) is generally more difficult due to the reduced continuum intensity contrast;
- observations in particular spectral lines and molecular bands, which are normally not used for Zeeman diagnostics;
- interest for the blue and near-UV region of the solar spectrum, where the scattering amplitudes are typically larger;
- simultaneous observation of lines with different sensitivities to the Hanle effect (differential Hanle effect techniques).

8.1 Investigation of the small-scale magnetism of the quiet solar photosphere

The quiet solar photosphere is permeated by small-scale (< 1 arcsec) magnetic fields, which interact strongly with the turbulent convection and typically evolve on a short time scale of tens of seconds, like the solar granulation. The individual quiet Sun magnetic structures are small in extent, compared to active regions, but ubiquitous on the Sun, and are therefore thought to play, in their entirety, a major role in solar activity and in the energetic coupling of the various height layers of the solar atmosphere. Some fundamental questions are related to these fields, like the existence and influence of a small-scale dynamo driven by the turbulent motions, or the question about the total amount of magnetic flux on the Sun. Multiple approaches can be taken to answer these questions making use of the fact that the Zeeman and Hanle effects probe magnetic fields in a complementary way. The Hanle effect, in particular, being sensitive to tangled magnetic fields showing opposite polarities below the resolution element (to which the Zeeman effect is practically blind), is a key tool for investigating this aspect of the solar magnetism (Stenflo 1982; Faurobert-Scholl et al. 1995; Trujillo Bueno et al. 2004).

One of the main difficulties of Hanle effect diagnostics is that the amplitude of a given scattering polarization signal depends on several factors, the most important one being the symmetry properties of the pumping radiation field. Such symmetry properties, on the other hand, strongly depend on the thermal and dynamic structure of the solar atmosphere, and they are very sensitive to the presence of horizontal inhomogeneities in the solar atmospheric plasma. As this kind of information is not known a priori, it is necessary to develop suitable techniques to disentangle the impact of the magnetic field from that of other “symmetry-breaking” causes.

There are basically two strategies to overcome this difficulty. The first one is to calculate theoretically the amplitude of a given scattering polarization signal by solving the radiative transfer problem, both taking into account and neglecting

the magnetic field, using three-dimensional (3D) numerical models of the solar atmosphere (Trujillo Bueno et al. 2004). The computational cost of 3D radiative transfer investigations is very significant, but they allow us to clearly identify and disentangle the impact of the various physical mechanisms affecting the observed scattering polarization, and in particular to obtain the zero-field reference signal. This method is necessarily model-dependent, but thanks to the availability of increasingly detailed and realistic simulations of the solar atmosphere, it is today a reliable approach. One of the most suitable and exploited spectral lines for this kind of diagnostics is the Sr I 4607 Å line. An OP focused on this line is described in Sect. 8.1.1.

A second approach is the so-called *differential Hanle effect technique* (Stenflo et al. 1998; Trujillo Bueno 2003). This approach consists in considering two (or more) spectral lines having similar formation properties but different sensitivities to the Hanle effect, and to consider amplitude ratios of the corresponding scattering polarization signals. Indeed, if the lines form under the same conditions, the dependence on several critical parameters cancels out in the ratios, and these latter can be used to directly infer information on the magnetic field, without the need of a theoretical zero-field reference signal. Spectral lines that are particularly suitable for the application of this technique are those produced by molecules, such as C₂ and MgH (Trujillo Bueno 2003). These are the main advantages:

- molecular lines are generally weak in the intensity spectrum and their radiative transfer modeling is relatively simple (which simplifies the application of the differential Hanle effect technique);
- molecular lines pertaining to the same branch have very similar formation properties and fall within a rather small frequency interval (they can be easily observed simultaneously);
- molecular lines pertaining to the same branch may have significantly different Landé factors, and therefore very different Hanle effect sensitivities;
- interestingly, several molecular lines produce clear scattering polarization signals in the Second Solar Spectrum.

In the past decades, differential Hanle effect diagnostics has been carried out exploiting various sets of molecular lines. Particular attention has been paid to lines of C₂ (Trujillo Bueno 2003; Trujillo Bueno et al. 2004; Berdyugina & Fluri 2004; Kleint et al. 2010, 2011), MgH (Asensio Ramos & Trujillo Bueno 2005), and CN (Shapiro et al. 2011). An OP focused on the C₂ lines at 5140 Å is described in Sect. 8.1.2

The differential Hanle effect technique can of course be applied also to atomic lines. The main disadvantage with respect to molecular lines is that atomic lines are generally stronger and their radiative transfer modeling is more complicated. A particularly nice example of a set of atomic lines suitable for differential Hanle effect diagnostics is a titanium multiplet at around 4530 Å (Manso Sainz et al. 2004). An OP dedicated to this multiplet is described in Sect. 8.1.3.

The possibility of performing differential Hanle effect diagnostics has been generally limited by the necessity of considering lines falling within the single and relatively narrow spectral windows of today's solar spectrographs. The new multi-wavelength capabilities of the EST instrumentation will significantly extend the wavelength range for differential diagnostics and thus increase the potentialities of this technique.

Another important aspect that has to be taken into account when applying Hanle effect diagnostics is the impact of depolarizing collisions with neutral perturbers (hydrogen and helium atoms). Such collisions sensibly affect the amplitude of the scattering polarization signals, and must be carefully taken into account. This is especially true in the lower layers of the solar atmosphere, where the plasma density is higher. Detailed theoretical calculations of the collisional rates for the Sr I 4607 Å line are today available (e.g. Faurobert-Scholl et al. 1995; Manso Sainz et al. 2014). The availability of precise estimates of these rates is crucial because relatively small variations of these quantities can have a clear impact on the amplitude of the synthetic scattering polarization profiles, and therefore on the inferred magnetic field (see del Pino Alemán et al. 2018). Unfortunately, the Sr I line is a kind of exception, and in most cases only very rough estimates of the depolarizing rates are available. Differential Hanle diagnostics can be suitably exploited to reduce the uncertainties related to collisional depolarization, especially concerning the number density of perturbers. This quantity, which is not known a priori, cancels out when combining scattering polarization signals observed in different spectral lines with similar formation properties. Such a multi-line observing technique applied with the EST will thus allow to reduce the uncertainties related to the interpretation of Hanle depolarization.

The Hanle effect has been successfully applied in the past for investigating the small-scale magnetism of the solar photosphere (e.g., Trujillo Bueno et al. 2006, and references therein). However, all such works were based on observing data with very low spatial and temporal resolution. On the other hand, sizable variations of the amplitude of the scattering polarization signals of photospheric lines are expected at sub-granular spatial scales (see Trujillo Bueno & Shchukina 2007; del Pino Alemán et al. 2018). Detecting such fluctuations would allow us to deepen our understanding of the various mechanisms affecting scattering polarization and, at the same time, to study the small-scale magnetic fields of the intergranular plasma. This is a typical goal for a 4m-class telescope like EST, and it is the basic objective of the OPs described in the following.

OP 8.1.1 Spatial fluctuations of scattering polarization in Sr I 4607 Å

See table on page 116.

The goal of this observing program is to study spatial fluctuations of the scattering polarization signal of the Sr I 4607.4 Å line, on subgranular spatial scales.

In observations with low spatio-temporal resolution, the Sr I line shows a relatively large scattering polarization amplitude (above 1% at $\mu = 0.1$, e.g. Stenflo & Keller 1997; Gandorfer 2002). Further, theoretical predictions (Trujillo Bueno & Shchukina 2007; del Pino Alemán et al. 2018) anticipate spatial fluctuations of the polarization signal in the order of several 0.1%, depending on limb distance and resolution of the instrument. The latter has a large effect on the signal amplitudes and so far observations are just at the limit of the critical combination of high spatial resolution and increased polarimetric sensitivity (e.g., Bianda et al. 2018; Zeuner, F. et al. 2018). The EST with its large telescope aperture will overcome those limitations and is therefore ideally suitable for this type of measurements. A key concept of this observing program is to refrain from strictly diffraction limited observations but to choose an optimum slightly larger spatial and temporal sampling, in order to reach the required signal-to-noise ratio in the best possible way, in the presence of solar evolution (cf. Appendix A.1).

OP 8.1.2 Spatial fluctuations of scattering polarization in C₂ molecular lines at 5140 Å

See table on page 116.

Molecular lines of particular interest for differential Hanle effect diagnostics are the C₂ lines of the Swan system ($d^3\Pi_u - a^3\Pi_g$), with their R and P branches (Trujillo Bueno 2003; Trujillo Bueno et al. 2006). Of particular interest are those around 5140 Å, proposed by Berdyugina & Fluri (2004). In particular, the three lines of the R branch are perfectly resolved and free of blends with other lines. They produce weak (10^{-3}) but clear signals in the Second Solar Spectrum (see Gandorfer 2002), and are characterized by the following critical fields for the onset of the Hanle effect (hereafter Hanle critical field): $B_H \approx 4$ G for the R₁ ($J = 14$) and R₃ ($J = 12$) lines and $B_H \approx 40$ G for the R₂ ($J = 13$) line. Their drawback with respect to the C₂ lines with $J > 20$ used by other authors, namely that a line with $B_H \approx 40$ G cannot really be considered as an ideal reference line given the significant magnetization of the quiet Sun photosphere, is compensated for by the fact that the R₂ ($J = 13$) and R₁ ($J = 14$) lines are not blended. These lines have been extensively used for synoptic programs aimed at monitoring possible long-term variations of weak photospheric magnetic fields with the solar cycle (Kleint et al. 2010, 2011). For the most recent results see Ramelli et al. (2019, in press).

Interestingly, the average magnetic field intensity inferred by means of C₂ molecular lines is sensibly smaller than that revealed by the Hanle effect in the Sr I 4607 Å line. A possible solution of this apparent disagreement was proposed by Trujillo Bueno (2003), who pointed out from numerical simulations that the scattering polarization signals produced by weak molecular lines should mainly come from the upflowing granular cell centers, while both granular and intergranular regions should contribute to the signal of the Sr I line (see also Trujillo Bueno et al. 2004). This further stresses the interest of observing scattering polarization in molecular and atomic lines to detect magnetic fields below the spatial resolution limit of the EST, where Zeeman diagnostics is known to suffer from potential signal cancellation. Apart from this particular aspect, the general detection of fluctuations in the scattering polarization signals produced by the above-mentioned C₂ lines at 5140 Å is the main goal of this OP.

OP 8.1.3 Spatial fluctuations of scattering polarization in the Ti I multiplet around 4530 Å

See table on page 117.

Some of the most remarkable features of the Second Solar Spectrum are the conspicuous spectral patterns produced by relatively low abundance atomic species (rare earths, transition metals — other than Fe) and molecules. This is important because it offers the possibility of studying tangled magnetic fields through the Hanle effect, at the lowest layers of the photosphere (e.g., Ce II forms barely above the continuum), just where magnetoconvection simulations can be best constrained.

Ti I is an important case in point; there are about 20 lines in the visible showing relatively large scattering polarization well above the continuum level. Several of them correspond to the singlet system of Ti I. They show some of the largest signals and are also relatively simple to model, which simplifies their analysis and the possibility of inversion. On the other hand, there are many multiplets from the quintet system. These are interesting mainly because lines of the same multiplet can be modeled similarly and consistently, which offers the possibility of differential analysis among different lines. Further, some of these multiplets lie on a relatively narrow spectral region, which allows observing if not all, at least many of them simultaneously. Probably, the most interesting case in this regard is the multiplet $a^5F - y^5F^0$ around 453 nm, which has a line (453.6 nm) which is completely unaffected by magnetic fields (the Landé factor of both levels is zero), and serves as a reference for Hanle effect depolarization, while 8 lines of the same multiplet lie within less than 1 nm, which allows observing all of them simultaneously (see Manso Sainz et al. 2004).

OP 8.1.4 Simultaneous observations in Sr I and C₂

The OPs described above are designed for observations in an individual line (Sr I) or in groups of lines located within a given spectrograph FOV (C₂, Ti I). The advantage of individual observations is the higher throughput of the instrument. For some science cases involving differential diagnostics we also want to perform observations in the C₂ molecular band and in Sr I simultaneously. In this case the requirements in terms of spectral resolution, FOV and sensitivity remain the same as in the case of individual observations. We accept a degradation of the spatio-temporal resolution though, as an inevitable consequence of the reduced instrument throughput. According to the EST photon budget the throughput will be decreased by about a factor 4 compared to individual observations, which results in a degradation of the optimum spatial sampling and cadence by about a factor $4^{1/3} \approx 1.6$ respectively (cf. Appendix A.1). The coarser sampling will be achieved by a post-facto numerical binning of the data and thus does not impose any additional requirements on the instrumentation.

8.2 Investigation of the magnetism of the solar chromosphere

The information on the magnetic field of the solar chromosphere is encoded in the polarization of the spectral line radiation that originates in a highly inhomogeneous and dynamic atmospheric region, where hydrodynamic and magnetic forces compete for dominance. Relatively few spectral lines whose core originates in the chromosphere can be observed with ground-based telescopes. Here we focus on the strongest lines of Ca I and Ca II and, in particular, on the diagnostic potential of their polarization signals produced by the joint action of scattering processes and the Hanle and Zeeman effects.

As mentioned above, the rarefied plasma of the solar chromosphere is very dynamic and highly inhomogeneous, and the magnetic field is thought to play a key role on the formation, evolution, and destruction of the observed plasma structures. We are still far from understanding the physics of the solar chromosphere, and precisely for this reason we need EST. This telescope will provide novel observations of the intensity and polarization of solar spectral lines, with unprecedented spatial, temporal, and spectral resolution. By confronting the Stokes profiles observed in chromospheric lines with those calculated in increasingly realistic numerical models of the solar chromosphere we may hope to achieve new breakthroughs in solar physics. By the time EST will be operative, such three-dimensional (3D) numerical models of the solar chromosphere will be much more realistic than nowadays, given that they will include a number of possibly key physical ingredients such as non-equilibrium ionization and ambipolar diffusion (interactions between ions and neutrals in the presence of magnetic fields). Likewise, ongoing synergistic efforts suggest that by the time EST will be available, we will have completed the development of a computer program capable of solving with massively parallel computers the (NLTE) problem of the generation and transfer of polarized radiation in such 3D models of the solar atmosphere, taking into account frequency correlation effects between the incoming and outgoing photons in the scattering events, as well as the joint action of the Hanle and Zeeman effects. Obviously, in order to validate or discard such numerical models we need to confront measured and calculated observable quantities sensitive to the thermal, dynamic, and magnetic time-dependent structure of the solar chromosphere (i.e., the Stokes profiles of chromospheric spectral lines). Comparing only the intensity of the spectral line radiation is not sufficient, since the Stokes *I* profile of spectral lines is practically insensitive to the strength and structure of the magnetic field.

OP 8.2.1 The Ca I 4227 Å resonance line

See table on page 117.

Outside sunspots, this spectral line shows very sizable linear polarization signals produced by scattering processes, of the order of a few percent when observing close to the limb (e.g., Gandorfer 2002) and of the order of 0.1% when observing moderately magnetized regions at the solar disk center (e.g., Bianda et al. 2011). Therefore, it is a very suitable line for observing with EST its scattering polarization with unprecedented spatial and temporal resolution.

A correct modeling of the linear polarization produced by scattering processes in this line requires taking into account the effects of correlations between the frequencies of the incoming and outgoing photons in the scattering events. For lines of sight (LOS) pointing to quiet regions located away from the solar disk center, such partial frequency redistribution (PRD) effects produce in the Ca I 4227 Å line a (typically) triple-peak fractional linear polarization pattern with a central peak and sizable lobes in the blue and red wings of the line.

In semi-empirical models of the solar atmosphere the height where the line-center optical depth is unity varies between about 900 km (disk center LOS) and 1200 km (close to the limb LOS), while the radiation of the extended wings of this line stems from increasingly deeper layers when going from the line center to the far wings (e.g., at ± 0.5 Å from line center the height where the ensuing optical depth is unity is about 200 km for a close to the limb line of sight).

Only one of the stable isotopes of calcium has non-zero nuclear spin, but its abundance is only 0.135%. Therefore, we can assume that hyperfine structure is a negligible physical ingredient for modeling the scattering polarization of calcium lines. The Ca I 4227 Å resonance line results from a transition with a lower level with angular momentum $J_l = 0$ and

and upper level with $J_u = 1$, and the Hanle critical field in this line is $B_H = 25$ G. Fortunately, a two-level model atom is a suitable approximation for modeling its polarization. This is good news, especially because the radiative transfer numerical modeling of lines for which PRD phenomena are important is complex and time-consuming. It is also good news that depolarization by elastic collisions with neutral hydrogen atoms is practically negligible.

Recent theoretical developments have shown that the polarization of this line, whose core originates in the lower chromosphere, is due to the following physical mechanisms (Alsina Ballester et al. 2018):

- the familiar Zeeman effect, which can produce measurable circular polarization for longitudinal field components of at least 10 G.
- the linear polarization caused by scattering processes in the line-core, along with its modification by the Hanle effect (which operates in the line core). Therefore, this line-core polarization is sensitive to the magnetism of the lower solar chromosphere.
- the linear polarization caused by scattering processes and PRD effects in the line wings, along with its magnetic sensitivity through the ρ_V magneto-optical (MO) terms of the Stokes-vector transfer equation. Such MO terms produce sizable U/I wing signals as well as a sensitivity of both the Q/I and U/I wings to the presence of magnetic fields in the region of formation of the line wings (the photosphere). Magnetic fields with strengths as low as 10 G are sufficient to produce a measurable impact on the wings of both Q/I and U/I .

In summary, the Ca I 4227 Å resonance line is of high scientific interest for EST because

- scattering processes produce very significant linear polarization signals, which are sensitive to the magnetization of the lower solar chromosphere (via the Hanle effect in the line core) and of the underlying photosphere (via the joint action of PRD and MO effects). Magnetic fields as weak as 10 G are sufficient to have a measurable impact on such linear polarization signals.
- the confrontation of the Stokes profiles observed by EST with those calculated by solving the problem of the generation and transfer of polarized radiation in 3D numerical models of the solar atmosphere, will allow us to probe the thermal, dynamic, and magnetic structure of the lower solar chromosphere and its coupling with the underlying layers.
- with the telescopes that are presently available spectropolarimetric observations of this line have always required to seriously sacrifice the spatio-temporal resolution.

OP 8.2.2 The Ca II H & K resonance lines

See table on page 118.

The Ca II H & K resonance lines at 3934 Å and 3969 Å are the strongest chromospheric lines that can be observed from ground-based facilities. In fact, in quiet regions of the solar atmosphere most of the calcium atoms are in the form of Ca II. In models of the quiet solar atmosphere the height where the line-center optical depth is unity is only about 200–300 km below the corrugated surface that delineates the chromosphere-corona transition region. Therefore, especially for close to the limb LOS the line-center radiation of these resonance lines stems mainly from the upper chromosphere.

The PRD effects are essential for understanding the Stokes profiles of these resonance lines. A rigorous modeling of their spectral line radiation requires using at least a 5-level model atom, with the H & K resonance lines and the IR triplet of Ca II. An additional important physical ingredient is quantum mechanical interference between the sublevels pertaining to the upper level of the K line, which has angular momentum $J_u = 3/2$, and those of the upper level of the H line whose $J_u = 1/2$. In spite of the fact that the H and K lines are separated by 35 Å, in the (optically thick) plasma of the solar atmosphere such J -state interference effects produce observable signatures in the scattering fractional linear polarization pattern, especially in their wings (Stenflo 1980). In semi-empirical models of the quiet solar atmosphere the Q/I pattern shows positive peaks in the blue wing of the K line and in the red wing of the H line (of the order of 1%, but with the K blue peak larger than the H red peak) and a negative peak between the K and H lines, in qualitative agreement with the observations of Stenflo (1980). Such wing signals are caused by the joint action of PRD and J -state interference, exactly as it occurs with the theoretical Q/I pattern of the Mg II h & k lines (see Belluzzi & Trujillo Bueno 2012).

In these resonance lines, the Hanle effect operates only in the core of the K line (critical Hanle field $B_H \approx 12$ G). On the other hand, given that the joint action of PRD and J -state interference produce sizable signals in the wings of the Q/I profiles, the MO terms of the Stokes-vector transfer equation already introduced in section 8.2.1 can produce significant signals in the wings of the U/I pattern and an interesting magnetic sensitivity in the wings of both Q/I and U/I (del Pino Alemán 2019; private communication), exactly as it happens with the wings of the Mg II k line (Alsina Ballester et al. 2016), with the wings of the Mg II h & k lines (del Pino Alemán et al. 2016), with the wings of the Ca I 4227 Å line (Alsina Ballester et al. 2018), and with the wings of the hydrogen Ly- α line (Alsina Ballester et al. 2019).

The photons of the far wings of the H & K lines stem from the solar photosphere, and via the MO effects photospheric fields as low as 20 G can therefore have a significant impact in the wings of Q/I and U/I . The same happens with the scattering polarization wings of the Mg II h & k lines (Alsina Ballester et al. 2016; del Pino Alemán et al. 2016). Finally, it should be noted that the circular polarization is dominated by the familiar Zeeman effect, which can produce measurable signals for longitudinal field components of at least 10 G.

In summary, the Ca II resonance lines are of high scientific interest for EST because

- Their line-core radiation stems from the upper solar chromosphere, only about 200–300 km below the corrugated surface that delineates the chromosphere-corona transition region.
- The line-core of the K line is sensitive to the Hanle effect. Therefore, it reacts to the presence of magnetic fields in the upper solar chromosphere. The critical Hanle field of the K line is 12 G, so we may expect good sensitivity to field strengths between 2 and 20 G, approximately.
- The wings of the fractional linear polarization profiles of the H and K lines are sensitive to the presence of magnetic fields as low as 20 G. This magnetic sensitivity in the wings of the Q/I and U/I profiles, which extends all through the solar atmosphere, is caused by the MO terms $\rho_V Q$ and $\rho_V U$ of the transfer equations for Stokes Q and U .
- the familiar Zeeman effect can produce measurable circular polarization if the longitudinal field component is sufficiently large (e.g., for a longitudinal field of 10 G the Stokes- V amplitude of the Ca II H line is about 0.2%).
- with the telescopes that are presently available spectropolarimetric observations of these lines have always required to dramatically sacrifice the spatio-temporal resolution.

OP 8.2.3 The Ca II IR triplet

See table on page 118.

The strongest line of the Ca II IR triplet is the 8542 Å line, followed by the 8662 Å and 8498 Å lines. Together, they encode information over a significant range of heights in the middle solar chromosphere. Manso Sainz & Trujillo Bueno (2003) demonstrated quantitatively that the physical origin of the enigmatic scattering polarization observed by Stenflo et al. (2000) in the solar disk radiation of the Ca II lines at 8662 Å and 8542 Å is “zero-field dichroism” (i.e., differential absorption of polarization components caused by the presence of a significant amount of atomic polarization in their metastable lower levels). The scattering polarization of the 8498 Å line, the weakest of the triplet, has contributions from the atomic polarization in its upper and lower levels.

The magnetic sensitivity of the scattering polarization in the Ca II IR triplet has been theoretically investigated by Manso Sainz & Trujillo Bueno (2010). They showed that the linear polarization profiles produced by scattering in the Ca II IR triplet have thermal and magnetic sensitivities potentially of great diagnostic value. The scattering polarization in the 8498 Å line shows a strong sensitivity to inclined magnetic fields with strengths between 0.001 and 10 G, while the scattering polarization in the 8542 Å and 8662 Å lines is mainly sensitive to magnetic fields with strengths between 0.001 and 0.1 G. The reason for this peculiar behavior is that the scattering polarization of the 8662 Å and 8542 Å lines, unlike the Ca II 8498 Å line, is controlled mainly by the lower level Hanle effect. Therefore, in regions with magnetic strengths sensibly larger than 1 G, their Stokes Q and U profiles are sensitive only to the orientation of the magnetic field vector. Manso Sainz & Trujillo Bueno (2010) also found that the sign of the emergent Stokes Q/I and U/I profiles of the 8662 Å and 8542 Å lines is rather insensitive to the chromospheric thermal structure, while the sign of the linear polarization profiles of the 8498 Å line turns out to be very sensitive to the thermal structure of the lower chromosphere. They concluded that spectropolarimetric observations providing information on the relative scattering polarization amplitudes of the Ca II IR triplet could be very useful to improve our empirical understanding of the thermal and magnetic structure of the quiet chromosphere. The required radiative transfer modeling for reliably inferring information from confrontations with future high-spatial resolution spectropolarimetric observations is however more complex. On the one hand, the line-core of the Ca II IR triplet originates in a shock-dominated region of the solar chromosphere and the ensuing macroscopic velocity gradients may have a significant impact on the anisotropy of the spectral line radiation and, therefore, on the shape and amplitude of the observed scattering line polarization (Carlin et al. 2012). On the other hand, the breaking of the axial symmetry of the pumping radiation field caused by the three-dimensional thermal, dynamical and magnetic structure of the solar chromosphere has to be taken into account (Štěpán & Trujillo Bueno 2016).

In general, the emergent Q/I and U/I profiles are produced by the joint action of atomic level polarization, and the Hanle and Zeeman effects. Atomic polarization and the Hanle effect dominate the emergent linear polarization profiles for inclined magnetic fields with strengths weaker than B_0 , where the B_0 value depends on the scattering geometry. In the forward scattering geometry of a disk center observation ($\mu = 1$), the linear polarization of the Ca II IR triplet is dominated by the Hanle effect if the magnetic field is weaker than about 10 G. In fact, while the contribution of the Zeeman effect to the linear polarization is negligible for $0 < B < 10$ G the Hanle effect creates weak but measurable fractional linear polarization signals already for horizontal magnetic fields as low as 0.1 G (Manso Sainz & Trujillo Bueno 2010; Štěpán

& Trujillo Bueno 2016). For magnetic strengths $10 \lesssim B \lesssim 100$ G the contribution of the transverse Zeeman effect to the linear polarization observed at the solar disk center should not be neglected. Detection of Q/I and U/I disk center signals caused either by the Hanle effect alone (if $B < 10$ G) or by the joint action of the Hanle and transverse Zeeman effects (if $10 \lesssim B \lesssim 100$ G) requires very high polarimetric sensitivity together with a spatial and temporal resolution sufficient to resolve the magnetic field azimuth. Finally, it is important to note that the circular polarization of the Ca II IR triplet is dominated by the Zeeman effect, and that the weak field approximation that relates the Stokes V profile with the wavelength derivative of the Stokes I profile can provide a reasonable estimation of the longitudinal component of the magnetic field vector (e.g., Centeno 2018).

8.3 Deepening our understanding of the physics of scattering polarization

OP 8.3.1 The physics and diagnostic potential of the Na I D₁ and D₂ lines

See table on page 119.

The theoretical interpretation of the rich variety of signals of the Second Solar Spectrum has played a key role in the development of new theoretical approaches for the description of scattering polarization. In particular, the interpretation of the signals produced by the Na I D₁ and D₂ lines at 5895.9 Å and 5890.0 Å respectively, has been challenging scientists for more than twenty years. Indeed, these signals show the signatures of several different physical mechanisms, and not by chance they have always represented a key test bench for the theory.

The Na I D₁ and D₂ lines originate from the transitions between the ground level of sodium, $^2S_{1/2}$, and the upper levels $^2P_{1/2}^o$ and $^2P_{3/2}^o$, respectively. Sodium has a single stable isotope, ^{23}Na , which has hyperfine structure (HFS; nuclear spin $I = 3/2$). The Na I D-lines are among the strongest spectral lines of the visible solar spectrum, and encode information on the physical properties of the solar atmosphere ranging from the low chromosphere to the photosphere. In semi-empirical models of the solar atmosphere, the line-core of these lines forms at about 900 km for an observation at $\mu = 1$, and at about 1200 km for an observation at $\mu = 0.1$.

In the Second Solar Spectrum, the D₂ line shows a peculiar triplet-peak Q/I profile. At $\mu = 0.1$, the central peak reaches an amplitude of about 0.35%, thus representing one of the largest signals of the Second Solar Spectrum (see Gandorfer 2000). The triplet-peak structure of this signal is ultimately due to frequency correlation effects between the incoming and outgoing photons in the scattering processes: PRD effects thus need to be taken into account for modeling this signal, as well as the depolarizing effect of HFS. The central peak is sensitive to the Hanle effect, the critical field being of the order of 5 G.

Besides the triplet-peak structure profile of the D₂ line, the scattering polarization signal produced by the sodium doublet shows an overall pattern across the two lines, with a sign reversal between D₁ and D₂, and an anti-symmetric structure across D₁, with a negative dip in the blue wing, and a positive bump in the red wing (Gandorfer 2000). This overall pattern is the result of quantum interference between the upper levels of D₁ and D₂ (J -state interference), and it is fully analogous to the one observed across the H and K lines of Ca II. Notably, these effects take place in the far wings of the spectral lines, where the emissivity is extremely low. This explains why the signatures of J -state interference cannot be observed in laboratory experiments, but only on the Sun, where the low emissivity is compensated by the optical thickness of the solar plasma.

Stenflo & Keller (1997) first observed a conspicuous scattering polarization signal also in the core of D₁ ($Q/I \approx 0.15\%$ at 5 arcsec from the limb). This signal was totally unexpected, as this line is produced by a $1/2 - 1/2$ transition, and it was considered therefore as intrinsically unpolarizable. A first possible explanation was proposed by Landi Degl'Innocenti (1998), who could reproduce the observed signal by taking PRD effects and HFS into account, and by assuming that a substantial amount of atomic polarization was present in the ground level of sodium. However, as pointed out by the same author, the required amount of lower level polarization is incompatible with the presence in the lower solar chromosphere of inclined magnetic fields stronger than 0.01 G, in apparent contradiction with the results obtained from other type of observations (e.g. Bianda et al. 1998; Stenflo et al. 1998). This circumstance opened a sort of “sodium paradox”, which is still largely debated. Recently, Belluzzi & Trujillo Bueno (2013) and Belluzzi et al. (2015) proposed a new mechanism that may explain the presence of such a signal, without the need of atomic polarization in the lower level. According to these works, the observed signals are ultimately due to small variations of the anisotropy of the chromospheric radiation, over spectral intervals as small as the separation among the HFS components of this line. This picture seems to be in agreement with the results of new observations carried out with the Zurich Imaging Polarimeter (ZIMPOL, see Ramelli et al. 2010), which have unequivocally confirmed the presence of such signals, and shown a rich diversity of profiles.

Up to now, the polarimetric accuracy and spectral resolution necessary to detect the polarimetric signals described above could only be reached by completely sacrificing the temporal and spatial resolution of the observation. The great challenge today is to observe these signals at high spatial and temporal resolution, so as to understand whether and how they change depending on the plasma structure that is observed. This is a scientific target for which the use of a 4 m aperture telescope, such as EST, is required. Observing the sodium doublet by combining high polarimetric accuracy and increased spectral, spatial, and temporal resolution will be of great scientific interest, both for getting more insights about the physics of

scattering polarization, and for investigating and possibly exploiting the diagnostic potential of the signals produced by these lines.

9 Solar science exploration between 350 and 400 nm

Authors: S. Mathews, J. Leenaarts, L. Bellot Rubio, A. Feller, T. Riethmueller, M. Mathioudakis

9.1 White-light emission from flares - Continuum diagnostics in vicinity of Balmer jump

White-light continuum emission in the visible and NUV carries a significant fraction of the solar flare radiative energy losses. The white-light continuum is usually attributed to hydrogen recombination radiation to $n=2$ and $n=3$ (Balmer and Paschen) and to a lesser extent negative hydrogen (H^-) bound-free and free-free emission with the local plasma conditions determining the processes that dominate. Radiative hydrodynamic (RHD) simulations of stellar and solar flares have shown that the 320nm – 400nm wavelength range contains important diagnostics that are very sensitive to the flare heating processes. If we consider flare energy transport by non-thermal electrons, the colour temperature of the NUV continuum and Balmer jump ratio (364.6nm), are very sensitive to the nonthermal electron flux. High resolution spectroscopy combined with RHD simulations can be used to determine the photospheric/chromospheric flare electron density and allow us to disentangle the atomic processes that contribute to form the observed shape of the flare white-light continua. The ratio of the relative strength of the higher Balmer lines (i.e. Balmer decrement) can also be used as a diagnostic for the optical depth, temperature and density of the flaring atmosphere. Despite a wealth of stellar flare observations in the visible and NUV, solar flare observations in vicinity of the Balmer jump are currently scarce (Kowalski et al. 2015, Kowalski et al. 2016). This is due to the lack of suitable solar instrumentation. A high-resolution spectrograph in the NUV will allow us to access important line and continuum diagnostics and hence determine the atmospheric conditions, dynamics and heating processes that lead to the formation of the white-light flare.

9.2 Coronal forbidden lines in the visible

Coronal forbidden lines offer many important diagnostic capabilities, including the potential to measure magnetic fields for those in the NIR (e.g. Penn (2014), Judge (1998)). As highlighted in Del Zanna and DeLuca (2017) they also offer diagnostic capability in the form of plasma densities, flows and line widths, with the latter in particular providing potential constraints for coronal heating models, as well as the prospect of detecting the presence of non-Maxwellian electron distributions (Dudík et al. 2014), in combination with measurements in the UV or EUV. The photospheric brightness is such that coronal lines in both visible and NIR wavelengths are most readily detected above the limb with coronagraphs. However, there are many lines in the visible range (see Table 2 in Del Zanna & DeLuca (2017)) including the Fe XIII line at 3388.1 Å, for which there has been a positive on disk detection during a flare on the M dwarfs CN Leo and LHS 2076, with the line displaying high variability (Fuhrmeister et al., 2003), confirming the detection by Schmitt & Wichmann (2001). Indeed, Fuhrmeister et al. find variability of the Fe XIII outside of the flare on timescales that they suggest may be consistent with microflaring. Table 2 of Del Zanna & DeLuca (2017) provides estimates of observed and predicted radiance for an AR at the limb, that indicate the Ca XIII 4087 Å, Ni XIII 5116 Å, Ca XV 5696 Å and Fe X 6374 Å lines have radiances of similar order of magnitude. The Fe XIV 5304 Å line is an order of magnitude brighter. A proper assessment of predicted on-disk intensities is needed, but a high-resolution spectrograph could potentially allow the measurement of the variability of these coronal lines (including measurement of velocities and line widths) during flares at the same time as chromospheric and white-light measurements, allowing complete tracking of the energy throughout the atmosphere.

9.3 Ca II H&K spectroscopy and spectropolarimetry

Recent high resolution imaging spectroscopy taking with CHROMIS at the SST shows the great potential of the Ca II H&K lines (Leenaarts et al. 2018)⁵, providing high spatial resolution owing to their short wavelength and great temperature sensitivity. Spectropolarimetric measurements in the same lines⁶ show Stokes V signal in sunspots, plage and flares of the order of 5%-10%, while quiet Sun signals appear weaker than 0.2%. The lines have a higher opacity than Ca II 854.2 and H α ⁷, and thus provide diagnostic information at larger heights in the chromosphere than any other line available from the ground (maybe except for the He I 587.6 and 1083.0 lines). The drawback of the lines are the low photon fluxes. At a total efficiency of 2% (=EST coude room) and $R=5e4$ one needs to integrate 60 s to reach a S/N of 1000 in a critically sampled, diffraction-limited pixel in the line core in the quiet sun. Putting an instrument in the Nasmyth focus

⁵ <http://iopscience.iop.org/article/10.3847/2041-8213/aa99dd/pdf>

⁶ <http://adsabs.harvard.edu/abs/1990ApJ...361L...81M>

⁷ <https://arxiv.org/pdf/1306.0671.pdf>, <https://arxiv.org/pdf/1712.01045.pdf>

would increase the total efficiency, perhaps as high as 10%-15%, which would lead to a factor 5-7 shorter integration time. Sampling at half the diffraction limit decreases integration time by another factor 4. High resolution imaging requires an AO system. A Fabry-Perot or 2D imaging spectropolarimeter or even a slit spectropolarimeter would provide a unique window on the solar chromosphere. DKIST first light instruments will not have a narrow-band imaging instrument in the Ca II H&K lines, only a slit spectropolarimeter.

9.4 Multi-line spectropolarimetry

Our knowledge of the lower solar atmosphere is mainly obtained from spectropolarimetric observations, which are often carried out in the red or infrared spectral range and almost always cover only a single or a few spectral lines. In the short-wavelength range, below 4300 Å, the line density but also the photon noise are considerably higher than in the red. This wavelength range is targeted for the next science flight of the balloon-borne solar telescope Sunrise. It is expected to be a rewarding science target for the EST as well. As a first step, the spectropolarimetric diagnostic potential of this spectral range has been studied based on simulations (Riethmüller & Solanki 2019). For an ensemble of state-of-the-art magneto-hydrodynamical atmospheres, exemplary spectral regions around, 4080 Å (328 lines), and, for comparison, around 6302 Å (111 lines) have been synthesised assuming a spectral resolving power of 150 000 as a reasonable trade-off between spectral FOV and spectral line resolution. The synthetic Stokes profiles are degraded with typical photon noise and then inverted. In the 4080 Å region the simulations show longitudinal Zeeman signal amplitudes up to 20% and transversal Zeeman amplitudes of order 1-3%. A polarimetric sensitivity of order 0.1% is thus recommended for Stokes Q, U. For Stokes V the required noise level is comparatively lower, but the instrument design shall be driven by the highest sensitivity requirement. The atmospheric parameters of the inversion are compared with the original noise-free MHD quantities. We find that from many-line inversions significantly more information can be obtained than from a traditionally used inversion of only a few lines. We further find that information about the upper photosphere can be significantly more reliably obtained at short wavelengths. In terms of sensitivity, in the mid and lower photosphere, the many-line approach at 4080 Å provides equally good results than at 6302 Å for the magnetic field strength and the line-of-sight (LOS) velocity, while the temperature determination is more precise by a factor of three. We conclude from our first theoretical modeling results that many-line spectropolarimetry at short wavelengths offers high potential in solar physics.

10 Tables for Observing Programmes

In the observing programme tables, we distinguish between the following instruments:

- (1) BBI: Broad Band Imagers, which take images with exposure times shorter than 1 ms. Images are taken at a fixed wavelength and can be centered at spectral continua or in spectral lines with pass bands as narrow as about 0.1 nm.
- (2) NBI: Narrow-Band Imagers, which scan a spectral range in wavelength. We assume that there will be three NBIs for three different wavelength regimes: blue visible, red visible, IR. For cost reasons, we assume a FOV of about 40 arcsec in diameter or 30 arcsec square side length.
- (3) SP: classical long-slit spectrograph, which scans the solar image. Slit length of 60 arcsec is assumed. Since IFUs are superior to SPs as image reconstruction techniques can be applied to their spectral images, only one SP from blue visible to IR is assumed.
- (4) IFU: Integral Field Units, which record spectrum and image simultaneously for a relatively small FOV of about 10 arcsec. We assume that IFUs are available for the spectral range from 392nm to 1600 nm. At this point of development it is unknown whether micro-lense systems (which are restricted to short wavelength ranges) or reflective image slicers (which can cover large wavelength regimes) are to be preferred.

All instruments, except for BBIs, are assumed to be operated in polarimetric mode, unless stated otherwise.

In this section we list the tables of a total 97 observing programmes that were addressed in the previous section. The electronic pdf of this document contains hyperlinks that relate the description of the science cases with the tables of the observing programme.

Spectral resolution is defined as $R = \delta\lambda/\lambda$.

The SNR (Signal-to-Noise) values in the OP tables correspond to the continuum of the intensity signal.

OP Table 1.1.1: Formation and evolution of intense flux tubes in the solar atmosphere

Duration of the observations: 1 hour, to cover the lifetime of newly formed magnetic flux tubes. The program should be repeated several times to build up a significant statistical sample. Light distribution: All instruments work simultaneously and receive 100% of the light at the indicated wavelengths.

Instrument 1	NBIs	
Goal	Detection of rapid flows and intensification of the field in photosphere and chromosphere over large FOVs, to build up statistically significant sample	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 617.3 nm	+Fe I 557.6 nm (Stokes I)
Chromosphere	Mg I 517.3 nm	+H α (Stokes I)
FOV	40'' \times 40''	60'' \times 60''
λ samples	10	20 for chromospheric lines
Spatial resolution	Diffraction limit	
SNR	500	
Cadence	30 s	10 s

Instrument 2	IFUs	
Goal	Detailed study of physical processes in and around flux tubes undergoing convective collapse	
	Requirement	Goal
Photosphere	Fe I 630.2 nm, Fe I 1565 nm	+Si I 1082.7 nm
Chromosphere	Ca II 854 nm	+Ca II H, He I 1083 nm
FOV	5'' \times 5''	As large as possible
Spatial resolution	0'.1	
SNR	3000	
Integration time	3 s	
Cadence	30 s	
Notes	IFU allows tiles to be reconstructed and minimizes differential refraction effects	

Instrument 3	BBIs	
Goal	Response of chromosphere to convective collapse and possible heating events. Context information	
	Requirement	Goal
Photosphere	CN bandhead	+ G-band
Chromosphere	Ca II K line core & line wing	
FOV	40'' \times 40''	As large as possible
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1 s

OP Table 1.2.1: Internal structure and evolution of magnetic elements

Duration of the observations: 1 hour, to cover the lifetime of magnetic flux tubes. The program should be repeated several times to build up a statistically significant sample.

Light distribution: All instruments work simultaneously and receive 100% of the light at the indicated wavelengths.

Instrument 1		
IFUs		
Goal	Detect spatial variations of field and flows across magnetic elements as a function of height in the atmosphere. Study temporal evolution of fields, flows, and waves	
	Requirement	
	Goal	
Photosphere	Fe I 525.02, Fe I 630, Fe I 1565 nm	+Fe I 709 (Stokes I), Si I 1082.7 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854	+Ca II H, He I 1083 nm
FOV	7" × 7"	15" × 15"
Spatial resolution	0'06	
SNR	2000	3000
Integration time/tile	3 s	
Cadence	15 s	
Notes	Four tiles of 3'8 × 3'8 needed to cover FOV	

Instrument 2		
BBIs		
Goal	Response of chromosphere to flows and waves. Context information (horizontal motions)	
	Requirement	
	Goal	
Photosphere	CN bandhead	+ G-band
Chromosphere	Ca II K line core & line wing	+ H α
FOV	20" × 20"	
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1 s
Notes	CN bandhead and Ca II K to allow IFU to observe Ca II H CN bandhead offers increased spatial resolution compared to G band	

OP Table 1.3.1: Magnetic bright points

Duration of the observations: Single co-temporal shots of several broad-band filtergrams to study morphologic appearance of MBPs from the photosphere to the chromosphere together with IFU for detailed magnetic field distribution characterization. The program should be repeated several times to build up a significant statistical sample.

Light distribution: IFU spectroplarimeters and BBIs work simultaneously with major (80%) fraction of light going to IFUs.

Instrument 1	IFUs	
Goal	Detailed study of the magnetic field strength distribution of MBPs at several atmospheric heights	
	Requirement	Goal
Photosphere	Fe I 630.2, Fe I 1565	+Si I 1082.7 nm
Chromosphere	Ca II 854 nm	+Ca II H, He I 1083 nm
FOV	5" × 5"	As large as possible
Spatial resolution	0.1	
SNR	1000	3000
Integration time/tile	3 s	
Cadence	not crucial	
Notes	IFU allows tiles to be reconstructed and minimizes differential refraction effects	

Instrument 2	BBIs	
Goal	Morphologic appearance of MBPs: size distribution and shape of MBPs at various atmospheric heights	
	Requirement	Goal
Photosphere	G band	+ CN bandhead
Chromosphere	Ca II K line core & line wing Mg I b ₂ 517.3	
FOV	40" × 40"	As large as possible
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	not crucial	not crucial

OP Table 1.3.2: Dynamic parameters and evolution of MBPs

Duration of the observations: 15 minutes minimum up to 1 hour, to cover the lifetime of newly formed MBPs. The program should be repeated several times to build up a significant statistical sample.

Light distribution: NBIs and broad-band imagers work simultaneously with 100% of the light going to each instrument at the indicated wavelength.

Instrument 1	NBIs	
Goal	Detection of physical parameters in and around MBP during its evolution.	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 617.3 nm	+Fe I 557.6 nm (Stokes I)
Chromosphere	Mg I 517.3 nm	+H α (Stokes I)
FOV	40'' \times 40''	60'' \times 60''
λ samples	10	20 for chromospheric lines
Spatial resolution	Diffraction limit	
SNR	500	
Cadence	10 s	5 s

Instrument 2	BBIs	
Goal	Intensity and shape of MBPs during their evolution in various heights	
	Requirement	Goal
Photosphere	G band	+ CN bandhead
Chromosphere	Ca II K line core & line wing	
FOV	40'' \times 40''	As large as possible
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1 s

OP Table 1.4.1: Emergence and evolution of magnetic fields in granular convection

Duration of the observations: 3-4 hours, to study temporal variation of appearance rates and build up a statistically significant sample.

Light distribution: All instruments work simultaneously and receive 100% of the light at the indicated wavelengths.

Instrument 1	NBIs	
Goal	Evaluate frequency of appearance of small-scale magnetic loops and unipolar flux features in quiet Sun. Estimate magnetic fluxes. Assess influence on chromospheric layers	
	Requirement	Goal
Photosphere	Fe I 630, Fe I 1565	+Si I 1082.7 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm, H α	+He I 1083 nm
λ samples	15	20 for chromospheric lines
FOV	40'' \times 40''	60'' \times 60''
Spatial resolution	0.2	
SNR	2000	
Integration time/wav	1 s	
Cadence	15 s	
Notes	FOV should cover at least one supergranule	

Instrument 2	BBI	
Goal	Resolve substructure of emerging flux concentrations. Detect small-scale chromospheric heating events. Context information (horizontal motions).	
	Requirement	Goal
Photosphere	CN bandhead	+G-band
Chromosphere	Ca II K line core & line wing	
FOV	40'' \times 40''	60'' \times 60''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1
Notes	Same FOV as narrow-band filtergraphs	

OP Table 1.4.2: Properties of magnetic fields emerging in the quiet Sun

Duration of the observations: 1 hour, to cover full emergence process and subsequent evolution and build up a statistically significant sample.

Light distribution: All instruments work simultaneously and receive 100% of the light at the indicated wavelengths.

Instrument 1	IFUs	
Goal	Determine magnetic field topology and dynamics of emerging loops as a function of height. Search for opposite polarities in unipolar flux emergence processes. Detailed study of the interaction of emerging fields with pre-existing flux in photosphere and chromosphere. Resolve polarity inversion lines.	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 630, Fe I 1565 nm	+Si I 1082.7 nm, Mn I 1526
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm	+H α , Ca II H, He I 1083 nm
FOV	7'' \times 7''	15'' \times 15''
Spatial resolution	0''.06	
SNR	2000	3000
Integration time/tile	10 s	
Cadence	40 s	30 s
Notes	Four tiles of 3''.8 \times 3''.8 with pixel size of 0''.03 needed to cover FOV Goal priority: FOV, cadence, SNR	

Instrument 2	BBIs	
Goal	Resolve internal structure of emerging flux concentrations. Detect small-scale chromospheric heating events. Context information (horizontal motions, large-scale magnetic topology at emergence site)	
	Requirement	Goal
Photosphere	CN bandhead	+G band
Chromosphere	Ca II K line core & line wing	
FOV	40'' \times 40''	As large as possible
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1 s
Notes	FOV should cover at least one full supergranule Ca II K instead of H to allow IFU to observe Ca II H	

OP Table 1.5.1: Magnetic field topology, dynamics and energy release at flux cancellation sites

Duration of the observations: 1 hour, to study evolution of cancelling features as they approach each other and post-cancellation effects. The program should be repeated several times to build up a statistically significant sample.

Instrument 1	IFUs	
Goal	Determine magnetic field topology of cancelling features as a function of height in the atmosphere. Detect magnetic field configuration at polarity inversion line. Study dynamical effects and possible energy release in upper atmospheric layers.	
	Requirement	Goal
Photosphere	Fe II 492.3, Fe I 630, Fe I 1565 nm	+H α (Stokes I)
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm Ca II K (intensity only)	
FOV	7'' \times 7''	15'' \times 15''
Spatial resolution	0'.06 (up to 854 nm)	
SNR	2000	3000
Integration time/tile	10 s	
Cadence	40 s	15 s
Notes	Four tiles of 3'.8 \times 3'.8 with pixel size of 0'.03 needed to cover FOV Goal priority: Cadence, SNR, FOV	

OP Table 1.6.1: Physical properties of internetwork magnetic fields

Duration of the observations: The time needed to take a single map. However, it would be useful to keep observing for 1-2 hours to have a few maps and study short-term variations within the same supergranular cell.

Instrument 1	IFUs	
Goal	Uncover linear polarisation signals. Determine distribution of field strengths, field inclinations, and flows in internetwork regions at highest spatial resolution. Determine height variation of field properties and flows. Resolve polarity inversion lines. Compare Zeeman and Hanle views of internetwork magnetism. Determine total flux content of supergranular cells.	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 630, Fe I 1565 nm, Sr I 460.7 nm	+ Mn I 553, Mn I 1526 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm	
FOV	40'' \times 40''	60'' \times 60''
Spatial resolution	0'.1	
SNR	10000	
Integration time/tile	25 s	
Cadence	20 min (not critical)	
Notes	FOV should cover at least one entire supergranule. Program can also be done with long-slit spectrograph, but IFU allows to reconstruct tiles and minimizes differential refraction effects. 44 tiles of 6'' \times 6'' with pixel size of 0'.05 needed to cover full FOV.	

OP Table 1.6.2: Short-term evolution of internetwork fields

Duration of the observations: 2 hours, to determine flux appearance and disappearance rates and their variations with time.

Light distribution: All instruments work simultaneously and receive 100% of the light at the indicated wavelengths.

Instrument 1		
Goal	Narrow-band filtergraphs Study appearance, evolution, and disappearance of IN fields. Determine connectivity between photosphere and chromosphere	
	Requirement	
	Goal	
Photosphere	Fe I 520.5, Fe I 630.2, Fe I 1565	+Ca II H (spectroscopy), He I 1083 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm, H α (spectroscopy)	
λ samples	10	20 for chromospheric lines
FOV	40'' \times 40''	70'' \times 70''
Spatial resolution	0:1	
SNR	2000	
Integration time/wav	3 s	
Cadence	30 s	
Notes	FOV should cover one full supergranule	

Instrument 2		
Goal	BBI's Detection of small-scale heating events at different heights in the atmosphere. Provide context information (horizontal motions, large scale structure of supergranule)	
	Requirement	
	Goal	
Photosphere	CN bandhead	
Chromosphere	Ca II K line core & line wing	
FOV	40'' \times 40''	70'' \times 70''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1 s
Notes	FOV should cover at least a supergranule. Ca II K to allow filtergraphs to observe Ca II H Really necessary when filtergraphs have wide-band cameras?	

OP Table 1.7.1: Structure of polar faculae

Duration of the observations: 1 hour, to cover lifetime of individual structures and build up a statistically significant sample.

Light distribution: All instruments work simultaneously and receive 100% of the light at the indicated wavelengths.

Instrument 1	IFUs	
Goal	Determine height variation of field properties and flows in isolated strong flux concentrations near the poles	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 630, Fe I 1565 nm	+Si I 1082.7, Mn I 1526 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm	+Ca II H (spectroscopy), He I 1083 nm
FOV	7" × 7"	As large as possible
Spatial resolution	0'06	
SNR	2000	
Integration time/tile	6 s	
Cadence	25 s	
Notes	Highest spatial resolution needed to partially compensate for projection effects. Four tiles of 3'8 × 3'8 with pixel size of 0'03 needed to cover FOV.	

Instrument 2	BBIs	
Goal	Resolve substructure of flux concentrations. Detect small-scale chromospheric events (jets, surges, waves). Context information	
	Requirement	Goal
Photosphere	CN bandhead	+ G band
Chromosphere	Ca II K line core & line wing	+ H α
FOV	40" × 40"	60" × 60"
Spatial resolution	Diffraction limit (0'025 at 397 nm)	
SNR	> 100	
Cadence	5 s	1 s
Notes	CN bandhead preferred over G band for poles because of higher spatial res	

OP Table 1.7.2: Properties, distribution and evolution of polar magnetic fields

Duration of the observations: 2 hour, to cover lifetime of individual structures and build up a statistically significant sample. If FOV is small, different pointings may be necessary to observe a wide range of latitudes.

Light distribution: All instruments work simultaneously and receive 100% of the light at the indicated wavelengths.

Instrument 1	NBIs	
Goal	Determine field strength, field inclination, and flux distribution as a function of latitude near the poles. Study temporal evolution of flux concentrations and interactions between them	
	Requirement	Goal
Photosphere	Fe I 630.2, Fe I 1565 nm	+ Fe I 525.02 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm	+ Ca II H (Stokes I), H α (Stokes I)
λ samples	15	20 for chromospheric lines
FOV	40'' \times 40''	80'' \times 80''
Spatial resolution	0':05	
SNR	1000	
Integration time/wav	2 s	
Cadence	30 s	
Notes	FOV should cover at least one full supergranule. Highest spatial resolution needed to partially compensate for projection effects	

Instrument 2	BBIs	
Goal	Detection of fast transient events produced by release of magnetic energy in photosphere/chromosphere. Context information	
	Requirement	Goal
Photosphere	CN bandhead	+ G band
Chromosphere	Ca II K line core & line wing	
FOV	40'' \times 40''	80'' \times 80''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1 s
Notes	CN bandhead preferred to G band because of higher spatial resolution. FOV matching that of filtergraphs.	

OP Table 1.8.1: Network evolution and dynamics

Instrument 1	BBIs	
Goal	Tracking of internetwork flux elements in supergranules	
	Requirement	Goal
Photosphere	G-band, blue and red continua	+CN bandhead
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm
FOV	60'' × 60''	120'' × 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1 s

Instrument 2	NBIs	
Goal	Detection of small-scale magnetic elements	
	Requirement	Goal
Photosphere	Fe 630.2 nm & Fe I 709.0 nm	+Fe I 1565 nm
Chromosphere	Ca II 854 nm	+He I 1083 nm
FOV	60'' × 60''	120'' × 120''
Spatial resolution	Diffraction limit	
λ samples	10 per line	31 per line
SNR	~ 500	
Cadence	30 s	10 s
Notes	The lines Fe I 630.25 and Ca II 854.2 nm should be observed both with NBI and IFU, but not simultaneously.	

Instrument 3	IFUs	
Goal	Detailed study of physical processes during the merging of internetwork elements with network elements	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 630.2, Fe I 1565 nm	+Si I 1082.7 nm & Mn I 1526 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854	+Ca II H, He I 1083 nm
FOV	5'' × 5''	As large as possible
Spatial resolution	0'.1	0'.05
SNR	500	1000
Integration time/tile	3 s	
Cadence	30 s	10 s
Notes	IFU allows to reconstruct tiles and minimizes differential refraction effects	

OP Table 2.1.1: Sausage and kink oscillations in MBPs

Instrument	BBI	
Goal	High-resolution imaging of MBPs at very high time cadence.	
	Requirement	Goal
Photosphere	G-band	80''×80''
FOV	50''×50''	
Spatial resolution	0.025''	
SNR	200	
Cadence	1 sec	
		0.5 sec

Instrument	NBIs	
Goal	High-resolution spectroscopy of MBPs.	
	Requirement	Goal
Photosphere	Na I D1	50''×50''
FOV	40''×40''	
Spatial resolution	0.05''	
SNR	200	
Wavelength samples	10	
Cadence	2 sec	1 sec

Instrument	IFU	
Goal	High-resolution spectropolarimetry of MBPs.	
	Requirement	Goal
Photosphere	Fe I 617.3 nm, Fe I 630.2 nm,	20''×20''
FOV	10''×10''	
Spatial resolution	0.025''	
SNR	1000	
Spectral resolution	150 000	
Cadence	2 sec	1 sec

OP Table 2.1.2: Magneto-acoustic waves in spicules on disk

Instrument	BBI	
Goal	High-resolution G-band imaging at very high cadence to determine wave signatures in RBEs/RREs.	
	Requirement	Goal
Photosphere	G-band	80''×80''
FOV	50''×50''	
Spatial resolution	0.025''	
SNR	200	
Cadence	1 sec	
		0.5 seconds

Instrument	NBIs	
Goal	High-resolution spectroscopy to determine wave signatures in RBEs/RREs.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Wavelength samples Cadence	Fe I 630.2 nm H α 656.3 nm, Ca II 854.2 nm He I 1083.0 nm 40'' \times 40'' 0.05'' 500 10 per line 10 sec	50'' \times 50'' 5 sec

Instrument	IFU	
Goal	High-resolution spectropolarimetry to determine wave signatures in RBEs/RREs.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Spectral Resolution Cadence	Fe I 617.3 nm, Fe I 630.2 nm, He I 1083.0 nm 10'' \times 10'' 0.025'' 1000 150 000 2 sec	20'' \times 20'' 2000 1 sec

OP Table 2.1.3: Magneto-acoustic waves in spicules at limb

Instrument	BBI	
Goal	High-resolution high cadence imaging to determine waves in spicules at the limb	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Cadence	H α , Ca II K 50'' \times 50'' 0.025'' 200 1 sec	80'' \times 80'' 0.5 sec

Instrument	NBIs	
Goal	High-resolution spectro-polarimetry to determine waves in spicules at the limb	
	Requirement	Goal
Chromosphere FOV Spatial resolution SNR Spectral Resolution Cadence	Ca II 854.2 nm, He I 1083.0 nm 40'' \times 40'' 0.05'' 500 50 000 10 sec	50'' \times 50'' 5 sec
Notes	Wavelength range: Doppler velocities in order of ± 100 km/s	

Instrument	IFU	
Goal	High-resolution spectro-polarimetry to determine waves in spicules at the limb	
	Requirement	Goal
Chromosphere	Ca II 854.2 nm, He I 1083.0 nm	
FOV	10''×10''	20''×20''
Spatial resolution	0.03''	
SNR	1000	2000
Spectral Resolution	70 000	
Cadence	5 sec	3 sec

OP Table 2.1.4: Torsional Alfvén wave (TAW) propagation in spicules

Instrument	IFU	
Goal	High-resolution, high cadence Fe I 630.2 and Ca II spectro-polarimetry to determine TAWs in spicules.	
	Requirement	Goal
Photosphere	Fe I 630.2 Stokes components	
Chromosphere	Ca II IR 854 Stokes components, Ca II K	
FOV	10''×10''	20''×20''
Spatial resolution	0.025'' (diffraction limit)	
Spectral resolution	150 000	
SNR	1000	
Cadence	2 second	

OP Table 2.2.1: Magnetic torsion and torsional oscillations of pores or sunspots

Instrument	IFU	
Goal	High-resolution, high cadence Fe I 630.2, Ca II and He I spectro-polarimetry to determine TAWs in pores (or sunspots).	
	Requirement	Goal
Photosphere	Fe I 630.2	
Chromosphere	Ca II IR, Ca II K, He I 1083.0 nm	
FOV	10''×10''	20''×20''
Spatial resolution	0.025'' (diffraction limit)	
Spectral resolution	150 000	
SNR	1000	
Cadence	1.5 second	

OP Table 2.2.2: Observation of magnetic vortexes and tornadoes

Instrument	BBI	
Goal	Statistical results in Liu et al. (2018) have shown that most (> 70%) swirls detected have lifetime less than 6 sec. High cadence is needed. Spatial resolution plays vital influence in the number and parameters of swirls detected.	
	Requirement	Goal
Photosphere	G-band	
Chromosphere	Ca II K	
FOV	40''×40''	
Spatial resolution	0.025''	
SNR	200	
Cadence	1 sec	
Instrument	NBIs	
Goal	Very high cadence observations with few line positions are needed to study the coupling and propagation of very short-lived intensity and magnetic swirls.	
	Requirement	Goal
Photosphere	Fe I 630 nm	
Chromosphere	H α , Ca II IR	
FOV	30''×30''	
Spatial resolution	0.05''	
Line positions	some 5 line positions	
SNR	300	
Cadence	2 sec	1 sec
Instrument	NBIs	
Goal	Extremely high cadence observations in Fe I line centre. More lines are desirable but high cadence is essential.	
	Requirement	Goal
Photosphere	Fe I 630 nm	
Chromosphere		
FOV	30''×30''	
Spatial resolution	0.05''	
SNR	300	
Line positions	1 line core position	
Cadence	0.1 sec	

OP Table 2.2.3: Formation of magnetic swirls in intergranular lanes

Instrument	IFU and simultaneous BBI imaging in G-Band	
Goal	High cadence and resolution observations of the intergranular lanes.	
	Requirement	Goal
Photosphere FOV Spatial resolution Spectral resolution SNR Cadence	G-Band (BBI), Fe I 630 nm 10''×10'' 0.025'' 150 000 1000 1.5 sec	

OP Table 2.2.4: Vortex flows in the lower solar atmosphere

Instrument 1	BBI (blue and red)	
Goal	Detection of vortex flows in the photosphere and tracking their evolution into the chromosphere	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Cadence	G-band, blue and red continua Ca II H line core & line wing 40'' × 40'' Diffraction limit > 100 2 s	+CN bandhead +Ca II 854 nm 90'' × 90'' 1 s
Instrument 2	NBIs	
Goal	Detection of vertical flows within vortex motions and swirls	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution Spectral resolution SNR Cadence	Fe I 630.2 nm, Fe I 709.0 nm (non-pol.) Ca II 854 nm 40'' × 40'' Diffraction limit 10 wavelength points per line ~ 500 20 s	+Fe I 1565 nm +He I 1083 nm 90'' × 90'' 31 wavelength points per line 10 s
Instrument 3	IFUs	
Goal	Detailed study of physical processes in and around vortex flows and their interaction with magnetic fields	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution Spectral resolution SNR Cadence	Fe I 525.02, Fe I 630.2, Fe I 1565 nm Mg I b ₂ 517.3, Ca II IR 854 10'' × 10'' 0.025'' (diffraction limit) 150 000 1000 1 s	+Si I 1082.7 nm, Mn I 1526 nm +Ca II H, He I 1083 nm As large as possible 2000
Notes	IFU allows to reconstruct tiles and minimizes differential refraction effects	

OP Table 2.3.1: Excitation mechanisms of sunspot waves

Instrument 1	NBIs	
Goal	Measure velocity and intensity oscillations in sunspots and nearby quiet Sun with high cadence, moderate spatial resolution, but large field of view. Long stable time series are required. Require simultaneously photospheric and chromospheric lines, spectroscopic measurements, not sensitive to magnetic field (non-polarimetric).	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution Spectral resolution SNR Cadence	Fe I 543.4, Fe I 557.6, Fe I 709.0 Ca II 854, He I 1083, H α , Ba I 455.4, K I 769.9 40'' \times 40'' 0.05'' 80 000 500 (SP mode) 20 sec	60'' \times 60'' 1000
Notes	Large field of view and simultaneous photospheric and chromospheric lines.	
Instrument 2	IFU	
Goal	Measure magnetic field vector in the photosphere and chromosphere to complement the velocity data	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution Spectral Resolution SNR Integration time Cadence	Fe I 1564.8 He I 1083, Ca II 854.2 40'' \times 40'' 0.05'' 150 000 2000 2 sec 45 sec	60'' \times 60''
Notes	Enough polarimetric precision to derive the magnetic field vector.	

OP Table 2.3.2: Alfvén waves in sunspots

Instrument 1	IFU	
Goal	Study oscillations of velocity and magnetic field in sunspots at several positions across the disc to determine if they are caused by Alfvén waves. Measurements in photospheric Fe I 1564 nm lines provide magnetic field in the deep layers, while He I 1083, and Ca II 854 nm provide magnetic field in the chromosphere. High polarimetric precision is crucial.	
	Requirement	Goal
Photosphere	Fe I 1564	40''×40''
Chromosphere	Ca II 854, He I 1083	
FOV	10''×10''	
Spatial resolution	0.05''	
SNR	2000	
Spectral resolution	150 000	
Integration time	2 sec	
Cadence	2 sec	
Instrument 2	NBI	
Goal	Bi-dimensional spectra to determine velocity oscillations at different heights from photosphere to chromosphere; phase speed of waves to determine their direction of propagation and heights of reflection. Spectroscopic (non-polarimetric mode).	
	Requirement	Goal
Photosphere	Si I 1080	40''×40''
Chromosphere	Ca II K , H α , Ba I 455.4, K I 769.9	
FOV	30''×30''	
Spatial resolution	0.05''	
Spectral resolution	80 000	
SNR	200	
Integration time	3 sec	
Cadence	30 sec	

OP Table 2.3.3: Magnetic field oscillations in sunspots

Instrument	IFU		
Goal	Measure precisely oscillations of magnetic field, together with velocity oscillations, across sunspot umbra and penumbra. Time series of 2D spectropolarimetric data in several lines with different temperature and magnetic field sensitivity simultaneously.		
	Requirement	Goal	
Photosphere	Fe I 1564, Fe I 525, Fe I 630, Fe I 613.7	20''×60''	
Chromosphere	Ca II 854, He I 1083		
FOV	10''×30''		
Spatial resolution	0.08''		
Spectral resolution	100 000		
SNR	1000		2000
Integration time	1 sec		
Cadence	5 sec		
Notes	High polarimetric precision and low noise is crucial.		

OP Table 2.3.4: Sunspot penumbral waves in the photosphere and above

Instrument	NBI		
Goal	Measure 2D velocity field over all sunspot simultaneously. Spectroscopic mode.		
	Requirement	Goal	
Photosphere	Fe I 543.4, Fe I 709.0, Fe I 630	60''×60''	
Chromosphere	Ca II 854, K I 769.9		
FOV	40''×40''		
Spatial resolution	0.05''		
Spectral resolution	100 000		
SNR	500		5000
Integration time	2 sec		
Cadence	1 minute		30 sec

Instrument 2	IFU		
Goal	Measure 2D magnetic field vector in the photosphere and chromosphere to complement the velocity data		
	Requirement	Goal	
Photosphere	Fe I 1564,	60''×60''	
Chromosphere	He I 1083, Ca II 854		
FOV	40''×40''		
Spatial resolution	0.05''		
Spectral resolution	120 000		
SNR	1000		2000
Integration time	2 sec		
Cadence	60 sec		
Notes	High polarimetric precision and low noise.		

OP Table 2.3.5: Sausage and kink oscillations in pores

Instrument 1	BBI, IFU	
Goal	High-resolution G-band images and LoS magnetic field to determine sausage and kink waves in pores.	
	Requirement	Goal
Photosphere	G-band, Fe I 1564	0.1 sec
Chromosphere	Na I D ₁ 589 nm	
FOV	10''×10''	
Spatial resolution	0.08''	
Spectral resolution	150 000	
SNR	800	
Integration time	1 sec	
Notes	Extremely high-cadence is desired.	

OP Table 2.3.6: (Torsional) Alfvén waves in pores

Instrument 1	IFU	
Goal	Study oscillations of velocity and magnetic field in sunspots at several positions across the disc to determine if they are caused by TAWs. Measurements in photospheric Fe I 1564 nm lines provide magnetic field in the deep layers, while He I 1083, and Ca II 854 nm provide magnetic field in the chromosphere.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Integration time Cadence	Fe I 1564 Ca II IR 854.5, He I 1083.0 10''×10'' 0.06'' 2000 3 sec 3 sec	40''×40''
Notes	High polarimetric precision is crucial	
Instrument 2	NBI	
Goal	Spectra to determine velocity oscillations at different heights from photosphere to chromosphere; phase speed of waves to determine their direction of propagation and heights of reflection.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Integration time Cadence	Si I 1080 Ca II K , H α , Ba I 455.4, K I 769.9 40''×40'' 0.05'' 300 3 sec < 20 sec	40''×40''
Notes	Bi-dimensional field of view	

OP Table 2.4.1: High-frequency wave propagation and dissipation from the photosphere to the chromosphere

Instrument 1	BBI, NBI	
Goal	Measure 2D velocity field over quiet and plage regions with simultaneous G-Band imaging.	
	Requirement	Goal
Photosphere	G-band, Fe I 543.4, Fe I 709.0	150 000 (with IFUs) 500
Chromosphere	H α , Ca II 854, K I 769.94, Ba I 455.5	
FOV	30'' \times 30''	
Spatial resolution	0.05''	
Spectral resolution	80 000	
SNR	300 (SP mode)	
Integration time	1 sec	
Cadence	3 sec	
Notes	High resolution and high cadence in 2D field of view in spectroscopic mode.	
Instrument 2	IFU	
Goal	Measure 2D magnetic field vector in the photosphere and chromosphere to complement the velocity data	
	Requirement	Goal
Photosphere	Fe I 1564,	3000 10 sec 90 sec
Chromosphere	He I 1083, Ca II IR 854	
FOV	30'' \times 30''	
Spatial resolution	0.06''	
SNR	2000	
Integration time	4 sec	
Cadence	45 sec	
Notes	High polarimetric precision and low noise.	

OP Table 2.4.2: Time-dependent behaviour of chromospheric jets

Instrument 1	NBI		
Goal	Measure 2D velocity and intensity oscillations over network magnetic field concentrations		
	Requirement	Goal	
Photosphere	Fe I 543.4, Fe I 709.0	40''×40''	
Chromosphere	Ca II H, H α , Ba I 455.5		
FOV	20''×20''		
Spatial resolution	0.05''		
SNR	500		1000
Integration time	2 sec		
Cadence	20 sec		6 sec
Notes	High resolution and high cadence in 2D field of view		

Instrument 2	IFU		
Goal	Measure 2D magnetic field vector in the photosphere and chromosphere to complement the velocity data		
	Requirement	Goal	
Photosphere	Fe I 1564,	3000	
Chromosphere	He I 1083, Ca II 854		
FOV	10''×10''		
Spatial resolution	0.06''		
Spectral Resolution	150 000		10 sec
SNR	2000		
Integration time	4 sec		
Notes	High polarimetric precision and low noise.		

OP Table 2.4.3: Network and plage oscillations

Instrument 1	NBI		
Goal	Measure long-period velocity oscillations in the network, high spatial resolution, medium-large field of view. Long stable time series are required. Require simultaneously photospheric and chromospheric lines best suited for velocity measurements.		
	Requirement	Goal	
Photosphere	Si I 1082.7, Fe I 709.0,	60''×60''	
Chromosphere	Ca II 854, He I 1083, H α , Ba I 455.4		
FOV	40''×40''		
Spatial resolution	0.05''		
SNR	500		1000
Cadence	30 sec		
Notes	Large field of view and simultaneous photospheric and chromospheric lines.		

Instrument 2	IFU	
Goal	Measure magnetic field vector in the photosphere and chromosphere to complement the velocity data	
	Requirement	Goal
Photosphere	Fe I 1564.8	60''×60''
Chromosphere	He I 1083, Ca II 854.2	
FOV	40''×40''	
Spatial resolution	0.05''	
Spectral resolution	150 000	
SNR	2000	
Integration time	3 sec	
Cadence	50 sec	
Notes	High polarimetric precision to derive the magnetic field vector in network.	

OP Table 3.1.1: Magnetic field structure in the quiet Sun chromosphere

Instrument	SP	
Goal	Determine chromospheric magnetic field on a supergranular scale	
	Requirement	Goal
Photosphere	Fe I 630.1/630.2, Si I 1082.7	0'1 10 s
Chromosphere	Na I 589.0/589.6, Ca II 854.2, He I 1083	
Spectral range	$\pm 75 \text{ km s}^{-1}$	
Spectral resolution	$\sim 3 \text{ km s}^{-1}$	
FOV	60'' \times 60''	
Spatial resolution	0'2	
SNR	10 ⁴	
Cadence per slit step	as long as needed for S/N	
Polarimetry	yes, all lines	
Notes	The spectrograph requires atmospheric refraction compensation to obtain strictly cospatial spectra. Needs 300 slit steps to cover the desired FOV	

OP Table 3.1.2: Fibrillar structure of the chromosphere

Instrument	NBI	
Goal	Observe fibrillar structure of the chromosphere at the highest resolution	
	Requirement	Goal
Photosphere	Fe I 630.1	He I 1083 as short as possible yes, in He I 1083
Chromosphere	Ca II 393.4, H I 656.3	
Wavelength samples	~ 10 per line	
Spectral range	$\pm 50 \text{ km s}^{-1}$	
Spectral resolution	3–6 km s^{-1}	
FOV	30'' \times 30''	
Spatial resolution	0'025 – 0'06	
SNR	$3 \times 10^2 - 10^3$	
Cadence	30 s	
Polarimetry	yes, in Fe I 630.1	

OP Table 3.2.1: Type II spicule acceleration on disk

Instrument	IFU	
Goal	Catch spicule acceleration on-disk at highest spatial and temporal resolution with polarimetry	
	Requirement	Goal
Photosphere	Fe I 630.1	Ca II 393.4
Chromosphere	Na I 589.0/589.6, H I 656.3, Ca II 854.2,	
Spectral range	$\pm 100 \text{ km s}^{-1}$	
Spectral resolution	3 – 6 km s^{-1}	
FOV	10'' \times 10''	
Spatial resolution	spectropolarimetry: 0'1, imaging: diffraction limit	
SNR	10 ³ (393.4, 656.3) – 3.3×10^3 (589.0/589.6, 630.1, 854.2)	
Cadence	4 s	
Polarimetry	yes, in NaI 589.0/589.6, Fe I 630.1 and Ca II 854.2	

OP Table 3.2.2: Type II spicule evolution on disk

Instrument	IFU	
Goal	Catch spicule evolution on disk at the highest spatial and temporal resolution with polarimetry	
	Requirement	Goal
Photosphere	Fe I 630.1	Ca II 393.4 diffraction limit
Chromosphere	HI 656.3, Ca II 854.2, He I 1083	
Spectral range	$\pm 100 \text{ km s}^{-1}$	
Spectral resolution	$3 - 6 \text{ km s}^{-1}$	
FOV	$10'' \times 10''$	
Spatial resolution	$0'.1$	
SNR	$10^3(393.4, 656.3) - 3.3 \times 10^3 (854.2, 1083)$	
Cadence	4 s	
Polarimetry	yes, in Fe I 630.1, Ca II 854.2, He I 1083	

OP Table 3.2.3: Type II spicule evolution off-limb

Instrument	IFU	
Goal	Catch spicule evolution off limb at highest spatial and temporal resolution with polarimetry	
	Requirement	Goal
Photosphere	HI 656.3, Ca II 854.2, He I 1083 $\pm 75 \text{ km s}^{-1}$ $3 - 6 \text{ km s}^{-1}$ $10'' \times 10''$ $0'.1$ $10^3(393.4, 656.3) - 3.3 \times 10^3 (854.2, 1083)$ 4 s yes, in Ca II 854.2 and He I 1083	Ca II 393.4 diffraction limit
Chromosphere		
Spectral range		
Spectral resolution		
FOV		
Spatial resolution		
SNR		
Cadence		
Polarimetry		

OP Table 3.2.4: Magnetic field of spicules

Instrument	SP	
Goal	Measure magnetic fields in spicules on disk using He I lines.	
	Requirement	Goal
Photosphere	He I 587.6, He I 1083 $\pm 100 \text{ km s}^{-1}$ $3 - 6 \text{ km s}^{-1}$ $60'' \times 60''$ $0'.1$ 3.3×10^3 5 s yes	10^4
Chromosphere		
Spectral range		
Spectral resolution		
FOV		
Spatial resolution		
SNR		
Cadence per slit step		
Polarimetry		
Notes	requires atmospheric refraction compensation	

OP Table 3.3.1: Small-scale chromospheric jets

Instrument	NBI	
Goal	Measure the fine structure of small-scale chromospheric jets	
	Requirement	Goal
Photosphere	Ca II 393.4, HI 656.3, Ca II 854.2 ~ 20 per line $\pm 100 \text{ km s}^{-1}$ 4 km s^{-1} $30'' \times 30''$ $< 0'.05$ 10^3 10-20 s no	
Chromosphere		
Wavelength samples		
Spectral range		
Spectral resolution		
FOV		
Spatial resolution		
SNR		
Cadence		
Polarimetry	no	
Notes	requires atmospheric refraction compensation	

OP Table 3.4.1: Alfvén waves

Instrument	NBI	
Goal	Study the propagation and damping of Alfvén waves in the chromosphere	
	Requirement	Goal
Photosphere	Fe I 630.1 Ca II 393.4, HI 656.3, Ca II 854.2 imaging: 1-5, spectropolarimetry: 10 $\pm 75 \text{ km s}^{-1}$ $3-6 \text{ km s}^{-1}$ $30'' \times 30''$ $0'.05 - 0'.15$ 10^3 imaging 1-5 s, spectropolarimetry: 10 s-15 s yes, in Fe I 630.1, Ca II 854.2	He I 1083 yes, in He I 1083
Chromosphere		
Wavelength samples		
Spectral range		
Spectral resolution		
FOV		
Spatial resolution		
SNR		
Cadence		
Polarimetry	yes, in Fe I 630.1, Ca II 854.2	

OP Table 3.4.2: Acoustic wave interaction with chromospheric field structure

Instrument	NBI	
Goal	Study the interaction of acoustic waves with the chromospheric magnetic field	
	Requirement	Goal
Photosphere	Fe I 630.1 Ca II 393.4, HI 656.3, Ca II 854.2 ~15 $\pm 50 \text{ km s}^{-1}$ $3-6 \text{ km s}^{-1}$ $30'' \times 30''$ $0'.05 - 0'.1$ 10^3 imaging 5 s, spectropolarimetry: 15 s yes, in Fe I 630.1, Ca II 854.2 and He I 1083	
Chromosphere		
Wavelength samples		
Spectral range		
Spectral resolution		
FOV		
Spatial resolution		
SNR		
Cadence		
Polarimetry	yes, in Fe I 630.1, Ca II 854.2 and He I 1083	

OP Table 3.5.1: Highly variable phenomena in the chromosphere

Instrument	NBI	
Goal	Study highly variable phenomena in the chromosphere	
	Requirement	Goal
Photosphere	Fe I 630.1	± 50 km s ⁻¹ diffraction limit
Chromosphere	Ca II 393.4, HI 656.3, Ca II 854.2	
Wavelength samples	1 – 5	
Spectral range	± 50 km s ⁻¹	
Spectral resolution	3 – 6 km s ⁻¹	
FOV	30'' × 30''	
Spatial resolution	0'.05	
SNR	10 ³	
Cadence	1 s	
Polarimetry	yes, in Fe I 630.1 at lower cadence	

OP Table 3.5.2: Highly variable phenomena in the chromosphere

Instrument	IFU	
Goal	Study highly variable phenomena in the chromosphere	
	Requirement	Goal
Photosphere	Fe I 630.1	He I 1083 10'' × 10'' diffraction limit
Chromosphere	Ca II 393.4, HI 656.3, Ca II 854.2	
Spectral range	± 100 km s ⁻¹	
Spectral resolution	3 – 6 km s ⁻¹	
FOV	5'' × 5''	
Spatial resolution	0'.05	
SNR	10 ³	
Cadence	1 s	
Polarimetry	yes, in Fe I 630.1 and Ca II 854.2	

OP Table 3.5.3: Reconnection at different heights

Instrument	IFU	
Goal	Study reconnection at different heights	
	Requirement	Goal
Photosphere	Fe I 630.1	
Chromosphere	Ca II 393.4, HI 656.3, Ca II 854.2, He I 1083	
Spectral range	± 150 km s ⁻¹	
Spectral resolution	3 – 6 km s ⁻¹	
FOV	10'' × 10''	
Spatial resolution	0'.05 – 0'.1	
SNR	imaging 10 ³ , polarimetry 3.3 × 10 ³	
Cadence	5 s	
Polarimetry	yes, in Fe I 630.1, Ca II 854.2 and He I 1083	

OP Table 3.6.1: Electric currents and heating in active regions

Instrument	NBI	
Goal	Determine electric currents in active regions	
	Requirement	Goal
Photosphere	Fe I 630.1	
Chromosphere	Na I 589.0/589.6, Ca II 849.8, Ca II 854.2, He I 1083	
Wavelength samples	15	
Spectral range	$\pm 70 \text{ km s}^{-1}$	
Spectral resolution	$3 - 6 \text{ km s}^{-1}$	
FOV	$30'' \times 30''$	
Spatial resolution	$0'.1$	
SNR	10^3	
Cadence	15 s	
Polarimetry	yes, in all lines	

OP Table 3.6.2: Electric currents and heating outside active regions

Instrument	SP	
Goal	Determine electric currents outside active regions	
	Requirement	Goal
Photosphere	Fe I 630.1	
Chromosphere	Na I 589.0/589.6, Ca II 849.8, Ca II 854.2, He I 1083	
Spectral range	$\pm 70 \text{ km s}^{-1}$	
Spectral resolution	$3 - 6 \text{ km s}^{-1}$	
FOV	$60'' \times 60''$	
Spatial resolution	$0'.1$	
SNR	3.3×10^3	
Cadence per slit step	5 s	
Polarimetry	yes, in all lines	

OP Table 3.6.3: Electric currents and heating outside active regions at small spatial scales

Instrument	IFU	
Goal	Determine time-varying electric currents outside active regions at small spatial scales	
	Requirement	Goal
Photosphere	Fe I 630.1	$10'' \times 10''$
Chromosphere	Na I 589.0/589.6, Ca II 849.8, Ca II 854.2, He I 1083	
Spectral range	$\pm 70 \text{ km s}^{-1}$	
Spectral resolution	$3 - 6 \text{ km s}^{-1}$	
FOV	$5'' \times 5''$	
Spatial resolution	$0'.1$	
SNR	3.3×10^3	
Cadence	5 s	
Polarimetry	yes, in all lines	

OP Table 3.7.1: Measurement of CO clouds

Instrument	SP	
Goal	Study properties of cool clouds in the chromosphere	
	Requirement	Goal
Photosphere Chromosphere Spectral range Spectral resolution FOV Spatial resolution SNR Cadence per slit step Polarimetry	4700 nm CO band 120 nm R=25,000 60'' × 60'' 0'3 10 ³ ? as long as needed for S/N yes	

OP Table 3.7.2: Temperature bifurcation diagnostics by scattering polarization in Ca II K

Instrument	SP	
Goal	Study properties of cool clouds in the chromosphere	
	Requirement	Goal
Photosphere Chromosphere Spectral range Spectral resolution FOV Spatial resolution SNR Cadence Polarimetry	Ca II 393.4 1 nm R = 100,000 0.3'' × 60'' 0'3 10 ³ 30 s yes	

OP Table 3.8.1: Magnetic field determination in plage including Ca II H&K

Instrument	SP	
Goal	Determine magnetic fields in plage, with a focus on the upper chromosphere.	
	Requirement	Goal
Photosphere Chromosphere Spectral range Spectral resolution FOV Spatial resolution SNR Cadence per slit step Polarimetry	Fe I 630.1/630.2 Ca II 393.4, Ca II 396.8,, Na I 589.0/589.6., Ca II 854.2, He I 1083 ± 100 km s ⁻¹ 3 km s ⁻¹ 60'' × 20'' 0'1 10 ³ 10 s yes, in all lines	60'' × 60'' 0'05 as high as possible
Notes	SNR is set for Ca II 393.4 and Ca II 396.8, other lines might have different requirements.	

OP Table 4.1.1: Stability of the umbra - interplay between the convection and magnetic forces

Instrument 1	IFU	
Goal	Spectropolarimetric observations of the umbral boundary to study the detailed processes in the magnetic field evolution.	
	Requirement	Goal
Spectral lines	Fe I 630.2 nm; Ca II 854.2 nm	+Fe I 630.15 nm
FOV	5'' × 5''	As large as possible
Spatial resolution	0'08	0'05
Spectral resolution R	200 000	
Spectral range	0.3 nm (for Ca II line) 0.1 nm (for Fe I line)	0.6 nm
SNR	2000	
Cadence	30 s	15 s
Instrument 2	NBI	
Goal	Context information about magnetic field configuration in a larger FOV.	
	Requirement	Goal
Spectral lines	Fe I 1565 nm; He I 1083 nm	+Si I 1082.7 nm
FOV	30'' × 30''	60'' × 60''
Spatial resolution	0'08	0'05
Spectral resolution R	50 000	
Spectral range	0.5 nm (for He I line) 0.1 nm (for Fe I line)	0.8 nm 0.2 nm
SNR	1000	2000
Cadence	120 s	30 s
Instrument 3	BBI	
Goal	Context information, for LCT and feature tracking.	
	Requirement	Goal
Spectral lines	G-band; Ca II H line core & line wing	
FOV	30'' × 30''	60'' × 60''
Spatial resolution	0'05	diffraction limit
SNR	100	200
Cadence	5 s	2 s

OP Table 4.2.1: Multi-wavelength analysis of umbral dots

Instrument 1	IFUs	
Goal	Spectropolarimetric observations of the umbral dots for detailed study of magnetic field configuration and temporal evolution.	
	Requirement	Goal
Spectral lines	Fe I 1565 nm; Ca II 854 nm	+Fe I 630.15 nm & 630.25 nm
FOV	5'' × 5''	As large as possible
Spatial resolution	0'08	0'05
Spectral resolution R	200 000	
Spectral range	0.3 nm (for Ca II line) 0.1 nm (for Fe I line)	0.6 nm 0.2 nm
SNR	2000	
Cadence	30 s	15 s
Instrument 2	NBI	
Goal	Context information about magnetic field configuration from other spectral lines.	
	Requirement	Goal
Spectral lines	Fe I 630.2 nm, He I 1083 nm, Si I 1082.7 nm	
FOV	30'' × 30''	60'' × 60''
Spatial resolution	0'08	0'05
Spectral resolution R	50 000	
Spectral range	0.5 nm (for He I line) 0.1 nm (for Fe I line)	0.8 nm
SNR	1000	2000
Cadence	120 s	60 s
Instrument 3	BBI	
Goal	Context information.	
	Requirement	Goal
Spectral lines	G-band; Ca II H line core & line wing	
FOV	30'' × 30''	60'' × 60''
Spatial resolution	diffraction limit	
SNR	100	200
Cadence	5 s	

OP Table 4.3.1: Probing the structure of cool sunspot umbrae

Instrument 1	IFU or SP	
Goal	High-precision spectropolarimetry of darkest umbral regions.	
	Requirement	Goal
Spectral lines FOV Spatial resolution Spectral resolution R Spectral range SNR Cadence	Fe I 630.15 nm & 630.25 nm; Fe I 1565 nm; Ti I 2.2 μ m 15'' \times 15'' 0'.1 200 000 0.1 nm (for Fe I lines) 12 nm (for Ti I lines) 3000 not crucial	0'.05
Notes	The observed FOV should cover the whole umbral region.	

OP Table 4.4.1: Umbral flashes

Instrument 1	IFUs	
Goal	Spectropolarimetric observations of the umbra to study the magnetic field configuration.	
	Requirement	Goal
Spectral lines FOV Spatial resolution Spectral resolution R Spectral range SNR Cadence	Fe I 1565 nm; Ca II 854 nm 15'' \times 15'' 0'.1 200 000 0.5 nm (for He I line) 0.1 nm (for Fe I line) 2000 60 s	+He I 1083 nm 30'' \times 30'' 0'.06 0.8 nm 0.2 nm 30 s
Instrument 2	NBI	
Goal	Additional spectropolarimetric observations using other spectral lines.	
	Requirement	Goal
Spectral lines FOV Spatial resolution Spectral resolution R Spectral range SNR Cadence	H α ; He I 1083 nm 30'' \times 30'' 0'.08 50000 0.5 nm (for He I line) 0.3 nm (for H α line) 1000 60 s	60'' \times 60'' 0'.05 0.5 nm 2000 30 s
Instrument 3	BBI	
Goal	Context information.	
	Requirement	Goal
Spectral lines FOV Spatial resolution SNR Cadence	Ca II H or K line core & line wing 30'' \times 30'' diffraction limit 100 5 s	60'' \times 60'' 200

OP Table 4.5.1: Penumbral and umbral micro-jets

Instrument 1	IFUs	
Goal	Determine the magnetic topology in and around micro-jets. High-cadence spectropolarimetric observations of a small FOV.	
	Requirement	Goal
Spectral lines	Fe I 630.15 nm; Ca II 854 nm	
FOV	10'' × 10''	as large as possible
Spatial resolution	0'.06	as good as possible
Spectral resolution R	200 000	
Spectral range	0.3 nm (for Ca II line) 0.1 nm (for Fe I line)	0.6 nm
SNR	1000	2000
Cadence	10 s	5 s
Notes	Core of this OP.	
Instrument 2	NBI	
Goal	Additional spectropolarimetric observations for context information.	
	Requirement	Goal
Spectral lines	Fe I 1565 nm; He I 1083 nm; Si I 1082.7 nm	
FOV	30'' × 30''	
Spatial resolution	0'.06	
Spectral resolution R	50000	
Spectral range	0.5 nm (for He I line) 0.1 nm (for Fe I line)	0.2 nm
SNR	500	
Cadence	20 s	10 s
Instrument 3	BBI	
Goal	Observe the dynamical evolution of micro-jets. G-band or continuum is for context information, for LCT and feature tracking.	
	Requirement	Goal
Spectral lines	G-band or continuum; Ca II H or K line core	
FOV	30'' × 30''	
Spatial resolution	diffraction limit	
SNR	100	200
Cadence	1 s	0.5 s
Notes	Extreme high cadence is needed to observe the onset and evolution of micro-jets.	

OP Table 4.6.1: Structure and dynamics of light-bridges

Instrument 1	IFU	
Goal	Determine the magnetic topology in and around light bridges. High-cadence spectropolarimetric observations of a small FOV. This program aims to study the details of magneto-convection and reconnection above light bridges.	
	Requirement	Goal
Spectral lines	Fe I 630.15 nm; Ca II 854 nm	
FOV	10'' × 10''	as large as possible
Spatial resolution	0'.06	as good as possible
Spectral resolution R	200 000	
Spectral range	0.3 nm (for Ca II line) 0.1 nm (for Fe I line)	0.6 nm
SNR	1000	2000
Cadence	10 s	5 s
Notes	Core of this OP.	
Instrument 2	NBI	
Goal	Additional spectropolarimetric observations for context information.	
	Requirement	Goal
Spectral lines	Fe I 1565 nm; He I 1083 nm	
FOV	30'' × 30''	
Spatial resolution	0'.06	
Spectral resolution R	50000	
Spectral range	0.5 nm (for He I line) 0.1 nm (for Fe I line)	0.2 nm
SNR	500	
Cadence	20 s	10 s
Instrument 3	BBI	
Goal	Observe the dynamical evolution of light bridges and jets in the chromosphere above light bridges. G-band or continuum is for context information and feature tracking.	
	Requirement	Goal
Spectral lines	G-band or continuum; Ca II H or K line core; H α line core	
FOV	30'' × 30''	
Spatial resolution	diffraction limit	
SNR	100	200
Cadence	1 s	0.5 s
Notes	Extreme high cadence is needed to observe the onset and evolution of jets above light bridges. Combination with UV/EUV imaging from space (e.g. Solar Orbiter, Solar-C) allows to study the impact of these jets on the Transition Region and Corona.	

OP Table 4.7.1: Evolution of an individual penumbral filament

Instrument 1	IFU	
Goal	Spectropolarimetric observations of the umbral boundary to study the detailed processes in the magnetic field evolution.	
	Requirement	Goal
Spectral lines	Fe I 1565 nm; Ca II 854 nm	As large as possible
FOV	10'' × 10''	
Spatial resolution	0'.06	
Spectral resolution R	200 000	
Spectral range	0.3 nm (for Ca II line) 0.1 nm (for Fe I line)	
SNR	1000	
Cadence	30 s	
		0.6 nm
		0.2 nm
		2000
		15 s

Instrument 2	NBI	
Goal	Additional spectropolarimetric observations using different spectral lines.	
	Requirement	Goal
Spectral lines	Fe I 630.2 nm; He I 1083 nm	
FOV	30'' × 30''	
Spatial resolution	0'.06	
Spectral resolution R	50000	
Spectral range	0.5 nm (for He I line) 0.1 nm (for Fe I line)	
SNR	1000	
Cadence	30 s	
		2000
		15 s

Instrument 3	BBI	
Goal	Context information, for LCT and feature tracking.	
	Requirement	Goal
Spectral lines	G-band; Ca II H line core & line wing	diffraction limit
FOV	30'' × 30''	
Spatial resolution	0'.05	
SNR	100	
Cadence	5 s	
		200
		2 s

OP Table 4.8.1: Capturing the formation and decay of penumbrae

Instrument 1	IFU	
Goal	Spectropolarimetric observations of the sunspot to study the magnetic field configuration.	
	Requirement	Goal
Spectral lines	Fe I 1565 nm; Ca II 854 nm	
FOV	30'' × 30'' (Mosaic)	60'' × 60''
Spatial resolution	0'.1	0'.06
Spectral resolution R	200 000	
Spectral range	0.3 nm (for Ca II line) 0.1 nm (for Fe I line)	0.6 nm 0.2 nm
SNR	1000	2000
Cadence	180 s	30 s
Notes	The goal is to cover the full process of penumbral formation and decay lasting several hours and the main requirement will be stable conditions on these time scales. In principle, short periods of worse seeing conditions can be tolerated as long as high quality spectral profiles are acquired every 3 minutes.	

Instrument 2	NBI	
Goal	Additional spectropolarimetric observations using other spectral lines.	
	Requirement	Goal
Spectral lines	Fe I 630.2 nm; He I 1083 nm	
FOV	60'' × 60'' (Mosaic)	
Spatial resolution	0'.08	0'.05
Spectral resolution R	50 000	
Spectral range	0.5 nm (for He I line) 0.1 nm (for Fe I line)	
SNR	1000	2000
Cadence	180 s	30 s

Instrument 3	BBI	
Goal	Context information, for LCT and feature tracking	
	Requirement	Goal
Spectral lines	G-band; Ca II H line core & line wing	
FOV	60'' × 60''	
Spatial resolution	diffraction limit	
SNR	100	200
Cadence	2 s	

OP Table 4.9.1: Observations of moat flow properties and its impact on sunspot decay

Instrument 1	IFUs	
Goal	Spectropolarimetric observations of a sunspot and its surroundings to study the magnetic field configuration.	
	Requirement	Goal
Spectral lines	Fe I 1565 nm; Ca II 854 nm	
FOV	120'' × 120'' (Mosaic)	
Spatial resolution	0'.1	
Spectral resolution R	200 000	
Spectral range	0.3 nm (for Ca II line) 0.1 nm (for Fe I line)	0.6 nm 0.2 nm
SNR	1000	2000
Cadence	300 s	150 s

Instrument 2	NBI	
Goal	Additional spectropolarimetric observations using other spectral lines.	
	Requirement	Goal
Spectral lines	Fe I 630.2 nm; He I 1083 nm	
FOV	120'' × 120'' (Mosaic)	
Spatial resolution	0'.08	
Spectral resolution R	50 000	
Spectral range	0.5 nm (for He I line) 0.1 nm (for Fe I line)	
SNR	1000	2000
Cadence	300 s	150 s

Instrument 3	BBI	
Goal	Context information, for LCT and feature tracking	
	Requirement	Goal
Spectral lines	G-band; Ca II H line core & line wing	
FOV	120'' × 120'' (Mosaic)	
Spatial resolution	diffraction limit	
SNR	100	200
Cadence	5 s	

OP Table 4.10.1: Dynamic magnetic fine structure of prominences and filaments

Instrument 1	IFU	
Goal	Spectropolarimetric observations of prominences/filaments to study the evolution of the magnetic field configuration and the dynamics at the smallest spatial scales possible, with the higher cadence possible. Type of observations: time series of a 20'' × 20'' area.	
	Requirement	Goal
Spectral lines	He I 1083 nm [1]; He I 587.6 nm [2]	
FOV	15'' × 15''	as large as possible
Spatial resolution	0'.1	0'.05
Spectral resolution R	300 000	
Spectral range	[1] ±0.11 nm; [2] ±0.06 nm, velocities as large as ±30 km s ⁻¹	
SNR	10000 (in line core)	20000
Exposure time	[1] 25 s; [2] 87 s	
Cadence	2 min	as high as possible
Notes	The use of the adaptive optics on an off-limb target is necessary in case of prominence observations. High stability is required to reduce spurious signals down to 10 ⁻⁵ , in terms of the continuum intensity. Pointing error of the order of 1''	

OP Table 4.10.2: Magnetic and dynamic fine structure of prominences and filaments

Instrument 1	IFU	
Goal	Spectropolarimetric observations of prominences/filaments to study the magnetic field configuration at the smallest spatial scales possible. Type of observations: Scan of a large area	
	Requirement	Goal
Spectral lines	He I 1083 nm [1]; He I 587.6 nm [2]	
FOV	40'' × 30'' (Mosaic)	as large as possible
Spatial resolution	0'.1	0'.05
Spectral resolution R	100 000	
Spectral range	[1] ±0.11 nm; [2] ±0.06 nm, velocities as large as ±30 km s ⁻¹	
SNR	10 000 (line core)	20000
Exposure time	[1] 7 s; [2] 30 s	
Cadence	30 min (assuming 10 tiles)	as high as possible
Notes	The use of the adaptive optics on an off-limb target is necessary in case of prominence observations. High stability is required to reduce spurious signals down to 10 ⁻⁵ , in terms of the continuum intensity. Pointing error of the order of 1''.	

OP Table 4.11.1: Magnetic field properties in QS prominences and AR filaments

Instrument 1	IFU	
Goal	Spectropolarimetric observations to study the magnetic field differences between QS prominences and AR filaments. Type of observations: Scan of a large area.	
	Requirement	Goal
Spectral lines	He I 1083 nm [1]; Ca II 854 nm [2]; He I 587.6 nm [3]	as large as possible 0'05
FOV	40'' × (10 – 30)''	
Spatial resolution	0'2	20000
Spectral resolution R	100 000	
Spectral range	[1] ±0.11 nm; [2] ±0.09 nm; [3] ±0.06 nm, velocities up to ±30 km s ⁻¹	
SNR	10000	
Exposure time	[1] 2s; [2] 15 s [3] 8 s	
Cadence	0.4 - 1.25 h (~ 30'' × 30'')	
Notes	The use of the adaptive optics on an off-limb target is necessary in case of prominence observations. High stability is required to reduce spurious signals down to 10 ⁻⁵ , in terms of the continuum intensity. Pointing error of the order of 1''.	

OP Table 4.12.1: Magnetism and dynamics of tornado prominences

Instrument 1	NBI	
Goal	Spectroscopic observations to study the dynamics of tornado prominences. Type of observations: Time series of a 50'' × 50'' area. These observations are simultaneous to the spectro-polarimetric ones detailed in next table.	
	Requirement	Goal
Spectral lines	He I 1083 nm [1]; Ca II 854 nm [2]; H _α [3]; H _β [4]	as large as possible 0'05
FOV	50'' × 50''	
Spatial resolution	0'2 in the IR	0.2 km s ⁻¹
Spectral sampling	1.8×10 ⁻³ nm [1]; 1.4×10 ⁻³ nm [2]; 1.1×10 ⁻³ nm [3]; 0.8×10 ⁻³ nm [4], resolve velocities of 0.5 km s ⁻¹	
Spectral range	± 0.11 nm; ± 0.09 nm [3]; ± 0.07 nm [3]; ±0.05 nm [4], equivalent to ± 30 km s ⁻¹	
SNR	3000	
Exposure time	[1] 300 ms; [2] 2.5 s; [3] 5 s; [4] 16 s, all measured at the core of the lines	
Cadence	20 s	
Notes	The adaptive optics has to be able to lock on the limb, where the tornado prominences are best observed. High stability is required to reduce spurious signals down to 10 ⁻⁵ , in terms of the continuum intensity. Pointing error of the order of 1''.	

Instrument 2	IFU	
Goal	Spectro-polarimetric observations to study the magnetic field configuration of tornado prominences. Type of observations: time series of a 20'' × 20'' area.	
	Requirement	Goal
Spectral lines	He I 1083 nm [1]; He I 587.6 nm [2]	
FOV	20'' × 20''	as large as possible
Spatial resolution	0'.2	0'.05
Spectral resolution R	300 000	
Spectral range	[1] ±0.11 nm; [2] ±0.06 nm, velocities as large as ±30 km s ⁻¹	
SNR	10000 (line core)	20000
Exposure time	[1] 6 s [2] 23 s	
Cadence	25 s	as high as possible
Notes	The use of the adaptive optics on an off-limb target is necessary in case of prominence observations. High stability is required to reduce spurious signals down to 10 ⁻⁵ , in terms of the continuum intensity. Pointing error of the order of 1''.	

OP Table 5.3.1: Determining which coronal upflows become outflows

Instrument 1	IFU	
Goal	Determine magnetic field topology and dynamics of the photosphere and chromosphere at active region and coronal hole boundaries as a function of height. Combine with UV/EUV imaging and spectroscopic observations of TR and coronal response using Solar Orbiter EUV/SPICE and/or Solar C EUVST.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Integration time/tile Cadence	Fe I 525.02, Fe I 630, Fe I 1565 nm Ca II H ₂ 396.8, Ca II 854, He I 1083 nm 10'' × 10'' 0.05'' 2000 5 s 60 s	+Si I 1082.7 nm +Na I D ₂ As large as possible 30 s
Notes	These observations need to be co-ordinated with space platforms to enable the connection to the TR and corona to be made. For Solar Orbiter this should be done at conjunction. This OP links to OP 3.3.1 on spicule acceleration on disk, so could potentially be merged, and is very similar to DKIST SUC-61 created by Louise Harra. Appropriate Solar Orbiter SOOPs, as listed in SUC-61 are L_BOTH_HIRES+LCAD_CH_Boundary_Expansion; L_SMALL_HIRES_HCAD_SlowWindConnection; L_SMALL_HIRES_HCAD_Fast_Wind; L_SMALL_HIRES_HCAD_SlowWindConenction; R_SMALL_HIRES_MCAD_PolarObservations. EST can potentially provide greater multi-height coverage.	
Instrument 2	BBI	
Goal	Detect small-scale photospheric and chromospheric activity. Provide context information on the surrounding AR/CH.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Cadence	G-band Ca II H line core & line wing 60'' × 60'' Diffraction limit > 100 5 s	+Ca II 854 nm, H α 120'' × 120'' 1 s
Notes	See above for co-ordination requirements.	

OP Table 5.4.1: Measuring the photospheric and chromospheric dynamics before flares

Instrument 1	IFU	
Goal	Determine magnetic field topology and dynamics of the photosphere and chromosphere within the vicinity of the neutral line in an active region as a function of height. Combine with UV/EUV imaging and spectroscopic observations of TR and coronal response using Solar Orbiter EUI/SPICE and/or Solar C EUVST.	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 630, Fe I 1565 nm	+Si I 1082.7 nm
Chromosphere	Ca II H 396.8, Ca II 854, He I 1083 nm	+Na I D ₂
FOV	30'' × 30''	As large as possible
Spatial resolution	0.05''	
SNR	2000	
Integration time/tile	10 s	
Cadence	100 s	30 s
Notes	These observations would have to be a TOO but need to be co-ordinated with space platforms to enable the connection to the TR and corona to be made. For Solar Orbiter this could be done either at conjunction or quadrature, with quadrature providing a 3D view of the overlying corona, particularly if combined with Solar C EUVST. Relevant SOOPs: L_FULL_HRES_HCAD_Coronal_Dynamics; R_SMALL_HRES_HCAD_AR_Dynamics; R_SMALL_HRES_HCAD_RSburst; L_FULL_HRES_HCAD_Eruption_Watch	
Instrument 2	BBI	
Goal	Detect small-scale photospheric and chromospheric activity around the neutral line, including flux emergence, cancellation. Provide context information on the surrounding AR.	
	Requirement	Goal
Photosphere	G-band	
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm, H α
FOV	60'' × 60''	120'' × 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	5 s	1 s
Notes	See above for co-ordination requirements.	

OP Table 5.5.1: Rising spicule signatures

Instruments	BBI, NBI, IFU		
Goal	High-resolution spectroscopy and spectropolarimetry combined with G-band imaging to determine the associated rising spicule signatures.		
	Requirement	Goal	
Photosphere	G-band, Fe I 630.2	30''×30''	
Chromosphere	H α , Ca II 8542, He I 10830		
FOV	20''×20''		
Spatial resolution	0.04''		
SNR	2000		
Integration time	0.2 sec		0.1 sec
Cadence	2 sec		1 seconds
Notes	The short lifetime of RBEs/RREs requires the highest cadence possible.		

OP Table 5.6.1: Ellerman bombs in the lower solar atmosphere

Instruments	BBI, NBI, IFU			
Goal	High-resolution spectro-polarimetry to determine signatures of EBs in the lower solar atmosphere			
	Requirement	Goal		
Photosphere	Ca II 854.2, H α , He I 1083.0, Ca II K	30''×20''		
Chromosphere				
FOV				
Spatial resolution				
SNR				
Integration time			1 sec	0.5 sec
Cadence			10 second	5 seconds
Notes	High-cadence is essential. Spectral range (km/s) \pm 100 for each line			

OP Table 6.1.1: Thermal structure of flare: line observations

Instrument 1		NBI	
Goal	Observe spectral lines formed at different heights in the chromosphere, from which temperature profile with column mass can be determined.		
	Requirement	Goal	
Photosphere	Fe I 543.4	+H α 60'' \times 60'' 500 1s	
Chromosphere	Ca II H 396.9 nm, Ca II 854 nm, He I 1083 nm		
Wavelength samples	0.01nm spectral resolution, across 0.3 nm spectral range		
FOV	40'' \times 40''		
Spatial resolution	0':04		
SNR	200		
Cadence	5 s		
Notes	The focus here is only on I, hence low SNR requirement. To capture dynamical changes need to prioritise cadence. This observing program requires stable observations with AO locked, at different μ values. Photospheric line chosen as it is formed quite high in photosphere, bearing in mind likely range in height of strong perturbations caused by flare heating.		

Instrument 2		BBI	
Goal	High temporal and spatial resolution context observations of the flare ribbons.		
	Requirement	Goal	
Photosphere	G-band	+CN Bandhead +H α 120'' \times 120'' 500 0.1 s	
Chromosphere	Ca II H line core and wing		
Wavelength samples	n/a		
FOV	60'' \times 60'' (100'' \times 100'')		
Spatial resolution	0':015		
SNR	> 100		
Cadence	1 s		
Notes	For FOV, the tradeoff is between having a good chance of catching a flare (FOV) and highest possible spatial and temporal resolutions. The programme requires stable observations with AO locked at different μ values.		

OP Table 6.2.1: Thermal structure of flare: continuum observations

Instrument 1	BBI	
Goal	High temporal and spatial resolution observations of a flare at several points in the continuum, to track the evolution of energy input via the hydrogen continuum emission (“white-light flare”).	
	Requirement	Goal
Photosphere Chromosphere	Paschen Continuum, G-band Continuum (Blue Arm, Channel 2) Brackett Continuum, H α continuum (Red Arm, Channel 3)	120'' \times 120''
Wavelength samples	n/a	
FOV	60'' \times 60'' (100'' \times 100''?)	
Spatial resolution	0''.015	
SNR	200	
500 Cadence	0.2 s	
Notes	For FOV, the tradeoff is between having a good chance of catching a flare (FOV) and highest possible spatial and temporal resolutions (the latter is necessary to catch sub-second ionisation evolution). The programme requires stable observations with AO locked at different μ values.	

OP Table 6.2.2: Thermal structure of flare: detection of the H Paschen Jump

Instrument 1	SP or NBI in spectroscopic mode	
Goal	Identification of the H Paschen jump at 815.1nm in flares.	
	Requirement	Goal
Photosphere Chromosphere Wavelength samples FOV Spectral range Imaging or polarimetry Spectral resolution Spatial resolution SNR Cadence	n/a 60'' × 60'' 800-810, 830-840 & 890-900 nm Narrowband images 10nm < 3''0 200 10 s	120'' × 120'' 800-810 & 890-900 nm Narrowband images < 10nm < 1''0 1000 1s
Notes	As an alternative to a spectrometer, 2 or 3 narrowband-filter imager could be used, capturing the continuum levels between the Ca and H Paschen spectral lines in the range 800-900nm. The filters would have to be designed in a way to avoid stray light from other spectral ranges, especially from regions densely populated by spectral lines.	

Instrument 2	NBI in spectroscopic mode	
Goal	Identification of the H Paschen jump at 815.1nm in flares.	
	Requirement	Goal
Photosphere Chromosphere Wavelength samples FOV Spectral range Spatial resolution SNR Cadence	Paschen Continuum (Blue Arm, Channel 2) Brackett Continuum (Red Arm, Channel 3) n/a 60'' × 60'' 800-850 nm < 3''0 200 10 s	120'' × 120'' 800-900 nm < 1''0 1000 1s
Notes	Ideally, a high-resolution imaging spectrometer in the range 800-900nm, with high spatial/temporal resolutions, should provide enough information about the continuum emission along flare ribbons to track the evolution of this emission and allow the detection of the Paschen jump (or lack thereof!). This setup would also provide observations of the Ca and H Paschen lines in the range.	

OP Table 6.3.1: Velocity structure of the flaring atmosphere

Instrument 1	NBI	
Goal	Observations to track flows in the chromosphere at a variety of optical depths.	
	Requirement	Goal
Photosphere Chromosphere Wavelength samples FOV Spatial resolution SNR Cadence	Fe I 557.6 nm H α 656.3 nm Ca II 854 nm 50mA spectral resolution, across 0.3 nm spectral range in all lines 40'' \times 40'' 0'.04 100 0.5 - 1 s	+ Fe I 543.4 nm +Ca II H 3969 nm 120'' \times 120'' 200 0.1s
Notes	The focus here is only on I, hence low SNR requirement. Based on observations e.g. with IRIS and in H α , can expect large flows, with also an optically thin chromospheric component possible in the nominally 'photospheric' lines. So a wide spectral range is necessary. This observing program requires stable observations with AO locked, at different μ values.	
Instrument 2	BBI	
Goal	High temporal and spatial resolution context observations of the flare ribbons.	
	Requirement	Goal
Photosphere Chromosphere Wavelength samples FOV Spatial resolution SNR Cadence	G-band Ca II H line core and wing n/a 60'' \times 60'' 0'.015 > 100 1 s	+CN Bandhead +H α 120'' \times 120'' 0.1 s
Notes	For FOV, the tradeoff is between having a good chance of catching a flare (FOV) and highest possible spatial and temporal resolutions. The programme requires stable observations with AO locked at different μ values.	

OP Table 6.4.1: Diagnostics for non-thermal particles

Instrument 1	NBI	
Goal	Linear polarization measurements in strong spectral lines, in a search for evidence of impact polarization generated by low-energy proton beams.	
	Requirement	Goal
Photosphere	n/a	
Chromosphere	H α 656.3 nm (I only), Ca II 854 nm, He I 1083 nm	+Na I 589
Wavelength samples	0.02nm spectral resolution, across 0.1 nm spectral range in both H α and Ca II 854 nm (i.e. fast but narrow spectral scans in line core)	
FOV	60'' \times 60''	120'' \times 120''
Spatial resolution	0'.04	
SNR	1000	500
Cadence	5 s	2s
Notes	The focus here is on I, Q, U in the core of the spectral lines only to try and isolate the upper part of the chromosphere where the particle beams would reach. FP is needed to capture different parts of a flare ribbon under different conditions. I, Q, U to be taken in as quick succession as possible. Note, it is expected that the spectral lines will brighten during a flare so short exposures should be possible. This observing program requires stable observations with AO locked.	
Instrument 2	BBI	
Goal	Detect flare ribbons at chromospheric heights. Context image information.	
	Requirement	Goal
Photosphere		
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm, H α
Wavelength samples		
FOV	60'' \times 60''	120'' \times 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	1 s	0.1 s
Notes		

OP Table 6.5.1: QPPs in different phases of flares

Instrument 1	NBI	
Goal	Detect QPPs during flares and verify whether those detected in different phases of the flare have different characteristics. Determine the magnetic field configuration of the flaring site, estimate magnetic fluxes and search for any variation correlated with the flare occurrence.	
	Requirement	Goal
Photosphere	Fe I 1565	+Si I 1082.7 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm, H α +He I 1083 nm	
Wavelength samples	15	20 for chromospheric lines
FOV	40'' \times 40''	120'' \times 120''
Spatial resolution	0'.05	
SNR	500	
Cadence	3 s	1 s
Notes		
Instrument 2	BBI	
Goal	Detect flare signatures at photospheric and chromospheric heights. Context information.	
	Requirement	Goal
Photosphere	G-band	+CN bandhead
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm, H α
Wavelength samples		
FOV	100'' \times 100''	120'' \times 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	1 s	0.1 s
Notes		

OP Table 6.5.2: QPPs and oscillatory reconnection

Instrument 1	IFU	
Goal	Determine magnetic field strength and topology of both pre-existing and emerging loops as a function of height. Detect signatures of (oscillating) magnetic reconnection due to the interaction of emerging fields with pre-existing flux in photosphere and chromosphere.	
	Requirement	Goal
Photosphere	Fe I 630	As large as possible
Chromosphere	Ca II 854, He I 1083 nm	
FOV	30'' × 30''	
Spatial resolution	0.1	
SNR	2000	
Integration time/tile	1 -3 s	
Cadence	30 s	10 s
Notes	The full active region FOV should be obtained by tiling the IFUs as appropriate.	
Instrument 2	BBI	
Goal	Detect flare emission in photosphere and chromosphere. Search for QPPs correlating with reconnection events.	
	Requirement	Goal
Photosphere	G-band	+CN bandhead
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm, H α
FOV	60'' × 60''	120'' × 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	1 s	0.1 s
Notes		

OP Table 6.6.1: Sunquakes initiated by impulsive heating during flares

Instrument 1	IFU	
Goal	Verify whether sunquakes are due to impulsive heating by means of simultaneous observations at different wavelengths: search for any time delay between photospheric and chromospheric emission that might shed light on the heating and energy transport mechanisms. Try to discriminate between the possible impulsive heating mechanisms: thick target, backwarming or wave heating (no polarimetry).	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 630, Fe I 1565 nm, Si I 1082.7 nm	+ Mn I 1526
Chromosphere	Mg I b ₂ 517.3, Ca II 854, He I 1083 nm	+Ca II H
FOV	60'' × 60''	As large as possible
Spatial resolution	0.1	
SNR	1000	
Integration time/tile	0.1 s	
Cadence	5 s	1 s
Notes	The full active region FOV should be obtained by tiling the IFUs as appropriate,	
Instrument 2	BBI	
Goal	Detect flare emission in photosphere and chromosphere. Context information.	
	Requirement	Goal
Photosphere	G-band	+CN bandhead
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm, H α
FOV	100'' × 100''	120'' × 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	1 s	fraction of s
Notes		

OP Table 6.6.2: Sunquakes initiated by changes in the Lorentz force during flares

Instrument 1	IFU	
Goal	Determine magnetic field configuration before and during flares. Detect signatures of sunquakes at photospheric and chromospheric levels.	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 630, Fe I 1565 nm	As large as possible
Chromosphere	Ca II 854, He I 1083 nm	
FOV	60'' × 60''	
Spatial resolution	0.1	
SNR	2000	
Integration time/tile	2 - 3 s	
Cadence	1.5 m	
Notes	The full active region FOV should be obtained by tiling the IFUs as appropriate.	
Instrument 2	BBI	
Goal	Detect flare emission in photosphere and chromosphere during sunquakes. Context information	
	Requirement	Goal
Photosphere	G-band	+CN bandhead
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm, H α
FOV	100'' × 100''	120'' × 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	1 s	fraction of s
Notes		

OP Table 6.7.1: Changes in the magnetic field configuration during/after flares

Instrument 1	NBI	
Goal	Detect changes in the sunspot configuration during/after flares. Determine the magnetic field configuration of the flaring site and search for any variation in the vector magnetic field correlated with the flare occurrence. Infer the time sequence leading to the magnetic field - sunspot variation.	
	Requirement	Goal
Photosphere	Fe I 1565	+Si I 1082.7 nm
Chromosphere	Mg I b ₂ 517.3, Ca II 854 nm, H α	+He I 1083 nm
Wavelength samples	15	20 for chromospheric lines
FOV	40'' \times 40''	120'' \times 120''
Spatial resolution	0'.1	
SNR	1000	
Cadence	3 - 5 s	1 s
Notes		
Instrument 2	IFU	
Goal	Context magnetic field at photospheric and chromospheric heights.	
	Requirement	Goal
Photosphere	Fe I 525.02, Fe I 630 nm	
Chromosphere	He I 1083 nm	
FOV	60'' \times 60''	As large as possible
Spatial resolution	0'.1	0'.05
SNR	2000	
Integration time/tile	1.3 s	
Cadence	40 s	10 s
Notes	The full active region FOV should be obtained by tiling the IFUs as appropriate. It may be desirable also to bin e.g. 4 \times 4 pixels to obtain higher SNR and hence increase the cadence.	
Instrument 3	BBI	
Goal	Detect flare signatures at photospheric and chromospheric heights. Context information on sunspots morphology before, during and after the flare.	
	Requirement	Goal
Photosphere	G-band	+CN bandhead
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm, H α
Wavelength samples		
FOV	60'' \times 60''	120'' \times 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	1 s	fraction of s
Notes		

OP Table 6.7.2: Determine magnetic helicity accumulation, dip and shear angle in flaring active regions

Instrument 1	SP	
Goal	Infer the magnetic field configuration and the photospheric flow field to determine the magnetic helicity accumulation in the active region. Determine the dip and shear angle along the main polarity inversion line.	
	Requirement	Goal
Photosphere	Fe I 1565	+Si I 1082.7 nm
Chromosphere	Ca II 854 nm, H α	+He I 1083 nm
Wavelength samples		
FOV	60'' \times 60''	120'' \times 120''
Spatial resolution	0'.05	
SNR	1000	
Cadence	140 s	
Notes	Raster scans with slit parallel to the magnetic neutral line.	
Instrument 2	BBI	
Goal	Detect flare signatures at photospheric and chromospheric heights. Context information on the active region morphology before, during and after the flare.	
	Requirement	Goal
Photosphere	G-band	+CN bandhead
Chromosphere	Ca II H line core & line wing	+Ca II 854 nm, H α
Wavelength samples		
FOV	60'' \times 60''	120'' \times 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	1 s	0.1 s
Notes		

OP Table 6.8.1: Filaments in flaring active regions

Instrument 1	SP	
Goal	Infer changes in the vector magnetic field and flows during the eruption of the filament. Scientific goal: identify the mechanisms which lead to flaring around filaments and eventually to the eruption of the filament. Predict future filament eruptions.	
	Requirement	Goal
Photosphere	Si I 1082.7	+Ca I 1083.9 nm
Chromosphere	He I 1083.0, Ca II 854.2	
FOV	10'' × 40''	20'' × 60''
Spatial resolution	0'.1	0'.05
Spectral range	±120 km s ⁻¹	±150 km s ⁻¹
SNR	1000	3000
Integration time		
Cadence	1 s	0.5 s
Notes	Two observing strategies: (1) Sit and stare: slit fixed and oriented along the filament. (2) Small raster scans (~10'') with slit parallel to filament covering more of the flaring area next to the filament.	
Instrument 2	SP	
	Requirement	Goal
Photosphere		+Fe I 630.2
Chromosphere	H α (only I), Na I 589, Ba II 455	
FOV	30'' × 30''	60'' × 60''
Spatial resolution	0'.1	0'.05
Spectral range	±80 km s ⁻¹	±120 km s ⁻¹
SNR	1000	3000
Integration time		
Cadence	5 s	1 s
Notes	Two observing strategies: (1) Sit and stare: slit fixed and oriented along the filament. (2) Small raster scans (~10'') with slit parallel to filament covering more of the flaring area next to the filament.	

OP Table 6.9.1: CME sources and temporal relation with flares

Instrument 1	SP	
Goal	Search for any brightening or flaring emission which could be spatially and temporally correlated with a CME. Infer the magnetic field configuration of the flux rope involved in the eruption leading to the coronal mass ejection.	
	Requirement	Goal
Photosphere	Fe I 1565	+Si I 1082.7 nm
Chromosphere	Ca II 854 nm, H α , Ba II 455	+He I 1083 nm
Wavelength samples		
FOV	60'' \times 60''	120'' \times 120''
Spatial resolution	0'.05	
SNR	2000	
Cadence	140 s	30 s
Notes	Raster scans with slit parallel to the magnetic flux rope axis to look for plasma motions and for brightenings in the surroundings.	
Instrument 2	BBI	
Goal	Detect flare signatures at photospheric and chromospheric heights. Context information on the active region morphology before, during and after the flare - CME occurrence.	
	Requirement	Goal
Photosphere	G-band	+CN bandhead
Chromosphere	Ca II H line core & line wing	H α
Wavelength samples		
FOV	120'' \times 120''	As large as possible
Spatial resolution	0'.1	
SNR	> 100	
Cadence	1 s	0.5 s
Notes	These observations should be complemented with data acquired by coronagraphs.	

OP Table 7.1.1: Waves in prominences observed in neutral and ionized spectral lines

Instrument 1	IFU in spectroscopic mode	
Goal	Very high-cadence velocity measurements strictly simultaneously, target on prominence. 2D field of view is essential to align spectral lines and correct for differential refraction effects.	
	Requirement	Goal
Photosphere Chromosphere	H α , Ca II 854.2, Ca II K , HI 397, He I 1083	+ HI 434, He I D ₃ 587.6
FOV	10'' \times 40''	20'' \times 80''
Spatial resolution	0.1''	0.05''
SNR	1000	2000
Integration time	0.5 sec	
Cadence	2 seconds	1 second
Notes	Extremely high-cadence in velocity is essential.	
Instrument 2	IFU with polarimetry	
Goal	Context magnetic field in prominence. Possibly, using the same IFU spectropolarimeter for both kinds of measurements and highest cadence possible (see above), then adding up polarimetric data to increase s/n.	
	Requirement	Goal
Photosphere Chromosphere	Ca II 854.2, He I 1083	
FOV	10'' \times 40''	20'' \times 80''
Spatial resolution	0.1''	0.05''
SNR	1000	2000
Integration time	1 sec	
Cadence	4 seconds	
Notes	Same field of view as for velocity measurements.	

OP Table 7.1.2: Prominence-corona transition region (PCTR) dynamics

Instrument 1	IFU in spectroscopic mode	
Goal	Very high spatial resolution velocity measurements strictly simultaneously, target on prominence. 2D field of view is essential to align spectral lines and correct for differential refraction effects.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Integration time Cadence	H α , Ca II 854.2, He I 1083 10'' \times 40'' 0.04'' 1000 1 second 4 seconds	20'' \times 80'' 2000 0.5 second 1 second
Notes	High spatial resolution is the priority	

OP Table 7.1.3: Draining of neutral material from prominences and filaments

Instrument 1	IFU with polarimetry	
Goal	High spatial resolution maps of filaments and prominences at different positions across the disc and at the limb. 2D field of view and polarimetry.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Integration time Cadence	H α , Ca II 854.2, He I 1083 40'' \times 100'' 0.04'' 2000 3 second	+ Ca I 422.7
Notes	High spatial resolution is the priority	

OP Table 7.2.1: Dynamics of spicules observed in neutral and ionized spectral lines

Instrument 1	IFU with polarimetry	
Goal	Very high-cadence velocity measurements strictly simultaneously, target on spicules. 2D field of view is essential to align spectral lines and correct for differential refraction effects. Spectropolarimetry in Ca II 854.2 and He I 1083.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Integration time Cadence	H α , Ca II 854.2, He I 1083 10'' \times 40'' 0.1'' 1000 1 sec 2 seconds	0.05'' 2000 0.5 sec 1 second
Notes	Extremely high-cadence in velocity is essential.	

OP Table 7.2.2: Alignment between the magnetic field in fibrils and disc counterparts of spicules

Instrument 1	IFU with polarimetry	
Goal	Very deep 2D Spectropolarimetry in Ca II 854.2. IFU to ensure full spectral line simultaneously.	
	Requirement	Goal
Photosphere Chromosphere FOV Spatial resolution SNR Integration time Cadence	Ca II 854.2 10''×40'' 0.1'' 10 ⁴ as long as needed for S/N not important	0.05'' 10 ⁴
Notes	Spectral integrity (IFU) comes at the expense of small FOV. Programs using FP should also be tried.	

OP Table 7.3.1: Neutral and ionized material Evershed flow in sunspots

Instrument 1	IFU in spectroscopic mode	
Goal	Very high-cadence velocity measurements strictly simultaneously, target on sunspot located at the limb	
	Requirement	Goal
Photosphere	Fe II 457.634, Fe I 457.422, Fe II 465.698, Fe I 465.758, Fe I 519.606, Fe II 519.758, Fe I 519.794, Fe I 519.871, Fe II 523.462, Fe I 523.620, Fe II 651.608, Fe I 651.837	
Chromosphere		
FOV	20''×20''	40''×40''
Spatial resolution	0.06''	
SNR	2000	
Integration time	4 sec	2 sec
Cadence	1 minute	30 seconds
Notes	High-cadence and simultaneous detection of all lines is essential.	

Instrument 2	SP with polarimetry	
Goal	Sunspot magnetic field vector accompanying velocity measurements	
	Requirement	Goal
Photosphere	Fe I 1564.8	
Chromosphere		
FOV	20''×20''	40''×40''
Spatial resolution	0.06''	
SNR	2000	
Integration time	0.5 sec	
Cadence	1 minute	30 seconds
Notes		

OP Table 7.4.1: Observation of magnetic swirls under 2-fluid condition

Instrument 1	IFU with polarimetry	
Goal	Extremely high cadence measurements of velocities in the photosphere and chromosphere. Spectropolarimetry in Fe I 630 and Ca II 854.2 at the same high cadence of velocity measurements. The polarimetric data will be averaged to increase s/n and determine magnetic field strength.	
	Requirement	Goal
Photosphere	Fe I 630	
Chromosphere	H α , Ca II 854.2	
FOV	20''×20''	30''×30''
Spatial resolution	0.04''	
SNR	500	1000
Integration time	0.5 sec	0.2 sec
Cadence	2 sec	1 sec
Notes	2D field of view and simultaneous measurements in all three spectral lines.	

OP Table 7.5.1: Observations of reconnection and plasmoids in partially ionized plasma

Instrument 1	IFU	
Goal	Detect spectroscopic signatures of plasmoids/evolution of tearing mode instability in solar flares, with accompanying magnetic field information	
	Requirement	Goal
Photosphere	Fe I 630	He I 1083 nm As large as possible 5 s
Chromosphere	Ca II 854	
FOV	10'' × 10''	
Spatial resolution	0.1''	
SNR	1000	
Integration time/tile	2 s	
Cadence	10 s	
Notes	You ideally want context imaging and spectroscopy in the TR and the corona as well, so co-ordination with Solar C EUVST is highly desirable.	
Instrument 2	BBI	
Goal	Detect fast moving blobs in the chromosphere that are consistent with being plasmoids	
	Requirement	Goal
Chromosphere	Ca II H line core & line wing, H α line core & line wing	+Ca II 854 nm
FOV	60'' × 60''	120'' × 120''
Spatial resolution	Diffraction limit	
SNR	> 100	
Cadence	1 s	fraction of s
Notes		

OP Table 7.5.2: Wave damping by ion-neutral friction as a possible cause of flare chromosphere heating, wight-light flares, and sunquakes

Instrument 1	IFU	
Goal	Measure oscillations of the magnetic field to detect the presence and the damping of Alfvén waves propagating downwards during a flare. Detect velocities and abundances of ionized and neutral species in the same region.	
	Requirement	Goal
Photosphere	Fe I 630 nm	
Chromosphere	Ca II 854.2, He I 1083 nm	
FOV	60'' × 60''	120'' × 120''
Spatial resolution	0.05''	0.04''
SNR	2000	
Integration time/tile	0.5 s	0.2 s
Cadence	2 s	1 s
Notes		
Instrument 2	BBI	
Goal	Detect flare signatures at photospheric and chromospheric heights. Context information on flare emission (chromospheric heating and evaporation, WL emission, sunquakes signatures).	
	Requirement	Goal
Photosphere	G-band	CN bandhead
Chromosphere	H α , Ca II H line core and wings	
Wavelength samples		
FOV	60'' × 60''	120'' × 120''
Spatial resolution	0.05''	
SNR	> 100	
Cadence	2 s	1 s
Notes		

OP Table 8.1.1: Spatial fluctuations of scattering polarization in Sr I 4607 Å

Instrument	NBI (Indiv. obs.)	
Goal	Detection of spatial fluctuations at sub-granular scales of scattering polarization in Sr I 4607 Å	
	Requirement	Goal
Wavelengths	Sr I 4607.4 Å, Fe I 4607.6 Å	
Polarimetry	Full Stokes	
Polarim. sensitivity	$3.3 \cdot 10^{-4}$ ($3\text{-}\sigma$ detect. of fluct. of 10^{-3})	
Spectral FOV	1 Å	
Spectral FWHM ⁽¹⁾	25 mÅ	20 mÅ
Spectral samples	3 samples: Sr I line center, Fe I line center, continuum	7 samples: 3 Sr I, 3 Fe I, 1 continuum
Spatial FOV	10'' × 10''	
Spatial sampling ⁽²⁾	0.04''/pix	0.04''/pix
Cadence ⁽²⁾	4 s	4 s
Notes	Stable observations with AO at different limb distances required, between $\mu = 1.0$ and $\mu = 0.1$ The above values for spatial sampling and cadence refer to disk center. For other limb distances, the values have to be scaled according to limb darkening.	

⁽¹⁾ For the approximate computation of spatial sampling and cadence as in this table, it is assumed that the FWHM of the NB transmission profile is equal to its equivalent width.

⁽²⁾ The values correspond to the optimum spatial sampling and cadence (cf. Appendix A.1), derived from the expected photon flux at line center (cf. EST photon budget), a solar evolution speed of 7 km/s (OP 8.1.1, 8.1.2, 8.1.3), 25 km/s (OP 8.2.1, 8.3.1), and 50 km/s (OP 8.2.2, 8.2.3), respectively, and the required spectral resolution and polarimetric sensitivity. An ideal balanced polarimetric efficiency $\varepsilon = 1/\sqrt{3}$ is assumed for Stokes Q , U , and V . The values have been computed using the photoncount code by J. Leenaarts, which also implements the equations of Appendix A.1.

OP Table 8.1.2: Spatial fluctuations of scattering polarization in C₂ molecular lines at 5140 Å

Instrument	SP (Indiv. obs.)	
Goal	Detection of spatial fluctuations at granular scales of scattering polarization in C ₂ molecular lines at 5140 Å	
	Requirement	Goal
Wavelengths	C ₂ 5140 Å	
Polarimetry	Full Stokes	
Polarim. sensitivity	$3.3 \cdot 10^{-5}$ ($3\text{-}\sigma$ detect. of fluct. of 10^{-4})	$1.7 \cdot 10^{-5}$ ($3\text{-}\sigma$ detect. of fluct. of $5 \cdot 10^{-5}$)
Spectral FOV	3 Å	
Spectral sampling	10 mÅ / pix	
Spatial FOV	10'' × 10''	
Spatial sampling ⁽¹⁾	0.23''/pix	0.35''/pix
Cadence ⁽¹⁾	23 s	36 s
Notes	Stable observations with AO at different limb distances required, between $\mu = 1.0$ and $\mu = 0.1$. The above values for spatial sampling and cadence refer to disk center. For other limb distances, the values have to be scaled according to limb darkening.	

⁽¹⁾ Same as footnote (2) in OP Table 8.1.1.

OP Table 8.1.3: Spatial fluct. of scattering polarization in the Ti I multiplet around 4530 Å

Instrument	SP (Indiv. obs.)	
Goal	Detection of spatial fluctuations at granular scales of scattering polarization in the Ti I $a^5F - y^5F^o$ multiplet around 4530 Å	
	Requirement	Goal
Wavelength	4530 Å	
Polarimetry	Full Stokes	
Polarim. sensitivity	$3.3 \cdot 10^{-5}$ ($3\text{-}\sigma$ detect. of fluct. of 10^{-4})	
Spectral FOV	10 Å	
Spectral sampling	10 mÅ / pix	
Spatial FOV	10'' × 10''	
Spatial sampling ⁽¹⁾	0.45'' / pix	
Cadence ⁽¹⁾	47 s	
Notes	Stable observations with AO at different limb distances required, between $\mu = 1.0$ and $\mu = 0.1$. The above values for spatial sampling and cadence refer to disk center. For other limb distances, the values have to be scaled according to limb darkening.	

⁽¹⁾ Same as footnote (2) in OP Table 8.1.1.

OP Table 8.2.1: The Ca I 4227 Å resonance line

Instrument	SP (Indiv. obs.)	
Goal	Detecting spatial and temporal variations of the scattering polarization signal of Ca I 4227 Å from the core to the wings	
	Requirement	Goal
Wavelength	4227 Å	
Polarimetry	Full Stokes	
Polarim. sensitivity	$3.3 \cdot 10^{-4}$ ($3\text{-}\sigma$ detect. of fluct. of 10^{-3})	
Spectral FOV	6 Å	
Spectral sampling	10 mÅ / pix	
Spatial FOV	1' × 1'	
Spatial sampling ⁽¹⁾	0.28'' / pix	
Cadence ⁽¹⁾	8 s	
Notes	Stable observations with AO at different limb distances required, between $\mu = 1.0$ and $\mu = 0.1$. The above values for spatial sampling and cadence refer to disk center. For other limb distances, the values have to be scaled according to limb darkening.	

⁽¹⁾ Same as footnote (2) in OP Table 8.1.1.

OP Table 8.2.2: The Ca II H & K resonance lines

Instrument	SP (Indiv. obs.)	
Goal	Detecting spatial fluctuations of scattering polarization in the line core and wings of the Ca II K line.	
	Requirement	Goal
Wavelength	3934 Å	
Polarimetry	Full Stokes	
Polarim. sensitivity	$3.3 \cdot 10^{-4}$ ($3\text{-}\sigma$ detect. of fluct. of 10^{-3})	
Spectral FOV	10 Å	
Spectral sampling	10 mÅ / pix	
Spatial FOV	1' × 1'	
Spatial sampling ⁽¹⁾	0.53 "/ pix	
Cadence ⁽¹⁾	7.8 s	
Notes	Stable observations with AO at different limb distances required, between $\mu = 1.0$ and $\mu = 0.1$. The above values for spatial sampling and cadence refer to disk center. For other limb distances, the values have to be scaled according to limb darkening.	

⁽¹⁾ Same as footnote (2) in OP Table 8.1.1.

OP Table 8.2.3: The Ca II IR triplet

Instrument	SP (Simult. obs.)	
Goal	Simultaneous observations of the 3 spectral lines of the Ca II IR triplet	
	Requirement	Goal
Wavelength	8498, 8542, 8662 Å	
Polarimetry	Full Stokes	
Polarim. sensitivity	$1 \cdot 10^{-4}$	
Spectral FOV	5 Å	
Spectral sampling	25 mÅ / pix	
Spatial FOV	1' × 1'	
Spatial sampling ⁽¹⁾	0.5 "/ pix	
Cadence ⁽¹⁾	7 s	
Notes	The spectral FOV refers to each line of the Ca II IR triplet separately and shall be centered on the respective line. Stable observations with AO at different limb distances required, between $\mu = 1.0$ and $\mu = 0.1$. The above values for spatial sampling and cadence refer to disk center. For other limb distances, the values have to be scaled according to limb darkening.	

⁽¹⁾ Same as footnote (2) in OP Table 8.1.1.

OP Table 8.3.1: The physics and diagnostic potential of the Na I D₁ and D₂ lines

Instrument	SP (Indiv. obs.)	
Goal	Investigate spatial and temporal variability of the “enigmatic” scattering polarization signal of the Na I D ₁ line	
	Requirement	Goal
Wavelength	5896 Å	
Polarimetry	Full Stokes	
Polarim. sensitivity	$3.3 \cdot 10^{-5}$ (3- σ detect. of fluct. of 10^{-4})	$1.7 \cdot 10^{-5}$ (3- σ detect. of fluct. of $5 \cdot 10^{-5}$)
Spectral FOV	10 Å	
Spectral sampling	10 mÅ / pix	5 mÅ / pix
Spatial FOV	1' \times 1'	
Spatial sampling ⁽¹⁾	0.9 "/ pix	1.8 "/ pix
Cadence ⁽¹⁾	27 s	53 s
Notes	Stable observations with AO at different limb distances required, between $\mu = 1.0$ and $\mu = 0.1$. The above values for spatial sampling and cadence refer to disk center. For other limb distances, the values have to be scaled according to limb darkening.	

⁽¹⁾ Same as footnote (2) in OP Table 8.1.1.

Part III

Discussion and Requirements

11 What are the particular strengths of the EST design?

11.1 EST preliminary design study 2008 - 2011

The conceptual design that was developed during the conceptual design study 2008 - 2011 defines EST to be a 4 m telescope. To assure highest polarimetric accuracy and compactness, it is carried out as an on-axis telescope mounted on an altitude-azimuth frame, with polarimetrically compensated transfer optics, i.e., which is free of introducing net polarisation to the light beam. The rotating transfer optics also compensates for image rotation such that the Coudé lab does not need to be rotated. This allows for a large room for the suite of EST science instruments.

On-axis versus off-axis: Both off-axis and on-axis designs were studied, and both designs have advantages and disadvantages. The off-axis design has the advantage of a clear aperture, such that spiders and obscurations can be avoided. This avoids stray-light and results in an ideal point-spread function. Spiders and a central obscuration are a necessity for on-axis telescopes. They cause the point-spread function of the telescope to have extended ‘wings’. While this can be corrected in terms of spatial resolution, it decreases the signal-to-noise ratio. It should be said that this negative effect is small compared to other aberration effects that are typically introduced by the train of telescope mirrors. The spider of an on-axis telescope rotates on the wavefront sensor and the measurement of the distorted wave front becomes complicated.

Off-axis telescopes at a given aperture are larger and heavier than on-axis telescopes. On-axis telescopes being more compact and having less weight can be integrated into higher towers, i.e., further away from ground-layer turbulence dominating the seeing, hereby improving the image quality. In contrast to off-axis telescopes, on-axis telescopes can be constructed such that the optical beam is polarimetrically compensated, thereby facilitating the calibration of polarimetric measurements. Another advantage of an on-axis telescope is that – at the same aperture – their cost is considerably smaller than for an off-axis telescope. Based on these reasons, the choice of an on-axis system was preferred during the conceptual design study.

Open versus closed dome: With the open design of an on-axis telescope, EST will have less weight than a telescope with a dome. EST will therefore allow for a high tower which elevates the telescope as high as possible above ground-layer seeing. As this is considered to be a major advantage, a closed dome should be avoided if possible.

Post focus instrumentation and light distribution: In the EST conceptual design from 2011, the post-focus instrument suite consists of

- 5 narrow-band imagers (NBIs) with wavelength ranges (WRs) in nm between 390 – 500 (WR1), 500 – 620 (WR2), 620 – 860 (WR3), 800 – 1100 (WR4), 1500 – 1800 (WR5).
- 4 spectrographs (SPs) with wavelength ranges in nm between 390 – 560, 560 – 1100, 700 – 1600, 1000 – 2300. These SPs were thought to be configured as long-slit spectrographs, as multi-slit multi-wavelength SPs with an integral field unit (IFU), or as double-pass imaging spectrographs (TUNIS or MSDP type of instruments).
- 3 Broad Band Imagers (BBIs), two with 380 – 500, and one with 600 - 900. They are associated to the NBIs of the corresponding wavelength, selecting the desired wavelength range with a suitable filter. Broad Band Imager provide diffraction limited context images with large FOVs at high cadence.

The light distribution system foresees exchangeable dichroic and partial beam splitters. This allows either simultaneous observations with all instruments by sharing light of a specific wavelength or observations where a subset of instruments is fed with all light of a specific wavelength.

11.2 Design developments 2011 – 2019

Instrumentation: Since the preliminary design phase, some 10 years ago, significant developments on post-focus instrumentation have happened. In recent years, prototypes of Integral-Field-Units (IFUs) with solar spectro-polarimetric measurements have successfully been developed and tested. Prototypes of micro-lense arrays and image slicer have proven to be most promising. With such IFUs, the 2D spatial information and the spectral dimension are recorded simultaneously, while traditional spectrographs and narrow band imagers either have long slits that scan the solar surface or

take very narrow-band images that sequentially scan through wavelength. This has fundamental consequences for those IFU systems that record coherent 2D images. *Post-factum* image restoration techniques can then be applied to the IFU data. Hereby, highest, diffraction-limited, spatial resolution can be achieved with full spectral integrity.

Spatially scanned maps with such long-slit spectrographs ask for a non-rotating solar image, since image rotation squeezes and stretches the solar scene. This effectively shrinks the field-of-view. With IFUs, the requirement of de-rotating the solar image is less stringent, since the larger fields-of-view are build up by mosaics. And since each tile of the mosaic has short exposure times, the image rotation is negligible, also for image restoration techniques.

Adaptive secondary mirror (ASM): Another crucial technical development during the last decade is that large-diameter deformable mirrors become available. If such a large-diameter deformable mirror could be placed as a secondary mirror, it appears feasible to integrate all AO and MCAO mirrors within the first seven mirrors of the telescope and its main axes. The transfer optics of the preliminary design that has seven additional mirrors becomes obsolete. Hence, with a deformable secondary mirror, the total number of mirrors could be reduced from fourteen to seven. This will substantially improve the optical quality and the photon flux in the science focus will increase by a factor of about 2 depending on wavelength⁸. For these reasons, scientists would prefer this solution, but it remains to be seen whether the involved technical risks are acceptable.

While the removal of the (rotating) transfer optics is of great advantage in terms of photon flux and image quality, it reintroduces image rotation. There are three options how to deal with this: (1) a rotating Coudé room, (2) a separate image de-rotator, or (3) accept image rotation. A rotating Coudé room is the straight-forward solution, but costly and may reduce the space for the instrument platform. Introducing a separate polarimetrically compensated image de-rotator with 5 additional mirrors appears to be a bad solution with respect to optical quality and photon flux. As mentioned above with instrumentation based on IFUs and NBIs, the problems of image rotation are much less severe and would be acceptable. Image rotation is less acceptable for long-slit spectrographs. They could be equipped with an internal image de-rotator.

Note on image rotation: Depending on the direction of the beam along the elevation axis, the image rotation rate differs between morning and afternoon. Since seeing conditions at Observatorio del Teide (Tenerife) and Roque de los Muchachos (La Palma) are typically better in the morning, the telescope design should allow for the smaller image rotation rate in the morning.

12 Discussion

The overall goal of EST is to understand the small-scale processes in the solar atmosphere. The telescope is designed to be a ‘microscope’ providing with best imaging and polarimetric quality and with the highest possible photon throughput. To achieve highest imaging quality, the telescope design contains Multi-Conjugate Adaptive Optics (MCAO). MCAO and image reconstruction techniques allow for diffraction limited imaging quality over the entire FOV, including FOVs at and off the limb. An optimal polarimetric performance is assured by assembling the mirrors such that they are polarimetrically compensated. The suite of post-focus instruments will change during its lifetime.

12.1 Discussion on photon flux

Science cases are presented and discussed in Part II (starting on page 8). Modelling the photon flux⁹, observing programmes were devised for each top-level science case. While doing this, it became obvious that compromises need to be made between spatial resolution, spectral resolution, and SNR. This is based on two fundamental properties: (i) The number of photons collected by pixels that sample the diffraction limit of a given aperture is independent of the aperture. I.e. the advantage of a larger aperture can either be used to increase the spatial resolution **or** to increase the number of photons (SNR). (ii) The solar scene evolves with time. This severely limits the exposure times that are needed to freeze a snapshot of the solar evolution. Hence, the top-level goal for the final design of the telescope is two-folded:

- (1) EST should be capable to reach the highest possible image quality and spatial resolution, i.e. diffraction limit. With the option to sacrifice spatial resolution in favour of collecting more photons.
- (2) The final design must be optimised for the highest possible photon flux, with the premises of securing polarimetric accuracy and sensitivity.

⁸A removal of 7 Ag-mirrors ($R@400\text{nm} \approx 0.85$, $R@1000\text{nm} \approx 0.95$) increases the photon flux by a factor of 3 at 400nm and a factor of 1.4 at 1000nm.

⁹*photoncount* app by J. Leenaarts.

12.2 Discussion on field-of-view (FOV)

EST is designed to study the small-scale structure of the solar atmosphere. The science cases presented in Part II request to resolve sub-structures from some 0.2 arcsec down to the diffraction limit (0.026 arcsec at 500 nm with a 4 m aperture). While scientists desire to have the FOV as large as possible, the reasons for limitations are manifold:

- (1) The size of optical surfaces in the telescope light beam increases with increasing FOV, leading to a substantial increase of costs. This cost increase is particularly high for NBIs. Large FOVs at highest spatial resolution are expensive.
- (2) The performance of the MCAO system degrades with increasing the wavefront-corrected FOV. The expectation is that MCAO will correct a FOV of less than 60 arcsec in diameter.

The FOVs of IFUs is expected to be limited to some 10 by 10 arcsec². These IFUs will be complemented by Narrow Band Imagers (NBIs). For the science cases in part II, a FOV of some 40 by 40 arcsec² is sufficiently large. The science cases require simultaneous observation in three or more different wavelength regions. I.e., rather than requesting a larger FOV, the science cases require a multitude of NBIs and IFUs that operate in different wavelengths simultaneously. If compromises need to be done, priority will be given to simultaneous multi-wavelength observation over larger FOVs.

If individual FOVs do not need to be larger than 40 by 40 arcsec², one could, in principle, relax the original requirement for the telescope FOV of 120 by 120 arcsec². Such large FOVs are traditionally required for context imaging. Instead of designing EST with such a large FOV, one could consider another telescope with a FOV of some 400 arcsec in diameter that delivers spectropolarimetric measurements with a spatial resolution of about 0.5 arcsec. This implies that such a telescope should be equipped with an adaptive optics system. This could also be an extended version of the auxiliary full-disk telescope (AFDT) that was foreseen in the preliminary EST design (cf. EST Conceptual Design Study, 2011).

12.3 Discussion on pointing requirements

For the science cases, the absolute pointing accuracy of the telescope on the sky is not relevant. Only the absolute pointing relative to the solar disk is relevant for the solar observer. In the World Coordinate System (Thomson 2005), the position on the solar disk can be determined by the Helioprojective-Cartesian coordinate system. These coordinates are given relative to the center of the solar disk and the separation between two points is given by the corresponding angle separation, i.e. in arcsec.

Absolute positioning is needed for simultaneous observations with other telescopes. Taking into account that the FOVs of IFUs are expected to be smaller than 10 by 10 arcsec², an absolute pointing on the solar disk should be on the order of 1 arcsec to guarantee the needed overlap with observations from other telescopes.

Mosaic with NBIs: For obtaining a large FOV mosaic with the NBIs, the tiling should be done by re-pointing the telescope. For this, the relative pointing is requested to have an accuracy of 1 arcsec. The re-pointing of the telescope including a closed-loop in the adaptive optics system at the new position should be performed as fast as possible. This will optimise the duty cycle. A time lapse of 2 seconds or less would allow for valuable time cadence of larger FOVs.

Mosaic with IFUs: In some science cases it is necessary to raster a larger FOV with IFUs. Since their FOV is smaller than those of NBIs, the positioning can not be done with the telescope, but needs an internal mechanism. To avoid redundant overlapping, the pointing accuracy of individual IFU tiles relative to each other should be in the order of 0.1 arcsec. For valuable time series of IFU mosaics the re-pointing time lapse should be as fast as possible. A time lapse of 0.1 sec or less is desirable to be able to acquire time series of mosaics at a reasonable cadence.

12.4 Discussion on wavelength range

The telescope should be optimised to cover the spectral range from Ca II K 393.4 to Fe I 1564.8 nm. Observations up to 2300 nm are desired. The choice of coatings should favour the transmission in the longer wavelengths between 800 and 1083 nm. This wavelength regime is considered to be more important to measure the magnetic coupling in the chromosphere. The optical train should be optimised to assure highest photon flux at the Ca II IR 854.2 nm as this line is crucial for the most important science cases, and since the time scales of the associated chromospheric signals are shorter than in the photosphere. Requirement for wavelength coverage beyond 1600 nm up to 2300 nm should not compromise performance for wavelengths smaller than 1600 nm.

12.5 Discussion on instrumentation and light distribution

The main science driver for EST consists in understanding the magnetic coupling of the solar atmosphere. This is reflected in many of the science cases described in Part II. In the corresponding observing programmes, measurements probing many different layers of the solar atmosphere are required to be taken co-temporally and co-spatially in many different wavelengths, as e.g., Ca II K 393.4 nm, Mg I 517.3 nm, Fe I 525.0 nm, Fe I 557.6 nm, He I 587.6 nm, Fe I 617.3 nm, Fe I 709.0 nm, H α 656.3 nm, Ca II IR triplet 849.8 nm, 854.2 nm and 866.2 nm, Si I 1082.7 nm, He I 1083.0 nm, Fe I 1564.8 nm, etc. For the observing programmes in Part II, typically three or more of these lines are required to probe the photosphere and chromosphere. Here again, time and photons are limited, such that it is desirable that multiple instruments operate at different wavelengths simultaneously. Ideally, each instrument would receive all available photons of a specific wavelength.

Efficient NBIs have low spectral resolution (between 30 000 and 100 000), meaning that science cases which request high spectral resolution (around 200 000) need to be served with IFUs or long-slit spectrographs.

The post-focus instrumentation and light distribution concept of the preliminary design from 2011 is powerful and sophisticated. With this concept, simultaneous observations as desired in the previous paragraph are possible. Yet, the old concept foresees to share light between SPs and NBIs. And IFUs are foreseen to replace classical long-slit spectrographs. As a consequence of the lesson learned during more recent developments, one should also consider to replace some of the NBIs by IFUs, or have both types of instruments for some wavelength ranges, depending on whether large FOV or short time cadence and high spectral resolution is needed for the science case. Different types of instruments for the same wavelength may be necessary to achieve the goals of different science cases, but simultaneous observation at a specific wavelength with different types of instruments, i.e. sharing the light at a specific wavelength, is not needed.

In this respect, the consensus of the present EST Science Advisory Group differs from the Preliminary Design Study: the light distribution should be such that each instrument receives all photons of a particular wavelength. Many of the top-level science goals need high SNRs to observe features that have short evolution time scales. Hence, the exposure time and the observing cadence must be short. Although IFUs have smaller FOVs than NBIs, they have higher spectral resolution and potentially can complete one measurement in a shorter time span.

Since photon flux and solar evolution time speed are critical, measurements should in general be done simultaneously rather than sequentially in 5 or more spectral lines. I.e., EST should be equipped with a multitude of exchangeable NBIs and IFUs. *However, in this document, we do not yet recommend how many NBIs and IFUs should observe at which wavelengths. Such a concept yet needs to be developed, and depends on the outcome of ongoing design studies of IFUs.*

13 Scientific Requirements

Which are the specifications that are relevant for the final design of the telescope, and for a preliminary planning of its instrumentation?

13.1 Telescope specifications

1. FOV:

- (a) Telescope FOV (Field stop in F1): Diameter of 125 arcsec corresponding to a square of 90 by 90 arcsec².
- (b) Seeing-corrected FOV: 40 by 40 arcsec². Seeing at the sites of OT and ORM are dominated by ground-layer turbulence. Therefore a high elevation of the telescope is to be preferred. The MCAO system shall also correct FOVs at and off the solar limb to observe spicules and prominences at the diffraction limit.
- (c) For context information larger FOVs are essential. Such context information can be supplied by another telescope that has spectropolarimetric capabilities.
- (d) Auxiliary full disk telescope (AFDT): At the site of EST a FDT is needed as a finder telescope for the orientation of the observer on the solar disk. This AFDT should provide full disk images (with FOV of 1 degree²) in Ca II H or Ca II K, H α 656.3 nm, possibly He I 1083.0 nm, and visible continuum light at a spatial resolution of 1.5 arcsec. The AFDT sketched in the EST Conceptual Design Study (2011) would serve the needs.

2. Pointing:

- (a) Pointing accuracy on solar disk: 1 arcsec.
- (b) Relative pointing accuracy for individual tiles of mosaic with NBIs: 1 arcsec
- (c) Telescope repointing time for tiles in NBI mosaic: 2 sec
- (d) Relative (internal) pointing accuracy for individual tiles of mosaic with IFUs (small FOVs): 0.1 arcsec

- (e) Repointing time (internal raster) for tiles in IFU mosaic: 0.1 sec
3. Optical quality: Optimised for high photon flux and best optical quality, i.e. optimised for a minimum amount of optical surfaces.
- (a) Diffraction-limited image quality
- (b) The Modulation Transfer Function (MTF) is required to be optimised for spatial scales that correspond to twice the diffraction-limit. At that scale a MTF value of about 70% relative to an ideal non-obscured aperture telescope is desired.¹⁰ The performance of the multi-conjugate adaptive optics (MCAO) system is required to retain this image quality under excellent Seeing conditions (Fried parameter $R_0 > 10$ cm and stable high altitude layers).
4. Secondary Mirror:
- (a) Design with Adaptive Secondary Mirror (ASM) is to be preferred if technically feasible.
- (b) In case of design with ASM: Image rotation is acceptable for NBIs and IFUs. Long-slit spectrographs benefit from an internal image de-rotator. Rotating Coudé platform to be preferred if it allows for sufficient space for instruments. The image rotation rate should be smallest during the morning¹¹.
5. The polarimetric properties of the solar light need to be measured with high sensitivity and accuracy. Polarimetric sensitivity refers to the ability to detect a signal above the noise and is therefore a requirement of the signal-to-noise ratio. The polarimetric sensitivity is defined as the ratio between the RMS noise value of a given Stokes parameter Q , U or V and the average intensity (Stokes I). Polarimetric accuracy quantifies the residual errors in establishing the zero polarisation levels and in removing the crosstalk between Stokes Q , U , and V , i.e., accuracy refers to the residual error after demodulating the measured Stokes parameters.
- (a) Measurements shall allow to be sensitive to polarimetric signals at the level of $3 \cdot 10^{-5}$.
- (b) For the errors, \mathcal{E} , of the demodulation matrix of the measured Stokes vectors, i.e. for the contamination, $\mathcal{E}_{i,j}$ due to cross talk between Stokes i and j , the requirements can be written in the following form¹²:

$$\mathcal{E} = \begin{pmatrix} 10^{-2} & 1 & 1 & 0.1 \\ 5 \cdot 10^{-4} & 10^{-2} & 5 \cdot 10^{-2} & 1 \cdot 10^{-3} \\ 5 \cdot 10^{-4} & 5 \cdot 10^{-2} & 10^{-2} & 1 \cdot 10^{-3} \\ 5 \cdot 10^{-3} & 5 \cdot 10^{-1} & 5 \cdot 10^{-1} & 10^{-2} \end{pmatrix}$$

This implies that

- i. the scaling accuracy needs to be better than 10^{-2} ,
 - ii. the residual polarisation measured in an unpolarised region must be smaller than $5 \cdot 10^{-4}$ (for Q , U) and $5 \cdot 10^{-3}$ (V), respectively,
 - iii. the cross talk $V \rightarrow (Q, U)$ is constrained to 0.1% to allow for scattering polarisation measurements, and
 - iv. if Q and U are 10% of V and V is 10% of I , the relative crosstalk contamination is evenly distributed among Q , U , and V , and amounts to 5%.
6. Wavelength coverage: 390 nm - 2300 nm. Transmission is required to be optimised for Ca II IR 854.2 nm. Requirement for wavelength coverage beyond 1600 nm should not compromise performance for wavelengths smaller than 1600nm.
7. Pointing off the solar disk: < 100 arcsec. The telescope is required to be able to point to the sun during day time.

13.2 Requirements for instrumentation and light distribution:

Many of the top-level science cases request to observe in many different layers of the solar atmosphere simultaneously. Up to three photospheric lines together with up to three chromospheric lines are requested. The selected lines for the Zeeman and Doppler diagnostics are one of the following: Ca II K 393.4 nm, Mg I 517.3 nm, Fe I 525.0 nm, Fe I 543.5 nm (non-magnetic), Fe I 557.6 nm (non-magnetic), He I 587.6 nm, Fe I 617.3 nm, Fe I 630.1 nm, Fe I 630.2 nm, Fe I 709.0 nm (non-magnetic), H α 656.3 nm, Ca II 854.2 nm, Si I 1082.7 nm, He I 1083.0 nm, Fe I 1564.8 nm. For the diagnostics using

¹⁰The EST baseline design foresees a central obscuration of 1.30 m, and 4 spiders with a width of 0.05 m. Relative to an open aperture, this results in a reduction of the MTF to 72% at a spatial frequency that corresponds to half of the diffraction limit.

¹¹The choice of direction of the beam in the elevation axis decides on image rotation rate being smaller or larger in the morning relative to the afternoon.

¹²If \mathbb{O} denotes the modulation matrix, \mathbb{D} the demodulation matrix, and \mathbb{E} the unity matrix, then $\mathcal{E} = \mathbb{D} \cdot \mathbb{O} - \mathbb{E}$.

scattering polarization the following lines are proposed: Ca II H , Ca II K , Ca I 422.7 nm, Ti I 453.0 nm, Sr I 460.7 nm, C₂ molecular line at 514.0 nm, Na I D₁ and D₂ 589 nm, Ca II IR triplet 849.8 nm, 854.2 nm and 866.2 nm.

Simultaneous observation in 6 different spectral lines are requested by the high-impact science cases. Some of those require larger FOVs, others require spectral integrity and smaller time cadences. I.e., 6 NBIs exchangeable with 6 IFUs for 6 different simultaneous wavelength regimes would suffice to serve science requirements.

The spatial resolution of NBIs and IFUs in some cases should be close to the diffraction limit. However many science cases ask for high SNR that will require to increase the collection area, i.e. reduce the spatial resolution. This could be achieved by a variable image scale or by binning of the recorded pixels.

Broad Band Imagers (BBIs) should be capable to image the solar evolution at the diffraction limit. Therefore a sequential observing procedure is not of advantage. Instead, one should foresee two BBIs for the blue visible and one BBI for the red visible light.

Part IV

Appendix, References, & Abbreviations

A Signal-to-noise ratio S/N

A.1 Optimum spatio-temporal resolution

(Alex Feller)

Number of photo-electrons, Z , in a resolution element (detector pixel):

$$Z = (S/N)^2 = \Phi \Delta\lambda \Delta x^2 \Delta t \quad (1)$$

S/N is the signal-to-noise ratio. Here we assume that the measurement noise is dominated by photon noise which is Poisson distributed. Φ is the photon-electron flux per unit wavelength, time, and resolution element. This quantity determines the actual S/N of the measurement and takes into account the overall transmission of the beam path from the solar surface through the Earth's atmosphere, the telescope optics, as well as the quantum efficiency of the detector. $\Delta\lambda, \Delta x, \Delta t$ denote the spectral, spatial and temporal sampling respectively.

If we assume that spatial and temporal resolution are coupled by a characteristic solar evolution speed v :

$$\Delta x = v\Delta t, \quad (2)$$

we can determine the optimum temporal sampling (cadence) in order to reach a given S/N :

$$\Delta t = \left(\frac{(S/N)^2}{\Phi \Delta\lambda v^2} \right)^{1/3} \quad (3)$$

The optimum spatial sampling is then given by Eq. (2). If the spatial sampling is lower than this optimum value the decreased photon flux per pixel leads to integration times which exceed the timescale of solar evolution at the given spatial sampling, leading to a smearing of the observed solar scene.

A.2 Number of photons at diffraction limit

(Rolf Schlichenmaier)

If one considers the signal-to-noise ratio at the diffraction limit of the telescope aperture, one needs to take into account that the size of the aperture increases the photon collection area, while it decreases the size of the diffraction-limited resolution element. The photon flux, Φ , is proportional to the square of the diameter, D , of the aperture: $\Phi = \tilde{\Phi} D^2$. Hence, Eq.(1) can be written as

$$(S/N)^2 = \tilde{\Phi} D^2 \Delta x^2 \Delta\lambda \Delta t$$

If the size of the resolution element is chosen such that it is given by the diffraction limit of the telescope, $\Delta x = \Delta x_{\text{limit}} = \lambda/D$, and if the exposure time is limited by the solar evolution speed (Eq. 2), $\Delta t = \Delta x_{\text{limit}}/v = \lambda/(Dv)$, then

$$(S/N)_{\text{limit}}^2 = \tilde{\Phi} D^2 \Delta x_{\text{limit}}^2 \Delta\lambda \Delta t_{\text{limit}} = \tilde{\Phi} D^2 \frac{\lambda^2}{D^2} \Delta\lambda \frac{\lambda}{Dv} = \tilde{\Phi} \frac{\lambda^3 \Delta\lambda}{v} \frac{1}{D}$$

This consideration demonstrates that bigger telescopes may give better spatial resolution, but at the diffraction limit they do not provide a higher signal-to-noise ratio. Even worse, the solar evolution speed, v , limits the allowed exposure time at the diffraction limit, such that the signal-to-noise ratio at the diffraction limit decreases with telescope aperture:

$$(S/N)_{\text{limit}}^2 \propto \frac{1}{D} \quad (4)$$

This somewhat counter-intuitive lesson stresses that high-resolution solar observations are limited by photons, and that solar telescopes must be optimised for photon flux.

The above equations are included in the `photoncount` code by Jorrit Leenaarts.

References

- Abramenko, V., Yurchyshyn, V., Goode, P., & Kilcik, A. 2010, *ApJL*, 725, L101
- Abramenko, V. I., Carbone, V., Yurchyshyn, V., et al. 2011, *ApJ*, 743, 133
- Allcock, M. & Erdélyi, R. 2017, *Solar Physics*, 292, 35
- Alsina Ballester, E., Belluzzi, L., & Trujillo Bueno, J. 2016, *ApJ*, 831, L15
- Alsina Ballester, E., Belluzzi, L., & Trujillo Bueno, J. 2018, *ApJ*, 854, 150
- Alsina Ballester, E., Belluzzi, L., & Trujillo Bueno, J. 2019, *ApJ*, 880, 85
- Anan, T., Ichimoto, K., & Hillier, A. 2017, *A&A*, 601, A103
- Arber, T. D., Haynes, M., & Leake, J. E. 2007, *ApJ*, 666, 541
- Arregui, I. 2015, *Philosophical Transactions of the Royal Society of London Series A*, 373, 20140261
- Asensio Ramos, A. 2009, *Astrophys. J.*, 701, 1032
- Asensio Ramos, A. & Martínez González, M. J. 2014, *A&A*, 572, A98
- Asensio Ramos, A. & Trujillo Bueno, J. 2005, *ApJ*, 635, L109
- Attie, R., Innes, D. E., & Potts, H. E. 2009, *A&A*, 493, L13
- Ayres, T. R. & Rabin, D. 1996, *ApJ*, 460, 1042
- Ballester, J. L., Alexeev, I., Collados, M., et al. 2018, *Space Science Reviews*, 214, 1
- Banerjee, D., Erdélyi, R., Oliver, R., & O'Shea, E. 2007, *Solar Physics*, 246, 3
- Bard, S. & Carlsson, M. 2010, *ApJ*, 722, 888
- Beck, C., Bellot Rubio, L. R., Schlichenmaier, R., & Sütterlin, P. 2007, *A&A*, 472, 607
- Beckers, J. M. & Tallant, P. E. 1969, *Solar Physics*, 7, 351
- Bellot Rubio, L. R., Collados, M., Ruiz Cobo, B., & Rodríguez Hidalgo, I. 2000a, *ApJ*, 534, 989
- Bellot Rubio, L. R. & Orozco Suárez, D. 2012, *Astrophys. J.*, 757, 19
- Bellot Rubio, L. R., Rodríguez Hidalgo, I., Collados, M., Khomeiko, E., & Ruiz Cobo, B. 2001, *ApJ*, 560, 1010
- Bellot Rubio, L. R., Ruiz Cobo, B., & Collados, M. 2000b, *ApJ*, 535, 489
- Bellot Rubio, L. R., Tritschler, A., & Martínez Pillet, V. 2008, *ApJ*, 676, 698
- Belluzzi, L. & Trujillo Bueno, J. 2012, *ApJ*, 750, L11
- Belluzzi, L. & Trujillo Bueno, J. 2013, *ApJ*, 774, L28
- Belluzzi, L., Trujillo Bueno, J., & Landi Degl'Innocenti, E. 2015, *ApJ*, 814, 116
- Berdyugina, S. V. & Fluri, D. M. 2004, *A&A*, 417, 775
- Berger, T. E., Rouppe van der Voort, L. H. M., Löfdahl, M. G., et al. 2004, *Astron. Astrophys.*, 428, 613
- Bharti, L., Hirzberger, J., & Solanki, S. K. 2013, *A&A*, 552, L1
- Bi, S. & Li, R. 1998, *A&A*, 335, 673
- Bianda, M., Benz, A. O., Stenflo, J. O., Küveler, G., & Ramelli, R. 2005, *A&A*, 434, 1183
- Bianda, M., Berdyugina, S., Gisler, D., et al. 2018, *A&A*, 614, A89
- Bianda, M., Ramelli, R., Anusha, L. S., et al. 2011, *A&A*, 530, L13
- Bianda, M., Solanki, S. K., & Stenflo, J. O. 1998, *A&A*, 331, 760

Bittencourt, J. A. 1986, *Fundamentals of plasma physics* (Oxford: Pergamon Press)

Bodnárová, M., Utz, D., & Rybák, J. 2013, *Solar Physics*

Buehler, D., Lagg, A., Solanki, S. K., & van Noort, M. 2015, *A&A*, 576, A27

Cally, P. S. 2007, *Astronomische Nachrichten*, 328, 286

Capparelli, V., Zuccarello, F., Romano, P., et al. 2017, *ApJ*, 850, 36

Carlin, E. S., Manso Sainz, R., Asensio Ramos, A., & Trujillo Bueno, J. 2012, *ApJ*, 751, 5

Carlsson, M. & Stein, R. F. 1997, *ApJ*, 481, 500

Casini, R. & Landi Degl'Innocenti, E. 2008, *Astrophysical Plasmas*, ed. T. Fujimoto & A. Iwamae, Vol. 44 (Springer-Verlag Berlin Heidelberg, ISBN 978-3-540-73586-1), 247

Casini, R., López Ariste, A., Paletou, F., & Léger, L. 2009, *ApJ*, 703, 114

Casini, R., López Ariste, A., Tomczyk, S., & Lites, B. W. 2003, *ApJ*, 598, L67

Casini, R., White, S. M., & Judge, P. G. 2017, *Space Science Reviews*, 210, 145

Cauzzi, G., Reardon, K., Rutten, R. J., Tritschler, A., & Uitenbroek, H. 2009, *A&A*, 503, 577

Centeno, R. 2018, *ApJ*, 866, 89

Centeno, R., Collados, M., & Trujillo Bueno, J. 2006, *ApJ*, 640, 1153

Centeno, R., Collados, M., & Trujillo Bueno, J. 2009, *ApJ*, 692, 1211

Centeno, R., Socas-Navarro, H., Lites, B., et al. 2007, *Astrophys. J. Lett.*, 666, L137

Centeno, R., Trujillo Bueno, J., & Asensio Ramos, A. 2010, *ApJ*, 708, 1579

Cirtain, J. W., Golub, L., Lundquist, L., et al. 2007, *Science*, 318, 1580

Danilovic, S. 2017, *A&A*, 601, A122

Danilovic, S., Schüssler, M., & Solanki, S. K. 2010a, *A&A*, 509, A76

Danilovic, S., Schüssler, M., & Solanki, S. K. 2010b, *A&A*, 513, A1

Danilovic, S., Solanki, S. K., Barthol, P., et al. 2017, *ApJ Supplements*, 229, 5

Danilovic, S., van Noort, M., & Rempel, M. 2016, *Astron. Astrophys.*, 593, A93

de la Cruz Rodríguez, J., Leenaarts, J., & Asensio Ramos, A. 2016, *ApJL*, 830, L30

de la Cruz Rodríguez, J., Rouppe van der Voort, L., Socas-Navarro, H., & van Noort, M. 2013, *A&A*, 556, A115

de la Cruz Rodríguez, J. & Socas-Navarro, H. 2011, *A&A*, 527, L8

de la Cruz Rodríguez, J., Socas-Navarro, H., Carlsson, M., & Leenaarts, J. 2012, *A&A*, 543, A34

De Pontieu, B., De Moortel, I., Martinez-Sykora, J., & McIntosh, S. W. 2017, *ApJL*, 845, L18

De Pontieu, B., Erdélyi, R., & De Moortel, I. 2005, *ApJL*, 624, L61

De Pontieu, B., Erdélyi, R., & de Wijn, A. G. 2003, *ApJ*, 595, L63

De Pontieu, B., Erdélyi, R. J., & Stewart, P. 2004, *Nature*, 430, Issue 6999, 536

De Pontieu, B. & Haerendel, G. 1998, *A&A*, 338, 729

De Pontieu, B., Hansteen, V. H., Rouppe van der Voort, L., van Noort, M., & Carlsson, M. 2007a, *ApJ*, 655, 624

De Pontieu, B., McIntosh, S., Hansteen, V. H., et al. 2007b, *PASJ*, 59, S655

De Pontieu, B., McIntosh, S. W., Carlsson, M., et al. 2007c, *Science*, 318, 1574

de Wijn, A. G., McIntosh, S. W., & De Pontieu, B. 2009, *ApJL*, 702, L168

- de Wijn, A. G., Rutten, R. J., Haverkamp, E. M. W. P., & Sütterlin, P. 2005, *A&A*, 441, 1183
- del Pino Alemán, T., Casini, R., & Manso Sainz, R. 2016, *ApJ*, 830, L24
- del Pino Alemán, T., Trujillo Bueno, J., Štěpán, J., & Shchukina, N. 2018, *ApJ*, 863, 164
- Deng, N., Liu, C., Yang, G., Wang, H., & Denker, C. 2005, *ApJ*, 623, 1195
- Díaz Baso, C. J., Martínez González, M. J., & Asensio Ramos, A. 2016, *ApJ*, 822, 50
- Donea, A. 2011, *Space Science Reviews*, 158, 451
- Drews, A. & Rouppe van der Voort, L. 2017, *A&A*, 602, A80
- Ellerman, F. 1917, *ApJ*, 46, 298
- Erdélyi, R. 2006, *Philosophical Transactions of the Royal Society of London Series A*, 364, 351
- Erdélyi, R. & Ballai, I. 2007, *Astronomische Nachrichten*, 328, 726
- Erdélyi, R. & Fedun, V. 2007, *Science*, 318, 1572
- Erdélyi, R., Malins, C., Tóth, G., & De Pontieu, B. 2007a, *A&A*, 467, 1299
- Erdélyi, R., Pintér, B., & Malins, C. 2007b, *Astronomische Nachrichten*, 328, 305
- Erdélyi, R. & Taroyan, Y. 2008, *A&A*, 489, L49
- Faurobert-Scholl, M., Feautrier, N., Machefert, F., Petrovay, K., & Spielfiedel, A. 1995, *A&A*, 298, 289
- Fawzy, D., Rammacher, W., Ulmschneider, P., Musielak, Z. E., & Stępień, K. 2002, *A&A*, 386, 971
- Fedun, V., Shelyag, S., & Erdélyi, R. 2011, *ApJ*, 727, 17
- Felipe, T., Khomenko, E., & Collados, M. 2011, *ApJ*, 735, 65
- Felipe, T., Khomenko, E., Collados, M., & Beck, C. 2010b, *ApJ*, 722, 131
- Felipe, T., Socas-Navarro, H., & Khomenko, E. 2014, *ApJ*, 795, 9
- Fischer, C. E., de Wijn, A. G., Centeno, R., Lites, B. W., & Keller, C. U. 2009, *A&A*, 504, 583
- Fisher, G. H., Bercik, D. J., Welsch, B. T., & Hudson, H. S. 2012, *Solar Physics*, 277, 59
- Fleck, B., Straus, T., Carlsson, M., et al. 2010, *Memorie della Societa Astronomica Italiana*, 81, 777
- Fossum, A. & Carlsson, M. 2006, *ApJ*, 646, 579
- Foullon, C., Verwichte, E., Nakariakov, V. M., & Fletcher, L. 2005, *A&A*, 440, L59
- Freij, N., Dorotovič, I., Morton, R. J., et al. 2016, *ApJ*, 817, 44
- Freij, N., Scullion, E. M., Nelson, C. J., et al. 2014, *ApJ*, 791, 61
- Fujimura, D. & Tsuneta, S. 2009, *ApJ*, 702, 1443
- Gandorfer, A. 2000, *The Second Solar Spectrum: A high spectral resolution polarimetric survey of scattering polarization at the solar limb in graphical representation. Volume I: 4625 Å to 6995 Å*
- Gandorfer, A. 2002, *The Second Solar Spectrum: A high spectral resolution polarimetric survey of scattering polarization at the solar limb in graphical representation. Volume II: 3910 Å to 4630 Å*
- Gandorfer, A. 2005, *The Second Solar Spectrum: A high spectral resolution polarimetric survey of scattering polarization at the solar limb in graphical representation. Volume III: 3160 Å to 3915 Å*
- Gilbert, H., Kilper, G., & Alexander, D. 2007, *ApJ*, 671, 978
- Gilbert, H. R., Hansteen, V. H., & Holzer, T. E. 2002, *ApJ*, 577, 464
- Giovanelli, R. G. 1972, *Solar Phys.*, 27, 71
- Gizon, L. & Birch, A. C. 2005, *Living Reviews in Solar Physics*, 2, 6

Goldreich, P. & Kumar, P. 1990, *ApJ*, 363, 694

Goldreich, P., Murray, N., & Kumar, P. 1994, *ApJ*, 424, 466

Gošić, M., Bellot Rubio, L. R., del Toro Iniesta, J. C., Orozco Suárez, D., & Katsukawa, Y. 2016, *ApJ*, 820, 35

Gošić, M., Bellot Rubio, L. R., Orozco Suárez, D., Katsukawa, Y., & del Toro Iniesta, J. C. 2014, *ApJ*, 797, 49

Gošić, M., de la Cruz Rodríguez, J., De Pontieu, B., et al. 2018, *ApJ*, 857, 48

Grigis, P. C. & Benz, A. O. 2005, *ApJL*, 625, L143

Hagenaar, H. J. & Shine, R. A. 2005, *ApJ*, 635, 659

Hansteen, V. H., Archontis, V., Pereira, T. M. D., et al. 2017, *ApJ*, 839, 22

Hasan, S. S. & Ulmschneider, P. 2004, *A&A*, 422, 1085

Hasan, S. S. & van Ballegooijen, A. A. 2008, *ApJ*, 680, 1542

Hegglund, L., Hansteen, V. H., De Pontieu, B., & Carlsson, M. 2011, *ApJ*, 743, 142

Hénoux, J. C. & Chambe, G. 1990, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 44, 193

Henriques, V. M. J., Kuridze, D., Mathioudakis, M., & Keenan, F. P. 2016, *ApJ*, 820, 124

Henriques, V. M. J., Scullion, E., Mathioudakis, M., et al. 2015, *A&A*, 574, A131

Hiei, E. 1982, *Solar Physics*, 80, 113

Holzreuter, R., Fluri, D. M., & Stenflo, J. O. 2006, *A&A*, 449, L41

Hou, Y., Zhang, J., Li, T., Yang, S., & Li, X. 2017, *ApJL*, 848, L9

Howard, R. & Stenflo, J. O. 1972, *Solar Physics*, 22, 402

Hurlburt, N. E. & Rucklidge, A. M. 2000, *MNRAS*, 314, 793

Iida, Y., Yokoyama, T., & Ichimoto, K. 2010, *ApJ*, 713, 325

Iijima, H. & Yokoyama, T. 2017, *ApJ*, 848, 38

Innes, D. E., Guo, L.-J., Huang, Y.-M., & Bhattacharjee, A. 2015, *ApJ*, 813, 86

Ishikawa, R., Tsuneta, S., Ichimoto, K., et al. 2008, *Astron. Astrophys.*, 481, L25

Isobe, H., Kubo, M., Minoshima, T., et al. 2007, *PASJ*, 59, S807

Ito, H., Tsuneta, S., Shiota, D., Tokumaru, M., & Fujiki, K. 2010, *ApJ*, 719, 131

Jahn, K. & Schmidt, H. U. 1994, *A&A*, 290, 295

Jefferies, S. M., McIntosh, S. W., Armstrong, J. D., et al. 2006, *ApJ*, 648, L151

Jess, D. B., Mathioudakis, M., Crockett, P. J., & Keenan, F. P. 2008, *ApJ*, 688, L119

Jess, D. B., Mathioudakis, M., Erdélyi, R., et al. 2009, *Science*, 323, 1582

Jess, D. B., Morton, R. J., Verth, G., et al. 2015, *Space Science Reviews*, 190, 103

Jess, D. B., Pascoe, D. J., Christian, D. J., et al. 2012, *ApJL*, 744, L5

Jess, D. B., Reznikova, V. E., Van Doorselaere, T., Keys, P. H., & Mackay, D. H. 2013, *ApJ*, 779, 168

Judge, P. G., Kleint, L., & Sainz Dalda, A. 2015, *ApJ*, 814, 100

Jurčák, J., Bello González, N., Schlichenmaier, R., & Rezaei, R. 2014a, *PASJ*, 66, S3

Jurčák, J., Bello González, N., Schlichenmaier, R., & Rezaei, R. 2015, *A&A*, 580, L1

Jurčák, J., Bello González, N., Schlichenmaier, R., & Rezaei, R. 2017, *A&A*, 597, A60

Jurčák, J., Bellot Rubio, L. R., & Sobotka, M. 2014b, *A&A*, 564, A91

Jurčák, J., Rezaei, R., González, N. B., Schlichenmaier, R., & Vomlel, J. 2018, *A&A*, 611, L4

Kaithakkal, A. J., Suematsu, Y., Kubo, M., et al. 2015, *ApJ*, 799, 139

Kato, Y., Steiner, O., Steffen, M., & Suematsu, Y. 2011, *ApJL*, 730, L24

Katsukawa, Y., Berger, T. E., Ichimoto, K., et al. 2007, *Science*, 318, 1594

Kerr, G. S., Fletcher, L., Russell, A. J. B., & Allred, J. C. 2016, *ApJ*, 827, 101

Keys, P. H., Mathioudakis, M., Jess, D. B., Mackay, D. H., & Keenan, F. P. 2014, *A&A*, 566, A99

Keys, P. H., Mathioudakis, M., Jess, D. B., et al. 2013, *Monthly Notices of the Royal Astronomical Society*, 428, 3220

Khomenko, E. & Collados, M. 2009, *A&A*, 506, L5

Khomenko, E. & Collados, M. 2012, *ApJ*, 747, 87

Khomenko, E. & Collados, M. 2015, *Living Reviews in Solar Physics*, 12, 6

Khomenko, E., Collados, M., Díaz, A., & Vitas, N. 2014a, *Physics of Plasmas*, 21, 092901

Khomenko, E., Collados, M., & Díaz, A. J. 2016, *ApJ*, 823, 132

Khomenko, E., Collados, M., Shchukina, N., & Díaz, A. 2015, *A&A*, 584, A66

Khomenko, E., Díaz, A., de Vicente, A., Collados, M., & Luna, M. 2014b, *A&A*, 565, A45

Khomenko, E. V., Collados, M., & Bellot Rubio, L. R. 2003, *ApJ*, 588, 606

Khomenko, E. V., Collados, M., Solanki, S. K., Lagg, A., & Trujillo Bueno, J. 2003, *Astron. Astrophys.*, 408, 1115

Kleint, L., Berdyugina, S. V., Shapiro, A. I., & Bianda, M. 2010, *A&A*, 524, A37

Kleint, L., Shapiro, A. I., Berdyugina, S. V., & Bianda, M. 2011, *A&A*, 536, A47

Kobanov, N. I., Chelpanov, A. A., & Kolobov, D. Y. 2013, *A&A*, 554, A146

Kontogiannis, I., Georgoulis, M. K., Park, S.-H., & Guerra, J. A. 2017, *Solar Physics*, 292, 159

Kontogiannis, I., Tsiropoula, G., Tziotziou, K., & Georgoulis, M. K. 2010, *A&A*, 524, A12

Kosovichev, A. G. 2006, *Solar Physics*, 238, 1

Krishna Prasad, S., Jess, D. B., & Khomenko, E. 2015, *ApJL*, 812, L15

Krishna Prasad, S., Jess, D. B., Van Doorselaere, T., et al. 2017, *ApJ*, 847, 5

Kubo, M., Chye Low, B., & Lites, B. W. 2014, *ApJL*, 793, L9

Kubo, M., Lites, B. W., Shimizu, T., & Ichimoto, K. 2008, *ApJ*, 686, 1447

Kubo, M., Low, B. C., & Lites, B. W. 2010, *Astrophys. J.*, 712, 1321

Kuckein, C., Martínez Pillet, V., & Centeno, R. 2012, *A&A*, 539, A131

Kuridze, D., Henriques, V., Mathioudakis, M., et al. 2015a, *ApJ*, 802, 26

Kuridze, D., Henriques, V., Mathioudakis, M., et al. 2017, *ApJ*, 846, 9

Kuridze, D., Mathioudakis, M., Simões, P. J. A., et al. 2015b, *ApJ*, 813, 125

Kuridze, D., Morton, R. J., Erdélyi, R., et al. 2012, *ApJ*, 750, 51

Kuridze, D., Verth, G., Mathioudakis, M., et al. 2013, *ApJ*, 779, 82

Kuridze, D., Zaqarashvili, T. V., Shergelashvili, B. M., & Poedts, S. 2009, *A&A*, 505, 763

Lagg, A., Lites, B., Harvey, J., Gosain, S., & Centeno, R. 2017, *Space Science Reviews*, 210, 37

Lagg, A., Solanki, S. K., van Noort, M., & Danilovic, S. 2014, *A&A*, 568, A60

Lamb, D. A., DeForest, C. E., Hagenaar, H. J., Parnell, C. E., & Welsch, B. T. 2010, *Astrophys. J.*, 720, 1405

Landi Degl'Innocenti, E. 1998, *Nature*, 392, 256

Leake, J. E., Lukin, V. S., Linton, M. G., & Meier, E. T. 2012, *ApJ*, 760, 109

Leenaarts, J., Carlsson, M., Hansteen, V., & Gudiksen, B. V. 2011, *A&A*, 530, A124

Leenaarts, J., de la Cruz Rodríguez, J., Danilovic, S., Scharmer, G., & Carlsson, M. 2018, *A&A*, 612, A28

Leroy, J. L. 1989, in *Astrophysics and Space Science Library*, Vol. 150, *Dynamics and Structure of Quiescent Solar Prominences*, ed. E. R. Priest, 77–113

Levens, P. J., Schmieder, B., Labrosse, N., & López Ariste, A. 2016, *ApJ*, 818, 31

Lighthill, M. J. 1952, *Proc. R. Soc. London A.*, 211, 564

Lin, Y., Engvold, O., & Rouppe van der Voort, L. H. M. 2012, *ApJ*, 747, 129

Lites, B. W., Kubo, M., Socas-Navarro, H., et al. 2008, *Astrophys. J.*, 672, 1237

Lites, B. W. & Socas-Navarro, H. 2004, *ApJ*, 613, 600

Liu, J., Nelson, C. J., & Erdélyi, R. 2018, *ArXiv e-prints*

Liu, J., Nelson, C. J., & Erdélyi, R. 2018, *ApJ*, To Be Published

Liu, Y. & Schuck, P. W. 2012, *ApJ*, 761, 105

Liu, Y., Xiang, Y., Erdélyi, R., et al. 2018, *ApJ*, 856, 17

Löhner-Böttcher, J. & Bello González, N. 2015, *A&A*, 580, A53

Löhner-Böttcher, J. & Schlichenmaier, R. 2013, *A&A*, 551, A105

López Ariste, A. & Casini, R. 2005, *A&A*, 436, 325

López Ariste, A., Martínez González, M. J., & Ramírez Vélez, J. C. 2007, *Astron. Astrophys.*, 464, 351

Louis, R. E., Beck, C., & Ichimoto, K. 2014, *A&A*, 567, A96

MacTaggart, D., Guglielmino, S. L., & Zuccarello, F. 2016, *ApJ*, 831, L4

Malins, C. & Erdélyi, R. 2007, *Solar Physics*, 246, 41

Manso Sainz, R., Landi Degl'Innocenti, E., & Trujillo Bueno, J. 2004, *ApJ*, 614, L89

Manso Sainz, R., Roncero, O., Sanz-Sanz, C., et al. 2014, *ApJ*, 788, 118

Manso Sainz, R. & Trujillo Bueno, J. 2003, *Physical Review Letters*, 91, 111102

Manso Sainz, R. & Trujillo Bueno, J. 2010, *ApJ*, 722, 1416

Martínez González, M. J. & Bellot Rubio, L. R. 2009, *Astrophys. J.*, 700, 1391

Martínez González, M. J., Asensio Ramos, A., Manso Sainz, R., et al. 2011, *ApJL*, 730, L37

Martínez González, M. J., Bellot Rubio, L. R., Solanki, S. K., et al. 2012, *ApJL*, 758, L40

Martínez González, M. J., Collados, M., Ruiz Cobo, B., & Beck, C. 2008, *Astron. Astrophys.*, 477, 953

Martínez González, M. J., Manso Sainz, R., Asensio Ramos, A., et al. 2015, *ApJ*, 802, 3

Martínez González, M. J., Manso Sainz, R., Asensio Ramos, A., & Hijano, E. 2012, *Astrophys. J.*, 755, 175

Martínez González, M. J., Pastor Yabar, A., Lagg, A., et al. 2016, *Astron. Astrophys.*, 596, A5

Martínez Pillet, V. 2002, *Astronomische Nachrichten*, 323, 342

Martínez Pillet, V., Garcia Lopez, R. J., del Toro Iniesta, J. C., et al. 1990, *ApJL*, 361, L81

Martínez-Sykora, J., De Pontieu, B., Carlsson, M., & Hansteen, V. 2016, *ApJL*, 831, L1

Martínez-Sykora, J., De Pontieu, B., & Hansteen, V. 2012, *ApJ*, 753, 161

Martínez-Sykora, J., De Pontieu, B., Hansteen, V. H., et al. 2017, *Science*, 356, 1269

Mather, J. F., Ballai, I., & Erdélyi, R. 2018, *A&A*, 610, A56

Mathioudakis, M., Jess, D. B., & Erdélyi, R. 2013, *Space Science Reviews*, 175, 1

McIntosh, S. W. & Jefferies, S. M. 2006, *ApJL*, 647, L77

McLaughlin, J. A., Nakariakov, V. M., Dominique, M., Jelínek, P., & Takasao, S. 2018, *Space Science Reviews*, 214, 45

Merenda, L., Trujillo Bueno, J., Landi Degl'Innocenti, E., & Collados, M. 2006, *ApJ*, 642, 554

Milligan, R. O., Kerr, G. S., Dennis, B. R., et al. 2014, *ApJ*, 793, 70

Moore, R. L., Sterling, A. C., & Falconer, D. A. 2015, *ApJ*, 806, 11

Moradi, H., Baldner, C., Birch, A., & et al. 2010, *Solar Physics*, 267, 1

Morton, R. J., Verth, G., Hillier, A., & Erdélyi, R. 2014, *ApJ*, 784, 29

Morton, R. J., Verth, G., Jess, D. B., et al. 2012, *Nature Communications*, 3, 1315

Muller, R., Hulot, J. C., & Roudier, T. 1989, *Solar Physics*, 119, 229

Muller, R., Roudier, T., Vigneau, J., & Auffret, H. 1994, *A&A*, 283, 232

Mumford, S. J. & Erdélyi, R. 2015, *mnras*, 449, 1679

Mumford, S. J., Fedun, V., & Erdélyi, R. 2015, *ApJ*, 799, 6

Murabito, M., Romano, P., Guglielmino, S. L., Zuccarello, F., & Solanki, S. K. 2016, *ApJ*, 825, 75

Murray, M. J., van Driel-Gesztelyi, L., & Baker, D. 2009, *A&A*, 494, 329

Musielak, Z. E., Rosner, R., Stein, R. F., & Ulmschneider, P. 1994, *ApJ*, 423, 474

Nagata, S., Tsuneta, S., Suematsu, Y., et al. 2008, *ApJL*, 677, L145

Nakariakov, V. M. & Melnikov, V. F. 2009, *Space Science Reviews*, 149, 119

Narayan, G. 2011, *A&A*, 529, A79

Neidig, D. F. & Wiborg, Jr., P. H. 1984, *Solar Physics*, 92, 217

Nelson, C. J., Scullion, E. M., Doyle, J. G., Freij, N., & Erdélyi, R. 2015, *ApJ*, 798, 19

Ni, L., Kliem, B., Lin, J., & Wu, N. 2015, *ApJ*, 799, 79

Nordlund, A&A. & Stein, R. F. 2001, *ApJ*, 546, 576

Nutto, C., Steiner, O., & Roth, M. 2012, *A&A*, 542, L30

Orozco Suárez, D., Asensio Ramos, A., & Trujillo Bueno, J. 2014, *A&A*, 566, A46

Orozco Suárez, D. & Bellot Rubio, L. R. 2012, *Astrophys. J.*, 751, 2

Orozco Suárez, D., Bellot Rubio, L. R., del Toro Iniesta, J. C., & Tsuneta, S. 2008, *Astron. Astrophys.*, 481, L33

Orozco Suárez, D., Katsukawa, Y., & Bellot Rubio, L. R. 2012, *ApJL*, 758, L38

Ortiz, A., Bellot Rubio, L. R., & Rouppe van der Voort, L. 2010, *ApJ*, 713, 1282

Parker, E. N. 1978, *ApJ*, 221, 368

Pascoe, D. J., Wright, A. N., & De Moortel, I. 2011, *ApJ*, 731, 73

Peter, H., Tian, H., Curdt, W., et al. 2014, *Science*, 346, 1255726

Pettit, E. 1932, *ApJ*, 76, 9

Pietarila, A., Hirzberger, J., Zakharov, V., & Solanki, S. K. 2009, *A&A*, 502, 647

Radziszewski, K., Rudawy, P., & Phillips, K. J. H. 2007, *A&A*, 461, 303

- Rajaguru, S. P., Couvidat, S., Sun, X., Hayashi, K., & Schunker, H. 2012, *Solar Physics*, 303
- Ramelli, R., Balemi, S., Bianda, M., et al. 2010, in *Proceedings of SPIE*, Vol. 7735, *Ground-based and Airborne Instrumentation for Astronomy III*, 77351Y
- Raouafi, N. E., Patsourakos, S., Pariat, E., et al. 2016, *Space Science Reviews*, 201, 1
- Reardon, K. P., Uitenbroek, H., & Cauzzi, G. 2009, *A&A*, 500, 1239
- Reep, J. W. & Russell, A. J. B. 2016, *ApJL*, 818, L20
- Rempel, M. 2011, *ApJ*, 729, 5
- Rempel, M. 2012, *ApJ*, 750, 62
- Rempel, M. 2014, *ApJ*, 789, 132
- Rempel, M. 2015, *ApJ*, 814, 125
- Requerey, I. S., Del Toro Iniesta, J. C., Bellot Rubio, L. R., et al. 2014, *ApJ*, 789, 6
- Requerey, I. S., Del Toro Iniesta, J. C., Bellot Rubio, L. R., et al. 2015, *ApJ*, 810, 79
- Rezaei, R. 2018, *A&A*, 609, A73
- Rezaei, R., Bello González, N., & Schlichenmaier, R. 2012, *A&A*, 537, A19
- Rezaei, R., Steiner, O., Wedemeyer-Böhm, S., et al. 2007, *A&A*, 476, L33
- Reznikova, V. E., Shibasaki, K., Sych, R. A., & Nakariakov, V. M. 2012, *ApJ*, 746, 119
- Riethmüller, T. L. & Solanki, S. K. 2019, *A&A*, 622, A36
- Riethmüller, T. L., Solanki, S. K., Berdyugina, S. V., et al. 2014, *A&A*, 568, A13
- Rijs, C., Rajaguru, S. P., Przybylski, D., et al. 2016, *ApJ*, 817, 45
- Rimmele, T. 2008, *ApJ*, 672, 684
- Robustini, C., Leenaarts, J., & de la Cruz Rodríguez, J. 2018, *A&A*, 609, A14
- Robustini, C., Leenaarts, J., de la Cruz Rodríguez, J., & Rouppe van der Voort, L. 2016, *A&A*, 590, A57
- Romano, P., Frasca, D., Guglielmino, S. L., et al. 2013, *ApJ*, 771, L3
- Romano, P., Guglielmino, S. L., Cristaldi, A., et al. 2014a, *ApJ*, 784, 10
- Romano, P., Zuccarello, F. P., Guglielmino, S. L., & Zuccarello, F. 2014b, *ApJ*, 794, 118
- Rouppe van der Voort, L., Bellot Rubio, L. R., & Ortiz, A. 2010, *ApJL*, 718, L78
- Rouppe van der Voort, L. & de la Cruz Rodríguez, J. 2013, *ApJ*, 776, 56
- Rouppe van der Voort, L., De Pontieu, B., Scharmer, G. B., et al. 2017, *ApJL*, 851, L6
- Rouppe van der Voort, L. H. M., Rutten, R. J., & Vissers, G. J. M. 2016, *A&A*, 592, A100
- Roy, J.-R. 1973, *Solar Physics*, 32, 139
- Ruderman, M. S. & Erdélyi, R. 2009, *Space Science Reviews*, 149, 199
- Russell, A. J. B. & Fletcher, L. 2013, *ApJ*, 765, 81
- Sainz Dalda, A. & Bellot Rubio, L. R. 2008, *A&A*, 481, L21
- Sainz Dalda, A., Martínez-Sykora, J., Bellot Rubio, L., & Title, A. 2012, *ApJ*, 748, 38
- Samanta, T., Tian, H., Banerjee, D., & Schanche, N. 2017, *ApJ*, 835, L19
- Sánchez Almeida, J. & Lites, B. W. 2000, *ApJ*, 532, 1215
- Santamaria, I. C., Khomenko, E., Collados, M., & de Vicente, A. 2016, *A&A*, 590, L3

- Sasso, C., Lagg, A., & Solanki, S. K. 2011, *A&A*, 526, A42
- Sasso, C., Lagg, A., & Solanki, S. K. 2014, *A&A*, 561, A98
- Sasso, C., Lagg, A., Solanki, S. K., Aznar Cuadrado, R., & Collados, M. 2007, in *Astronomical Society of the Pacific Conference Series*, Vol. 368, *The Physics of Chromospheric Plasmas*, ed. P. Heinzel, I. Dorotović, & R. J. Rutten, Heinzel
- Schaffenberger, W., Wedemeyer-Böhm, S., Steiner, O., & Freytag, B. 2006, in *Astronomical Society of the Pacific Conference Series*, Vol. 354, *Solar MHD Theory and Observations: A High Spatial Resolution Perspective*, ed. J. Leibacher, R. F. Stein, & H. Uitenbroek, 345
- Schlichenmaier, R. 2002, *Astronomische Nachrichten*, 323, 303
- Schlichenmaier, R., González, N. B., & Rezaei, R. 2011, in *IAU Symposium*, Vol. 273, *Physics of Sun and Star Spots*, ed. D. Prasad Choudhary & K. G. Strassmeier, 134–140
- Schlichenmaier, R., Rezaei, R., Bello González, N., & Waldmann, T. A. 2010, *A&A*, 512, L1
- Schlichenmaier, R., von der Lühe, O., Hoch, S., et al. 2016, *A&A*, 596, A7
- Schrijver, C. J., Title, A. M., van Ballegooijen, A. A., Hagenaar, H. J., & Shine, R. A. 1997, *ApJ*, 487, 424
- Schüssler, M. & Vögler, A. 2006, *ApJ*, 641, L73
- Schüssler, M. & Vögler, A. 2008, *Astron. Astrophys.*, 481, L5
- Scullion, E., Engvold, O., Lin, Y., & Rouppe van der Voort, L. 2015, *ApJ*, 814, 123
- Scullion, E., Erdélyi, R., Fedun, V., & Doyle, J. G. 2011, *ApJ*, 743, 14
- Shapiro, A. I., Fluri, D. M., Berdyugina, S. V., Bianda, M., & Ramelli, R. 2011, *A&A*, 529, A139
- Sharykin, I. N. & Kosovichev, A. G. 2014, *ApJL*, 788, L18
- Sheeley, Jr., N. R. 1972, *Sol. Phys.*, 25, 98
- Shelyag, S., Cally, P. S., Reid, A., & Mathioudakis, M. 2013, *ApJL*, 776, L4
- Shelyag, S., Fedun, V., Keenan, F. P., Erdélyi, R., & Mathioudakis, M. 2011, *Ann. Geophys.*, 29, 883
- Shelyag, S., Khomenko, E., de Vicente, A., & Przybylski, D. 2016, *ApJL*, 819, L11
- Shimizu, T., Ichimoto, K., & Suematsu, Y. 2012, *ApJ*, 747, L18
- Shimizu, T., Katsukawa, Y., Kubo, M., et al. 2009, *ApJL*, 696, L66
- Shimojo, M., Narukage, N., Kano, R., et al. 2007, *PASJ*, 59, S745
- Simões, P. J. A., Kerr, G. S., Fletcher, L., et al. 2017, *A&A*, 605, A125
- Simon, G. W. & Weiss, N. O. 1970, *Sol. Phys.*, 13, 85
- Singh, K. A. P., Hillier, A., Isobe, H., & Shibata, K. 2015, *PASJ*, 67, 96
- Smitha, H. N., Anusha, L. S., Solanki, S. K., & Riethmüller, T. L. 2017, *Astrophys. J. Suppl. Ser.*, 229, 17
- Snow, B., Fedun, V., Gent, F. A., Verth, G., & Erdélyi, R. 2018, *ApJ*, 857, 125
- Socas-Navarro, H. 2005, *ApJL*, 633, L57
- Socas-Navarro, H. & Manso Sainz, R. 2005, *ApJL*, 620, L71
- Socas-Navarro, H., McIntosh, S. W., Centeno, R., de Wijn, A. G., & Lites, B. W. 2009, *ApJ*, 696, 1683
- Solanki, S. K., Lagg, A., Woch, J., Krupp, N., & Collados, M. 2003, *Nature*, 425, 692
- Solanki, S. K. & Montavon, C. A. P. 1993, *A&A*, 275, 283
- Soler, R., Carbonell, M., Ballester, J. L., & Terradas, J. 2013, *ApJ*, 767, 171
- Soler, R., Díaz, A. J., Ballester, J. L., & Goossens, M. 2012, *ApJ*, 749, 163

- Soler, R., Oliver, R., & Ballester, J. L. 2010, *A&A*, 512, A28
- Spruit, H. C. 1979, *Solar Physics*, 61, 363
- Stangalini, M., Solanki, S. K., Cameron, R., & Martínez Pillet, V. 2013, *A&A*, 554, A115
- Stein, R. 1967, *Solar Physics*, 2, 385
- Stein, R. F. & Nordlund, A&A. 2001, *ApJ*, 546, 585
- Stein, R. F. & Nordlund, Å. 2006, *ApJ*, 642, 1246
- Steiner, O., Grossmann-Doerth, U., Knölker, M., & Schüssler, M. 1998, *ApJ*, 495, 468
- Steiner, O., Hauschildt, P. H., & Bruls, J. 2001, *A&A*, 372, L13
- Steiner, O., Rezaei, R., Schaffenberger, W., & Wedemeyer-Böhm, S. 2008, *Astrophys. J. Lett.*, 680, L85
- Stenflo, J. O. 1973, *Solar Phys.*, 32, 41
- Stenflo, J. O. 1980, *A&A*, 84, 68
- Stenflo, J. O. 1982, *Sol. Phys.*, 80, 209
- Stenflo, J. O. & Keller, C. U. 1997, *A&A*, 321, 927
- Stenflo, J. O., Keller, C. U., & Gandorfer, A. 1998, *A&A*, 329, 319
- Stenflo, J. O., Keller, C. U., & Gandorfer, A. 2000, *A&A*, 355, 789
- Stenflo, J. O., Solanki, S., Harvey, J. W., & Brault, J. W. 1984, *Astron. Astrophys.*, 131, 333
- Sterling, A. C., Moore, R. L., Falconer, D. A., et al. 2016, *ApJ*, 821, 100
- Su, J. T., Shen, Y. D., Liu, Y., Liu, Y., & Mao, X. J. 2012, *ApJ*, 755, 113
- Tandberg-Hanssen, E. 1995, *Science*, 269, 111
- Taroyan, Y. & Erdélyi, R. 2009, *Space Science Reviews*, 149, 229
- Taroyan, Y., Erdélyi, R., Doyle, J. G., & Bradshaw, S. J. 2005, *A&A*, 438, 713
- Thibault, K., Charbonneau, P., & Béland, M. 2014, *ApJ*, 796, 19
- Thornton, L. M. & Parnell, C. E. 2011, *Solar Phys.*, 269, 13
- Tian, H., Yurchyshyn, V., Peter, H., et al. 2018, *ApJ*, 854, 92
- Tiwari, S. K., van Noort, M., Lagg, A., & Solanki, S. K. 2013, *A&A*, 557, A25
- Tomczyk, S., McIntosh, S. W., Keil, S. L., et al. 2007, *Science*, 317, 1192
- Trujillo Bueno, J. 2003, in *Astronomical Society of the Pacific Conference Series*, Vol. 307, *Solar Polarization*, ed. J. Trujillo-Bueno & J. Sanchez Almeida, 407
- Trujillo Bueno, J. 2009, in *Astronomical Society of the Pacific Conference Series*, Vol. 405, *Solar Polarization 5: In Honor of Jan Stenflo*, ed. S. V. Berdyugina, K. N. Nagendra, & R. Ramelli, 65
- Trujillo Bueno, J. 2010, *Astrophysics and Space Science Proceedings*, 19, 118
- Trujillo Bueno, J., Asensio Ramos, A., & Shchukina, N. 2006, in *Astronomical Society of the Pacific Conference Series*, Vol. 358, *Solar Polarization 4*, ed. R. Casini & B. W. Lites, 269
- Trujillo Bueno, J., Landi Degl'Innocenti, E., & Belluzzi, L. 2017, *Space Science Reviews*, 210, 183
- Trujillo Bueno, J., Landi Degl'Innocenti, E., Collados, M., Merenda, L., & Manso Sainz, R. 2002, *Nature*, 415, 403
- Trujillo Bueno, J. & Shchukina, N. 2007, *ApJ*, 664, L135
- Trujillo Bueno, J., Shchukina, N., & Asensio Ramos, A. 2004, *Nature*, 430, 326
- Tsuneta, S., Ichimoto, K., Katsukawa, Y., et al. 2008, *ApJ*, 688, 1374

Ulmschneider, P. 1989, *A&A*, 222, 171

Utz, D., del Toro Iniesta, J. C., Bellot Rubio, L. R., et al. 2014, *ApJ*, 796, 79

Utz, D., Hanslmeier, A., Muller, R., et al. 2010, *A&A*, 511, A39+

Utz, D., Jurčák, J., Hanslmeier, A., et al. 2013, *A&A*, 554, A65

Štěpán, J. & Trujillo Bueno, J. 2016, *ApJ*, 826, L10

Van Doorselaere, T., De Groof, A., Zender, J., Berghmans, D., & Goossens, M. 2011, *ApJ*, 740, 90

Van Doorselaere, T., Kupriyanova, E. G., & Yuan, D. 2016, *Solar Physics*, 291, 3143

Van Doorselaere, T., Nakariakov, V. M., & Verwichte, E. 2008, *ApJL*, 676, L73

van Noort, M. 2012, *A&A*, 548, A5

Vargas Domínguez, S., Rouppe van der Voort, L., Bonet, J. A., et al. 2008, *ApJ*, 679, 900

Vecchio, A., Cauzzi, G., & Reardon, K. P. 2009, *A&A*, 494, 269

Velli, M. & Liewer, P. 1999, *Space Sci. Rev.*, 87, 339

Verma, M., Denker, C., Balthasar, H., et al. 2018, *A&A*, 614, A2

Verth, G., Terradas, J., & Goossens, M. 2010, *ApJL*, 718, L102

Vigeesh, G., Fedun, V., Hasan, S. S., & Erdélyi, R. 2012, *ApJ*, 755, 18

Vigeesh, G., Hasan, S. S., & Steiner, O. 2009, *A&A*, 508, 951

Vissers, G. J. M., Rouppe van der Voort, L. H. M., Rutten, R. J., Carlsson, M., & De Pontieu, B. 2015, *ApJ*, 812, 11

Vitas, N. 2011, PhD thesis, Utrecht University

Vögler, A., Shelyag, S., Schüssler, M., et al. 2005, *A&A*, 429, 335

Wang, H., Liu, C., Qiu, J., et al. 2004, *ApJ*, 601, L195

Wang, Y., Noyes, R. W., Tarbell, T. D., & Title, A. M. 1995, *Astrophys. J.*, 447, 419

Watanabe, H., Kitai, R., & Otsuji, K. 2014, *ApJ*, 796, 77

Wedemeyer-Böhm, S., Scullion, E., Steiner, O., et al. 2012, *Nature*, 486, 505

Wentzel, D. G. 1992, *ApJ*, 388, 211

Woods, M. M., Harra, L. K., Matthews, S. A., et al. 2017, *Solar Physics*, 292, 38

Wyper, P. F., Antiochos, S. K., & DeVore, C. R. 2017, *Nature*, 544, 452

Xu, Y., Cao, W., Jing, J., & Wang, H. 2012a, *ApJL*, 750, L7

Xu, Z., Lagg, A., Solanki, S., & Liu, Y. 2012b, *ApJ*, 749, 138

Yan, L., He, J., Xia, L., & Jiao, F. 2015, *ApJ*, 804, 69

Yang, S., Zhang, J., Erdélyi, R., et al. 2017, *ApJL*, 843, L15

Yang, Z., Tian, H., Peter, H., et al. 2018, *ApJ*, 852, 79

Yelles Chaouche, L., Solanki, S. K., & Schüssler, M. 2009, *A&A*, 504, 595

Yuan, D., Sych, R., Reznikova, V. E., & Nakariakov, V. M. 2014, *A&A*, 561, A19

Yurchyshyn, V., Abramenko, V., Kosovichev, A., & Goode, P. 2014, *ApJ*, 787, 58

Zaqarashvili, T. V., Carbonell, M., Ballester, J. L., & Khodachenko, M. L. 2012, *A&A*, 544, A143

Zeuner, F., Feller, A., Iglesias, F. A., & Solanki, S. K. 2018, *A&A*, 619, A179

Zhao, J., Felipe, T., Chen, R., & Khomenko, E. 2016, *ApJL*, 830, L17

- Zharkova, V. V. & Zharkov, S. I. 2007, ApJ, 664, 573
- Zirin, H. & Stein, A. 1972, Astrophys. J. Lett., 178, L85
- Zirker, J. B., Engvold, O., & Martin, S. F. 1998, Nature, 396, 440
- Zsámberger, N. K., Allcock, M., & Erdélyi, R. 2018, ApJ, 853, 136
- Zuccarello, F., Guglielmino, S. L., & Romano, P. 2014, ApJ, 787, 57
- Zwaan, C. 1987, Annual Review of Astron and Astrophys, 25, 83

C List of Abbreviations

- ASM → Adaptive Secondary Mirror
- AFDT → Auxiliary full-disk telescope
- BBI → Broad Band Imager (cf. page 53)
- EAST → European Association for Solar Telescopes:
<http://www.est-east.eu/est/index.php/people/>
- EST → European Solar Telescope: <http://www.est-east.eu>
- FoV → Field-of-view
- IFU → Integral field unit: Instrument that measures 2D-spatial and spectral dimensions simultaneously. The big advantage of such devices is that image restoration techniques can be applied to enable full spectral integrity at highest spatial resolution. An IFU includes a grating spectrograph, but in contrast to SP which has a long slit, its FOV is 2D. (cf. page 53)
- MCAO → Multi-Conjugate Adaptive Optics: EST will be equipped with a ground-layer adaptive optics system (one deformable mirror in the pupil and one TipTilt mirror). In addition, EST will be equipped with a MCAO system that incorporates 4 deformable mirrors conjugated to four different atmospheric heights aiming to correct a FoV of up to 60 arcsec in diameter.
- NBI → Narrow Band Imager (Fabry Perot system): 2D images sequentially sample spectral positions (cf. page 53).
- PSF → Point-Spread-Function
- R → Spectral Resolution $R = \lambda/\delta\lambda$
- SAG → Science Advisory Group
- SNR → Signal-to-noise ratio: Definition see page 126.
- SP → Spectropolarimeter based on a long-slit spectrograph. To obtain spatial maps the slit must scan across the image (cf. page 53).
- SRD → Science Requirement Document