

CLINICAL IMAGE

A 48-years-old patient with non–ST-elevation myocardial infarction caused by a narrowing in the anomalous circumflex artery

Artur Pawlik¹, Rafał Januszek¹, Dariusz Dudek², Stanisław Bartuś³

¹ 2nd Department of Cardiology and Cardiovascular Interventions, University Hospital, Kraków, Poland

² Department of Interventional Cardiology, Jagiellonian University Medical College, Kraków, Poland

³ 2nd Department of Cardiology, Jagiellonian University Medical College, Kraków, Poland

Coronary artery anomalies are rare findings with an incidence of 0.3% to 1.5%.¹ The circumflex branch of the left coronary artery (Cx) arising from the right coronary artery or from the right Valsalva sinus is the most common type of such malformations.¹ They are considered to be asymptomatic, although cases of exertional pain, myocardial infarction, or even sudden cardiac death have been reported.^{2,3} We present a case of a patient with non–ST-elevated myocardial infarction caused by critical stenosis of the ectopic Cx.

A 48-years-old man was admitted to the emergency department with acute onset of typical angina. He did not have any risk factors for atherosclerosis except for being a heavy smoker for 30 years. After the confirmation of non–ST-elevated myocardial infarction, the patient was redirected to the catheterization laboratory. On admission,

the patient had no chest pain, his blood pressure was 139/76 mm Hg, and heart and lung auscultation was normal. Electrocardiography did not reveal any abnormalities. Lateral and inferior wall hypokinesia of the left ventricle with preserved ejection fraction were observed on the echocardiographic examination. Coronary artery angiography revealed no significant atherosclerotic stenoses in the right or left coronary arteries (**FIGURE 1A** and **1B**). Lack of Cx led to the suspicion of coronary anomaly. Therefore, computed tomography of the coronary arteries (angio-CT) was performed and exposed Cx originating in the right sinus of Valsalva with a possible narrowing within (**FIGURE 1C** and **1D**). The second coronary angiography confirmed ectopic Cx with significant stenosis, and percutaneous coronary angioplasty with drug eluting stent implantation was performed




FIGURE 1 A, B – initial coronary angiogram: no significant narrowing and lack of the circumflex branch are noticeable. Abbreviations: see **FIGURE 1** on the next page

Correspondence to:

Artur Pawlik, MD, 2nd Department of Cardiology and Cardiovascular Interventions, University Hospital in Kraków, ul. Kopernika 17, 31-501 Kraków, Poland, phone: +48 12 424 71 70, email: arturo.pawlik@gmail.com

Received: February 16, 2019.

Revision accepted: March 21, 2019.

Published online: March 21, 2019.

Pol Arch Intern Med. 2019;





129 (6): 424–425

doi:10.20452/pamw.4491

Copyright by Medycyna Praktyczna, Kraków 2019

FIGURE 1 C – angio-CT showing abnormal origin of the circumflex branch from the right Valsalva sinus with possible narrowing in the proximal segment; **D** – critical lesion in the proximal segment of ectopic circumflex branch originating from the right Valsalva sinus (**a**); **E** – final coronary angiogram with the circumflex branch successfully treated by drug eluting stent implantation

Abbreviations: Cx, circumflex coronary artery; Dg, diagonal artery; LAD, left anterior descending artery; RC, right coronary artery



(FIGURE 1E and 1F). After a few days of recovery, the patient was discharged home.

The current follow-up is 4 months from the time of percutaneous coronary intervention and none of the coronary artery disease symptoms were reported. An electrocardiographic exercise treadmill test was negative with 14.2 metabolic equivalents achieved.

Although anomalies of the coronary arteries are usually accidentally found on coronary angiography or autopsy, they may be the cause of serious medical conditions. Also in cases of anomalous Cx, which is considered a benign malformation, life-threatening cardiac events may occur due to the compression of the ectopic artery by the great vessels. Another possible pathophysiological mechanism is accelerated atherosclerosis in the anomalous artery as in the presented patient.⁴

Ectopic coronary arteries may cause numerous technical problems even for greatly experienced operators. They may depart not only from different parts of the aortic bulb and ascending aorta but also from the pulmonary trunk.⁵ In these situations, additional imaging techniques such as angio-CT or magnetic resonance scans are valuable diagnostic methods. When abnormal artery origination is suspected, we should strive for its visualization at all costs because it

may completely change the diagnosis, and may save lives by using proper treatment.

ARTICLE INFORMATION

CONFLICT OF INTEREST None declared.

OPEN ACCESS This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (CC BY-NC-SA 4.0), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, provided the original work is properly cited, distributed under the same license, and used for noncommercial purposes only. For commercial use, please contact the journal office at pamw@mp.pl.

HOW TO CITE Pawlik A, Januszek R, Dudek D, Bartuś S. A 48-years-old patient with non-ST-elevated myocardial infarction caused by narrowing in the anomalous circumflex artery. *Pol Arch Intern Med.* 2019; 129: 424-425. doi:10.20452/pamw.4491

REFERENCES

- Alexander RW, Griffith GC. Anomalies of the coronary arteries and their clinical significance. *Circulation.* 1956; 14: 800-805. [↗](#)
- Taylor AJ, Byers JP, Cheitlin MD, Virmani R. Anomalous right or left coronary artery from the contralateral coronary sinus: "high-risk" abnormalities in the initial coronary artery course and heterogeneous clinical outcomes. *Am Heart J.* 1997; 133: 428-435. [↗](#)
- Piovesana P, Corrado D, Verlato R, et al. Morbidity associated with anomalous origin of the left circumflex coronary artery from the right aortic sinus. *Am J Cardiol.* 1989; 63: 762-763. [↗](#)
- Samarendra P, Kumari S, Hafeez M, et al. Anomalous circumflex coronary artery: benign or predisposed to selective atherosclerosis. *Angiology.* 2001; 52: 521-526. [↗](#)
- Cowles RA, Berdon WE. Bland-White-Garland syndrome of anomalous left coronary artery arising from the pulmonary artery (ALCAPA): A historical review. *Pediatric Radiology.* 2007; 37: 890-895. [↗](#)