



Research Article

Evaluation of Ferritin and Nitric Oxide Levels in Breast Cancer

Rakesh Dhankhar¹, Adarsh Chandramouleswar¹, Kiran Dahiya^{2*}, Veena Singh Ghalaut², Nityasha Nara³, Anil Khurana¹, Prashanta Saha Roy²

¹ Department of Radiotherapy, Pandit Bhagwat Dayal Sharma Postgraduate Institute of Medical Sciences, Rohtak, Haryana, India

² Department of Biochemistry, Pandit Bhagwat Dayal Sharma Postgraduate Institute of Medical Sciences, Rohtak, Haryana, India

³ Department of Surgery, Pandit Bhagwat Dayal Sharma Postgraduate Institute of medical Sciences, Rohtak, Haryana, India

Abstract

Breast carcinoma is one of the commonest malignancies in females. It is curable, if detected, at an early stage. Levels of serum ferritin and nitric oxide are found raised in various cancers including breast carcinoma. These parameters are sufficiently interlinked to be used as markers for breast cancer. In this study, the levels of ferritin and nitric oxide were estimated in 30 patients of early stage (stage I and II) and 30 patients of advanced breast cancer (stage III and IV). These levels were compared with 30 healthy females as controls. Serum ferritin and nitric oxide were found to be raised ($p < 0.001$) in all breast cancer patients as compared to controls. The rise in their levels was significantly more in advanced stage as compared to early stage carcinoma ($p < 0.001$). Treatment had a curative effect on these parameters also as shown by a decrease in their levels in both the groups. Thus, estimation of ferritin and nitric oxide may aid in diagnosis, assessment of severity and monitoring of breast cancer patients though results will be highly reliable in conjunction with other tumor markers.

Keywords: Ferritin; nitric oxide; breast carcinoma

Peer Reviewer: Jaeku Kang, PhD, College of Medicine, Konyang University, Republic of Korea; Anupam Bishayee, PhD, Department of Pharmaceutical Sciences, American University of Health Sciences (AUHS), United States; Kimitoshi Kohno, MD, PhD, University of Occupational & Environmental Health, Japan; Amro Amara, PhD, Pharmaceutics Department, Collage of Pharmacy, King Saud University, Egypt

Received: September 29, 2013; **Accepted:** December 13, 2013; **Published:** February 11, 2014

Competing Interests: The authors have declared that no competing interests exist.

Copyright: 2014 Dahiya K *et al.* This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

***Correspondence to:** Kiran Dahiya, Department of Biochemistry, Pandit Bhagwat Dayal Sharma Postgraduate Institute of Medical Sciences, Rohtak, Haryana, India

Email: kirandahiya_2002@yahoo.com

Introduction

The most important function of ferritin is in iron storage but it performs a number of other significant biological functions in the body. It has, recently, been implicated in the pathogenesis of a number of diseases including cancer. A number of mechanisms such as pro-oxidant and pro-inflammatory pathways are responsible for this [1]. Secretion of ferritin is stimulated by cytokines. Cytokines play an important role in causation of cancer and ferritin plays a prominent role in cytokine response [2].

Oxidants induce ferritin transcription by directly targeting conserved region of ferritin genes. These oxidants, including nitric oxide (NO), may release iron from ferritin either directly or through heme oxygenase [3,4] Cytokines may also affect ferritin translation indirectly through their ability to induce the enzyme inducible nitric oxide synthase (iNOS) and increase NO which, in turn, causes the activation of iron regulatory proteins 1 and 2 (IRP 1 and IRP 2) [2,4]. It has been reported that high levels of NO may be cytostatic or cytotoxic for tumor cells while low levels may have the opposite effect and promote tumor growth. NO has genotoxic and angiogenic properties and also modulates tumor deoxyribonucleic acid (DNA) repair mechanisms. It may mediate induction of ferritin synthesis by selective IRP 2 downregulation [5,6].

Breast cancer is the most common malignant disease and is the chief cause of cancer related mortality among women worldwide [7]. In India, breast cancer is the second most common cancer overall (both sexes) and second most common cancer in incidence and mortality next only to

cervix carcinoma [8]. It accounts for around 9% of all cancers seen in Department of Radiotherapy of our institute [9]. The standard management of breast cancer is by means of combined modality which includes locoregional treatment (surgery and radiotherapy) and systemic treatment (chemotherapy, hormonal therapy and targeted therapy) [10]. The tumor markers in breast carcinoma like CA 15-3 and carcinoembryonic antigen (CEA) are generally useful in the follow-up of the patients with metastatic disease in combination with other diagnostic techniques. This disease is still in the need of more precise biomarkers which might help in early detection, assessment of severity and for prediction of therapy response [7].

Therefore, this study was planned to estimate the levels of ferritin and NO in patients of breast cancer as both these parameters are interrelated and are markers of inflammation, iron metabolism, cell proliferation, angiogenesis and vasomotor activity which are generally found disturbed in malignancies.

Material and methods

Ninety females were enrolled for this study. Out of these, 60 were patients of newly diagnosed/untreated histopathologically proven breast cancer and 30 were healthy, non-anaemic females of age more than 15 years. Informed consent from all the subjects and approval from institutional board of postgraduate studies was obtained beforehand. The pre-treatment evaluation included complete history, general physical examination, routine biochemical and hematological investigations and appropriate radiological assessment (chest X-ray, ultrasonography of abdomen and pelvis for all

patients and computed tomography scan of chest and abdomen whenever indicated). The patients were staged according to American Joint Committee of Cancer (AJCC) staging 2010 (TNM). Pregnant or lactating females and those with any associated chronic medical condition or hemoglobin <10g% were excluded from the study. The subjects were divided into three groups:

Group I: 30 randomly selected apparently healthy females.

Group II: 30 patients with histopathologically proven breast cancer in early stage disease (stage I and II, AJCC-TNM stage).

Group III: 30 patients with histopathologically proven breast cancer in advanced stage disease (stage III and IV, AJCC-TNM stage).

5 mL of venous blood was collected before and 3 weeks after completion of treatment in group II and group III females and only once in group I females from antecubital vein under all aseptic conditions. Treatment modalities included surgery alone, surgery followed by adjuvant chemotherapy, surgery followed by chemoradiotherapy, neoadjuvant chemotherapy followed by adjuvant treatment in stage I, II and III or palliative treatment in stage IV disease. The average time interval for sampling before and after treatment in groups I and II was six months. Serum was separated within an hour and kept at -20 °C for subsequent analysis possible at the earliest.

Ferritin was estimated by chemiluminescence immunoassay (Advia Centaur CP, Siemens, Switzerland, USA) and NO was estimated colorimetrically by Griess reaction [11,12].

The results were analysed by standard statistical techniques which included ANOVA test for comparison of pre-treatment levels in all the

three groups and paired 't' test for comparison in groups II and III before and after treatment.

Results

The mean age of presentation in group I was 47.5 years (25-75 years), group II was 48.2 years (29-65 years) and in group III was 47.9 years (23-75 years). The ratio premenopausal: postmenopausal females in group I was 14%: 16%, group II 12%: 18% and in group III was 16%: 14%. In group III, one patient was unmarried and nulliparous, otherwise all other females were married and parous. All the patients (group II and III) were having Karnofsky Performance Status (KPS) 70 and above. Overall 50% patients had tumor in upper outer quadrant, 10% in upper inner, 13.4% in lower outer, 3.3% in lower inner quadrant and 23.3% in central part of the breast. In group II, 4 patients presented in stage I and 26 patients in stage II. In group III, 25 patients presented in stage III and 5 patients in stage IV. Histopathologically, 29 patients in group II and all 30 in group III were having infiltrating ductal carcinoma and 1 patient in group II presented with DCIS, Paget's disease. Overall 90% (54) of patients underwent modified radical mastectomy and 10% (6) were unsuitable for surgical intervention; 56.7% (34) received neoadjuvant chemotherapy, 88.3% (53) adjuvant and 8.3% (5) received salvage chemotherapy; 75% (45) received radical, 5% (3) palliative and 20% (12) received no radiotherapy. About hormonal therapy, 48.3% (29) were administered tamoxifen, 33.3% (20) letrozole, 3.3% (2) anastrozole and 15% (9) patients were given no hormonal treatment. The levels of ferritin and NO before (all the three groups) and after treatment (group II and group III) are shown in Table 1.

Table 1 Serum ferritin and NO (mean \pm SD) in healthy controls (group I) and early (group II) and advanced stage (group III) breast carcinoma before and after treatment

Group I	Group II		Group III		
	Before treatment	After treatment	Before treatment	After treatment	
Ferritin (μ g/L)	33.26 \pm 16.34 (18.7-41.8)	73.73 \pm 42.7* (35.0-116.4)	61.6 \pm 25.8*# (35.8-87.4)	103.4 \pm 83.1* (20.3-186.5)	84.7 \pm 38.4*# (46.3-123.1)
NO (μ mol/L)	21.6 \pm 8.67 (11.3-29.6)	40.8 \pm 16.2* (24.6-57.0)	27.5 \pm 12.4*§ (15.1-39.9)	70.7 \pm 69.0* (1.7-139.7)	27.2 \pm 12.4*§ (14.8-39.6)

* p value <0.001 as compared to group I.

p value <0.001 as compared to levels before treatment.

§ p value <0.05 as compared to levels before treatment.

Discussion

The levels of ferritin were found to be significantly raised ($p < 0.001$) in breast cancer patients (in both groups) as compared to controls. After treatment, these levels were found to be decreased significantly in groups II and III. It has been reported in literature that ferritin levels are raised in breast and other solid cancers [13,14]. This increase may either be due to increased expression of a tumor derived protein which interferes with iron metabolism or due to non-specific effect of malignancy on reticulo-endothelial iron metabolism [14, 15]. The rise in ferritin levels is reported to be linked with more advanced stage breast cancer and high ferritin levels are associated with more advanced malignant phenotype [15-17]. In our study also, the levels in group III were significantly higher as compared to group II ($p < 0.001$).

Rise in serum ferritin levels may be attributed to increased iron requirement by malignant cells for growth and for modulation of transferrin receptor. Transferrin receptors on proliferating and malignant

cells are well documented and are considered potential markers for identifying cells undergoing divisional activity and requiring the incorporation of additional iron [18]. Ferritin levels have been found raised in breast cancer tissue also [19]. In addition to increased synthesis by malignant cells, other causes of raised serum levels include presence of inflammation, hepatic necrosis due to metastasis and reduced hepatic clearance of ferritin [20]. All these factors might have been responsible for higher ferritin levels in advanced stage as compared to early breast cancer.

The levels of nitric oxide were found raised ($p < 0.001$) in both groups II and III as compared to group I. The difference in NO levels between group II and group III was also statistically significant ($p < 0.001$). NO influences various aspects of tumor biology like modulation of cell growth, apoptosis, differentiation, angiogenesis and metastatic capability. Mechanism of action of nitric oxide involves inhibition of DNA synthesis, mitochondrial respiration and enhancement of cellular oxidative injury. This

increased oxidative stress may be implicated in increasing ferritin secretion by breast tissue [21]. After treatment, levels were found to decline significantly in both group II ($p=0.002$) and group III (0.001).

Hypoxia, often associated with neoplastic tissues, is another factor which may induce ferritin as well as NO production [22]. NO, acting as an oxidant, may induce ferritin synthesis [4]. Thus, both these markers are sufficiently interlinked in malignant conditions including carcinoma breast. The tumor markers being presently used for breast carcinoma like CA 15-3 and CEA are also not specific and are found raised in other diseases too. The utility of these markers is more if analysed in conjunction with other markers rather than relying on a single one [7].

Therefore, it is concluded that ferritin and NO may help in assessing the severity and monitoring of breast cancer patients. As both these markers are non-specific and their mechanism of action is multifactorial, these might prove more useful if combined with other biomarkers for breast carcinoma.

References

1. Thiel EC. The ferritin family of iron storage proteins. *Adv Enzymol Relat Areas Mol Biol.* 1990, 63: 421-449
2. Tran TN, Eubanks SK, Schaffer KJ, Zhou CY, Linder MC. Secretion of ferritin by rat hepatoma cells and its regulation by inflammatory cytokines and iron. *Blood.* 1997, 90: 4979-4986
3. Pantopoulos K, Hentze MW. Activation of iron regulatory protein-1 by oxidative stress in vitro. *Proc Natl Acad Sci USA.* 1998, 95: 10559-10563
4. Orino K, Lehman L, Tsuji Y, Ayaki H, Torti SV, Torti FM. Ferritin and the response to oxidative stress. *Biochem J.* 2001, 357: 241-247
5. Xu W, Liu LZ, Loizidou M, Ahmed M, Charles IG. The role of nitric oxide in cancer. *Cell Res.* 2002, 12: 311-320
6. Fahmy M, Young SP. Modulation of iron metabolism in monocyte cell line U937 by inflammatory cytokines: changes in transferrin uptake, iron handling and ferritin mRNA. *Biochem J.* 1994, 296: 175-181
7. Ćujić D, Stefanoska I, Golubović S. Serum Ferritin in healthy women and breast cancer patients. *J Med Biochem.* 2011, 30: 33-37
8. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer.* 2010, 127: 2893-2917
9. Das BP. Cancer patterns in Haryana: Twenty-one years' experience. *Radiation Oncology.* 2005, 5: 22-32
10. Sauer R. Adjuvant radiotherapy after breast conserving surgery for breast cancer. *Pro Eur J Cancer.* 2000, 36: 1073-1078
11. Dupuy AM, Debarge L, Poulain M, Badiou S, Rossi M, Cristol JP. Determination of serum ferritin using immunoturbidimetry or chemiluminescent detection in comparison with radioimmunoassay- a compendium of a methodological juxtaposition. *Clin Lab.* 2009, 55: 207-215
12. Dimitrios T. Analysis of nitrite and nitrate in biological fluids by assays based on Griess reaction: appraisal of the Griess reaction in the L-arginine/nitric oxide area of research. *J Chromatography.* 2007, 851: 51-70
13. Wang W, Knovich MA, Coffman LG, Torti FM, Torti SV. Serum ferritin: past, present and future. *Biochim Biophys Acta.* 2010, 1800: 760-769
14. Jacobs A, Jones B, Ricketts C, Bulbrook RD, Wang DY. Serum ferritin concentration in early breast cancer. *Br J Cancer.* 1976, 34: 286-289
15. Shpyleva SI, TryndyakVP, Kovalchuk O, Starlard-Davenport A, Chekhun VF, Beland FA,

- Pogribny IP. Role of ferritin alterations in human breast cancer cells. *Breast Cancer Res Treat.* 2011, 126: 63-71
16. Alkhateeb AA, Han B, Connor JR. Ferritin stimulates breast cancer cells through an iron-independent mechanism and is localized within tumor-associated macrophages. *Breast Cancer Res Treat.* 2013, 137: 733-744
 17. Mishra S, Sharma DC, Sharma P. Studies of biochemical parameters in breast cancer with and without metastasis. *Indian J Clin Biochem.* 2004, 19: 71-75
 18. Elliot RL, Elliot MC, Wang F, Head JF. Breast carcinoma and the role of iron metabolism. A cytochemical tissue, culture and ultrasonic study. *Ann NY Acad Sci.* 1993, 698: 159-166
 19. Weinstein RE, Bond BH, Silberg BK. Tissue ferritin concentration in carcinoma of breast. *Cancer.* 1982, 50: 2406-2409
 20. Toyokuni S. Iron and carcinogenesis: from fenton reaction to target genes. *Redox Rep.* 2002, 7: 189-197
 21. Sayed-Ahmad MM, Shouman SA, Ramadan LA. Contribution of Nitric oxide in doxorubicin-related cytotoxicity in murine tumor model. *J Egyptian Nat Cancer Inst.* 2002, 14: 111-116
 22. Qi Y, Jamindar TM, Dawson GJ. Hypoxia alters iron homeostasis and induces ferritin synthesis in oligodendrocytes. *Neurochem.* 1995, 64: 2458-2464