

Swarthmore College

Works

Biology Faculty Works

Biology

12-5-2019

Pharmacological Or Genetic Targeting Of Transient Receptor Potential (TRP) Channels Can Disrupt The Planarian Escape Response

Ziad Sabry , '21

A. Ho

D. Ireland

C. Rabeler

O. Cochet-Escartin

See next page for additional authors

Follow this and additional works at: <https://works.swarthmore.edu/fac-biology>



Part of the [Biology Commons](#), and the [Systems Biology Commons](#)

[Let us know how access to these works benefits you](#)

Recommended Citation

Ziad Sabry , '21; A. Ho; D. Ireland; C. Rabeler; O. Cochet-Escartin; and Eva-Maria S. Collins. (2019). "Pharmacological Or Genetic Targeting Of Transient Receptor Potential (TRP) Channels Can Disrupt The Planarian Escape Response". *PLoS ONE*. Volume 14, Issue 12. e0226104 DOI: 10.1371/journal.pone.0226104
<https://works.swarthmore.edu/fac-biology/592>



This work is licensed under a [Creative Commons Attribution 4.0 License](#).

This work is brought to you for free and open access by . It has been accepted for inclusion in Biology Faculty Works by an authorized administrator of Works. For more information, please contact myworks@swarthmore.edu.

Authors

Ziad Sabry , '21; A. Ho; D. Ireland; C. Rabeler; O. Cochet-Escartin; and Eva-Maria S. Collins

RESEARCH ARTICLE


Pharmacological or genetic targeting of Transient Receptor Potential (TRP) channels can disrupt the planarian escape response

Ziad Sabry¹, Alicia Ho², Danielle Ireland^{1,2}, Christina Rabeler¹, Olivier Cochet-Escartin^{3*}, Eva-Maria S. Collins^{1,2,3*}

1 Department of Biology, Swarthmore College, Swarthmore, Pennsylvania, United States of America, **2** Section of Cell and Developmental Biology, Division of Biological Sciences, University of California San Diego, La Jolla, California, United States of America, **3** Department of Physics, University of California San Diego, La Jolla, California, United States of America

* Current address: Institut Lumière Matière, UMR5306, Université Lyon 1-CNRS, Université de Lyon, Villeurbanne, France

* ecollin3@swarthmore.edu



OPEN ACCESS

Citation: Sabry Z, Ho A, Ireland D, Rabeler C, Cochet-Escartin O, Collins E-MS (2019) Pharmacological or genetic targeting of Transient Receptor Potential (TRP) channels can disrupt the planarian escape response. PLoS ONE 14(12): e0226104. <https://doi.org/10.1371/journal.pone.0226104>

Editor: Shang-Zhong Xu, University of Hull, UNITED KINGDOM

Received: August 23, 2019

Accepted: November 19, 2019

Published: December 5, 2019

Copyright: © 2019 Sabry et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: The following grants provided support for this study: Burroughs Wellcome CASI award (E-MS), NSF CAREER award (E-MS). Partial salary support was provided by the Marye Anne Fox Endowed Fellowship (DI). The funders had no role in study design, data collection and analysis,

Abstract

In response to noxious stimuli, planarians cease their typical ciliary gliding and exhibit an oscillatory type of locomotion called scrunching. We have previously characterized the bio-mechanics of scrunching and shown that it is induced by specific stimuli, such as amputation, noxious heat, and extreme pH. Because these specific inducers are known to activate Transient Receptor Potential (TRP) channels in other systems, we hypothesized that TRP channels control scrunching. We found that chemicals known to activate TRPA1 (allyl isothiocyanate (AITC) and hydrogen peroxide) and TRPV (capsaicin and anandamide) in other systems induce scrunching in the planarian species *Dugesia japonica* and, except for anandamide, in *Schmidtea mediterranea*. To confirm that these responses were specific to either TRPA1 or TRPV, respectively, we tried to block scrunching using selective TRPA1 or TRPV antagonists and RNA interference (RNAi) mediated knockdown. Unexpectedly, co-treatment with a mammalian TRPA1 antagonist, HC-030031, enhanced AITC-induced scrunching by decreasing the latency time, suggesting an agonistic relationship in planarians. We further confirmed that TRPA1 in both planarian species is necessary for AITC-induced scrunching using RNAi. Conversely, while co-treatment of a mammalian TRPV antagonist, SB-366791, also enhanced capsaicin-induced reactions in *D. japonica*, combined knock-down of two previously identified *D. japonica* TRPV genes (*DjTRPVa* and *DjTRPVb*) did not inhibit capsaicin-induced scrunching. RNAi of *DjTRPVa/DjTRPVb* attenuated scrunching induced by the endocannabinoid and TRPV agonist, anandamide. Overall, our results show that although scrunching induction can involve different initial pathways for sensing stimuli, this behavior's signature dynamical features are independent of the inducer, implying that scrunching is a stereotypical planarian escape behavior in response to various noxious stimuli that converge on a single downstream pathway. Understanding which aspects of nociception are conserved or not across different organisms can provide insight into the underlying regulatory mechanisms to better understand pain sensation.

decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Normal locomotion of freshwater planarians, termed gliding, is achieved through synchronous beating of cilia in a layer of secreted mucus [1–3]. Gliding planarians display a smooth motion without major body shape changes, except for turning movements of the anterior end. However, when exposed to certain noxious stimuli (e.g. low pH, high temperature, or amputation), planarians switch to a muscular-based escape gait that is characterized by oscillatory body length changes [4]. We termed this gait scrunching and showed that it has a characteristic set of 4 quantifiable parameters: 1. frequency of body length oscillations, 2. relative speed, 3. maximum amplitude, and 4. asymmetry of body elongation and contraction [4]. Moreover, scrunching is conserved among different planarian species, with each species demonstrating a characteristic frequency and speed. Although scrunching shares similarities with peristalsis, another muscle-based oscillatory gait that occurs when cilia beating is disrupted [2,5–7], scrunching is cilia-independent, can be induced in animals performing peristalsis, and is distinguishable from peristalsis based on the 4 parameters listed above [4], demonstrating that scrunching and peristalsis are distinct gaits. Because scrunching is such a stereotypical response involving many steps of neuronal communication (sensation, processing, neuromuscular communication), scrunching in response to noxious heat has proven to be a useful and sensitive readout of neuronal function in planarian toxicological studies [8,9]. However, which molecular targets and neuronal circuits regulate scrunching remain an open question.

Recently, it has been shown using RNA interference (RNAi) that the Transient Receptor Potential (TRP) channel, TRP ankyrin 1 (TRPA1), is required for avoidance behaviors in *Schmidtea mediterranea* in response to noxious heat and the pungent ingredient in mustard oil, allyl isothiocyanate (AITC) [10]. The authors also showed that the noxious heat response was mediated by H₂O₂ and/or reactive oxygen species which directly activate SmTRPA1, causing the planarian to avoid hot regions. Moreover, in response to physical injury (tail snips), *Smed-TRPA1*-knockdown worms scrunched with a significantly reduced amplitude [10]. Based on these results, the authors hypothesized that SmTRPA1 signaling, induced through H₂O₂ upregulation at the site of wounding, may regulate amputation-induced-scrunching in *S. mediterranea*. Whether H₂O₂ exposure alone induces scrunching or whether SmTRPA1 also plays a role in triggering scrunching in response to other stimuli is still unknown.

TRPA1 is a member of the TRP superfamily, comprised of widely conserved transmembrane, nonselective cation channels [11]. TRP channels mediate responses to almost all classes of external stimuli, including various nociceptive stimuli such as extreme temperatures, ultraviolet light, and specific chemical irritants, and as such mediate the initial steps of pain sensation [11–13]. TRPs are classified into sub-families depending on their main mode of activation (mechanical, thermal, chemical. . .), but are often polymodal, integrating different stimuli in the same channel [11,13,14].

In addition to TRPA1, TRPV channels are also good candidates for possibly regulating scrunching. Scrunching is activated by low pH and noxious heat [4], stimuli which are known to activate members of the TRP vanilloid (TRPV) sub-family, named after their sensitivity to vanilloid compounds such as capsaicin, in vertebrates and invertebrates [15–20]. TRPV channels are activated by a diverse range of stimuli and exhibit a high level of species-dependent functional differences [20,21]. For example, while human and rat TRPV1 are highly sensitive to capsaicin, rabbit and bird have greatly reduced sensitivities [21–23]. Historically it was thought that, like fruit flies and nematodes [24–26], most invertebrates were also insensitive to this chili pepper irritant [27]. However, medicinal leech was recently found to contain a capsaicin-sensitive TRPV channel [16]. Interestingly, although no TRPV homologs exist in the parasitic flatworm *Schistosoma mansoni*, it was shown that TRPA1 in this species mediates the

behavioral response to capsaicin [28,29]. Previous analysis of a *Dugesia japonica* transcriptome has estimated that at least 25 TRP homologs may be present in this planarian species [30]. Thus far, partial sequences for seven TRP genes have been identified and their expression profiles characterized [31]. DjTRP_{Ma} channels have been shown to regulate thermotaxis behavior at lower temperatures (0–25°C) [31], but the function of DjTRPA and DjTRPV channels has not yet been studied in this species.

Thus, based on previous literature in planarians and the known activators of scrunching, we hypothesized that planarian TRPA1 and TRPV channels control scrunching to specific stimuli. To test this hypothesis, we first assayed a variety of chemicals for their ability to induce scrunching in two freshwater planarian species, *S. mediterranea* and *D. japonica*. We focused on chemical compounds that have been shown to activate TRPA1 and/or TRPV, such as AITC, an agonist of planarian [10] and mammalian [32,33] TRPA1, and capsaicin, a specific agonist of mammalian TRPV1 [34,35].

We found that scrunching is a specific response to known modulators of mammalian TRPA1 or TRPV channels, including AITC and capsaicin. These findings were substantiated by knocking down either *SmTRPA1*, *DjTRPAa* or *DjTRPVa/DjTRPVb* using RNAi and evaluating the behaviors of these worms when exposed to a subset of the confirmed scrunching inducers. We found that planarian TRPA1, and partially TRPV, modulate scrunching in response to different triggers.

The observation that scrunching is a stereotypical response that is the same for different stimuli and sensing mechanisms suggests the existence of a single convergent pathway that regulates scrunching downstream of planarian TRP activation. TRP channels are involved in various chemical and physical sensing capacities across eukaryotes, from yeast to humans, yet exhibit a high level of diversity, both within the superfamily and across species [11,13]. By understanding how these channels are used in different species, such as planarians, we gain better insight into their regulatory mechanisms, with the potential to reveal elements important to control pain sensation.

Materials and methods

Animal care

Asexual freshwater planarians of the species *Dugesia japonica* and *Schmidtea mediterranea* were used for all experiments. The animals were fed organic chicken or beef liver 1–2 times per week, cleaned twice per week, and starved for 5–7 days prior to experimentation. Planarians were stored in a temperature-controlled Panasonic incubator in the dark at 20°C with *D. japonica* in dilute (0.5 g/L) Instant Ocean (IO) water (Spectrum Brands, Blacksburg, VA, USA) and *S. mediterranea* in 1X Montjuïc Salts (MS) water [36].

Behavioral assays

Pharmacological perturbations. All chemicals used are listed in Table 1. Chemicals were stored according to supplier specifications. All stock solutions were made directly in IO or MS water or in 100% dimethyl sulfoxide (DMSO, Sigma-Aldrich, St. Louis, MO, USA), depending on chemical solubility. For chemicals prepared in DMSO, the final DMSO concentrations were kept ≤1%, which does not induce scrunching (S1 Fig). Specific conditions for each chemical experiment were determined empirically by qualitative observation of planarian scrunching behavior (Table 1). The lowest exposure concentration tested which induced the most straight-line scrunches in wildtype worms was used for each experiment. The pH of all exposure solutions (except for hydrogen chloride) was checked and adjusted with NaOH to fall between 6.90–7.10, to ensure the observed scrunching behavior was not due to low pH

Table 1. Overview of chemicals used to induce scrunching.

Chemical	CAS	Provider	Purity	Exposure concentration	Exposure method	Type/action and references
Allyl isothiocyanate (AITC)	57-06-7	Sigma-Aldrich	95%	Dj, Sm—50, 75, 100 μ M	25 mL bath	Planarian and mammalian TRPA1 agonist [10,32,37]
Hydrogen peroxide (H_2O_2)	7722-84-1	Sigma-Aldrich	30%	Dj, Sm—40 mM	25 mL bath	Planarian TRPA1 agonist [10]
Capsaicin	404-86-4	Cayman Chemicals	\geq 95%	Dj, Sm—33, 82.5, 165 μ M	25 mL bath	TRPV1 agonist (various species) [15,20,32,34,35]
Anandamide	94421-68-8	Sigma-Aldrich	\geq 97%	Dj— 100, 125 μ M Sm— 100 μ M	25 mL bath	Endocannabinoid and mammalian TRPV1 agonist [35,38]
HC-030031	349085-38-7	Cayman Chemicals	\geq 98%	Dj, Sm— 100 μ M	25 mL bath	Human, rat, mouse, medicinal leech TRPA1 antagonist [37,39,40]
SB-366791	472981-92-3	Cayman Chemicals	\geq 98%	Dj, Sm— 1, 10 μ M	25 mL bath	Rat, parasitic flatworm and medicinal leech TRPV1 antagonist [16,41–43]
Hydrogen chloride (HCl)	7647-01-0	Sigma-Aldrich	36.5–38.0%	Dj, Sm—pH to 2.7	10 μ L pipette	Low pH, planarian scrunching inducer [4]

<https://doi.org/10.1371/journal.pone.0226104.t001>

conditions. Planarians were exposed to the chemicals either in a bathing solution or by pipetting a fixed volume directly onto a worm. Pipetting allowed for the usage of small volumes of locally higher chemical concentrations and was used in cases of poor chemical solubility or when baths failed to produce sufficiently long stretches of straight-line scrunching that could be used for quantitative analysis. Working solutions for chemicals were made fresh immediately prior to starting experiments. Planarians were individually placed into 100 x 15-mm or 60 x 15-mm petri dishes (Celltreat Scientific Products, Shirley, MA, USA) depending on whether experiments were conducted in baths or by pipetting, respectively. The behavior of each planarian was recorded, starting immediately after initial exposure to the chemical, for up to five minutes at 10 frames per second (fps) using a charge-coupled device camera (PointGrey Flea3 1.3MP Mono USB 3.0) with a 16-mm lens (Tamron M118FM16 Megapixel Fixed-focal Industrial Lens) attached to a ring stand.

Amputation experiments. Individual *S. mediterranea* planarians were placed into 100 x 15-mm petri dishes containing 15 mL of MS water. Using a razor blade, planarians were amputated just above the pharynx. The behavior of each planarian was recorded at 10 fps using the same setup as in the chemical assays. The number of scrunches of the resulting head piece was counted for each amputation with the scrunching sequence beginning after the first immediate contraction.

High temperature experiments. 60 mm x 15-mm petri dishes were filled with 5 mL of either IO or MS water. Individual planarians were placed in the dishes and induced to scrunch by pipetting 100 μ L 65°C IO/MS water as in [4]. Additionally, we tested the effects of a heated water bath using an automated set-up for screening individual planarians in a 48-well plate [8]. To induce scrunching, the plate was placed on a warmed peltier plate (TE Technology Inc., Traverse City, MI), whose temperature was computer controlled to heat the water in the wells. The temperature of the peltier was initially set to 65°C for the first 30 seconds to quickly heat up the plate from room temperature and then gradually decreased to 43°C to stabilize the aquatic temperature across the plate at around 32°C for 4 minutes. The plate was imaged from above and the movies were analyzed using a custom, automated MATLAB (MathWorks, Natick, MA, USA) script to detect instances of scrunching, as previously described [8].

Scrunching quantification. Recordings of planarian behavioral responses to the noxious stimuli were processed using ImageJ (National Institutes of Health, Bethesda, MD, USA). The

background-subtracted image sequences were cropped to capture the first set of at least four consecutive straight-line scrunches or oscillations. An ellipse was then fit to the sequence of binary images of the worm to track and quantify the major axis (length of the worm) over time. From these data, the parameters frequency (number of scrunching/oscillation cycles per second), maximum elongation (difference between longest and shortest elongations/contractions as a fraction of the longest), relative speed (product of maximum elongation and frequency), and fraction of time spent elongating were then quantified in MATLAB as in [4]. Unless stated otherwise, all values denote mean \pm standard deviation (SD). Statistical significance for each scrunching parameter (or number of scrunches for amputation experiments) was calculated using a student's t-test comparing to either previously published values for amputation for wild-type animals or to the *control* RNAi population for RNAi experiments.

Behavior scoring. Recordings of worms in chemical baths were scored by 2 blind reviewers. For every 15 s interval in the first 90 s of recording, worms were scored as either scrunching, exhibiting a non-scrunching reaction, or not reacting. Worms were scored as scrunching if they scrunched at least once in a given 15 s interval, based on the definitions set in [4]. Worms were scored as exhibiting a non-scrunching behavior if they performed other behaviors, such as head shaking, frequent turning or abnormally long body elongation. Planarians that glided unhindered throughout the 15 s interval were scored to have no reaction. Three experimental replicates were carried out for each condition, with $N = 8$ *S. mediterranea* and $N = 10$ *D. japonica* used per replicate. The mean of the scored responses from the two reviewers and across the experimental replicates for each 15 s interval are shown in the respective figures. To compare the timing of the initiation of scrunching in AITC or capsaicin alone or with the addition of HC-030031 or SB-366791, respectively, a Fisher's exact test was performed by comparing the number of planarians scrunching or not scrunching (no reaction or a non-scrunching reaction) in two early time periods (15–30, and 31–45 s), using the averaged numbers from the two reviewers. A Fisher's exact test was also performed to compare the number of planarians reacting (scrunching or exhibiting a non-scrunching reaction) or not reacting in *control* versus *DjTRPVab* or *DjTRPAa* RNAi populations.

Mucus staining

Staining procedures were performed using fluorescently labeled *Vicia villosa* (VVA) lectins as described previously [4]. *D. japonica* planarians were individually placed in wells and induced to scrunch atop a glass coverslip using baths of 33 μ M capsaicin or 50 μ M AITC. The same staining procedure was followed for *S. mediterranea* planarians with scrunching being induced by a bath of 33 μ M capsaicin or 50 μ M AITC. Mucus trails were imaged in 4x under GFP fluorescence using a Nikon Eclipse Ci microscope (Nikon Corporation, Minato, Tokyo, Japan). Images were stitched together using Fiji [44] and the MosaicJ plugin [45].

Cilia imaging

To view cilia beating, *D. japonica* and *S. mediterranea* planarians were incubated for five minutes in baths of 100 μ M anandamide before mounting between a glass slide and a 22 \times 22' coverslip. Imaging procedures were performed as previously described in (4).

Partial cloning of *Dugesia japonica* TRP genes

Partial mRNA sequences for TRPA1 (*DjTRPAa*) and TRPV (*DjTRPVa* and *DjTRPVb*) homologs in *D. japonica* have been previously published [31]. Primers were designed using Primer3 [46] from these templates for *DjTRPVa* and *DjTRPVb* to generate 213 and 430 bp fragments, respectively (Table 2). For *DjTRPAa*, using the published partial sequence as a starting point,

Table 2. Primers used in this study.

Gene	Fragment length (bp)	Forward primer	Reverse primer
<i>DjTRPAa</i>	895	GCAATTAATGACCGAGCAAAC	AACCGATTTCGTTCAAAGTGG
<i>DjTRPVa</i>	213	TATTGAGTGCGCCAATGAAA	AATCACCGCGAACCATTTTA
<i>DjTRPVb</i>	430	TCCATTACTTTGGATGGGTTTAC	TTTTGCCCAAATTGCTATCC
<i>DjTRPAa-qPCR</i>	109	TCGAGGGGAAATGCCAATG	ACTTGAGCTTCAGATGAGCC
<i>DjTRPVa-qPCR</i>	89	ATTTCGCGAAGATGAACACGG	GCCCCTCTTTGGTCAATGTC
<i>DjTRPVb-qPCR</i>	142	ATAAGTGCCTCCAATCATTGC	TCTCGGTGAATCAAGCTGC
<i>SmTRPA1-qPCR</i>	99	CCTCGTGTGGAAATAGTGCG	TGGGACTACAGACTAACGCG

<https://doi.org/10.1371/journal.pone.0226104.t002>

we blasted against a *D. japonica* transcriptome (dd_Djap_v4) on PlanMine [47] to identify the full coding sequence (transcript dd_Djap_v4_9060_1_1). An 895 bp fragment was identified from this transcript and cloned using the primers in Table 2.

These fragments were cloned into the pPR244-TRP vector using ligase independent cloning [48]. The *Smed-TRPA1-pGEMt* plasmid was a gift from Dr. Marco Gallio [10].

Sequence alignments

Full coding sequences for *DjTRPVa* and *DjTRPVb* were obtained by blasting the published partial sequences [31] on PlanMine [47]. Predicted protein sequences for SmTRPA1, DjTRPA1, DjTRPVa, and DjTRPVb were obtained using the National Center for Biotechnology Information Open Reading Frame Finder (<https://www.ncbi.nlm.nih.gov/orffinder/>). Alignments were created with JalView [49] using T-Coffee [50].

RNAi feedings and injections

Expression of *SmTRPA1* and *DjTRPAa* were knocked down separately in their respective species. Expression of *DjTRPVa* and *DjTRPVb* were knocked down in combination and referred to as *DjTRPVab* RNAi. The respective TRP genes of interest were knocked down by injecting *S. mediterranea* or *D. japonica* worms on four consecutive days with *in vitro* transcribed dsRNA to a final concentration of at least 1 µg/µL as in [51]. Negative control populations of both species, denoted as *control* RNAi, were injected with *unc22* dsRNA, a non-homologous *C. elegans* gene. Approximately 180 nL dsRNA were injected into each worm per day of injection using a standard dissection microscope and Pneumatic PicoPump Model PV 820 (World Precision Instruments, Sarasota, FL, USA). Needles were made by pulling 1-mm/0.58-mm OD/ID Kwik-Fill borosilicate glass capillaries through a two-stage program on a P-1000 micropipette puller (Sutter Instrument Company, Novato, CA, USA). On the fourth day of injection, after the fourth injection had been administered, worms were fed organic chicken liver mixed with at least 1 µg/µL dsRNA. Worms were then starved for six days prior to experiments.

qRT-PCR

RNA was extracted from ten worms for each RNAi population using TRIzol (Invitrogen, Carlsbad, CA, USA) then purified using an RNeasy Mini Kit (QIAGEN, Germantown, MD, USA) including a DNase treatment. cDNA was synthesized from each RNA pool using the SuperScript® III First-Strand Synthesis System for RT-PCR (Invitrogen, Carlsbad, CA, USA), following the manufacturer's protocol and priming with random hexamers. Primers for qPCR were designed using Primer3 [46] and are listed in Table 2.

DjGAPDH and *SmedGAPDH* were used as housekeeping genes for their respective species. qPCR was performed on an MJ Research PTC-200 thermocycler equipped with a Chromo4 Real-Time PCR Detector (Bio-Rad Laboratories, Hercules, CA, USA), using PerfeCTa® SYBR® Green FastMix® (Quantabio, Beverly, MA, USA). Technical triplicates were run for all reactions within an experiment, and two biological replicates were performed. To analyze primer efficiency, standard curves were obtained using a 1:1:1:1 mix of all cDNA pools for each species, serially diluted. The efficiency for each primer pair was found to be between 87–116%. Analysis of relative expression for the genes targeted by RNAi was performed using the $\Delta\Delta C_t$ method, where reported values are the mean of all replicates.

Results and discussion

TRPA1 and TRPV agonists induce scrunching in *S. mediterranea* and *D. japonica*

Based on the known inducers of scrunching in *D. japonica* and *S. mediterranea* (noxious heat, pH, amputation) ([4] and S2 Fig), and the recent work by Arenas et al. suggesting that TRPA1 mediates scrunching in response to amputation in *S. mediterranea* [10], we tested known chemical agonists of planarian and/or other species' TRPA1 and TRPV (Table 1) for their ability to trigger scrunching in *D. japonica* and *S. mediterranea*. The two oscillatory planarian gaits, scrunching and peristalsis, can be hard to distinguish qualitatively by eye. Thus, if oscillatory motion was observed, we distinguished peristalsis and scrunching by quantifying the 4 characteristic parameters (frequency, speed, maximum elongation, asymmetry of elongation/contraction) and comparing with published reference values for these gaits [4]. Notably, asymmetry between contraction and elongation cycles is the most distinct feature of scrunching that is conserved among different planarian species [4]. We previously found that each planarian species exhibits a characteristic scrunching frequency and speed, with *D. japonica* scrunching at higher speeds and with almost double the frequency of *S. mediterranea* planarians [4]. Therefore, all comparisons are done with references in the same species.

Under normal conditions, planarians glide, maintaining a constant body length over time (Fig 1A and 1B). When exposed to 50 μM of the TRPA1 activator, AITC, planarians scrunched showing oscillations of body length elongation and contraction (Fig 1C) with quantitative parameters consistent with those previously determined for *D. japonica* and *S. mediterranea* using amputation (Table 3).

Previous studies in *S. mediterranea* demonstrated that TRPA1 is directly activated by H_2O_2 [10]; therefore, we assayed whether H_2O_2 exposure could induce scrunching. As expected from the results of AITC exposure, 40 mM H_2O_2 elicited scrunching in both planarian species (Fig 1D and Table 3). Notably, while *S. mediterranea* scrunching parameters were statistically insignificant from those induced by amputation, *D. japonica* scrunched at slightly (but statistically significant) lower speeds in 40 mM H_2O_2 compared to the reference values for amputation-induced scrunching in this species (Table 3). However, all other scrunching parameters were consistent with the previous reference values, demonstrating the overall characteristics of scrunching. This decrease in speed may be due to negative health effects of the exposure, since we found that *D. japonica* but not *S. mediterranea* disintegrated within a day following the 5 min H_2O_2 exposure unless excessively washed post H_2O_2 exposure. Even after washing in three separate 50 mL baths of IO water, 1/12 *D. japonica* planarians died within 24 hours. Additional range-finding tests were unable to determine a concentration of H_2O_2 that induced scrunching without negative health effects in *D. japonica*. Lower concentrations (10 and 20 mM) only induced head wiggling within 5 min of exposure. It is currently unclear why *D. japonica* show such an increased sensitivity to H_2O_2 .




Fig 1. TRPA1 and TRPV agonists induce scrunching in planarians. (A) Single frames of gliding *S. mediterranea* (left) and *D. japonica* (right). (B-F) Representative length versus time plots for *S. mediterranea* (left) and *D. japonica*

(right) planarians during (B) gliding or (C-F) chemically induced scrunching, when exposed to (C) 50 μM AITC, (D) 40 mM H₂O₂, (E) 165 μM capsaicin, or (F) 100 μM anandamide. *D. japonica* planarians only scrunched temporarily in capsaicin, as shown in the representative plot in (E), where scrunching occurs from 2–9 seconds. Length was normalized by the average gliding length for all plots. (G-H) Example *S. mediterranea* (i) and *D. japonica* (ii) mucus trails stained with fluorescein-conjugated VVA lectin (see Methods) for worms exposed to (G) 50 μM AITC or (H) 165 μM capsaicin. Mucus trail images were black/white inverted. Scale bars: 1 mm.

<https://doi.org/10.1371/journal.pone.0226104.g001>

We have previously demonstrated that scrunching is induced by high heat and low pH [4], which are known activators of TRPV1 in other species [15–20]. Thus, we tested whether the classical TRPV1 agonist, capsaicin, and the endocannabinoid anandamide, known to directly activate TRPV1 in mammalian systems [35,38,52–54], were able to induce scrunching (Fig 1E and 1F). Although not all organisms are equally sensitive to these mammalian TRPV1 modulators, some invertebrates such as medicinal leech [16] and parasitic flatworms [28] have been shown to exhibit behavioral effects when exposed to capsaicin, suggesting that planarians may also be sensitive to TRPV1 modulators.

Indeed, both *S. mediterranea* and *D. japonica* scrunched with stereotypical parameters when exposed to 165 μM capsaicin (Fig 1E and Table 3). Additionally, in both species, scrunching in capsaicin was often accompanied by vigorous head shaking and jerking (S1 Movie). In contrast, exposure to 100 μM anandamide elicited scrunching in *D. japonica* but not in *S. mediterranea* (Fig 1F and Table 3). The slightly reduced maximum body elongation in *D. japonica* is likely because anandamide also induced other behaviors, such as increased head lifting or head wiggling, but all other parameters are consistent with scrunching. *S. mediterranea* worms displayed oscillatory locomotion (Fig 1F), but a quantitative analysis of the parameters shows that *S. mediterranea* performed peristalsis (Table 3), which we have previously demonstrated to be a distinct gait from scrunching [4]. Peristalsis is induced when cilia beating is disrupted, whereas scrunching is cilia-independent [2,4–7]. Using cilia imaging, we confirmed that cilia beating was disrupted in *S. mediterranea* but not *D. japonica* exposed to 100 μM anandamide (S3 Fig). A possible explanation for this finding is that in addition to being a low potency agonist of TRPV1, anandamide also activates cannabinoid receptor 1 (CB-1) [35,38]. Although the cannabinoid receptor(s) have not yet been directly identified in

Table 3. Quantification of scrunching parameters in response to TRPA1 and TRPV chemical agonists in *D. japonica* and *S. mediterranea*.

Species	Stimulus	Conc.	Frequency (cycles s ⁻¹)	Maximum elongation	Speed (body length s ⁻¹)	Fraction of time spent elongating	Gait	N
<i>D. japonica</i>	AITC	50 μM	0.72 ± 0.08	0.52 ± 0.04	0.37 ± 0.04	0.59 ± 0.03	S	9
<i>D. japonica</i>	H ₂ O ₂	40 mM	0.54 ± 0.09	0.44 ± 0.04	0.23 ± 0.05*	0.60 ± 0.02	S	8
<i>D. japonica</i>	Capsaicin	165 μM	0.80 ± 0.14	0.48 ± 0.07	0.38 ± 0.05	0.59 ± 0.04	S	8
<i>D. japonica</i>	Anandamide	100 μM	0.63 ± 0.08	0.43 ± 0.05*	0.27 ± 0.05	0.57 ± 0.03	S	8
<i>D. japonica</i>	Amputation ^a	–	0.70 ± 0.27	0.50 ± 0.08	0.34 ± 0.12	0.60 ± 0.12	S	15
<i>S. mediterranea</i>	AITC	50 μM	0.33 ± 0.02	0.43 ± 0.05	0.14 ± 0.02	0.58 ± 0.02	S	5
<i>S. mediterranea</i>	H ₂ O ₂	40 mM	0.37 ± 0.03	0.39 ± 0.05	0.14 ± 0.02	0.52 ± 0.02	S	10
<i>S. mediterranea</i>	Capsaicin	165 μM	0.44 ± 0.04	0.49 ± 0.06	0.21 ± 0.03	0.57 ± 0.03	S	8
<i>S. mediterranea</i>	Anandamide	100 μM	0.28 ± 0.03**	0.30 ± 0.05**	0.08 ± 0.01**	0.49 ± 0.03**	P	7
<i>S. mediterranea</i>	Amputation ^a	–	0.40 ± 0.09	0.44 ± 0.09	0.17 ± 0.09	0.62 ± 0.18	S	77
<i>S. mediterranea</i>	Peristalsis ^a	–	0.26 ± 0.07	0.23 ± 0.19	0.06 ± 0.04	0.50 ± 0.07	P	14

Values denote mean ± SD. S: scrunching; P: peristalsis

^aAmputation and peristalsis data are previously published values [4], provided for reference.

* p < 0.05 and

** p < 0.01 from student t-tests, conducted by comparing against the same parameter in the published amputation data for each species as the control group.

<https://doi.org/10.1371/journal.pone.0226104.t003>

planarians, pharmacological experiments with specific cannabinoid receptor agonists and antagonists in the planarian *Dugesia gonocephala* suggest the presence of functional cannabinoid receptor homologs in planarians [55,56]. Complicating matters, in other systems, cross-talk with the cannabinoid system can modulate the responsiveness of TRPV1 [38]. Furthermore, the efficiency of anandamide binding to TRPV1 appears to be species-specific [35]. Thus, these factors could interact to produce different manifestations of similar, yet distinct, oscillatory gaits in the two species, resulting in scrunching in *D. japonica*, but peristalsis in *S. mediterranea*. This dissimilarity in behavioral phenotypes, together with the sensitivity differences to H₂O₂ exposure, emphasize that care needs to be taken when attempting to extrapolate findings from pharmacological studies from one planarian species to another. Similar pharmacological differences have also been found in parasitic flatworms as AITC was shown to activate TRPA in *S. mansoni*, but not in the closely related *S. haematobium* [28].

Finally, we also visualized the mucus trails of worms of both species exposed to either AITC or capsaicin (Fig 1G and 1H) and saw the characteristic profiles of triangular anchor points that we have previously demonstrated to be associated with scrunching [4].

In summary, we found that archetypal agonists of TRPA1 and TRPV channels induce scrunching in planarians, supporting our hypothesis that induction of this gait is mediated by TRPA1 and TRPV activation.

Increasing concentrations of AITC and the TRPA1 antagonist HC-030031 enhance scrunching

It has been previously shown that increasing the concentration of AITC decreases the latency to initiate the nocifensive response in the medicinal leech [40]. We observed this same trend in both planarian species, as at higher concentrations of AITC more planarians scrunched earlier (Fig 2A), with more pronounced differences seen in *S. mediterranea*. However, in all concentrations of AITC tested, even when planarians did not scrunch initially, they still reacted to the AITC as evidenced by vigorous head turning (shown as percent worms reacting in Fig 2A), thus demonstrating that the initial sensation of AITC does not appear to be affected by concentration within the range tested here. Interestingly, the scrunching parameters were also dependent on AITC concentration, with increasing concentrations causing significantly increased maximum elongation and speed for both *D. japonica* and *S. mediterranea* when compared to 50 μ M AITC (S1 Table). Whereas for *D. japonica*, these values, as well as frequency in 100 μ M AITC, were also significantly different from the scrunching parameters in response to amputation, the parameters for *S. mediterranea* exposed to the different concentrations of AITC were not significantly different from amputation-induced scrunching parameters, indicating these differences were within the range observed in this species (S1 Table).

A striking behavioral difference was observed between *S. mediterranea* and *D. japonica* when exposed to AITC. In all tested AITC concentrations, the majority (at least ~80% in all tested concentrations) of scrunching *D. japonica* ceased scrunching by 90 seconds and began gliding, as seen by the decrease in both the percent worms scrunching and reacting in Fig 2A. This apparent desensitization was concentration-dependent; *D. japonica* planarians at higher AITC concentrations started and ceased scrunching earlier than those at lower AITC concentrations. *S. mediterranea* did not share this desensitization behavior and showed longer periods where all worms were scrunching (Fig 2A, compare 100 μ M AITC between the two species). Consistent with this observed desensitization to prolonged scrunching activation in *D. japonica*, the continuous application of high concentrations of AITC completely desensitizes currents in the dorsal root ganglion neurons of mice [57]. AITC is thought to activate TRPA1 through covalent modification of conserved cysteine residues [58], causing seemingly




Fig 2. HC-030031 decreases scrunching latency induced by AITC. (A, B) Behavior scoring plots for *S. mediterranea* (left) and *D. japonica* (right) showing the percentage of worms scrunching (black lines) or reacting (behaviors other than scrunching, see [Methods](#); red lines) every 15 s over 90 s when exposed to (A) AITC or (B) AITC + 100 μM HC-030031. Markers and shading represent the mean and standard deviation of 3 technical replicates, respectively.

<https://doi.org/10.1371/journal.pone.0226104.g002>

irreversible activation on the time scale of the mice electrophysiological experiments (15–60 mins) [57]. Since acute or repeated exposure to lower concentrations of AITC were not shown to induce desensitization of mouse TRPA1 [57], it is possible that differences in sensitivity between the two planarian species, which could be due to differences in chemical uptake and/or receptor sensitivities, could explain the lack of desensitization observed at the used concentrations in *S. mediterranea*.

HC-030031 is a specific TRPA1 antagonist that has been shown to block nocifensive responses to AITC in other systems, including rat and the medicinal leech [39,40]. Therefore, we tested whether HC-030031 could block or at least attenuate planarian scrunching. During initial tests with multiple concentrations of HC-030031, we unexpectedly found that 200 μM HC-030031 induced scrunching in 10/10 *D. japonica* planarians at some point within 2 minutes of exposure, whereas it did not have that effect on *S. mediterranea* (S4A Fig). At 100 μM HC-030031, neither planarian species scrunched, but *D. japonica* displayed a mild reaction including vigorous head turning, which was not observed in *S. mediterranea*. However, since scrunching was absent at this concentration in both species, 100 μM HC-030031 was used for further experiments.

When co-administered with 50, 75, or 100 μM AITC, 100 μM HC-030031 (Fig 2B) decreased the latency to induce scrunching in the majority of planarians at 50 and 75 μM

AITC compared to AITC treatment alone (Fig 2A) in both planarian species (S2 and S3 Movies). This decreased latency was confirmed by Fisher's exact tests showing a statistically significant increase in the number of worms scrunching at 15–30 and/or 31–45 seconds in 100 μ M HC-030031 compared to AITC alone (S2 Table). These data suggest a cooperative interaction between AITC and HC-030031, which mimicked the trend seen in increasing concentrations of AITC alone (Fig 2A). This effect was not as pronounced at 100 μ M, especially in *S. mediterranea* where no significant differences were found, suggesting that the maximal activity may have already been reached with 100 μ M AITC alone.

Together, these findings suggest that increasing concentrations of AITC or cooperative actions of AITC and HC-030031 enhance scrunching, further supporting the idea that TRPA1 is involved in mediating the scrunching response.

Genetic modulation of TRPA1 expression disrupts scrunching in response to AITC, H₂O₂, and amputation

Our chemical experiments suggest that TRPA1 activation in *S. mediterranea* and *D. japonica* induces scrunching. Sequence analysis of the putative proteins encoded by *SmTRPA1* and *DjTRPA1* further support that they would be sensitive to AITC as 2 of the 3 cysteines important for covalent modification by AITC [58] are conserved (S5 Fig). To confirm this, we knocked down *SmTRPA1* and *DjTRPA1* using RNAi and evaluated how this affected scrunching in response to TRPA1 modulators. Gene knockdown was confirmed by qRT-PCR showing 73.0% knockdown in *SmTRPA1* RNAi populations and 51.4% knockdown in *DjTRPA1* RNAi populations compared to expression in the species-specific control RNAi populations (S6A and S6B Fig).

When exposed to 100 μ M AITC for 90 s, none of the *SmTRPA1* RNAi or *DjTRPA1* RNAi planarians scrunched, while all control RNAi animals of each species scrunched under the same conditions (Fig 3A and 3B, S4 and S5 Movies). Similarly, scrunching in response to 200 μ M HC-030031 alone or to 100 μ M HC-030031 + 100 μ M AITC was completely lost in *DjTRPA1* RNAi animals (S4B and S4C Fig), demonstrating that planarian TRPA1 is essential for AITC-induced scrunching and that HC-030031 activates TRPA1. Our results suggest a cooperative, rather than antagonistic, action of HC-030031 on planarian TRPA1. Different organisms have been shown to have different sensitivities to the antagonistic effects of HC-030031. For example, divergence of a single amino acid (N855 in human TRPA1) in frog and zebrafish TRPA1 is responsible for their insensitivity to the inhibitor [59]. Although the mechanism of human TRPA1 inhibition could not be resolved structurally [60], it has been suggested that HC-030031 causes a conformational change in human TRPA1 which disrupts ligand binding [59]. Thus, it is possible that in planarians HC-030031 may cause a different conformational change in TRPA1 to instead potentiate AITC activation.

Scrunching was also completely lost in both *SmTRPA1* RNAi and *DjTRPA1* RNAi populations exposed to 40 mM H₂O₂ for either 270 s or 60 s, respectively, during which times all control RNAi planarians of both species scrunched (Fig 3C and 3D). These data are consistent with previous reports of H₂O₂ as a direct activator of *S. mediterranea* TRPA1 [10]. Together, these results show that planarian TRPA1 is essential to induce scrunching with either AITC or H₂O₂.

One of the most robust but unspecific inducers of scrunching is amputation [4]. Arenas et al. found that when doing tail snips on filter paper, *Smed-TRPA1* RNAi animals exhibited a decreased scrunching amplitude compared to control RNAi animals [10]. Because dry environments alone can induce scrunching (S7A Fig and [4]), we did not perform amputation experiments on filter paper, but in an aqueous environment instead. Under these experimental




Fig 3. TRPA1 is necessary for AITC- and H₂O₂-induced scrunching. (A-D) Representative length versus time plots for RNAi treated *S. mediterranea* (left) and *D. japonica* (right) exposed to (A-B) 100 μ M AITC or (C-D) 40 mM H₂O₂. Note that the scrunching frequencies in the *control* RNAi populations differ between the two inducers because a higher (100 μ M) concentration of AITC was used (compare values in Table 3 and S1 Table). Plots are representative of the total number of worms tested, as indicated in the respective panels for each condition.

<https://doi.org/10.1371/journal.pone.0226104.g003>

conditions, we found that knockdown of *SmTRPA1* caused reduced scrunching compared to *control* RNAi animals after amputation, evidenced by fewer total scrunches (S6B Fig). Thus, *SmTRPA1* appears to partially mediate scrunching in response to amputation. These results are consistent with the work of Arenas et al., who also found attenuated rather than completely abolished scrunching in *Smed-TRPA1* RNAi animals after amputation [10].

In *S. mediterranea*, it has been shown that amputation leads to a burst of H₂O₂ production at the wound site [61]. Thus, because H₂O₂ directly activates *SmTRPA1*, it has been suggested that mechanical injury (such as amputation) indirectly activates *SmTRPA1* through H₂O₂

production [10]. While our results confirm that H₂O₂ activation of TRPA1 induces scrunching, H₂O₂ activation of TRPA1 is likely not the only mechanism mediating amputation-induced scrunching since scrunching is not completely abolished in amputated *SmTRPA1* RNAi planarians. We were unable to perform these same experiments with the *D. japonica* RNAi populations as even in *control* RNAi animals, amputation only induces few scrunches robustly.

Together, our results confirm that TRPA1 in both *S. mediterranea* and *D. japonica* is necessary to induce scrunching in response to AITC and H₂O₂ and is partially involved in amputation-induced scrunching in *S. mediterranea*.

TRPV antagonist SB-366791 enhances scrunching

As we did for TRPA1, we similarly dissected the role of TRPV in scrunching. Because anandamide did not elicit scrunching in *S. mediterranea* and because two *D. japonica* TRPV genes have previously been characterized [31], we carried out all further experiments in *D. japonica* only. When comparing the behavioral responses to different concentrations of capsaicin, we found that at all tested concentrations, the planarians initially reacted by vigorously turning and shaking their heads and then transitioned to a scrunching phenotype over time. Because of this transitional nature of the behavior, it was often difficult to confidently distinguish non-scrunching reactions from scrunching by eye alone. Increasing the concentration of capsaicin decreased the latency time to switch to a scrunching reaction, similarly to AITC (Fig 4A). As with AITC (Fig 2A), after 45 s in 165 μM capsaicin, *D. japonica* scrunching behavior began to cease (Fig 4A). However, unlike in AITC where *D. japonica* began gliding again, *D. japonica* continued to react in capsaicin by maintaining a contracted body length and exhibiting minor oscillations (S6 Movie). Many nociceptors, including TRPA1 and TRPV, have been shown to become desensitized following prolonged activation. This desensitization is why certain TRP agonists, such as capsaicin, have been used therapeutically as analgesics [62]. In rat neuronal cell culture, it was shown that prolonged capsaicin exposure causes rat TRPV1 channels to be removed from the membrane through endocytosis and lysosomal degradation [63]. A similar desensitization to scrunching induction appears to be present in *D. japonica*, though the underlying mechanism remains to be determined.

We then tested SB-366791, a selective antagonist of human and rat TRPV1 [41,42]. Initial experiments using a range of different concentrations of SB-366791 showed that *D. japonica* began vigorous head turning at a concentration of 10 μM SB-366791 but did not scrunch. No abnormal behaviors were observed at 1 μM (S7 Movie). Similarly to AITC co-administration with HC-030031, co-administration of capsaicin with 10 μM SB-366791 decreased the scrunching latency compared to exposure to capsaicin alone (compare Fig 4A and 4B for the same capsaicin concentrations). Statistically significant differences were seen in the proportion of worms scrunching at 16–30 or 31–45 seconds in 33 and 82.5 μM capsaicin with or without 10 μM SB-366791 (S3 Table). However, unlike the trends seen with HC-030031, co-administration with SB-366791 decreased the number of worms which stopped scrunching over time, creating a more prolonged scrunching reaction compared to capsaicin alone. Together these results suggest that planarians are sensitive to a known agonist and antagonist of human and rat TRPV1, though the identity of this purported “capsaicin-receptor” remains to be verified. While SB-366791 enhances capsaicin-induced scrunching it does not have the same potentiation effects seen with HC-030031 and AITC.

DjTRPVab modulates scrunching behavior in response to anandamide

To assay their roles in mediating scrunching in response to the TRPV modulators, we knocked down both known *DjTRPV* genes (*DjTRPVa* and *DjTRPVb*) [31] in combination via RNAi




Fig 4. Scrunching in capsaicin is enhanced by SB-366791. (A-B) Behavior scoring plots for *D. japonica* showing the percentage of worms scrunching (black lines) or reacting (non-scrunching behaviors, red lines) every 15 s over 90 s when exposed to (A) capsaicin or (B) capsaicin + 10 μ M SB-366791. Markers and shading represent the mean and standard deviation of 3 technical replicates, respectively.

<https://doi.org/10.1371/journal.pone.0226104.g004>

(referred to as *DjTRPVab* RNAi). Gene knockdown was confirmed by qRT-PCR showing a 41.2% decrease in *DjTRPVa* and 83.3% decrease in *DjTRPVb* in the *DjTRPVab* RNAi population compared to expression in *control* RNAi planarians (S6C and S6D Fig). Because TRPA1 has been found to modulate sensitivity to capsaicin in the parasitic flatworm *S. mansoni*, which does not have any TRPV homologs [28,29,43], we also evaluated the reactions of *DjTRPAa* RNAi worms to the mammalian TRPV agonists. Neither *DjTRPVab* nor *DjTRPAa* RNAi significantly attenuated scrunching in response to 165 μ M capsaicin (Fig 5A and S6 Movie). Although there was a slight decrease in the percentage of worms scrunching in the *DjTRPVab* and *DjTRPAa* RNAi populations compared to *control* RNAi, all *DjTRPVab* and *DjTRPAa* RNAi planarians either reacted or scrunched when exposed to 165 μ M capsaicin. Sequence comparisons between TRPV1s which are sensitive (human, rat) or insensitive (chicken, rabbit) to capsaicin have revealed 3 residues (Y511, S512, and T550 using human TRPV1 numbering) which are important for capsaicin sensitivity [22,23]. None of these

residues are conserved in DjTRPVa, while only S512 is conserved in DjTRPVb (S8 Fig), consistent with our RNAi results showing DjTRPVa and b are not required for capsaicin-induced scrunching. Similarly, for *S. mediterranea*, pipetting 165 μ M capsaicin onto *control* or *SmTRPA1* RNAi populations induced scrunching in all tested animals (N = 10). A TRPV homolog has not yet been identified in this species. Thus, none of these channels are solely responsible for capsaicin-induced scrunching.

Although mammalian TRPV1 was originally identified as the “capsaicin receptor”, capsaicin-sensing ability and the responsible receptor varies dramatically across invertebrates. Several invertebrate species, including fruit flies and nematodes, are insensitive to capsaicin [25,26]. In *Caenorhabditis elegans*, capsaicin potentiates the thermal avoidance response, but this effect is not dependent on OSM-9, the purported *C. elegans* TRPV1 homolog, suggesting another unknown receptor is involved [25]. A similar situation appears to be present in *D. japonica*, where scrunching is not dependent on two previously identified TRPV homologs, DjTRPVa and DjTRPVb (Fig 5A). Our results also show that, unlike in *S. mansoni*, TRPA1 in both planarian species is not responsible for capsaicin sensing, suggesting evolutionary divergence. While our data show that DjTRPVa/b and DjTRPA are not essential for capsaicin sensing in planarians, it is possible that other planarian TRP channels may be involved. Transcriptomic analysis suggests *D. japonica* may have at least 25 TRPs [30], of which only 7 have been characterized so far [31]. What receptor is responsible for capsaicin sensing, whether it be another unidentified TRPV, a different TRP channel, or some unrelated protein, in freshwater planarians remains to be determined.

Next, we tested whether scrunching induced by anandamide could be affected by knock-down of either *DjTRPVab* or *DjTRPAa*. Even in *control* RNAi planarians, scrunching in response to 125 μ M anandamide was complicated as non-scrunching behaviors, such as vigorous head turning or other body shape contortions (Fig 5B, S8 Movie), were also prevalent. Thus, we focused our comparisons on the number of worms reacting by either clearly scrunching or exhibiting these mixed behavioral phenotypes. Most (75%) *control* RNAi planarians began reacting within 30 seconds, whereas in *DjTRPVab* RNAi planarians, anandamide-induced scrunching was attenuated, evidenced by an increase in the latency time to induce scrunching (Fig 5B, S8 Movie). This increased latency was confirmed by a Fisher’s exact showing a significant decrease in the number of *DjTRPVab* RNAi planarians reacting during 16-30s compared to *control* RNAi planarians at the same time (p-value: 0.02). No significant differences were found in the behaviors of *DjTRPAa* RNAi planarians. These data suggest that DjTRPVa/b partially mediate anandamide-sensing in *D. japonica*, though other receptor(s) are likely also involved.

Anandamide and other cannabinoids have complicated relationships with TRP channels. Both endogenous and synthetic cannabinoids act through the canonical cannabinoid receptors, CB-1 and CB-2, but some have been found to activate TRPV and TRPA1 channels as well [54,64]. Additionally, because of the extensive crosstalk between the endocannabinoid system and TRPV1, leading to sensitization of TRPV1 to other endogenous ligands, it has been suggested that even when anandamide treatment mimics the physiological outcomes of TRPV agonists, the effects are not necessarily due to direct activation of TRPV1 [38]. Thus, it is unclear from our RNAi results whether anandamide’s role in scrunching is due to direct activation of DjTRPVab or indirectly through its role as an endocannabinoid.

Finally, we assayed the scrunching response of all RNAi populations (*SmTRPA1*, *DjTRPA1*, and *DjTRPVab*) to noxious heat and low pH exposure, which are known to affect TRPV in other species [15,16,18–20]. We observed scrunching in all populations (S7 Fig), indicating that none of these 3 genes are involved in the scrunching response to these stimuli. Strengthening this conclusion is the observation that, in addition to pipetting experiments, when




Fig 5. DjTRPV mediates the behavioral response to anandamide. (A-B) Behavior of control RNAi, *DjTRPAa* RNAi, and *DjTRPVab* RNAi planarians in (A) 165 μ M capsaicin and (B) 125 μ M anandamide. Markers and shading represent the mean and standard deviation of 3 technical replicates, respectively.

<https://doi.org/10.1371/journal.pone.0226104.g005>

scrunching was induced by heating the aquatic environment, scrunching was observed in all RNAi populations with no statistically significant differences, as determined by a Fisher's exact test ($p > 0.05$). Scrunching was found in 24/34 *DjTRPAa* and 20/28 *DjTRPVab* RNAi planarians, similar to control RNAi worms (25/36). Consistent results were also found for *S. mediterranea* as similar proportions of animals scrunched in the control (9/22) or *SmTRPA1* (7/21) RNAi populations. The finding that in this assay *S. mediterranea* planarians across RNAi populations scrunched much less than *D. japonica* planarians may be a consequence of the experimental setup being optimized for *D. japonica* [8]. While previous reports have shown that *SmTRPA1* is involved in mediating heat avoidance behaviors via direct activation by H_2O_2 [10], our data suggest that other channels may be involved in triggering scrunching in response to high temperatures.

Taken together, these data demonstrate that TRPA1 is required for scrunching in response to AITC and H_2O_2 in both planarian species, whereas DjTRPVab partially regulates

anandamide-induced scrunching in *D. japonica*. It remains to be determined which other receptor(s) are involved in regulating the anandamide response and are responsible for the other scrunching inducers, including capsaicin, low pH and noxious heat. Importantly, because scrunching in response to capsaicin, heat or pH were not affected by knockdown of either *DjTRPA* or *DjTRPVab*, these genes are likely not responsible for the general scrunching response but rather mediate sensing of specific stimuli.

Conclusions

Combining the results presented here with our previous studies of scrunching allowed us to partially decipher the molecular mechanisms responsible for sensing noxious stimuli in planarians. In this and our previous work, we found that planarian TRPA1 and TRPV channels, as well as the planarian Big Potassium ion channel SLO-1 which mediates ethanol-induced scrunching [65], are involved in inducing scrunching in response to specific stimuli. Using RNAi, we found that some inducers are specific to one of these pathways, such as AITC and H₂O₂ to TRPA1, while others, such as anandamide and amputation, rely on potential epistatic or overlapping functions of other unidentified channels. Lastly, the responsible receptors mediating the response to some scrunching inducers, including capsaicin, low pH, and noxious heat remain elusive and it remains to be determined which receptor(s) are responsible for sensing these stimuli. Based on our results, it is likely that multiple receptors could have epistatic or overlapping functions. Further complicating matters, we found species differences with several of the scrunching inducers. For example, while anandamide induced scrunching which was partially mediated by DjTRPVa/b in *D. japonica*, it induced peristalsis in *S. mediterranea*. These observations and the differences in sensitivity to H₂O₂ and HC-030031 between the two species demonstrate that although these two species are closely related, caution must be used when extrapolating the pharmacological effects of one species to another.

It is striking that despite the existence of multiple induction routes, the dynamic features of scrunching are independent of the inducer, and hence of the sensing pathway. This suggests some form of signal integration occurring downstream of these receptors, as illustrated graphically in Fig 6, which represents our current understanding of the molecular mechanisms of scrunching induction.

Signal integration could occur at the neuronal level, raising the question of which parts of the planarian nervous system are involved. Our previous results [4] have shown that tail pieces which lack a brain are still capable of scrunching in response to some stimuli, albeit much more rarely. This would suggest that the ventral nerve cords are sufficient for scrunching execution but that the brain plays an important role in achieving consistent induction.

Our pharmacological studies revealed that several agonist-antagonist pairs (AITC/HC-030031 and capsaicin/SB-366791) either both triggered scrunching and/or were unable to pharmacologically rescue the scrunching phenotype. These results were surprising given that in other systems, including invertebrates such as the medicinal leech and schistosomes, the antagonists work as expected to inhibit the action of the agonists [16,40,43]. In contrast, in the two planarian species studied here, we found that both a TRPA1 agonist and antagonist induced scrunching, with potentiating effects when co-exposed in both planarian species. Similarly, although the mammalian TRPV antagonist SB-366791 did not induce scrunching alone, it also potentiated capsaicin-induced scrunching in *D. japonica*. Only RNAi against the target genes allowed us to suppress scrunching in response to specific chemical inducers. One possible explanation for these findings is that the planarian sensory system is highly sensitive to any deviation from normal and that scrunching is a default downstream response to system perturbations or stress. However, the observed species differences demonstrate that scrunching is




Fig 6. Overview of our current understanding of mediators of scrunching induction. Solid lines indicate that direct connections have been experimentally shown. Dotted lines indicate inducers which were only found to induce scrunching in one of the two species. Dashed lines are hypothesized connections.

<https://doi.org/10.1371/journal.pone.0226104.g006>

not always triggered, in agreement with our previous findings that ethanol, but not methanol, trigger scrunching [65]. Furthermore, knockdown of a single receptor, such as SLO-1 [65] or TRPA1, abolishes the scrunching response to specific inducers (ethanol and AITC/H₂O₂, respectively), without perturbing scrunching in response to other inducers. Together, these observations argue that scrunching is a specific response, whose regulation, despite the progress made in this work, remains poorly understood.

Understanding the molecular mechanisms controlling the scrunching gait, from initiation to execution will require systematic studies of these different aspects using chemical and/or molecular approaches as presented here. The observed complexity and myriad of pathways involved in scrunching initiation reported here may explain why scrunching is a sensitive read-out of neurotoxicity [8] and gaining a deeper understanding of its regulation will allow for more mechanistic studies of potential toxicants in the future using the planarian system. Moreover, by understanding the extent that aspects of nociception are conserved (or not) across species will provide better informative context to understand species-specific sensitivity differences and provide insight into the important mechanisms regulating noxious stimuli and pain sensation.

Supporting information

S1 Table. Maximum elongation and speed of scrunching are dependent on AITC concentration. Scrunching parameters for *D. japonica* and *S. mediterranea* exposed to 50, 75, or 100 μ M AITC, denoted as mean \pm standard deviation. For each concentration, planarians were observed to scrunch with the parameters listed within the first minute in the bath. * denotes $p < 0.05$ and ** denotes $p < 0.01$ significance level compared to 50 μ M AITC given by a two-tailed t-test. ^ denotes $p < 0.05$ and ^^ denotes $p < 0.01$ significance level compared to amputation given by a two-tailed t-test. ^aAmputation data are previously published values [4], provided for reference. (PDF)

S2 Table. P-values comparing scrunching rate in AITC treatment with or without HC-030031. A Fisher's exact test was used to compare the number of worms scrunching vs not

scrunching (no reaction or non-scrunching reaction) at each listed time point in different concentrations of AITC alone or co-exposed with 100 μ M HC-030031. * denotes $p < 0.05$ and ** denotes $p < 0.01$ significance level.

(PDF)

S3 Table. P-values comparing scrunching rate in capsaicin with or without SB-366791. A Fisher's exact test was used to compare the number of worms scrunching vs not scrunching (no reaction or non-scrunching reaction) at each listed time point in different concentrations of capsaicin alone or co-exposed with 10 μ M SB-366791. * denotes $p < 0.05$ and ** denotes $p < 0.01$ significance level.

(PDF)

S1 Fig. 1% DMSO does not induce scrunching in either *S. mediterranea* or *D. japonica*.

Representative length versus time plot for wildtype (A) *S. mediterranea* or (B) *D. japonica* planarians in 1% dimethyl sulfoxide (DMSO) (N = 10). Planarians were exposed to 1% DMSO by directly pipetting 100 μ L.

(TIF)

S2 Fig. *S. mediterranea* scrunch in response to low pH exposure. Representative length versus time plot for wildtype *S. mediterranea* planarians in Instant Ocean water at neutral pH (red, N = 5) and pH 2.7 (black, N = 10). Scrunching was induced by directly pipetting 100 μ L onto planarians.

(TIF)

S3 Fig. Anandamide impairs cilia beating in *S. mediterranea* but not *D. japonica*. Representative (N = 3/3) 1 s kymograph of cilia beating for *S. mediterranea* (left) and *D. japonica* (right). (A) Controls in planarian water and (B) when exposed to 100 μ M anandamide for 5 minutes. Notice that cilia beating is almost completely lost in *S. mediterranea* while cilia beat normally in *D. japonica*. Scale bar shows 0.1 s horizontally and 1 μ m vertically.

(TIF)

S4 Fig. 200 μ M HC-030031 induction of scrunching in *D. japonica* is dependent on TRPA1. (A) Representative length versus time plot for wildtype (i) *S. mediterranea* and (ii) *D. japonica* planarians in 100 μ M (black) and 200 μ M (red) HC-030031. Plots are representative of N = 10. (B) Representative length versus time plot for *D. japonica control* RNAi (N = 13) and *DjTRPAa* RNAi (N = 11) planarians in 200 μ M HC-030031. (C) Representative length versus time plot for *D. japonica control* RNAi (N = 8) and *DjTRPAa* RNAi (N = 9) planarians in 100 μ M HC-030031 + 100 μ M AITC.

(TIF)

S5 Fig. Planarian TRPA1s have conserved AITC-responsive cysteines. Alignment of human and mouse TRPA1 with predicted protein sequences for planarian TRPA1. Darkness of purple color-coding represents levels of shared identity. Cysteines shown to be involved in AITC sensitivity are shown in orange. Two of the three cysteines are conserved in both SmTRPA1 and DjTRPA1.

(TIF)

S6 Fig. Confirmation of RNAi knockdown by qRT-PCR. (A-D) Relative expression of (A) *SmTRPA1*, (B) *DjTRPAa*, (C) *DjTRPVa* and (D) *DjTRPVb* in the respective RNAi populations compared to the *control* RNAi population in that species. Data are shown as the mean of two biological replicates (each including 3 technical replicates). Error bars represent the standard error.

(TIF)

S7 Fig. SmTRPA1 mediates scrunching in response to amputation. (A) Representative length versus time plot for wildtype *S. mediterranea* planarians in an aqueous (red, N = 5) and dry (black, N = 10) environment, created by placing the planarians on wet filter paper as in (10). (B) Distribution showing median and quartiles of the number of scrunches directly following amputation in *control* RNAi (N = 21) and *SmTRPA1* RNAi (N = 24) planarians. * denotes $p < 0.01$ significance from *control* RNAi given by a two-tailed t-test. (TIF)

S8 Fig. Residues important for capsaicin binding are not conserved in DjTRPVa/b. Fragment of a sequence alignment of capsaicin-sensitive (human, rat) and capsaicin-insensitive (rabbit) TRPVs with predicted protein sequences for DjTRPVa and DjTRPVb. Darkness of purple color-coding represents levels of shared identity. Residues important for capsaicin-sensitivity are in orange. The only residue found in the planarian TRPVs is S512 in DjTRPVb. (TIF)

S9 Fig. Heat and low pH sensing are not significantly impaired in DjTRPAa or DjTRPVab RNAi planarians. (A-B) Representative oscillation plots for *control*, *DjTRPAa*, and *DjTRPVab* RNAi planarians exposed to (A) pH 2.7 and (B) 65°C IO water via pipette. No significant differences in scrunching induction are seen in any of the conditions. N = 10 for all conditions. (TIF)

S1 Movie. *D. japonica* and *S. mediterranea* planarians in 165 µM capsaicin. Comparison of *D. japonica* and *S. mediterranea* behaviors in 165 µM capsaicin. Planarians exhibit vigorous head shaking and turning as well as scrunching. Movie is recorded and played at 10 fps. (AVI)

S2 Movie. *D. japonica* exposed to 50 µM AITC alone and with 100 µM HC-030031. First 50 s of exposure of *D. japonica* to 50 µM AITC alone or co-administered with 100 µM HC-030031. Movie is recorded and played at 10 fps. Scale bar: 1 cm. (AVI)

S3 Movie. *S. mediterranea* exposed to 50 µM AITC alone and with 100 µM HC-030031. First 50 s of exposure of *S. mediterranea* planarians to 50 µM AITC alone or co-administered with 100 µM HC-030031. Movie is recorded and played at 10 fps. Scale bar: 1 cm. (AVI)

S4 Movie. *D. japonica* control and *DjTRPAa* RNAi planarians in 100 µM AITC. Behavior of *D. japonica* *control* RNAi and *DjTRPAa* RNAi in 100 µM AITC during the first 30 seconds of exposure. *Control* RNAi planarians scrunch whereas *DjTRPAa* RNAi planarians exhibit rapid head turning but lack a scrunching response. Movie is recorded and played at 10 fps. Scale bar: 1 cm. (AVI)

S5 Movie. *S. mediterranea* control and *SmTRPA1* RNAi *S. mediterranea* planarians in 100 µM AITC. Behavior of *S. mediterranea* *control* RNAi and *SmTRPA1* RNAi planarians in 100 µM AITC during the first 30 s of exposure. *Control* RNAi planarians scrunch whereas *SmTRPA1* RNAi planarians glide. Movie is recorded and played at 10 fps. Scale bar: 1 cm. (AVI)

S6 Movie. *D. japonica* control, *DjTRPAa*, and *DjTRPVab* RNAi planarians in 165 µM capsaicin. First 60 seconds of *D. japonica* RNAi populations when exposed to 165 µM capsaicin. Movie is recorded and played at 10 fps. Scale bar: 1 cm. (AVI)

S7 Movie. *D. japonica* behavior in 1 μ M and 10 μ M SB-366791. Movie showing *D. japonica* behavior in SB-366791. *D. japonica* planarians exhibit no abnormal behaviors in 1 μ M SB-366791 but display vigorous head turning at 10 μ M SB-366791. Movie is recorded and played at 10 fps. Scale bar: 1 cm.

(AVI)

S8 Movie. *D. japonica* control, *DjTRPAa*, and *DjTRPVab* RNAi planarians in 125 μ M anandamide. First 60 seconds of *D. japonica* RNAi populations when exposed to 125 μ M anandamide. *Control* RNAi *D. japonica* planarians exhibit either scrunching or a non-scrunching behavior. Conversely, both *DjTRPAa* and *DjTRPVab* RNAi planarians show a significant decrease in both reactions. Movie is recorded and played at 10 fps. Scale bar: 1 cm.

(AVI)

Acknowledgments

The authors thank Cambria Neal, Arya Dadhania, Kelson Kaj, Angel Leu, Addam Debebe, Eileen Tsai, and Yingtian He for help with planarian care and experiments, Dr. Marco Gallio for the TRPA1 plasmid and discussions, Oscar Arenas for discussions of the AITC experiments, and Dr. William Kristan for discussions and comments on the manuscript.

Author Contributions

Conceptualization: Danielle Ireland, Olivier Cochet-Escartin, Eva-Maria S. Collins.

Formal analysis: Ziad Sabry, Alicia Ho, Christina Rabeler.

Funding acquisition: Eva-Maria S. Collins.

Investigation: Ziad Sabry, Alicia Ho, Christina Rabeler, Eva-Maria S. Collins.

Methodology: Ziad Sabry, Alicia Ho, Danielle Ireland, Olivier Cochet-Escartin, Eva-Maria S. Collins.

Project administration: Eva-Maria S. Collins.

Resources: Eva-Maria S. Collins.

Supervision: Danielle Ireland, Olivier Cochet-Escartin, Eva-Maria S. Collins.

Validation: Ziad Sabry.

Writing – original draft: Ziad Sabry, Alicia Ho, Danielle Ireland, Eva-Maria S. Collins.

Writing – review & editing: Ziad Sabry, Alicia Ho, Danielle Ireland, Christina Rabeler, Olivier Cochet-Escartin, Eva-Maria S. Collins.

References

1. Martin GG. A new function of rhabdites: Mucus production for ciliary gliding. *Zoomorphologie*. 1978; 91: 235–248. <https://doi.org/10.1007/BF00999813>
2. Rompolas P, Patel-King RS, King SM. An outer arm dynein conformational switch is required for metachronal synchrony of motile cilia in planaria. *Mol Biol Cell*. 2010; 21: 3617–3759. <https://doi.org/10.1091/mbc.E10-03-0246>
3. Elgeti J, Gompper G. Emergence of metachronal waves in cilia arrays. *Proc Natl Acad Sci*. 2013; 110: 4470–4475. <https://doi.org/10.1073/pnas.1218869110> PMID: 23487771
4. Cochet-Escartin O, Mickolajczk KJ, Collins E-MS. Scrunching: a novel escape gait in planarians. *Phys Biol*. 2015; 12: 055001. <https://doi.org/10.1088/1478-3975/12/5/055001>

5. Rompolas P, Azimzadeh J, Marshall WF, King SM. Analysis of ciliary assembly and function in planaria. In: Marshall WF, editor. *Methods in Enzymology*. 2013. pp. 245–264. <https://doi.org/10.1016/B978-0-12-397944-5.00012-2>
6. Azimzadeh J, Wong ML, Downhour DM, Alvarado AS, Marshall WF. Centrosome loss in the evolution of planarians. *Science* (80-). 2012; 335: 461–463. <https://doi.org/10.1126/science.1214457> PMID: [22223737](https://pubmed.ncbi.nlm.nih.gov/22223737/)
7. Patel-King RS, Gilberti RM, Hom EFY, King SM. WD60/FAP163 is a dynein intermediate chain required for retrograde intraflagellar transport in cilia. *Mol Biol Cell*. 2013; 24: 2593–2763. <https://doi.org/10.1091/mbc.E12-08-0617>
8. Zhang S, Hagstrom D, Hayes P, Graham A, Collins E-MS. Multi-behavioral endpoint testing of an 87-chemical compound library in freshwater planarians. *Toxicol Sci*. 2019; 167: 26–44. <https://doi.org/10.1093/toxsci/kyf145> PMID: [29893936](https://pubmed.ncbi.nlm.nih.gov/29893936/)
9. Zhang S, Ireland D, Sipes NS, Behl M, Collins E-MS. Screening for neurotoxic potential of 15 flame retardants using freshwater planarians. *Neurotoxicol Teratol*. 2019; 73: 54–66. <https://doi.org/10.1016/j.ntt.2019.03.003> PMID: [30943442](https://pubmed.ncbi.nlm.nih.gov/30943442/)
10. Arenas OM, Zaharieva EE, Para A, Vásquez-Doorman C, Petersen CP, Gallio M. Activation of planarian TRPA1 by reactive oxygen species reveals a conserved mechanism for animal nociception. *Nat Neurosci*. 2017; 20: 1686–1693. <https://doi.org/10.1038/s41593-017-0005-0> PMID: [29184198](https://pubmed.ncbi.nlm.nih.gov/29184198/)
11. Venkatachalam K, Montell C. TRP channels. *Annu Rev Biochem*. 2007; 76: 387–417. <https://doi.org/10.1146/annurev.biochem.75.103004.142819> PMID: [17579562](https://pubmed.ncbi.nlm.nih.gov/17579562/)
12. Birkholz TR, Beane WS. The planarian TRPA1 homolog mediates extraocular behavioral responses to near-ultraviolet light. *J Exp Biol*. 2017; 220: 2616–2625. <https://doi.org/10.1242/jeb.152298> PMID: [28495872](https://pubmed.ncbi.nlm.nih.gov/28495872/)
13. Nilius B, Owsianik G. The transient receptor potential family of ion channels. *Genome Biol*. 2011; 12. <https://doi.org/10.1186/gb-2011-12-3-218> PMID: [21401968](https://pubmed.ncbi.nlm.nih.gov/21401968/)
14. Li H. TRP Channel Classification. In: Wang Y, editor. *Transient Receptor Potential Canonical Channels and Brain Diseases*. Springer, Dordrecht; 2017. pp. 1–8. https://doi.org/10.1007/978-94-024-1088-4_1
15. Dhaka A, Uzzell V, Dubin AE, Mathur J, Petrus M, Bandell M, et al. TRPV1 Is activated by both acidic and basic pH. *J Neurosci*. 2009; 29: 153–158. <https://doi.org/10.1523/JNEUROSCI.4901-08.2009> PMID: [19129393](https://pubmed.ncbi.nlm.nih.gov/19129393/)
16. Summers T, Holec S, Burrell BD. Physiological and behavioral evidence of a capsaicin-sensitive TRPV-like channel in the medicinal leech. *J Exp Biol*. 2014; 217: 4167–73. <https://doi.org/10.1242/jeb.110049> PMID: [25324339](https://pubmed.ncbi.nlm.nih.gov/25324339/)
17. Caterina MJ, Rosen TA, Tominaga M, Brake AJ, Julius D. A capsaicin-receptor homologue with a high threshold for noxious heat. *Nature*. 1999; 398: 436–441. <https://doi.org/10.1038/18906> PMID: [10201375](https://pubmed.ncbi.nlm.nih.gov/10201375/)
18. King RS, Newmark PA. In situ hybridization protocol for enhanced detection of gene expression in the planarian *Schmidtea mediterranea*. *BMC Dev Biol*. 2013; 13: 8. <https://doi.org/10.1186/1471-213X-13-8> PMID: [23497040](https://pubmed.ncbi.nlm.nih.gov/23497040/)
19. Aneiros E, Cao L, Papakosta M, Stevens EB, Phillips S, Grimm C. The biophysical and molecular basis of TRPV1 proton gating. *EMBO J*. 2011; 30: 994–1002. <https://doi.org/10.1038/emboj.2011.19> PMID: [21285946](https://pubmed.ncbi.nlm.nih.gov/21285946/)
20. Gunthorpe MJ, Benham CD, Randall A, Davis JB. The diversity in the vanilloid (TRPV) receptor family of ion channels. *Trends in Pharmacological Sciences*. 2002. pp. 183–191. [https://doi.org/10.1016/S0165-6147\(02\)01999-5](https://doi.org/10.1016/S0165-6147(02)01999-5) PMID: [11931994](https://pubmed.ncbi.nlm.nih.gov/11931994/)
21. Zheng J. Molecular mechanism of TRP channels. *Compr Physiol*. 2013; 3: 221–42. <https://doi.org/10.1002/cphy.c120001> PMID: [23720286](https://pubmed.ncbi.nlm.nih.gov/23720286/)
22. Gavva NR, Klionsky L, Qu Y, Shi L, Tamir R, Edenson S, et al. Molecular determinants of vanilloid sensitivity in TRPV1. *J Biol Chem*. 2004; 279: 20283–20295. <https://doi.org/10.1074/jbc.M312577200> PMID: [14996838](https://pubmed.ncbi.nlm.nih.gov/14996838/)
23. Jordt S-E, Julius D. Molecular basis for species-specific sensitivity to “hot” chili peppers. *Cell*. 2002; 108: 421–430. [https://doi.org/10.1016/S0092-8674\(02\)00637-2](https://doi.org/10.1016/S0092-8674(02)00637-2) PMID: [11853675](https://pubmed.ncbi.nlm.nih.gov/11853675/)
24. Tobin DM, Madsen DM, Kahn-Kirby A, Peckol EL, Moulder G, Barstead R, et al. Combinatorial expression of TRPV channel proteins defines their sensory functions and subcellular localization in *C. elegans* neurons. *Neuron*. 2002; 35: 307–18. [https://doi.org/10.1016/S0896-6273\(02\)00757-2](https://doi.org/10.1016/S0896-6273(02)00757-2) PMID: [12160748](https://pubmed.ncbi.nlm.nih.gov/12160748/)
25. Wittenburg N, Baumeister R. Thermal avoidance in *Caenorhabditis elegans*: an approach to the study of nociception. *Proc Natl Acad Sci U S A*. 1999; 96: 10477–82. <https://doi.org/10.1073/pnas.96.18.10477> PMID: [10468634](https://pubmed.ncbi.nlm.nih.gov/10468634/)

26. Vriens J, Owsianik G, Voets T, Droogmans G, Nilius B. Invertebrate TRP proteins as functional models for mammalian channels. *Pflügers Arch—Eur J Physiol.* 2004; 449: 213–226. <https://doi.org/10.1007/s00424-004-1314-1> PMID: 15480752
27. Holzer P. Capsaicin: cellular targets, mechanisms of action, and selectivity for thin sensory neurons. *Pharmacol Rev.* 1991; 43.
28. Bais S, Berry CT, Liu X, Ruthel G, Freedman BD, Greenberg RM. Atypical pharmacology of schistosome TRPA1-like ion channels. *PLoS Negl Trop Dis.* 2018; 12: e0006495. <https://doi.org/10.1371/journal.pntd.0006495> PMID: 29746471
29. Bais S, Greenberg RM. TRP channels in schistosomes. *Int J Parasitol Drugs Drug Resist.* 2016; 6: 335–342. <https://doi.org/10.1016/j.ijpddr.2016.07.002> PMID: 27496302
30. Chan JD, Zhang D, Liu X, Zarowiecki M, Berriman M, Marchant JS. Utilizing the planarian voltage-gated ion channel transcriptome to resolve a role for a Ca²⁺ channel in neuromuscular function and regeneration. *Biochim Biophys Acta—Mol Cell Res.* 2017; 1864: 1036–1045. <https://doi.org/10.1016/j.bbamcr.2016.10.010> PMID: 27771293
31. Inoue T, Yamashita T, Agata K. Thermosensory Signaling by TRPM Is Processed by Brain Serotonergic Neurons to Produce Planarian Thermotaxis. *J Neurosci.* 2014; 34: 15701–15714. <https://doi.org/10.1523/JNEUROSCI.5379-13.2014> PMID: 25411498
32. Andrade EL, Luiz AP, Ferreira J, Calixto JB. Pronociceptive response elicited by TRPA1 receptor activation in mice. *Neuroscience.* 2008; 152: 511–520. <https://doi.org/10.1016/j.neuroscience.2007.12.039> PMID: 18272293
33. Calixto JB, Kassuya CAL, André E, Ferreira J. Contribution of natural products to the discovery of the transient receptor potential (TRP) channels family and their functions. *Pharmacol Ther.* 2005; 106: 179–208. <https://doi.org/10.1016/j.pharmthera.2004.11.008> PMID: 15866319
34. Yang F, Xiao X, Cheng W, Yang W, Yu P, Song Z, et al. Structural mechanism underlying capsaicin binding and activation of the TRPV1 ion channel. *Nat Chem Biol.* 2015; 11: 518–524. <https://doi.org/10.1038/nchembio.1835> PMID: 26053297
35. Ross RA. Anandamide and vanilloid TRPV1 receptors. *Br J Pharmacol.* 2003; 140: 790–801. <https://doi.org/10.1038/sj.bjp.0705467> PMID: 14517174
36. Cebria F. Planarian homologs of netrin and netrin receptor are required for proper regeneration of the central nervous system and the maintenance of nervous system architecture. *Development.* 2005; 132: 3691–3703. <https://doi.org/10.1242/dev.01941> PMID: 16033796
37. McNamara CR, Mandel-Brehm J, Bautista DM, Siemens J, Deranian KL, Zhao M, et al. TRPA1 mediates formalin-induced pain. *Proc Natl Acad Sci.* 2007; 104: 13525–13530. <https://doi.org/10.1073/pnas.0705924104> PMID: 17686976
38. Tóth A, Blumberg PM, Boczán J. Anandamide and the Vanilloid Receptor (TRPV1). *Vitamins and Hormones.* Elsevier Inc.; 2009. pp. 389–419. [https://doi.org/10.1016/S0083-6729\(09\)81015-7](https://doi.org/10.1016/S0083-6729(09)81015-7)
39. Eid SR, Crown ED, Moore EL, Liang HA, Choong K-C, Dima S, et al. HC-030031, a TRPA1 selective antagonist, attenuates inflammatory- and neuropathy-induced mechanical hypersensitivity. *Mol Pain.* 2008; 4. <https://doi.org/10.1186/1744-8069-4-48> PMID: 18954467
40. Summers T, Wang Y, Hanten B, Burrell BD. Physiological, pharmacological and behavioral evidence for a TRPA1 channel that can elicit defensive responses in the medicinal leech. *J Exp Biol.* 2015; 218: 3023–3031. <https://doi.org/10.1242/jeb.120600> PMID: 26254323
41. Varga A, Németh J, Szabó Á, McDougall JJ, Zhang C, Elekes K, et al. Effects of the novel TRPV1 receptor antagonist SB366791 in vitro and in vivo in the rat. *Neurosci Lett.* 2005; 385: 137–142. <https://doi.org/10.1016/j.neulet.2005.05.015> PMID: 15950380
42. Gunthorpe MJ, Rami HK, Jerman JC, Smart D, Gill CH, Soffin EM, et al. Identification and characterisation of SB-366791, a potent and selective vanilloid receptor (VR1/TRPV1) antagonist. *Neuropharmacology.* 2004; 46: 133–149. [https://doi.org/10.1016/s0028-3908\(03\)00305-8](https://doi.org/10.1016/s0028-3908(03)00305-8) PMID: 14654105
43. Bais S, Churgin MA, Fang-Yen C, Greenberg RM. Evidence for novel pharmacological sensitivities of Transient Receptor Potential (TRP) channels in *Schistosoma mansoni*. Keiser J, editor. *PLoS Negl Trop Dis.* 2015; 9: e0004295. <https://doi.org/10.1371/journal.pntd.0004295> PMID: 26655809
44. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. *Nat Methods.* 2012; 9: 676–682. <https://doi.org/10.1038/nmeth.2019> PMID: 22743772
45. Thévenaz P, Unser M. User-friendly semiautomated assembly of accurate image mosaics in microscopy. *Microsc Res Tech.* 2007; 70: 135–146. <https://doi.org/10.1002/jemt.20393> PMID: 17133410
46. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3—new capabilities and interfaces. *Nucleic Acids Res.* 2012; 40: e115–e115. <https://doi.org/10.1093/nar/gks596> PMID: 22730293

47. Rozanski A, Moon H, Brandl H, Martín-Durán JM, Grohme MA, Hüttner K, et al. PlanMine 3.0—improvements to a mineable resource of flatworm biology and biodiversity. *Nucleic Acids Res.* 2019; 47: D812–D820. <https://doi.org/10.1093/nar/gky1070> PMID: 30496475
48. Hagstrom D, Zhang S, Ho A, Tsai ES, Radić Z, Jahromi A, et al. Planarian cholinesterase: molecular and functional characterization of an evolutionarily ancient enzyme to study organophosphorus pesticide toxicity. *Arch Toxicol.* 2018; 92: 1161–1176. <https://doi.org/10.1007/s00204-017-2130-7> PMID: 29167930
49. Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics.* 2009; 25: 1189–1191. <https://doi.org/10.1093/bioinformatics/btp033> PMID: 19151095
50. Notredame C, Higgins DG, Heringa J. T-coffee: a novel method for fast and accurate multiple sequence alignment. *J Mol Biol.* 2000; 302: 205–217. <https://doi.org/10.1006/jmbi.2000.4042> PMID: 10964570
51. Takano T, Pulvers JN, Inoue T, Tarui H, Sakamoto H, Agata K, et al. Regeneration-dependent conditional gene knockdown (Readyknock) in planarian: Demonstration of requirement for Djsnap-25 expression in the brain for negative phototactic behavior. *Dev Growth Differ.* 2007; 49: 383–394. <https://doi.org/10.1111/j.1440-169X.2007.00936.x> PMID: 17547648
52. Fenwick AJ, Fowler DK, Wu S-W, Shaffer FJ, Lindberg JEM, Kinch DC, et al. Direct anandamide activation of TRPV1 produces divergent calcium and current responses. *Front Mol Neurosci.* 2017; 10. <https://doi.org/10.3389/fnmol.2017.00200> PMID: 28680392
53. Zygmunt PM, Petersson J2, Andersson DA2, Chuang H-H3, Sürga M, Rd Ê2, et al. Vanilloid receptors on sensory nerves mediate the vasodilator action of anandamide. *Nature.* 1999. Available: www.nature.com
54. Muller C, Morales P, Reggio PH. Cannabinoid ligands targeting TRP channels. *Front Mol Neurosci.* 2019; 11: 487. <https://doi.org/10.3389/fnmol.2018.00487> PMID: 30697147
55. Buttarelli FR, Pontieri FE, Margotta V, Palladini G. Cannabinoid-induced stimulation of motor activity in planaria through an opioid receptor-mediated mechanism. *Prog Neuro-Psychopharmacology Biol Psychiatry.* 2002; 26: 65–68. [https://doi.org/10.1016/S0278-5846\(01\)00230-5](https://doi.org/10.1016/S0278-5846(01)00230-5)
56. Buttarelli FR, Pellicano C, Pontieri FE. Neuropharmacology and behavior in planarians: Translations to mammals. *Comp Biochem Physiol—C Toxicol Pharmacol.* 2008; 147: 399–408. <https://doi.org/10.1016/j.cbpc.2008.01.009> PMID: 18294919
57. Raisinghani M, Zhong L, Jeffry JA, Bishnoi M, Pabbidi RM, Pimentel F, et al. Activation characteristics of transient receptor potential ankyrin 1 and its role in nociception. *Am J Physiol Cell Physiol.* 2011; 301: C587–600. <https://doi.org/10.1152/ajpcell.00465.2010> PMID: 21653898
58. Macpherson LJ, Dubin AE, Evans MJ, Marr F, Schultz PG, Cravatt BF, et al. Noxious compounds activate TRPA1 ion channels through covalent modification of cysteines. *Nature.* 2007; 445: 541–545. <https://doi.org/10.1038/nature05544> PMID: 17237762
59. Gupta R, Saito S, Mori Y, Itoh SG, Okumura H, Tominaga M. Structural basis of TRPA1 inhibition by HC-030031 utilizing species-specific differences. *Sci Rep.* 2016; 6: 37460. <https://doi.org/10.1038/srep37460> PMID: 27874100
60. Paulsen CE, Armache J-P, Gao Y, Cheng Y, Julius D. Structure of the TRPA1 ion channel suggests regulatory mechanisms. *Nature.* 2015; 520: 511–517. <https://doi.org/10.1038/nature14367> PMID: 25855297
61. Pirotte N, Stevens A-S, Fraguas S, Plusquin M, Van Roten A, Van Belleghem F, et al. Reactive oxygen species in planarian regeneration: An upstream necessity for correct patterning and brain formation. *Oxid Med Cell Longev.* 2015; 2015: 392476. <https://doi.org/10.1155/2015/392476> PMID: 26180588
62. Cortright D, Szallasi A. TRP channels and pain. *Curr Pharm Des.* 2009; 15: 1736–1749. <https://doi.org/10.2174/138161209788186308> PMID: 19442187
63. Sanz-Salvador L, Andrés-Bordería A, Ferrer-Montiel A, Planells-Cases R. Agonist- and Ca²⁺-dependent desensitization of TRPV1 channel targets the receptor to lysosomes for degradation. *J Biol Chem.* 2012; 287: 19462–71. <https://doi.org/10.1074/jbc.M111.289751> PMID: 22493457
64. Akopian AN, Ruparel NB, Patwardhan A, Hargreaves KM. Cannabinoids desensitize capsaicin and mustard oil responses in sensory neurons via TRPA1 activation. *J Neurosci.* 2008; 28: 1064–1075. <https://doi.org/10.1523/JNEUROSCI.1565-06.2008> PMID: 18234885
65. Cochet-Escartin O, Carter JA, Chakraverti-Wuerthwein M, Sinha J, Collins EMS. Slo1 regulates ethanol-induced scrunching in freshwater planarians. *Phys Biol.* 2016; 13: 1–12. <https://doi.org/10.1088/1478-3975/13/5/055001> PMID: 27609598