

Copyright The University of Chicago 2019. Preprint (not copyedited or formatted).
Please use DOI when citing or quoting. DOI: 10.1086/704248

Title:

Validating the demethylating effects of 5-aza-2'-deoxycytidine in insects requires a whole-genome approach (A reply to Ellers *et al.*)

Authors: Nicola Cook*¹, Darren J. Parker*^{2,3}, Eran Tauber⁴, Bart A. Pannebakker,⁵ and David M. Shuker¹.

*these authors contributed equally to the paper

1 School of Biology, University of St Andrews, St Andrews, UK.

2 Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland.

3 Swiss Institute of Bioinformatics, Lausanne, Switzerland

4 Faculty of Natural Sciences, University of Haifa, Haifa, Israel.

5 Department of Plant Sciences, Laboratory of Genetics, Wageningen University, Wageningen, The Netherlands.

Keywords: sex ratio, 5-aza-2'-deoxycytidine, DNA methylation, *Nasonia vitripennis*

No. words in main text: 4815 (excluding references)

Note: This article is a reply to a comment by Ellers *et al.* 2019 (manuscript number: 58628).

Elements: Title Page, Abstract, Main Text, Acknowledgements, References, Figure legends

ABSTRACT

We previously demonstrated that treatment with the demethylating agent 5-aza-2'-deoxycytidine (5-aza-dC) alters the offspring sex ratios produced by females of the parasitoid wasp *Nasonia vitripennis* (Cook *et al.* 2015). Females allocate offspring sex ratio in line with Local Mate Competition theory, producing more or less female-biased sex ratios as the number of other females laying eggs on a patch varies, thereby reducing competition amongst their sons for mates. Interestingly, treatment with 5-aza-dC did not ablate the facultative sex allocation response. Instead, sex ratios became less female-biased, a shift in the direction of the optimum sex ratio for paternally-inherited alleles according to genomic conflict theory. This was the first (albeit indirect) experimental evidence for genomic conflict over sex allocation. Ellers *et al.* (2019) have since assayed the effects of 5-aza-dC on DNA methylation in ten *Nasonia* genes, finding no evidence of demethylation in these 10 genes, from which they conclude that 5-aza-dC has no demethylating capability in *Nasonia vitripennis*. Quantifying the efficacy of 5-aza-dC in terms of demethylation is indeed crucial to in-depth interpretation of studies using 5-aza-dC to link phenotypes to epigenetic regulation. Here, we outline the mode of action of 5-aza-dC and demonstrate that determining the efficacy of 5-aza-dC in insect systems requires a whole-genome approach.

Main Text

DNA methylation, sex allocation and 5-aza-2'-deoxycytidine

In a previous study, we demonstrated that treatment with the demethylating agent 5-aza-2'-deoxycytidine (5-aza-dC) alters the sex allocation response of *Nasonia vitripennis* females (Cook *et al.* 2015). As with all Hymenoptera, *N. vitripennis* is a haplodiploid species, with males arising from unfertilized, haploid eggs and females developing from fertilized, diploid eggs. Females are in putative control of sex allocation, either laying unfertilized eggs that will become sons or releasing sperm to fertilize eggs that will become daughters. Female *Nasonia* allocate sex broadly in line with Local Mate Competition (LMC) theory (Hamilton 1967, Burton-Chellew *et al.* 2008, West 2009); females laying eggs alone will produce very female-biased sex ratios to minimize competition amongst their sons for mates. As the number of laying females (foundresses) on a patch increases though, the level of LMC consequently declines and less female-biased sex ratios are produced.

Recent theory, however, has demonstrated that maternally- and paternally- inherited alleles should have differing patterns of optimal sex allocation under LMC, with maternally-inherited alleles selected to produce more female-biased sex ratios (see Wild and West 2009 for a detailed explanation). Importantly, parent-of-origin information must be associated with each allele, for instance via epigenetic marks such as DNA methylation. We demonstrated that disrupting DNA methylation with 5-aza-dC did not ablate the facultative sex allocation response, but instead found that sex ratios were shifted in the direction of the paternal optimum (sex ratios became more less female-biased; Cook *et al.* 2015). This result was consistent with predictions from genomic conflict over sex allocation theory (Wild and West 2009), suggesting that unmanipulated sex ratios are closer to the optimum for maternally-inherited alleles (Figure 1). Ellers *et al.* (2019), in their valuable comment on this study, raised the valid point that the

efficacy of 5-aza-dC in insect systems should be tested, and have suggested from their own experimentation that 5-aza-dC does not disrupt methylation in *Nasonia vitripennis*. To fully address their comments, we will begin by briefly reviewing the mechanism by which 5-aza-dC brings about the demethylation of DNA, before showing that assessing the role of 5-az-dC requires a whole genome approach.

Originally synthesized in 1964 by Piskala and Sorm (and also known as 5-AZA-CdR, DAC, decitabine, ZdCyD and AzaD amongst other names), 5-aza-dC is an analog of the natural nucleoside 2'-deoxycytidine (the carbon at the 5-position of the pyrimidine ring is replaced by a nitrogen) and was repurposed as a demethylating agent in 1980 (Jones and Taylor 1980). Much of the current attention surrounding 5-aza-dC is attributable to its success in the treatment of cancers, particularly acute myeloid leukaemia and myelodysplastic syndrome (Seelan 2018, Bryan *et al.* 2011, Momparler 2012). In mammals, DNA methylation acts as a transcriptional silencer and the clinical efficacy of 5-aza-dC results from its ability to reactivate tumor suppressor genes silenced by aberrant DNA methylation (Karahoca and Momparler 2013).

DNA methylation for the vast majority of cases occurs at the C5 position of the cytosine ring in CpG dinucleotides and is catalysed by DNA methyltransferase enzymes (Dnmts), of which 3 types are known in eukaryotes. Dnmt3 establishes DNA methylation patterns *de novo*, Dnmt1 maintains these patterns, and Dnmt2 is involved in tRNA methylation (Lyko 2018). DNA methyltransferase genes have been found across the Insecta, varying in presence/absence and copy number across species. The *Nasonia vitripennis* genome encodes a full DNA methylation toolkit including 3 copies of Dnmt1 (Werren *et al.* 2010). DNA methylation itself is also present across the Insecta and is largely restricted to the transcribed regions of genes, in contrast to DNA methylation in mammals, which is found throughout the genome (Bewick *et al.* 2017, Glastad *et al.* 2011). In addition, genomic levels of DNA methylation in insects is much lower (0-14%; Bewick *et al.* 2017) as compared to 60-90% in mammals (Glastad *et al.* 2011). Regardless of

the level of DNA methylation present within a genome, or whether methylation signals transcriptional suppression (as in mammals) or acts to maintain transcript integrity and/or initiate/regulate splicing (as has been suggested for the insects: reviewed Glastad *et al.* 2011), the epigenetic modification itself is, to our knowledge, chemically identical.

The introduction of 5-aza-dC to a cell results in a cascade of chemical reactions to cause an eventual reduction in DNA methylation. Upon its introduction to a cell 5-aza-dC is activated *in situ* via phosphorylation by deoxycytidine kinase, producing the biologically active triphosphate form of 5-aza-dC, 5-aza-dCTP (Seelan *et al.* 2018). This active form is an excellent substrate for DNA polymerase and is readily incorporated into DNA during the S-phase of the cell cycle (Momparler 2005, Christman 2002, Christman *et al.* 1983, Bouchard and Momparler 1983). In the resulting daughter strands produced from a DNA replication event, hemi-methylated CpG sites are remethylated by Dnmt1 to fully restore the patterns of methylation present in the parent strand. However, where 5-aza-dCTP is incorporated into the daughter DNA strand, a covalent bond is formed between 5-aza-dCTP and a cysteine residue in the active site of Dnmt1. Since the carbon at the 5-position in dCTP is replaced with a nitrogen in 5-aza-dCTP, the methyl transfer reaction cannot take place and Dnmt1 becomes irreversibly bound to the DNA (Seelan *et al.* 2018, Creusot *et al.* 1982, Taylor and Jones 1982). Thus, treatment with 5-aza-dC results in a diminishing pool of Dnmt1, resulting in the passive loss of DNA methylation: this is the key effect of 5-aza-dC. Importantly, there is evidence that the effects of 5-aza-dC are unlikely to be random across methylated CpGs in a given genome, as genomic context can influence the extent to which its action is effective. DNA sequence context plays a role, with demethylation of CpGs outside of CpG islands occurring at a higher frequency (Hagemann *et al.* 2011, Mossmann *et al.* 2010). The distribution of transcription factor binding sites is also influential with, for example, “demethylation-sensitive” genes enriched for Forkhead box (Fox) binding sites and “demethylation-resistant” genes enriched for basic Helix-Loop-Helix (bHLH) binding sites. Indeed, some genes are never demethylated and these are enriched for binding motifs not present

in demethylated genes. More broadly, 5-aza-dC activity appears to be targeted to transcriptionally active euchromatin in mammalian cells (Ramos *et al.* 2015). Given that the mode of action of 5-aza-dC is passive, these results indicate that specific loci remain demethylated after replication, whereas others at first appear “resistant” because they become preferentially remethylated by the remaining pool of DNA methyltransferase, leading to non-random patterns of demethylation (Hagemann *et al.* 2011).

We agree with Ellers *et al.* (2019) that quantifying the demethylation efficacy of orally-administered 5-aza-dC in *N. vitripennis* will permit a more complete understanding of the relationship between DNA methylation and the sex allocation phenotype in *Nasonia* (see Cook *et al.* 2015). More broadly, this will allow us to determine the extent to which genome-wide demethylation can be altered with 5-aza-dC in insect systems. In their analysis of the involvement of DNA methylation in the photoperiodic diapause response in *Nasonia*, Pegoraro *et al.* (2016) demonstrated that consumption of 5-aza-dC led to demethylation at specified CpG sites on a small scale. Ellers *et al.* (2019) have examined the demethylating effect of 5-aza-dC at 155 CpG sites spread across 9 candidate genes in a slightly more comprehensive analysis. The authors found no significantly differentially methylated CpGs at any of the 155 sites in association with 5-aza-dC exposure and conclude from this result that 5-aza-dC has no demethylating effect. To our knowledge, there is no evidence to suggest that the candidate genes chosen by Ellers *et al.* (2019) are more (or less) likely to be influenced by 5-aza-dC – i.e. to have genomic contexts which favor or disfavor its action. It would also be the first time that 5-aza-dC would have been shown to be inert in an organism in terms of influencing DNA methylation. As such, to get a more complete picture and to more accurately quantify the demethylation efficacy of 5-aza-dC, we suggest that a whole-genome approach is required.

In the first instance, it is important to note that genomic DNA methylation in *Nasonia* is relatively low, with estimates ranging from 0.63% to 1.6 % of all CpG sites methylated (Beeler *et al.* 2014, Wang *et al.* 2013). These estimates are comparable with other hymenopterans such as the honeybee, which has

approximately 0.7% methylated CpGs (Lyko *et al.* 2010). Therefore, a demethylation effect will be inherently difficult to detect just by looking across a small number of CpGs. Ellers *et al.* (2019) examined fragments of 9 genes to examine for evidence of demethylation due to 5-aza-dC treatment. Two genes were chosen because of a heavy methylation pattern (aa and rnapol) and five (acc, fabd, fasn1, fasn3, and mcd) are part of the fatty acid metabolic pathway, purported to be epigenetically regulated. Two additional genes were selected from the study of Pegoraro *et al.* (2016; perq and wdr36) based on the incorrect assumption that these genes are heavily methylated under the conditions tested by Ellers *et al.* In fact, perq and wdr36 display elevated methylation only under long photoperiods. Whilst these genes may be heavily methylated under certain conditions and/or epigenetically regulated, there is no confirmed evidence in insects that the genomic context of these particular genes predisposes them to demethylation by 5-aza-dC. As outlined above, demethylation via 5-aza-dC occurs due to a lack of functional Dnmt1 to restore methylation to CpG sites post-replication. Given the contextual nature of the action of 5-aza-dC, there is a risk of false negatives in terms of assessing its genome-wide action. Therefore, a lack of demethylation in the gene fragments examined by Ellers *et al.* (2019) cannot be used to conclude that 5-aza-dC does not affect DNA methylation across the genome more broadly.

Quantifying the efficacy of 5-aza-dC

Recently we carried out a genome-wide analysis of the demethylating capacity of 5-aza-dC in *Nasonia vitripennis* (Cook *et al.* 2018). Briefly, female wasps were exposed to one of three 5-aza-dC exposure regimes: 1) 20% sucrose for 24h, 2) 20% sucrose supplemented with 10 μ M 5-aza-dC for 24h and 3) 20% sucrose supplemented with 10 μ M 5-aza-dC for 48h. For each of these three regimes, females were harvested at 0, 24 and 48h post-exposure and, prior to DNA extraction, heads were excised from the bodies as a nod in the direction of a tissue-specific analysis. Whole-genome sequencing of bisulphite-

treated DNA samples was carried out to determine the methylation status of individual CpG sites across the genome (proportion of methylated reads/total reads per CpG). Median coverage depth across the whole experiment was approximately 900X across 4 million CpGs after quality filtering.

To examine the data as a whole, we summed proportion of methylated reads for each CpG by gene, to broadly account for contextual similarity of CpGs within the same genic region. Data were available for over 11,000 genes comprising 85% of the protein-coding genes in the *Nasonia vitripennis* genome. Distinct clustering of samples was associated with 5-aza-dC exposure regime, clearly demonstrating that 5-aza-dC exposure has a quantifiable effect on DNA methylation status on a genome-wide scale (Figure 2A). To explore DNA methylation changes at the gene level, we used a generalized linear modelling approach to determine which genes had significantly different proportions of methylation across the gene with respect to 5-aza-dC exposure regime, time-point after exposure, and tissue. To encapsulate these results, we found that approximately 8500 genes showed significantly altered methylation in response to treatment with 5-aza-dC. Interestingly, it appears that methylation in some genes initially increases in response to treatment with 5-aza-dC, an effect that is noticeable after 24h of exposure and then less prevalent after 48h of exposure (Figure 2B). In our data, two of the genes tested by Ellers *et al.*, *aa* and *rnapol*, do not exhibit significant differential methylation in association with “feeding regime” in our experiments. However, four of their tested genes (*perq*, *acc*, *fasn1*, and *fasn3*) exhibit significantly altered methylation levels in association with an overall effect of “feeding regime”. Interestingly a decrease in methylation levels of these genes (relative to controls) is noted in wasps that were exposed to 5-aza-dC for a 48-hour period but not in wasps exposed to 5-aza-dC for a 24-hour period. This indicates that 5-aza-dC may take longer to exert its effects on these four genes. We also note that Ellers *et al.* have used newly emerged *N. vitripennis* females, whereas we used two-day old mated females; the effects of 5-aza-dC are known to differ with life-stage (see below). We are unable to draw comparisons between our analysis and that of Ellers *et al.* for their remaining three genes; for *wdr36* and

fabd we do not find any CpGs in the Ensembl *Nasonia vitripennis* annotation (v 2.1), and mcd was excluded from our analysis as genes with zero coverage for all samples in any particular treatment group were removed prior to statistical analysis.

Hyper-methylation has been observed previously in response to treatment with 5-aza-dC. Using “amplification of intermethylated sites” (AIMS) in honeybees, Amarasinghe *et al.* (2014) observed hypomethylation in 10 and hypermethylation in 9 of 62 loci in 5-aza-dC exposed individuals relative to controls. Also, it looks as though Ellers *et al.* (2019), using bisulphite amplicon sequencing, may have uncovered some CpGs that exhibit hypermethylation in 5-aza-dC treated samples, particularly in *perq* (Ellers *et al.* 2019: Figure 3). It is also noteworthy that two of the nine genes reported by Ellers *et al.* (2019; in their Table 3 and Figure 3) show marginally significant Treatment X Site interaction (fabd, $p=0.055$; *fasn1*, $p=0.079$). This reflects the fact that 5-aza-dC affects CpG sites in a context-specific manner (and possibly in different directions), an effect which is likely masked in their data by other sites that do not differ. The mechanism by which hypermethylation is brought about by 5-aza-dC is as yet unclear. Current understanding of the DNA methylation machinery suggests that there are a range of Dnmt-including complexes and that there is cross-talk between them in terms of their roles in *de novo* and maintenance methylation (Hervouet *et al.* 2018). It is possible that drug-induced demethylation in one region of the genome may have altered the regulation of the DNA methylation machinery itself, such that hypermethylation occurs. Nonetheless, it is clear that 5-aza-dC alters methylation patterns in the *N. vitripennis* genome and that this is associated with the allocation of offspring sex ratio (Cook *et al.* 2018, 2015).

Possible confounding effects

Ellers *et al.* (2019) raised the valid concern that the anti-metabolic activity of 5-aza-dC could result in a decline in the physical condition of *Nasonia vitripennis* females, indirectly producing observed phenotypic

differences through a mechanism independent of DNA demethylation. Whilst we have shown that 5-aza-dC at the dosage used in our original experiment (Cook *et al.* 2015) produces altered methylation patterns (above; Cook *et al.* 2018), the possibility of confounding side-effects should not be ignored. We do note, however, that an effect of condition-dependent sex allocation has yet to be elucidated in *Nasonia vitripennis*, although pathological side-effects of any given chemical exposure cannot be discounted. We also note that sub-lethal exposure to neurotoxic neonicotinoid pesticides does also influence sex allocation, albeit in the opposite direction to that seen with 5-aza-dC; i.e. more female-biased sex ratios are seen than expected: Whitehorn *et al.* 2015, Cook *et al.* 2016.

In humans, the major side effect of 5-aza-dC treatment is myelosuppression (Karahoca and Momparler 2013), a condition in which bone marrow activity is decreased resulting in fewer red and white blood cells and platelets. It is difficult of course to extrapolate to what the major side effect of the drug in insects might be, if any. In mammals, high-dose schedules result in an increased presence of DNA-Dnmt1 adducts which impede DNA polymerase function, leading to growth arrest and apoptosis. However, low-dose schedules are known to capitalize on the hypomethylating capability of the drug and minimize these additional cytotoxic effects (Jones and Taylor 1980, Jüttermann *et al.* 1994, Issa *et al.* 2004, Yang *et al.* 2006). In trialling the cancer drug 5-aza-dC as a tool for molecular ecology and evolution studies, researchers have also used low doses to achieve DNA demethylation (Uçkan *et al.* 2007, Amarasinghe *et al.* 2014, Cook *et al.* 2015, Pegoraro *et al.* 2016, Sak 2017). In their 2014 study, examining DNA methylation differences between different reproductive worker castes in *Bombus terrestris*, Amarasinghe *et al.* used 5-aza-dC (<10 μ M) on both callows (bees less than one day old) and adults (bees more than one day old). This is less than half of the lowest concentration (25 μ M) used to test for genotoxic effects of 5-aza-dC in *Drosophila melanogaster* at a range of concentrations (Cunha *et al.* 2002). Their experimentation showed no effect of 5-aza-dC on the adult bees in contrast to the callows, which developed larger ovaries, became more aggressive, and showed altered methylation patterns (at 19 of 62 tested loci) relative to controls of

the same age. The authors hypothesized that cells in adult bees may be largely post-mitotic thus decreasing the efficacy of 5-aza-dC in adults vs callows. Importantly, neither callows nor adults showed reduced activity or reproduction, indeed callows became more reproductive. If the effects of 5-aza-dC were merely toxic it is reasonable to assume that reduced fitness would be observed in both age groups even if altered methylation patterns occurred only in callow bees.

Ellers *et al.* quantified the lipid reserves of both control wasps and those treated with 5-aza-dC as a general measure of body condition. The authors showed that lipid reserves were significantly lower as a result of 5-aza-dC treatment and that reserves in both treatment and control groups declined over the duration of the experiment. The authors conclude that these differences in lipid reserves can only be attributed to a detrimental effect of 5-aza-dC on wasp health given that they observed no alterations in DNA methylation in a total of 6 genes involved in fatty acid metabolism. Given demonstrations that 5-aza-dC does produce altered methylation patterns (Amarasinghe *et al.* 2014, Pegoraro *et al.* 2016, Cook *et al.* 2018) it is possible that hypo- or hyper-methylation in genes untested in their experiment could have induced a decrease in lipid reserves; the six genes tested cannot be regarded as representative of the entire fatty acid metabolism pathway. Interestingly, Amarasinghe *et al.* (2014) noted increased reproductive capability as a result of 5-aza-dC so it is possible that the observed reduction in lipid reserves is due to the diversion of resources elsewhere. It is also unclear whether the decline in lipid reserves was sufficient to produce a true reduction in fitness in terms of offspring production or longevity as this information is not available.

5-aza-dC and evolutionary ecology

Research has determined that 5-aza-dC acts to alter methylation patterns in insects and has assisted evolutionary biologists in elucidating a role for epigenetics in a range of phenotypic traits, including sex

allocation, worker reproduction, and diapause (Cook *et al.* 2018, Pegoraro *et al.* 2016, Amarasinghe *et al.* 2014). However, a greater understanding of how 5-aza-dC brings about altered methylation could improve the precision of its use in evolutionary ecology studies. As a starting point, careful selection of the life-stage of experimental individuals can prove important. 5-aza-dC had no demonstrable effect on DNA methylation when administered to bees more than one day old but resulted in significant hypo- and hyper-methylation in younger bees (Amarasinghe *et al.* 2014). This may be due to the passive mode of action of 5-aza-dC and a higher percentage of cells in adult bees being post-mitotic. This effect is also likely to be dose-dependent and species-specific. Therefore, if no effect of 5-aza-dC is initially observed in a given species, it is worthwhile investigating the efficacy at different life-stages.

Upon introduction to the cell, 5-aza-dC is converted to 5-aza-dCTP but, in addition, 5-aza-dC is also subject to deamination resulting in an inactive form of the drug being produced alongside the active form. This is due to very high levels of cytidine deaminase in the human liver and spleen (Momparker *et al.* 1985, Chabot *et al.* 1983). Ellers *et al.* (2019) called attention to the fact that cytidine deaminase is also present in insects. Indeed, there is a “cytidine deaminase-like” gene in the *Nasonia vitripennis* genome (NCBI: LOC107980937) which may act to degrade some of the introduced 5-aza-dC to its inactive form. To counteract this effect, the use of an inhibitor of cytidine deaminase, such as tetrahydrouridine or zebularine has been suggested (Eliopoulos 1998). When administered in combination with 5-aza-dC in preclinical trials, the cytidine deaminase inhibitor zebularine was shown to increase the clinical efficacy of 5-aza-dC in cell lines by reducing the degradation of 5-aza-dC (Lemaire *et al.* 2009). Pilot experiments to determine whether the use of a cytidine deaminase inhibitor increases the efficacy of 5-aza-dC in insect study systems would be beneficial.

Concluding remarks

Here, we have discussed the mechanisms by which the demethylating agent 5-aza-dC exerts its effect and illustrated that a whole-genome approach is required to determine the efficacy of this agent in *Nasonia vitripennis*. Having used drug-induced demethylation to validate a role for epigenetic regulation in sex allocation (Cook *et al.* 2015, 2018), the desirable next step would be to ascertain which CpGs are involved in the expression of the sex allocation phenotype; i.e. the causal CpGs. As outlined above, research has shown that drug-induced demethylation is non-random and reproducible. Experimentation is warranted to determine the reproducibility and the genomic context underlying the methylation patterns produced by 5-aza-dC in *N. vitripennis* and other insect systems.

Acknowledgements

Firstly, we would like to thank Mark Lammers and Jacintha Ellers for the collegiate and constructive engagement with our previous research and the editorial board for the opportunity to respond to their critique. Secondly, we thank Jade Green for assistance in the laboratory with practical work referred to in this manuscript. We are also grateful to Matthew Arno and the Edinburgh Genomics team for their sequencing and downstream support with the data. This work was initiated as part of Natural Environment Research Council grant NE/J024481/1.

Literature Cited

- Amarasinghe, H. E., C. I. Clayton, and E. B. Mallon. 2014. Methylation and worker reproduction in the bumble-bee (*Bombus terrestris*). *Proceedings of the Royal Society B* 281: 20132502.
- Beeler, S. M., G. T. Wong, J. M. Zheng, E. C. Bush, E. J. Remnant, B. P. Oldroyd, and R. A. Drewell. 2014. Whole-genome DNA methylation profile of the Jewel Wasp (*Nasonia vitripennis*). *G3: Genes|Genomes|Genetics* 4: 383-388.
- Bewick, A. J., K. J. Vogel, A. J. Moore, and R. J. Schmitz. 2017. Evolution of DNA methylation across Insects. *Molecular Biology and Evolution* 34:654-665.
- Bouchard, J., and R. L. Mompalmer. 1983. Incorporation of 5-aza-2'-deoxycytidine 5'-triphosphate into DNA: Interactions with mammalian DNA polymerase and DNA methylase. *Molecular Pharmacology* 24:109-114.
- Bryan, J., H. Kantarjian, G. Garcia-Manero, and E. Jabbour. 2011. Pharmacokinetic evaluation of decitabine for the treatment of leukemia. *Expert Opinion on Drug Metabolism and Toxicology* 7:661-672.
- Burton-Chellew, M. N., T. Koevoets, B. K. Grillenberger, E. M. Sykes, S. L. Underwood, K. Bijlsma, J. Gadau, L. van de Zande, L. W. Beukeboom, S. A. West, and D. M. Shuker. 2008. Facultative sex ratio adjustment in natural populations of wasps: cues of local mate competition and the precision of adaptation. *American Naturalist* 172:393-404.
- Chabot, G., J. Bouchard, and R. L. Mompalmer. 1983. Kinetics of deamination of 5-aza-2'-deoxycytidine and cytosine arabinoside by human liver cytidine deaminase and its inhibition by 3-deazauridine, thymidine or uracil arabinoside. *Biochemical Pharmacology* 32:1327-1328.
- Christman, J. K. 2002. 5-Azacytidine and 5-aza-2'-deoxycytidine as inhibitors of DNA methylation: mechanistic studies and their implications for cancer therapy. *Oncogene* 21:5483-5495.
- Christman, J. K., N. Mendelsohn, D. Herzog, and N. Schneiderman. 1983. Effect of 5-Azacytidine on Differentiation and DNA Methylation in Human Promyelocytic Leukemia Cells (HL-60). *Cancer Research* 43:763-769.
- Cook, N., D. J. Parker, E. Tauber, B. A. Pannebakker, and D. M. Shuker. 2018. 5-aza-2'-deoxycytidine alters methylation patterns in *Nasonia vitripennis* on a genome-wide scale. *bioRxiv*: <https://doi.org/10.1101/437202>.

- Cook, N., J. Green, D. M. Shuker, and P. W. Whitehorn. 2016. Exposure to the neonicotinoid imidacloprid disrupts sex allocation cue use during superparasitism in the parasitoid wasp *Nasonia vitripennis*. *Ecological Entomology* 41:693-697.
- Cook, N., B. A. Pannebakker, E. Tauber, and D. M. Shuker. 2015. DNA methylation and sex allocation in the parasitoid wasp *Nasonia vitripennis*. *American Naturalist* 186:513-518.
- Creusot, F., G. Acs, and J. K. Christman. 1982. Inhibition of DNA methyltransferase and induction of Friend erythroleukemia cell differentiation by 5-azacytidine and 5-aza-2'-deoxycytidine. *The Journal of Biological Chemistry* 257:2041-2048.
- Cunha, K. S., M. L. Reguly, U. Graf, and H. H. Rodrigues de Andrade. 2002. Somatic recombination: a major genotoxic effect of two pyrimidine antimetabolite chemotherapeutic drugs in *Drosophila melanogaster*. *Mutation Research* 514:95-103.
- Eliopoulos, N., C. Denis, and R. L. Momparler. 1998. Drug resistance to 5-aza-2'-deoxycytidine, 2',2'-difluorodeoxycytidine, and cytosine arabinoside conferred by retroviral-mediated transfer of human cytidine deaminase cDNA into murine cells. *Cancer Chemotherapy and Pharmacology* 42:373-378.
- Ellers, J., M. Visser, J. Mariën, K. Kraaijeveld, and M. Lammers. 2019. The importance of validating the demethylating effect of 5-aza-dC in model species. *American Naturalist* 194:____ - ____.
- Glastad, K. M., B. G. Hunt, S. V. Yi, and M. A. D. Goodisman. 2011. DNA methylation in insects: on the brink of the epigenomic era. *Insect Molecular Biology* 20:553-565.
- Hagemann, S., O. Heil, F. Lyko, and B. Brueckner. 2011. Azacytidine and decitabine induce gene-specific and non-random DNA demethylation in human cancer cell lines. *PLOS ONE* 6:e17388.
- Hamilton, W. D. 1967. Extraordinary sex ratios. *Science* 156:477-488.
- Hervouet, E., P. Peixoto, R. Delage-Mourroux, M. Boyer-Guittaut, and P. Cartron. 2018. Specific or not specific recruitment of DNMTs for DNA methylation, an epigenetic dilemma. *Clinical Epigenetics* 10:17.
- Issa, J. J., G. Garcia-Manero, F. J. Giles, R. Mannari, D. Thomas, S. Faderl, E. Bayar, J. Lyons, C. S. Rosenfeld, J. Cortes, and H. M. Kantarjian. 2004. Phase I study of low-dose prolonged exposure schedules of the hypomethylating agent 5-aza-2'-deoxycytidine (decitabine) in hematopoietic malignancies. *Blood* 103:1635-1640.

- Jones, P. A., and S. M. Taylor. 1980. Cellular differentiation, cytidine analogs and DNA methylation. *Cell* 20:85-93.
- Juttermann, R., E. Li, and R. Jaenisch. 1994. Toxicity of 5-aza-2'-deoxycytidine to mammalian cells is mediated primarily by covalent trapping of DNA methyltransferase rather than DNA demethylation. *Proceedings of the National Academy of Sciences of the USA* 91:11797-11801.
- Karahoca, M., and R. L. Momparler. 2013. Pharmacokinetic and pharmacodynamic analysis of 5-aza-2'-deoxycytidine (decitabine) in the design of its dose-schedule for cancer therapy. *Clinical Epigenetics* 5:3.
- Lemaire, M., L. F. Momparler, N. J. M. Raynal, M. L. Bernstein, and R. L. Momparler. 2009. Inhibition of cytidine deaminase by zebularine enhances the antineoplastic action of 5-aza-2'-deoxycytidine. *Cancer Chemotherapy and Pharmacology* 63:411-416.
- Lyko, F., S. Foret, R. Kucharski, S. Wolf, C. Falckenhayn, and R. Maleszka. 2010. The honey bee epigenomes: differential methylation of brain DNA in queens and workers. *PLoS Biology* 8:e1000506.
- Lyko, F. 2018. The DNA methyltransferase family: a versatile toolkit for epigenetic regulation. *Nature Reviews Genetics* 19:81-92.
- Momparler, R. L. 1985. Molecular, cellular and animal pharmacology of 5-aza-2'-deoxycytidine. *Pharmacology and Therapeutics* 30:287-299.
- Momparler, R. L. 2005. Epigenetic therapy of cancer with 5-aza-2'-deoxycytidine (decitabine). *Seminars in Oncology* 32:443-451.
- Momparler, R. L. 2012. A perspective on the comparative antileukemic activity of 5-Aza-2'-deoxycytidine (Decitabine) and 5-Azacytidine (Vidaza). *Pharmaceuticals* 5:875-881.
- Mossman, D., K. Kim, and R. J. Scott. 2010. Demethylation by 5-aza-2'-deoxycytidine in colorectal cancer cells targets genomic DNA whilst promoter CpG island methylation persists. *BMC Cancer* 10:366.
- Pegoraro, M., A. Bafna, N. J. Davies, D. M. Shuker, and E. Tauber. 2016. DNA methylation changes induced by long and short photoperiods in *Nasonia*. *Genome Research* 26:203-210.
- Piskala, A., and F. Sorm. 1964. Nucleic acid components and their analogues. LI. Synthesis of 1-glycosyl derivatives of 5-azauracil and 5-azacytosine. *Collection of Czechoslovak Chemical Communications* 29:2060-2076.

- Ramos, M., N. A. Wijetunga, A. S. McLellan, M. Suzuki, and J. M. Greally. 2015. DNA demethylation by 5-aza-2'-deoxycytidine is imprinted, targeted to euchromatin and has limited transcriptional consequences. *Epigenetics and Chromatin* 8:11.
- Sak, O. 2017. The effects of 5-aza-2'-deoxycytidine on total lipid and fatty acid composition of *Apanteles galleriae* Wilkinson (Hymenoptera: Braconidae) and on its parasitized host. *Journal of the Entomological Research Society* 19:105-120.
- Seelan, R. S., P. Mukhopadhyay, M. M. Pisano, and R. M. Greene. 2018. Effects of 5-Aza-2'-deoxycytidine (decitabine) on gene expression. *Drug Metabolism Reviews* 50:193-207.
- Taylor, S. M., and P. A. Jones. 1982. Mechanism of action of eukaryotic DNA methyltransferase: use of 5-Azacytosine-containing DNA. *Journal of Molecular Biology* 162:679-692.
- Uçkan, F., S. Hepçorman Şengül, O. Sak, and M. Korkmaz. 2007. Effects of 5-aza-2'-deoxycytidine on biological parameters of larval endoparasitoid *Apanteles galleriae* (Hymenoptera: Braconidae) and on its host *Achoria grisella* (Lepidoptera: Pyralidae). *Annals of the Entomological Society of America* 100:265-269.
- Wang, X., D. Wheeler, A. Avery, A. Rago, J. Choi, J. K. Colbourne, A. G. Clark, and J. H. Werren. 2013. Function and evolution of DNA methylation in *Nasonia vitripennis*. *PLOS Genetics* 9:e1003872.
- Werren, J. H., S. Richards, C. A. Desjardins, O. Niehuis, J. Gadau, J. K. Colbourne, and *Nasonia* Genome Working Group. 2010. Functional and evolutionary insights from the genomes of three parasitoid *Nasonia* species. *Science* 327:343-348.
- West, S. A. 2009. Sex allocation. Princeton University Press, Princeton NJ.
- Whitehorn, P. W., N. Cook, C. V. Blackburn, S. M. Gill, J. Green, and D. M. Shuker. 2015. Sex allocation theory reveals a hidden cost of neonicotinoid exposure in a parasitoid wasp. *Proceedings of the Royal Society B* 282:20150389.
- Wild, G., and S. A. West. 2009. Genomic imprinting and sex allocation. *American Naturalist* 173:E1-14.
- Yang, A. S., K. D. Doshi, S. Choi, J. B. Mason, R. K. Mannari, V. Gharybian, R. Luna, A. Rashi, L. Shen, M. R. H. Estecio, H. M. Kantarjian, G. Garcia-Manero, and J. J. Issa. 2006. DNA methylation changes after 5-aza-2'-deoxycytidine therapy in patients with leukemia. *Cancer Research* 66:5495-5503.

Figure Legends

Figure 1. Reproduced from Cook *et al.* 2015. A, The optimal sex ratios under Local Mate Competition (LMC) for maternally-inherited alleles (dark gray) and paternally-inherited alleles (light gray) for a haplodiploid species with female control of sex allocation. Shown in black is the LMC prediction with no genomic conflict. Equations shown in Wild and West 2009. B, Females treated with 5-aza-dC (light gray bars) produce slightly less female-biased sex ratios than controls (dark gray bars). Both treated and untreated females vary sex ratio with foundress number as predicted. Data are presented as residuals after controlling for experimental replicate. Error bars are binomial confidence intervals.

Figure 2. Based on data from Cook *et al.* 2018. A PCA of the proportion of methylated reads per gene ($n=11,582$), showing the effects of exposure regime and tissue type. Samples cluster by exposure regime with no obvious effect of tissue type. B Log odds of methylation for each of the two 5-aza-dC exposure regimes (24-hour or 48-hour exposure period) vs the sucrose control treatment group ($n = 8,556$ genes).

Figure 1






Figure 2

B



A

