


# Anaerobic Transformation Kinetics and Mechanism of Steroid Estrogenic Hormones in Dairy Lagoon Water

Dairy products are a staple of the American diet. The main ingredient of the dairy industry, milk, has a byproduct even before it reaches the pasteurizing factory. For cows to produce milk, they need to have given birth recently. Pregnancy and lactation hormones include estrogen compounds. These hormones are excreted through urine and feces. Liquid manure and wastewater from precipitation and washing processes of the herd end up in a series of dairy farm lagoons designed to remove solids from the water. What happens to the estrogen compounds? How can we prevent hormones from contaminating the environment and causing detrimental effects in aquatic species? One research query is to determine if the lagoons have long enough retention times for hormones to degrade.


Lab experiments conducted by mixing dairy lagoon water with deionized water at a 1:1 ratio, and then spiking that solution with starting concentrations of 5 mg/L estrogen compounds showed that the estrogen compounds mainly biodegraded, and little to no abiotic degradation occurred at various temperatures. In addition, degradation rates increased with rising temperatures (15, 25, and 35°C), but declined at 45°C. These results indicate that the well-known optimum temperature for biological growth, ~37°C, would be effective at degrading estrogen compounds.

Estrogen Compounds



Additionally, degradation rates varied based on the structure of the estrogen compound. The compounds 17β-estradiol and 17α-estradiol degraded 35 and 12 times more than estrone when incubated at 25°C, respectively. This difference is most likely due to their slightly differing physical structure, and the bacterium's ability to use one compound more efficiently than the other. Over 52 days at an incubation temperature of 35°C, the total removal of 17α-estradiol was 85%; 17β-estradiol was 78%; and estrone was 77%.

Lab experiments at relevant starting concentrations (5 µg/L) of estrogen compounds were also conducted, showing similar results at higher concentrations. To achieve these similar results with lower concentrations, however, ITC scientists had to develop a new analytical detection method to accommodate the low concentrations of estrogen compounds.



## Energy

## Pollutants

### Aquatic Plastic Debris

### Metals

### Metalworking Fluids

### Per- and Polyfluoroalkyl Substances (PFASs)

### Agricultural Chemicals

### PPCPs in the Environment

### Pilot Study on PPCPs at Champaign and Urbana Wastewater Treatment Plants (WWTPs)

### PPCPs: Extending Knowledge and Mitigation Strategies

### Fate and Transport of Steroid Hormones and Veterinary Antibiotics Derived from Cattle Farms

### Uptake, Translocation, and Accumulation of Pharmaceutical and Hormone Contaminants in Vegetables

### Fate of Pharmaceutical and Personal Care Products in Irrigated Wastewater Effluent

### Karst Groundwater Contaminants in Western Illinois

### PPCPs in Karst Groundwater in Southwestern Illinois

### Triclosan in Illinois Rivers and Streams

### Tunable Luminescent Carbon Nanospheres with Well-Defined Nanoscale Chemistry for Synchronized Imaging and Therapy

### Occurrence and Removal of Pharmaceutical and Hormone Contaminants in Rural Wastewater Treatment Lagoons

### Degradation Kinetics and Mechanism of Antibiotic Cefitiofur in Recycled Water Derived from a Beef Farm

### Anaerobic Transformation Kinetics and Mechanism of Steroid Estrogenic Hormones in Dairy Lagoon Water

### Nano-CarboScavengers

### Medicine Collection Boxes

### 2008 PPCPs Symposium

2016 PPCPs in the Environment Conference

2017 Emerging Contaminants in the Aquatic Environment Conference

2018 Emerging Contaminants in the Aquatic Environment Conference

2019 Emerging Contaminants in the Environment Conference

2016 Teacher Workshop on Pharmaceutical and Personal Care Products in the Environment

PPCPs Videos

Emerging Contaminants Consortium

PCBs & PBDEs

Polycyclic Aromatic Hydrocarbons (PAHs)

Waste Utilization

Water

Instruments & Equipment

Hazardous Waste Research Fund

## Meet the Scientists

Wei Zheng

## Publications

Anaerobic Transformation Kinetics and Mechanism of Steroid Estrogenic Hormones in Dairy Lagoon Water



One Hazelwood Drive, MC-676  
Champaign, IL 61820  
p: 217-333-8940  
**Email us**

**Home of Illinois' State Scientific Surveys**  
Illinois Natural History Survey  
Illinois State Archaeological Survey  
Illinois State Geological Survey  
Illinois State Water Survey  
Illinois Sustainable Technology Center