

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications in Food Science and  
Technology

Food Science and Technology Department

---

11-24-2013

## Lifestyle, dietary factors and antibody levels to oral bacteria in cancer-free participants of a European cohort study

Dominique S. Michaud

Jacques Izard

Zachary Rubin

Ingegerd Johansson

Elisabete Weiderpass

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/foodsciefacpub>

 Part of the [Food Science Commons](#)

---

This Article is brought to you for free and open access by the Food Science and Technology Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in Food Science and Technology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

## Authors

Dominique S. Michaud, Jacques Izard, Zachary Rubin, Ingegerd Johansson, Elisabete Weiderpass, Anne Tjønneland, Anja Olsen, Kim Overvad, Marie Christine Boutron-Ruault, Françoise Clavel-Chapelon, Laure Dossus, Rudolf Kaaks, Verena A. Katzke, Heiner Boeing, Jana Foerster, Antonia Trichopoulou, Androniki Naska, Giana Ziara, Paolo Vineis, Sara Grioni, Domenico Palli, Rosario Tumino, Amalia Mattiello, Petra HM Peeters, Peter D. Siersema, Aurelio Barricarte, José-María Huerta, Esther Molina-Montes, Miren Dorronsoro, J. Ramón Quirós, Eric J. Duell, Bodil Ohlsson, Bengt Jeppsson, Anders Johansson, Pernilla Lif, Kay-Tee Khaw, Nick Wareham, Ruth C. Travis, Tim J. Key, Heinz Freisling, Talita Duarte-Salles, Magdalena Stepien, Elio Riboli, and H. Bas Bueno-de- Mesquita

---



Published in final edited form as:

*Cancer Causes Control*. 2013 November ; 24(11): . doi:10.1007/s10552-013-0265-2.

## Lifestyle, dietary factors and antibody levels to oral bacteria in cancer-free participants of a European cohort study

Dominique S. Michaud<sup>1,2</sup>, Jacques Izard<sup>3,4</sup>, Zachary Rubin<sup>1</sup>, Ingegerd Johansson<sup>5,6</sup>, Elisabete Weiderpass<sup>6,7,8,9</sup>, Anne Tjønneland<sup>10</sup>, Anja Olsen<sup>10</sup>, Kim Overvad<sup>11</sup>, Marie Christine Boutron-Ruault<sup>12,13</sup>, Françoise Clavel-Chapelon<sup>12,13</sup>, Laure Dossus<sup>12,13</sup>, Rudolf Kaaks<sup>14</sup>, Verena A. Katzke<sup>14</sup>, Heiner Boeing<sup>15</sup>, Jana Foerster<sup>15</sup>, Antonia Trichopoulou<sup>16,17</sup>, Androniki Naska<sup>16</sup>, Giana Ziara<sup>17</sup>, Paolo Vineis<sup>2,18</sup>, Sara Grioni<sup>19</sup>, Domenico Palli<sup>20</sup>, Rosario Tumino<sup>21</sup>, Amalia Mattiello<sup>22</sup>, Petra HM Peeters<sup>23</sup>, Peter D. Siersema<sup>24</sup>, Aurelio Barricarte<sup>26,27</sup>, José-María Huerta<sup>27,28</sup>, Esther Molina-Montes<sup>27,29</sup>, Miren Dorronsoro<sup>30</sup>, J. Ramón Quirós<sup>31</sup>, Eric J. Duell<sup>32</sup>, Bodil Ohlsson<sup>33</sup>, Bengt Jeppsson<sup>34</sup>, Anders Johansson<sup>5</sup>, Pernilla Lif<sup>5</sup>, Kay-Tee Khaw<sup>35</sup>, Nick Wareham<sup>36</sup>, Ruth C. Travis<sup>37</sup>, Tim J. Key<sup>37</sup>, Heinz Freisling<sup>38</sup>, Talita Duarte-Salles<sup>38</sup>, Magdalena Stepien<sup>38</sup>, Elio Riboli<sup>2</sup>, and H. Bas Bueno-de-Mesquita<sup>24,39</sup>

<sup>1</sup>Department of Epidemiology, Brown University, Providence, RI, USA <sup>2</sup>School of Public Health, Imperial College London, London, UK <sup>3</sup>The Forsyth Institute, Cambridge, MA, USA <sup>4</sup>Department of Oral Medicine, Infection and Immunity, Harvard School of Dental Medicine, Boston, MA, USA <sup>5</sup>Department of Odontology/Cariology, Umeå University, Umeå, Sweden <sup>6</sup>Samfundet Folkhälsan, Helsinki, Finland <sup>7</sup>Department of Community Medicine, Faculty of Health Sciences, University of Tromsø, Tromsø, Norway <sup>8</sup>Department of Research, Cancer Registry of Norway, Oslo, Norway <sup>9</sup>Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden <sup>10</sup>Institute of Cancer Epidemiology, Danish Cancer Society, Copenhagen, Denmark <sup>11</sup>Department of Public Health, Aarhus University, Denmark <sup>12</sup>Inserm, Centre for Research in Epidemiology and Population Health, U1018, Institut Gustave Roussy, F-94805, Villejuif, France <sup>13</sup>Paris South University, UMRS 1018, F-94805, Villejuif, France <sup>14</sup>Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany <sup>15</sup>Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany <sup>16</sup>WHO Collaborating Center for Food and Nutrition Policies, Department of Hygiene, Epidemiology and Medical Statistics, University of Athens Medical School, Athens, Greece <sup>17</sup>Hellenic Health Foundation, Athens, Greece <sup>18</sup>Center for Cancer Prevention (CPO-Piemonte) and Human Genetic Foundation (HuGeF), Torino, Italy <sup>19</sup>Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy <sup>20</sup>Molecular and Nutritional Epidemiology Unit, ISPO- Cancer Research and Prevention Institute, Florence Italy <sup>21</sup>Cancer Registry and Histopathology Unit, "Civile - M.P.Arezzo" Hospital, ASP 7 Ragusa, Italy <sup>22</sup>Department of Clinical and Experimental Medicine, Federico II University, Naples, Italy <sup>23</sup>Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, the Netherlands <sup>24</sup>Department of Gastroenterology and Hepatology, University Medical Centre Utrecht (UMCU), Utrecht, The Netherlands <sup>25</sup>Institute of Community Medicine, University of Tromsø, Tromsø, Norway <sup>26</sup>Public Health Institute of Navarra, Pamplona, Spain <sup>27</sup>CIBER Epidemiología y Salud Pública (CIBERESP), Spain <sup>28</sup>Department of Epidemiology, Murcia Regional Health Authority, Murcia, Spain <sup>29</sup>Andalusian School of Public Health, Granada, Spain <sup>30</sup>Public Health Division of Gipuzkoa, Basque Regional Health Department, San Sebastian, Spain <sup>31</sup>Public Health and Participation Directorate, Health and

**Correspondence to:** Dominique Michaud, ScD Department of Epidemiology Brown University Box G-S121-2 Providence, RI 02912  
Dominique\_Michaud@brown.edu.

The authors have no conflicts of interest.

Health Care Services Council, Asturias, Spain <sup>32</sup>Unit of Nutrition, Environment and Cancer, Cancer Epidemiology Research Programme, Catalan Institute of Oncology (ICO-IDIBELL), Barcelona, Spain <sup>33</sup>Department of Clinical Sciences, Division of Gastroenterology, Skåne University Hospital, Malmö, Lund University, Sweden <sup>34</sup>Department of Surgery, Lund University and Skåne University Hospital, Malmö, Sweden <sup>35</sup>School of Clinical Medicine, University of Cambridge, Cambridge, UK <sup>36</sup>MRC Epidemiology Unit, Cambridge, UK <sup>37</sup>Cancer Epidemiology Unit, University of Oxford, Nuffield Department of Clinical Medicine, Oxford, UK <sup>38</sup>International Agency for Research on Cancer (IARC-WHO), Lyon, France <sup>39</sup>National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

## Abstract

**Background**—Increasing evidence suggests that oral microbiota play a pivotal role in chronic diseases, in addition to the well-established role in periodontal disease. Moreover, recent studies suggest that oral bacteria may also be involved in carcinogenesis; periodontal disease has been linked several cancers. In this study, we examined whether lifestyle factors have an impact on antibody levels to oral bacteria.

**Methods**—Data on demographic characteristics, lifestyle factors, and medical conditions were obtained at the time of blood sample collection. For the current analysis, we measured antibody levels to 25 oral bacteria in 395 cancer-free individuals using an immunoblot array. Combined total immunoglobulin G (IgG) levels were obtained by summing concentrations for all oral bacteria measured.

**Results**—IgG antibody levels were substantially lower among current and former smokers (1697 and 1677 ng/mL, respectively) than never smokers (1960 ng/mL; p-trend = 0.01), but did not vary by other factors, including BMI, diabetes, physical activity, or by dietary factors, after adjusting for age, sex, education, country and smoking status. The highest levels of total IgG were found among individuals with low education (2419 ng/mL).

**Conclusions**—Our findings on smoking are consistent with previous studies and support the notion that smokers have a compromised humoral immune response. Moreover, other major factors known to be associated with inflammatory markers, including obesity, were not associated with antibody levels to a large number of oral bacteria.

## Introduction

Within the oral cavity, more than 700 different bacterial species or phylotypes inhabit a complicated ecosystem that remains in dynamic equilibrium<sup>1, 2</sup>. This ecological equilibrium is marked by the stable and diverse composition of the bacterial community and maintains the health of the oral cavity<sup>3, 4</sup>. Disruptions of this balance can promote the growth of opportunistic strains and disrupt the relationship between the microbes and the human host<sup>5</sup>. Increasing evidence indicates that oral microbiota play a pivotal role in chronic diseases; poor oral health (including periodontal disease and tooth loss) has been associated with respiratory disease<sup>6</sup>, cardiovascular diseases<sup>7</sup>, stroke<sup>8, 9</sup>, rheumatoid arthritis<sup>10</sup>, diabetes<sup>11, 12</sup>, head and neck cancer<sup>12, 13</sup>, pancreatic cancer<sup>14, 15</sup>, colon cancer<sup>16, 17</sup>, and orodigestive cancers<sup>18</sup>.

Lifestyle, diet and disease status influence the host in different ways. Studies have consistently shown that smokers have lower immunoglobulin G (IgG) levels to oral bacteria, suggesting that smoking suppresses the humoral immune response<sup>19, 20</sup>. While obesity and diabetes can impact the innate immune response<sup>21, 22</sup>, it is less clear whether these conditions can impact the humoral immune response, and specifically the immune response

to oral bacteria. No study has examined the relationship between dietary factors and IgG levels to oral bacteria in blood. A few studies observed that diet can influence bacteria in the mouth suggesting that it may impact the mounted immune response to these bacteria; for example, coffee and wine have strong anti-bacterial and anti-adhesive properties<sup>23</sup>, cocoa can reduce bacterial polysaccharide production<sup>24</sup>, sugars and acidic foods favor aciduric bacterial species<sup>25</sup>, and yogurt may inhibit periodontal pathogens<sup>26</sup>. Furthermore, certain foods are rich in antioxidants and may influence the immune response by lowering levels of oxidative stress, a significant modifier of immune mechanisms. In one study, high levels of systemic oxidative stress were strongly associated with lower total IgG levels, after controlling for a large number of known cardiovascular risk factors<sup>27</sup>.

Understanding the association between lifestyle factors and antibodies to oral bacteria may provide new insights into the interaction between the environment and the host immune response to pathogens. We measured antibodies to 25 oral bacteria and examined their association with lifestyle factors, including smoking, physical activity, obesity and diabetes, and dietary factors previously linked to oral bacteria, in a population study of cancer-free individuals.

## Materials and Methods

### Study Population

The European Prospective Investigation into Cancer and Nutrition (EPIC) includes 519,978 participants, mostly aged 35–70 years, who were recruited in 23 centers within 10 European countries (Denmark, France, Germany, Greece, Italy, the Netherlands, Norway, Spain, Sweden, and the United Kingdom) between 1992 and 2000. Detailed descriptions of the study design, population, and baseline data collection of the cohort can be found in a previous report<sup>28</sup>. Each participant provided informed consent. The local ethical review committees approved the EPIC cohort study as well as the current project. For this analysis, we included 395 cancer-free control subjects (i.e., no known medical conditions at the time of blood draw and no major chronic diseases diagnosed during follow-up) who had available antibody measurements for 25 oral bacteria (see previous study<sup>15</sup>). In all but three countries (Denmark, Norway and Sweden), blood samples were collected based on a standardized protocol and aliquoted in plastic straws (plasma, serum, erythrocytes, and buffy coat for DNA). The aliquoted specimens were then stored in a central biorepository (IARC, France) in liquid nitrogen (−196°C). In Sweden, all samples were stored locally in freezers at −80°C and in Denmark in nitrogen vapour (−150°C). Due to small numbers (n=2), samples from Norway were excluded from this analysis.

### Antibody detection

Plasma content of IgG antibodies against a panel of 25 bacterial species/strains selected for a prior study examining the relation between periodontal disease and pancreatic cancer<sup>15</sup>. Measurements were conducted using an immunoblot array as described previously<sup>15, 29</sup>. This method has the advantage of using a very small amount of primary sample (less than 10 µl). Briefly, whole-cell formalin fixed bacteria suspensions and protein A from *Staphylococcus aureus* (control) were used as antigens. Plasma samples (diluted 1:125), and serial dilutions of human IgG immunoglobulin (250, 125, 62.5, 30, 15, and 7.5 ng/ml) were applied to the perpendicular lanes of the immunoblotter. Bound IgG was detected by anti-human IgG antibody. Plasma IgG concentrations were estimated by using the human IgG immunoglobulin reference curve. The lower detection limit of the method was 3.0 ng/ml and interassay correlations were previously reported to be good (percent concordance between 0.67 and 0.84<sup>15</sup>).

## Diet and lifestyle questionnaires

Dietary and lifestyle questionnaires were completed by participants at enrolment. Diet was assessed using country-specific, validated questionnaires designed to capture habitual consumption of food over the preceding year<sup>30</sup>. Extensive quantitative dietary questionnaires with up to 260 food items were used in northern Italy, The Netherlands, Germany, Greece, France and Spain. Semi-quantitative food frequency questionnaires were used in Denmark, Norway, Umeå (Sweden), and Naples (Italy). Combined dietary methods of food records and questionnaires were used in the UK and Malmö (Sweden)<sup>28</sup>.

Lifestyle questionnaires included questions on education, occupation, medical history, lifetime history of consumption of tobacco, alcoholic beverages and physical activity<sup>28</sup>. Height and weight were measured directly in most centres at recruitment; self-reported height and weight were obtained in France and at one centre in the United Kingdom. Body mass index (BMI) was calculated as kg/m<sup>2</sup>. Waist circumference was measured either at the narrowest torso circumference (Italy; Cambridge, United Kingdom; and Utrecht, the Netherlands) or at the midpoint between the lower ribs and iliac crest (Bilthoven, the Netherlands; Potsdam, Germany; Malmö, Sweden; and Oxford, United Kingdom). In Spain, Greece, Denmark, and Heidelberg, Germany, a combination of methods was used, although the majority of participants were measured at the narrowest circumference. Hip circumference was measured at the widest circumference (Italy; Spain; Bilthoven, the Netherlands; Greece; and Malmö, Sweden) or over the buttocks (United Kingdom; Utrecht, the Netherlands; Germany; and Denmark).

## Statistical Analysis

Total IgG level was estimated by summing together the antibody concentration to 25 oral bacteria. A transformation (square root) was conducted to obtain a normal distribution of this variable. We used generalized linear models (GLM) to estimate marginal means (LS Means) and p-values are based on the Type III *F*-test; major demographic, lifestyle and dietary factors were modelled simultaneously to examine the independent contribution of each. We created quartiles for dietary variables and for the waist-to-hip ratio. Mean values were squared to obtain interpretable measures (total IgG in ng/ml). We used a generalized linear model with binomial distribution (PROC GENMOD, LOGIT LINK) to obtain proportions of individuals with elevated antibodies (defined as >200ng/ml; >50 ng/ml; or >7.5 ng/ml, depending on prevalence of values in the higher categories) within different strata of smoking (never, former, current) controlling for age, sex, country of origin and education. This analysis was repeated for education, controlling for smoking status. *P*-values were based on tests for trend (ordered categorical variables modelled as continuous). All statistical analyses were conducted using the Statistical Analysis System (SAS) software package, Version 9.2 (SAS Institute Inc., Cary, North Carolina, USA).

## Results

In this cancer-free population, we observed a statistically significant association between smoking status and total IgG ( $p = 0.01$ ); never smokers had higher total IgG levels than former and current smokers (Table 1) after adjusting for potential confounding variables (i.e., sex, age, country and education). Males had higher antibody level than females ( $p=0.02$ ), and an inverse relationship was observed between education and total IgG levels ( $p=0.003$ ). No statistically significant association was observed for total IgG levels and age, BMI, waist-to-hip ratio, diabetes, hypertension, hyperlipidemia, or physical activity (Table 1). Mean total IgG levels were higher among subjects with low education who were never smokers (2536 ng/ml), compared with subjects with low education who were ever smokers

(2056 ng/ml); the difference between smokers and nonsmokers was similar across educational level (data not shown).

Three of the five oral bacterial pathogens previously associated with periodontal disease were significantly associated with smoking status (*P. gingivalis* ATCC 33277,  $p = 0.03$ ; *Aggregatibacter actinomycetemcomitans* ATCC 29523,  $p=0.02$ ; *Tannerella forsythia*,  $p=0.003$ ; Table 2) after adjusting for potential confounders; antibody levels to these oral pathogens were lower in smokers than in never smokers. For *A. actinomycetemcomitans* ATCC 43718, an inverse association was observed between IgG levels and smoking status, but the association did not reach statistical significance ( $p=0.17$ ). Half of the antibodies to Gram-negative oral bacterial species measured in this study were associated with smoking status (in addition to the pathogenic strains: *Fusobacterium polymorphum*,  $p = 0.004$ ; *Prevotella intermedia*,  $p=0.02$ ; *Prevotella melaninogenica*,  $p=0.01$ ; *Prevotella nigrescens*,  $p < 0.001$ ). None of the antibodies to the Gram-positive bacteria were associated with smoking status after adjusting for potential confounders (Table 2).

After adjusting for potential confounders, including smoking, statistically significant associations were observed between education and IgG levels, such that having lower education was associated with higher levels of antibodies to *P. gingivalis* ATCC 33277, *P. intermedia*, *P. melaninogenica*, and *P. nigrescens* (Supplemental Table).

Quartiles of consumption of vegetables, fruits, vitamin C, dairy products, alcohol, and coffee were evaluated in relation to total IgG levels (Table 3); no associations were noted for these dietary factors. Additionally, total IgG levels for the consumption of tea, wine, yogurt, sugar and meats were examined separately, but no associations were observed.

Total IgG levels were not associated with BMI in this population. However, given that *T. forsythia* levels were reported to be significantly higher in subgingival biofilm of obese individuals in a recent study<sup>31</sup>, we examined the association between these antibodies and BMI separately. While the association was not statistically significant ( $p\text{-value} = 0.18$ ), overweight subjects had more elevated antibodies to *T. forsythia* ( $>50$  ng/ml; 17.9%) compared to those with BMI $<25$  kg/m<sup>2</sup> (12.8%), adjusting for potential confounders.

## Discussion

In this large cross-sectional analysis of cancer-free individuals living throughout Europe, we observed lower overall antibody levels to 25 oral bacteria in current smokers compared to never smokers, women compared to men, and individuals with higher education compared to lower education. Antibody levels associated with smoking status included antibodies to a number of periodontal pathogens and other Gram-negative bacteria. Antibody levels associated with education were similarly associated with Gram-negative bacteria. No differences in overall antibody levels were noted by age, obesity, diabetes, physical activity, and a number of dietary factors, including fruit and vegetable consumption.

An association between oral bacteria and systemic antibody response is well-established in individuals who have different degrees of periodontal disease; serum antibody levels to oral pathogens, such as *P. gingivalis*, correlate with severity of periodontal disease<sup>32</sup>. However, in the general population, the association between serum antibody levels to oral pathogenic bacteria and periodontal disease status is inconsistent; other factors, such as smoking status, can influence the immune response (i.e., IgG levels)<sup>19, 20</sup>. Additionally, most studies on antibody response to oral bacteria have been performed on small numbers of patients with periodontitis, which limits interpretation to the general population. To date, two large population studies examined the relation between antibody levels to a number of oral bacteria and periodontal disease status. In a study of 8153 individuals (NHANES III), *P.*

*gingivalis* antibodies were the only antibodies (out of 19 species) that were significantly correlated to periodontal disease status (positive associations measured with two different definitions of periodontitis)<sup>19</sup>. In the other study, conducted in the Atherosclerosis Risk in Communities study (ARIC) and including 4,717 individuals, higher total IgG levels to 17 commensal oral bacteria were observed in those with more severe periodontal disease<sup>27</sup>.

Smoking has been consistently shown to suppress the humoral immune response<sup>20</sup>. Total IgG levels have been observed to be lower among current smokers<sup>33</sup>, and approximately half of IgG levels to specific oral bacteria were also lower in current smokers in a large population based study<sup>19</sup>. In the present study, we also observed that about half of IgG levels to oral bacteria were significantly lower among current smokers.

Dietary intake has been associated with oral bacteria measured directly in saliva and plaques of healthy subjects<sup>23</sup>. Individuals consuming large quantities of coffee, tea or wine, had lower levels of pathogenic bacteria in saliva and plaque than those drinking primarily water<sup>23</sup>; however, the same study made no statistical adjustment for potential confounders, including smoking status. Studies on diet and periodontal disease suggest that whole grains<sup>34</sup> may lower risk of periodontal disease, but overall findings with diet are inconsistent<sup>35</sup>. Dietary factors in our study were not associated with total antibody levels or with antibodies to pathogenic bacteria.

The relation between oral bacteria and obesity was previously examined in one study. Levels of *T. forsythia* measured in subgingival biofilm of subjects with periodontally healthy gums, or exhibiting gingivitis (n=120), were significantly higher among those who were obese (r=0.43, p<0.001), but other bacteria were not associated with obesity<sup>31</sup>. No association was noted for other pathogenic or commensal bacteria. While we did observe higher levels of antibodies to *T. forsythia* among overweight individuals, compared to normal weight (BMI <25 kg/m<sup>2</sup>), the association was not statistically significant. Unlike smoking, total antibody IgG levels were not associated with BMI in our population.

Systemic oxidative stress has been shown to be more strongly associated with total oral bacteria antibodies than periodontal disease status, smoking or other demographic and lifestyle factors in a large population study<sup>27</sup>. This finding highlights the need for more studies on modulators of the immune response to bacterial pathogens. Although we did not have any measure of oxidative stress in our study population to examine this question, intake of foods high in antioxidants, including fruits and vegetables, was not related to antibody levels. More studies are also needed to understand factors that impact oxidative stress markers and the humoral immune response.

The major limitation of our study was the lack of data on periodontal disease status. Another limitation was the large differences in antibody levels by country which were not attributable to major lifestyle and medical factors examined. The differences therefore remain unexplained and may have influenced our results. However, the observed association between antibodies for periodontal pathogens and education, and the findings for smoking status, provide some indirect validation of the assays we conducted, given consistency with prior studies.

Our data are consistent with previous studies demonstrating suppressed immune response among smokers. Although obesity and diabetes can modulate the innate immune response, we observed no association with antibodies to oral bacteria in this study; however, as the number of diabetics in this study was low (n=17) more studies will need to confirm this finding. Similarly, consumption of antioxidant rich foods, or foods known to modulate bacteria colonization in the mouth, were not related to plasma levels of oral bacteria antibodies.



## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## References

1. Zarco MF, Vess TJ, Ginsburg GS. The oral microbiome in health and disease and the potential impact on personalized dental medicine. *Oral Dis.* Mar; 2012 18(2):109–120. [PubMed: 21902769]
2. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, et al. The human oral microbiome. *J Bacteriol.* Oct; 2010 192(19):5002–5017. [PubMed: 20656903]
3. Segata N, Haake SK, Mannon P, Lemon KP, Waldron L, Gevers D, et al. Composition of the adult digestive tract bacterial microbiome based on seven mouth surfaces, tonsils, throat and stool samples. *Genome Biol.* Jun 14.2012 13(6):R42. [PubMed: 22698087]
4. Structure, function and diversity of the healthy human microbiome. *Nature.* Jun 14; 2012 486(7402): 207–214. [PubMed: 22699609]
5. Hajishengallis G, Lamont RJ. Beyond the red complex and into more complexity: the polymicrobial synergy and dysbiosis (PSD) model of periodontal disease etiology. *Mol Oral Microbiol.* Dec; 2012 27(6):409–419. [PubMed: 23134607]
6. Awano S, Ansai T, Takata Y, Soh I, Akifusa S, Hamasaki T, et al. Oral health and mortality risk from pneumonia in the elderly. *J Dent Res.* Apr; 2008 87(4):334–339. [PubMed: 18362314]
7. Blaizot A, Vergnes JN, Nuwwareh S, Amar J, Sixou M. Periodontal diseases and cardiovascular events: meta-analysis of observational studies. *Int Dent J.* Aug; 2009 59(4):197–209. [PubMed: 19774803]
8. Sfyroeras GS, Roussas N, Saleptsis VG, Argyriou C, Giannoukas AD. Association between periodontal disease and stroke. *J Vasc Surg.* Apr; 2012 55(4):1178–1184. [PubMed: 22244863]
9. Joshipura KJ, Hung HC, Rimm EB, Willett WC, Ascherio A. Periodontal disease, tooth loss, and incidence of ischemic stroke. *Stroke.* Jan; 2003 34(1):47–52. [PubMed: 12511749]
10. Chen HH, Huang N, Chen YM, Chen TJ, Chou P, Lee YL, et al. Association between a history of periodontitis and the risk of rheumatoid arthritis: a nationwide, population-based, case-control study. *Ann Rheum Dis.* Aug 31.2012
11. Taylor GW, Burt BA, Becker MP, Genco RJ, Shlossman M, Knowler WC, et al. Severe periodontitis and risk for poor glycemic control in patients with non-insulin-dependent diabetes mellitus. *J Periodontol.* Oct; 1996 67(10 Suppl):1085–1093. [PubMed: 8910827]
12. Tezal M, Sullivan MA, Hyland A, Marshall JR, Stoler D, Reid ME, et al. Chronic periodontitis and the incidence of head and neck squamous cell carcinoma. *Cancer Epidemiol Biomarkers Prev.* Sep; 2009 18(9):2406–2412. [PubMed: 19745222]
13. Meyer MS, Joshipura K, Giovannucci E, Michaud DS. A review of the relationship between tooth loss, periodontal disease, and cancer. *Cancer Causes Control.* May 14.2008
14. Michaud DS, Joshipura K, Giovannucci E, Fuchs CS. A prospective study of periodontal disease and pancreatic cancer in US male health professionals. *J Natl Cancer Inst.* Jan 17; 2007 99(2):171–175. [PubMed: 17228001]
15. Michaud DS, Izard J, Wilhelm-Benartzi CS, You DH, Grote VA, Tjonneland A, et al. Plasma antibodies to oral bacteria and risk of pancreatic cancer in a large European prospective cohort study. *Gut.* Sep 18.2012
16. Kostic AD, Gevers D, Pedamallu CS, Michaud M, Duke F, Earl AM, et al. Genomic analysis identifies association of *Fusobacterium* with colorectal carcinoma. *Genome Res.* Feb; 2012 22(2): 292–298. [PubMed: 22009990]
17. Castellarin M, Warren RL, Freeman JD, Dreolini L, Krzywinski M, Strauss J, et al. *Fusobacterium nucleatum* infection is prevalent in human colorectal carcinoma. *Genome Res.* Feb; 2012 22(2): 299–306. [PubMed: 22009989]
18. Ahn J, Segers S, Hayes RB. Periodontal disease, *Porphyromonas gingivalis* serum antibody levels and orodigestive cancer mortality. *Carcinogenesis.* May; 2012 33(5):1055–1058. [PubMed: 22367402]

19. Vlachojannis C, Dye BA, Herrera-Abreu M, Pikdoken L, Lerche-Sehm J, Pretzl B, et al. Determinants of serum IgG responses to periodontal bacteria in a nationally representative sample of US adults. *J Clin Periodontol*. Aug 1; 2010 37(8):685–696. [PubMed: 20561113]
20. Barbour SE, Nakashima K, Zhang JB, Tangada S, Hahn CL, Schenkein HA, et al. Tobacco and smoking: environmental factors that modify the host response (immune system) and have an impact on periodontal health. *Crit Rev Oral Biol Med*. 1997; 8(4):437–460. [PubMed: 9391754]
21. Milner JJ, Beck MA. The impact of obesity on the immune response to infection. *Proc Nutr Soc*. May; 2012 71(2):298–306. [PubMed: 22414338]
22. Knapp S. Diabetes and Infection: Is There a Link? - A Mini-Review. *Gerontology*. Nov 24.2012
23. Signoretto C, Burlacchini G, Bianchi F, Cavalleri G, Canepari P. Differences in microbiological composition of saliva and dental plaque in subjects with different drinking habits. *New Microbiol*. Oct; 2006 29(4):293–302. [PubMed: 17201096]
24. Paolino VJ, Kashket S. Inhibition by cocoa extracts of biosynthesis of extracellular polysaccharide by human oral bacteria. *Arch Oral Biol*. 1985; 30(4):359–363. [PubMed: 3857906]
25. Bradshaw DJ, Marsh PD. Analysis of pH-driven disruption of oral microbial communities in vitro. *Caries Res*. 1998; 32(6):456–462. [PubMed: 9745120]
26. Zhu Y, Xiao L, Shen D, Hao Y. Competition between yogurt probiotics and periodontal pathogens in vitro. *Acta Odontol Scand*. Sep; 2010 68(5):261–268. [PubMed: 20491536]
27. Singer RE, Moss K, Beck JD, Offenbacher S. Association of systemic oxidative stress with suppressed serum IgG to commensal oral biofilm and modulation by periodontal infection. *Antioxid Redox Signal*. Dec; 2009 11(12):2973–2983. [PubMed: 19496702]
28. Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, Fahey M, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr*. Dec; 2002 5(6B):1113–1124. [PubMed: 12639222]
29. Sakellari D, Socransky SS, Dibart S, Eftimiadi C, Taubman MA. Estimation of serum antibody to subgingival species using checkerboard immunoblotting. *Oral Microbiol Immunol*. Oct; 1997 12(5):303–310. [PubMed: 9467384]
30. Kaaks R, Slimani N, Riboli E. Pilot phase studies on the accuracy of dietary intake measurements in the EPIC project: overall evaluation of results. *European Prospective Investigation into Cancer and Nutrition*. *Int J Epidemiol*. 1997; 26(Suppl 1):S26–36. [PubMed: 9126531]
31. Haffajee AD, Socransky SS. Relation of body mass index, periodontitis and *Tannerella forsythia*. *J Clin Periodontol*. Feb; 2009 36(2):89–99. [PubMed: 19207883]
32. Hayman L, Steffen MJ, Stevens J, Badger E, Temprow P, Fuller B, et al. Smoking and periodontal disease: discrimination of antibody responses to pathogenic and commensal oral bacteria. *Clin Exp Immunol*. Apr; 2011 164(1):118–126. [PubMed: 21303363]
33. Quinn SM, Zhang JB, Gunsolley JC, Schenkein JG, Schenkein HA, Tew JG. Influence of smoking and race on immunoglobulin G subclass concentrations in early-onset periodontitis patients. *Infect Immun*. Jul; 1996 64(7):2500–2505. [PubMed: 8698472]
34. Merchant AT, Pitiphat W, Franz M, Joshupura KJ. Whole-grain and fiber intakes and periodontitis risk in men. *Am J Clin Nutr*. Jun; 2006 83(6):1395–1400. [PubMed: 16762952]
35. Kaye EK. Nutrition, dietary guidelines and optimal periodontal health. *Periodontol* 2000. Feb; 2012 58(1):93–111. [PubMed: 22133369]

**Table 1**

Total antibody measured to 25 oral bacteria in apparently healthy individuals in the EPIC sub-cohort (n=395) in relation to baseline characteristics and lifestyle factors.

	Numbers (%)	Total IgG (ng/mL)*	P-values**
<b>Sex</b>			0.02
Males	187 (47)	1892	
Females	208 (53)	1663	
<b>Age (years)</b>			0.79
<50	53 (13)	1716	
50–59	166 (42)	1801	
60	176 (45)	1811	
<b>Smoking status</b>			0.01
Never	174 (44)	1960	
Former	134 (34)	1677	
Current	87 (22)	1697	
<b>BMI (kg/m<sup>2</sup>)</b>			0.41
<25	171 (43)	1693	
25–<30	162 (41)	1809	
30	62 (16)	1827	
<b>Waist-to-hip ratio <sup>§</sup></b>			0.61
Quartile 1	83 (23)	1863	
Quartile 2	97 (27)	1734	
Quartile 3	91 (25)	1811	
Quartile 4	90 (25)	1688	
<b>Education</b>			0.003
None	17 (4)	2419	
Primary school completed	147 (37)	1662	
Technical/professional school	104 (26)	1839	
Secondary school	54 (14)	1520	18
Higher education	73 (19)	1511	
<b>Country</b>			<0.001
Denmark	82 (21)	1020	
France	15 (4)	2314	
Germany	52 (13)	1803	
Greece	22 (6)	1972	
Italy	38 (10)	2009	
The Netherlands	38 (10)	1548	
Spain	28 (7)	2095	
Sweden	82 (21)	1529	

	Numbers (%)	Total IgG (ng/mL)*	P-values**
United Kingdom	38 (10)	1876	
<b>Physical activity</b> #			0.90
Inactive	106 (30)	1686	
Moderately inactive	123 (34)	1761	
Moderately active	66 (18)	1781	
Active	65 (18)	1784	
<b>Diabetes</b> §			0.20
No	362 (96)	1799	
Yes	17 (4)	1525	
<b>Hypertension</b> §			0.94
No	205 (70)	1829	
Yes	80 (27)	1836	
<b>Hyperlipidemia</b> §			0.72
No	183 (65)	1823	
Yes	66 (35)	1739	

\* Sum of IgG measured for 25 oral bacteria.

\*\* *p*-values mutually adjusted for other variables in this table (physical activity, waist-to-hip ratio, and diabetes, hypertension and hyperlipidemia not adjusted for to maintain sample size of 395).

§ Quartiles: males, <0.90, 0.90 to <0.93, 0.94 to <1.0, and 1.0; females, <0.76, 0.76 to <0.81, 0.79 to <0.85, and 0.85; 36 with missing data for waist-to-hip ratio.

# Among 360 individuals with data on physical activity.

§ Missing data: 16 for diabetes; 110 for hypertension; 146 for hyperlipidemia.

Table 2

Percent individuals with elevated antibody levels to oral bacteria by smoking status \*.

Bacteria	% with high IgG level <sup>§</sup>			P-values
	Never smokers	Past smokers	Current smokers	
<b>Oral periodontal pathogens ** (Gram negative)</b>				
<i>Porphyromonas gingivalis</i> ATCC 33277	26.2	14.6	13.8	0.03
<i>Porphyromonas gingivalis</i> ATCC 53978 <sup>‡</sup>	38.4	30.7	36.7	0.17
<i>Aggregatibacter actinomycetemcomitans</i> ATCC 29523	31.9	22.3	17.2	0.02
<i>Aggregatibacter actinomycetemcomitans</i> ATCC 43718	31.9	26.4	21.8	0.17
<i>Tannerella forsythia</i> ATCC 43037 <sup>‡</sup>	27.2	11.5	12.7	0.003
<b>Oral bacterial species of the human microbiome (Gram negative)</b>				
<i>Fusobacterium nucleatum</i> ATCC 25586 <sup>‡</sup>	37.9	27.0	35.2	0.52
<i>Fusobacterium periodonticum</i> ATCC 33693 <sup>^</sup>	5.5	3.1	4.5	0.39
<i>Fusobacterium polymorphum</i> ATCC 10953 <sup>^</sup>	48.3	34.2	26.5	0.004
<i>Prevotella intermedia</i> ATCC 25611 <sup>‡</sup>	60.3	40.1	46.0	0.02
<i>Prevotella melaninogenica</i> ATCC 25845 <sup>‡</sup>	73.7	62.1	57.3	0.01
<i>Prevotella nigrescens</i> ATCC 33563 <sup>‡</sup>	81.3	70.8	59.4	<0.001
<i>Veillonella atypica</i> ATCC 17744 <sup>‡</sup>	9.8	6.9	8.1	0.55
<i>Veillonella parvula</i> ATCC 10790 <sup>‡</sup>	23.1	17.5	20.2	0.49
<b>Oral bacterial species of the human microbiome (Gram positive)</b>				
<i>Bifidobacterium dentium</i> ATCC 27534 <sup>^</sup>	48.8	39.7	40.0	0.20
<i>Corynebacterium matruchotii</i> ATCC 14266 <sup>^</sup>	56.0	47.8	42.9	0.09
<i>Enterococcus faecalis</i> ATCC 29212 <sup>‡</sup>	8.7	7.0	5.7	0.34
<i>Parvimonas micra</i> ATCC 33270	63.2	48.1	63.5	0.69
<i>Peptostreptococcus anaerobius</i> ATCC 27337 <sup>^</sup>	7.5	5.4	4.9	0.35

Bacteria	% with high IgG level <sup>§</sup>			P-values
	Never smokers	Past smokers	Current smokers	
<i>Streptococcus intermedius</i> ATCC 27335 <sup>‡</sup>	36.5	28.5	35.4	0.68
<i>Streptococcus mitis</i> ATCC 49456	48.1	45.4	57.3	0.27
<i>Streptococcus salivarius</i> ATCC 7073	10.9	7.7	10.2	0.73

Note: *Captonocytophaga ochracea*; *Eikenella corrodens*; *Actinomyces naeslundii*; *Finegoldia magna* are not listed due to insufficient variation after adjusting for country.

\* Percentages and p-values for trend are adjusted for sex, age, country, and education.

\*\* Oral bacterial species that have been previously associated with periodontal disease.

<sup>§</sup> "High" defined as >200ng/mL, unless otherwise indicated with footnote.

<sup>‡</sup> due to small numbers with antibodies above 200ng/mL, the cutpoint was set at >50 ng/ml

<sup>^</sup> due to small numbers with antibodies above 50ng/mL, the cutpoint was set at >7.5 ng/ml.

**Table 3**

Total antibody to 25 oral bacteria in apparently healthy individuals in the EPIC sub-cohort (n=395) in relation to dietary factors \*

	Total IgG (ng/mL)	P-values
<b>Vegetable intake</b>		0.61
Quartile 1 (lowest)	1888	
Quartile 2	1748	
Quartile 3	1760	
Quartile 4 (highest)	1836	
<b>Fruit intake</b>		0.49
Quartile 1	1869	
Quartile 2	1892	
Quartile 3	1760	
Quartile 4	1837	
<b>Dairy intake</b>		0.59
Quartile 1	1704	
Quartile 2	1794	
Quartile 3	1890	
Quartile 4	1826	
<b>Vitamin C</b>		0.50
Quartile 1	1840	
Quartile 2	1918	
Quartile 3	1739	
Quartile 4	1884	
<b>Coffee intake</b>		0.86
Quartile 1	1834	
Quartile 2	1748	
Quartile 3	1783	
Quartile 4	1847	
<b>Alcohol intake</b>		0.93
Quartile 1	1820	
Quartile 2	1788	
Quartile 3	1761	
Quartile 4	1734	

\* Means and p-values for trend are adjusted for sex, age, country, education, BMI and smoking status.