

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

8-28-2019

Mitogenome and Nuclear-encoded Fungicide-target Genes of Thecaphora frezii - Causal Agent of Peanut Smut

R. S. Arias

USDA-ARS National Peanut Research Laboratory (NPRL), renee.arias@ars.usda.gov

Luis I. Cazon

CIAP-INTA

Alicia N. Massa

USDA-ARS National Peanut Research Laboratory (NPRL), alicia.massa@usda.gov

Brian E. Scheffler

USDA-ARS Genomics and Bioinformatics Research Unit

Victor S. Sobolev

USDA-ARS National Peanut Research Laboratory (NPRL)

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

Arias, R. S.; Cazon, Luis I.; Massa, Alicia N.; Scheffler, Brian E.; Sobolev, Victor S.; Lamb, Marshall C.; Duke, Mary V.; Simpson, Sheron A.; Conforto, Cinthia; Paredes, Juan A.; Buteler, Mario I.; and Rago, Alejandro M., "Mitogenome and Nuclear-encoded Fungicide-target Genes of Thecaphora frezii - Causal Agent of Peanut Smut" (2019). *Publications from USDA-ARS / UNL Faculty*. 2198.

<https://digitalcommons.unl.edu/usdaarsfacpub/2198>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

R. S. Arias, Luis I. Cazon, Alicia N. Massa, Brian E. Scheffler, Victor S. Sobolev, Marshall C. Lamb, Mary V. Duke, Sheron A. Simpson, Cinthia Conforto, Juan A. Paredes, Mario I. Buteler, and Alejandro M. Rago

Mitogenome and Nuclear-encoded Fungicide-target Genes of *Thecaphora frezii* - Causal Agent of Peanut Smut

Renee S. Arias^{1*}, Luis I. Cazon², Alicia N. Massa¹, Brian E. Scheffler³, Victor S. Sobolev¹, Marshall C. Lamb¹, Mary V. Duke³, Sheron A. Simpson³, Cinthia Conforto², Juan A. Paredes², Juan H. Soave⁴, Mario I. Buteler⁴ and Alejandro M. Rago²

¹USDA-ARS National Peanut Research Laboratory (NPRL), Dawson, Georgia, USA; ²Instituto de Patologia Vegetal, CIAP-INTA, Cordoba, Argentina; ³USDA-ARS Genomics and Bioinformatics Research Unit, Stoneville, Mississippi, U.S.A.; ⁴El Carmen S.A., General Cabrera, Cordoba, Argentina

ABSTRACT

Background: *Thecaphora frezii* Carranza and Lindquist causes smut disease in peanut (*Arachis hypogaea* L.) resulting in up to 35% yield losses. Fungicides have shown ineffective in controlling the disease; whereas research on the molecular basis of that fungicide resistance has been hindered because of the lack of genetic information about *T. frezii*. The goal of this work was to provide molecular information about fungicide-target loci in *T. frezii*, including its mitochondrial genome (mitogenome) and critical nuclear-encoded genes.

Results: Here we report the complete annotated mitogenome of *T. frezii*, a 123,773 bp molecule containing the standard 14 genes that form part of mitochondrial complexes I, III, IV and V, 22 transfer RNAs, small and large subunits of ribosomal RNA, DNA polymerase, ribonuclease P, GII-reverse transcriptase/maturase, nine hypothetical open-reading frames and homing endonucleases (LAGLIDADG, GIY-YIG, HEG). In addition, we report the full-length cDNA sequence of *T. frezii* cytochrome *b* (*cob*) and cytochrome oxidase 1 (*cox1*) genes; as well as partial sequences of *T. frezii* succinate dehydrogenase (*sdhb*), ergosterol biosynthesis (*Erg4*), cytochrome P450 (*cyp51*), and beta tubulin (β -tubulin) genes, which are respective targets of strobilurins, quinone oxidation inhibitors, triazoles and beta-tubulin inhibitor fungicides commonly used in the peanut crop. Translation of *cob* and *sdhb* genes in this particular *T. frezii* isolate suggests potential resistance to strobilurin and carboxamide fungicides.

Conclusion: The mitogenome and nuclear-encoded gene sequences presented here provide the molecular tools to research *T. frezii* fungicide-target loci.

Keywords: Mitochondrial genome; Pathogens; Groundnut; Carbon del mani; Fungicides

INTRODUCTION

Currently endemic to Argentina, peanut smut is an emerging disease that constitutes a threat to the peanut industry around the world [1]. The etiological agent is the Basidiomycete fungus *T. frezii*, originally described on wild peanuts in Brazil in 1962 [1] and detected in the peanut production area of Argentina in the early 1990s [2]. Argentina is the seventh peanut producer in the world but is the first exporter in the peanut market [3]. *T. frezii* invades peanuts when the pegs touch the ground [4], and upon infection, seeds are replaced by a dark-brown powder of fungal teliospores (Figure 1) [5,6]. Without proper crop rotations *T. frezii* teliospores accumulate in soil, resulting in up to 51% of disease incidence [7] and up to 35% losses [8]. Much concern has spread throughout the peanut community, thus, in May 2017 the Australian peanut growers asked for a ban of peanut imports from Argentina [9], and

in October 2017, the United States Department of Agriculture (USDA) - Animal and Plant Health Inspection Service (APHIS) restricted imports of raw peanut from Argentina and Brazil due to peanut smut [10]. Though peanut smut disease has not been reported in the United States, trade, travel and climate change may increase the risk of the disease being introduced and spread. Currently, United States is the fourth largest producer of peanut in the world (2.6 Mt) and Argentina is the eighth [11].

Mitochondria in fungi participate in virulence, pathogenicity, survival and drug resistance, but they are essentially involved in energy metabolism [12]. Consequently, fungicides commonly target the energy-production protein complexes located in the inner membrane of the mitochondria (Figure 2) [13]. For example, diflumetorim (Pyricut) from the group of NADH inhibitors is a fungicide that targets complex I by inhibiting the NADH oxidoreductase activity

Correspondence to: Renee S. Arias, USDA-ARS National Peanut Research Laboratory (NPRL), Dawson, Georgia, USA, Telephone: +1 229-995-4441; E-mail: renee.arias@ars.usda.gov

Received: June 18, 2019; **Accepted:** August 21, 2019; **Published:** August 28, 2019

Citation: Arias RS, Cazon LI, Massa AN, Scheffler BE, Sobolev VS, Lamb MC, et al. (2019) Mitogenome and nuclear-encoded fungicide-target genes of *Thecaphora frezii* causal agent of peanut smut. *Fungal Genom Bio* 9: 160. doi: 10.35248/2165-8056.19.9.160

Copyright: © 2019 Arias RS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Figure 1: Symptoms caused by peanut smut, *Thecaphora frezii*, the dark powder is a mass of fungal teliospores developed inside the peanut pods after infection. Left: peanut pods still attached to the plant, Right: close view of infection.




Figure 2: Schematic representation of a mitochondrion and a close view of the five complexes I, III, II, IV and V, in the order described by Kuhlbrandt, 2015 [13] that constitute the electron transport chain (ETC). In the matrix, nutrients are metabolized and their sub-products converted to energy in the form of adenosine triphosphate (ATP) through the ETC. The mitochondrial DNA is located in the matrix, and it contributes gene products to complexes I, III, IV and V. Many fungicides target complexes I, II, III and V to prevent respiration in fungal pathogens of plants and humans.

[14]. Carboxamides (succinate dehydrogenase inhibitors, SDHI group) target complex II [15]. Strobilurins (Quinone oxidation inhibitors, QoI) target complex III, by inhibiting the electron transfer in the cytochrome bc1 and paralyzing ATP production [16]. Lastly, complex V is targeted by fentin hydroxide, or triphenyltin hydroxide [17], listed by the Fungicide Resistance Action Committee (FRAC) as an inhibitor of the oxidative phosphorylation (ATP synthase). Carboxamides and strobilurins are the most common fungicides used in the peanut crop along with triazoles and phtalonitriles, [1,18-20] (Supplementary Table 1).

Fungicide resistance frequently results from mutations in mitochondrial genes [21-23] and other fungicide-target loci [24-27]. To better plan strategies to control peanut smut it is essential to know the genetic makeup of *T. frezii* at those loci, but information about this species had been very limited. Until May 2019, there were merely 10 entries (607 to 722 bp long) for *T. frezii* in GenBank (NCBI database) corresponding to ribosomal RNA sequences and internal transcribed spacer (ITS) of *T. frezii* [28,29]. The objective of this work was to generate basic genetic information and molecular tools to investigate *T. frezii* and its response to fungicides. We report the complete annotated mitochondrial genome of *T. frezii*, as well as cDNA sequences of nuclear-encoded genes that are target of fungicides used in the peanut crop.

Table 1: Primers used to amplify *Thecaphora frezii* genes that are target of fungicides, plus primers T3 and T7 that were added to their 5' ends (forward and reverse, respectively).

TF_CYTB_Fw1	GGAAATAATATGCGACTACTTAAATCACATCC
TF_CYTB_Rv1	TACAGCTGTTTCGTAGTAGAGGTGCTGTAGG
TF_COX1_Fw1	ATGACTCGATGGCTTTTTTCAACAAACGC
TF_COX1_Rv1	TAAGATTGACGTTAAAGCTACCCTCACTGGTG
TF_SDH_Fw	ATGACGATGTCGCTCTTCGCCGGCAACG
TF_SDH_Rv	ACGATCGCCAGCACCCCTCACAGCTAGC
TF_Beta_Fw	CGCCGGTAACTACGAGGGCAGC
TF_Beta_Rv	TGGCGTATAGCTTCTTCAGG
TF_CYP51_Fw	CACCACCCTCACCCCTACACGC
TF_CYP51_Rv	AGGACCACCATGGAAATCCGGC
TF_Erg4_Fw	CACTTTGAGTTCTGACTGACCGC
TF_Erg4_Rv	CTGACGATGCACCAGCAGGATCGC
T3	ATTAACCCTCACTAAAGGGA
T7	TAATACGACTCACTATAGG

MATERIALS AND METHODS

DNA extraction and sequencing

Peanut pods containing kernels with teliospores were collected from General Deheza (32° 45' 08.54"S, 63° 46' 06.5"W), Cordoba,

Argentina, in 2016. Pods were surface disinfected in 0.26% NaOCl with continued agitation for 15 min, then 200 mg of teliospores were surface disinfected in 0.63% NaOCl for 5 min, rinsed twice with sterile distilled water and resuspended in 2 mL sterile distilled water [30]. Disinfected teliospores were plated on potato-dextrose agar (PDA) (Sigma-Aldrich, St. Louis, MO) supplemented with the extract of 50 g·L⁻¹ peanut-seeds and incubated in the dark at 25°C [31]. DNA was extracted from the mycelial growth using the CTAB method [32], and identified as *T. frezii* by 18S ribosomal RNA sequencing [28]. DNA of an isolate identified as “A” was sent to the USDA-ARS-National Peanut Research Laboratory (NPRL) (Dawson, GA, U.S.A.), and the isolate was stored in the culture collection of the Instituto de Patología Vegetal (IPAVE) at the Instituto Nacional de Tecnología Agropecuaria (INTA), Córdoba, Argentina. Genomic DNA of isolate A was sequenced using Illumina HiSeq 2500 at LC_Sciences (Houston, TX) and at the USDA-ARS-Genomics and Bioinformatics Research Unit (GBRU) (Stoneville, MS). Reads from those runs were trimmed and assembled de novo at the NPRL using CLC_Bio Genomics Workbench 12, Qiagen (Redwood City, CA). No contaminating sequences were detected. A contig identified by BLAST analysis [33] as mitochondrial DNA was preliminary annotated with MAKER [34,35]. Additional open reading frames (ORFs) were predicted using Clone Manager Professional 9 (Sci-Ed Software, Denver, CO), and a map was built using Seqbuilder followed by GenVision in DNASTAR Lasergene 11 Core Suite (DNASTAR, Inc. Madison, WI). ORFs and hypothetical protein genes were translated by Clone Manager software and protein sequences subjected to search of motifs in Pfam and PROSITE databases using MOTIF (Kyoto University Bioinformatics Center, <https://www.genome.jp/tools/motif/>) to further characterize both predicted and intron encoded proteins. Mycelium preserved in RNAlater solution (Ambion, Life Technologies, Austin, TX) was used to extract RNA from the same *T. frezii* isolate using Direct-zol RNA MiniPrep Plus (Zymo Research, Irvine, CA), and two cDNAs were synthesized, one with oligo dT and the other with oligo DT+random hexamers using Superscript III reverse transcriptase (Invitrogen, Life Technologies, Carlsbad, CA).

Gene amplification and annotation

To validate the annotation of *T. frezii* mitogenome, primers were designed to amplify transcripts of *cytochrome b* (*cob*) and *cytochrome c oxidase 1* (*cox1*); both genes were PCR amplified from *T. frezii* cDNA libraries by Platinum Taq DNA Polymerase High Fidelity (Invitrogen, Life Technologies, CA), using the following PCR conditions for amplification: 94°C 2 min; then 36 cycles of 94°C 30 s, 60°C 30 s, 68°C 1 min; then 68°C 5 min and kept at 4°C. The sequences of the universal forward primers of the T3 and T7 bacteriophages (Table 1) were added to the forward and reverse primers of the specific genes to increase the length of quality-sequencing obtained. PCR products were sequenced on SimpleSeq reactions at Eurofins Genomics LLC (Louisville, KY), the sequences were assembled using SeqMan Pro in DNASTAR Lasergene 11 Core Suite (DNASTAR, Inc. Madison, WI). Exon and intron annotation of *cox1* and *cob* were revised according to the results, their corresponding proteins were translated using Clone Manager Professional 9 (Sci-Ed Software, Denver, CO) with the genetic code of yeast mitogenome [36].

Nuclear-encoded fungicide-target genes of *T. frezii*

We performed several iterations of mapping and de novo assembly

and the final assemblies were screened by BLASTn in NCBI. Primers were designed to amplify nearly the full length of each gene from cDNAs of *T. frezii*. Gene coding regions were PCR amplified by the same conditions described for genes *cob* and *cox1*, using as template cDNA and DNA from *T. frezii*. Universal primers T3 and T7 were added to the 5' end of forward and reverse primers, respectively, to increase the length of quality-sequencing obtained as indicated for *cob* and *cox1* genes (Table 1). Putative *T. frezii succinate dehydrogenase B subunit* (*sdhb*), a *lanosterol 14β-demethylase cytochrome P450 monooxygenase 51* (*cyp51*), *ergosterol biosynthesis gene 4* (*Erg4*) and *β-tubulin* partial sequences were obtained by the described iterations of mapping reads, de novo assembly in CLC_Bio Workbench 11, and BLAST analysis. These sequences were submitted to the NCBI Database.

RESULTS

Thecaphora frezii mitogenome summary

T. frezii genomic DNA was sequenced. The total number of reads in two Illumina sequencing runs after trimming was 8,442,832 and 53,186,795, respectively, with an average length per read of 165 bp. After de novo assembly, one contig of 123,895 bp with a 10,664 x average coverage was identified as mitochondrial DNA by BLAST analysis. A 122 bp overlap of 5' and 3' ends allowed circularization into a 123,773 bp mitochondrial DNA molecule that constitutes the *T. frezii* mitogenome. The sequence was submitted to GenBank and has Accession number MH392473.

The *T. frezii* mitogenome contained genes identified as: seven units of nicotinamide-adenine dinucleotide reduced-form (NADH) reductase genes (*nad*) (1, 2, 3, 4, 4L, 5, 6); small ribosomal protein 3 (rpS3); partial large subunit of ribosomal RNA (rRNA); small subunit ribosomal RNA (rns); Ribonuclease P (EC 3.1.26.5) or ribozyme (rnpB); three units of ATP synthase subunit genes: *atp6*, *atp8* and *atp9*; a putative DNA Polymerase (DNAPol); three hypothetical proteins; another 7 open reading frames (ORFs); plus IEND, HEG, GIY genes, all represented in dark-blue arrows in Figure 3. Two ORFs (ORF255 and ORF186) were translated, and searches using MOTIF identified ORF 255 [70 amino acids (aa), pI: 10.13, MW: 7,592] as the N-terminal domain of a reverse transcriptase/maturase of group II (Expected value 5.4×10^{-16}). ORF186 (186aa, pI:10.82, MW: 21,880) Group II intron maturase specific domain (E-value: 6×10^{-16}), thus they were annotated as GII_MAT in the diagram. The mitogenome also had 23 transfer RNA (tRNA) that included all 20 amino acids plus two copies of transfer RNAs for methionine *trnM* (cat), arginine *trnR* (tcg, tct) and serine *trnS* (tga, gct), indicated as dark-red arrows in Figure 3. There were three subunits of cytochrome oxidase (*cox*), *cox1*, *cox2*, and *cox3*; only *cox1* presented introns (light-green lines in Figure 3) for which MAKER predicted 14 introns and one more intron was identified by PCR amplification of cDNA and Sanger sequencing to a total of 15 introns. Also encoded in the *T. frezii* mitogenome was the *cytochrome oxidase B* gene (*cob*) (8,477 bp), which contained five introns, this was confirmed by PCR amplification and sequencing of cDNA. A total of 34,055 bp of the 123,773 bp *T. frezii* mitogenome (27.5%) corresponded to introns. In the reported orientation, *T. frezii* mitogenome had a 67.5% (A+T) and a 32.5% (G+C) content, color coded red/yellow for AT and blue/green for GC in Figure 3.

The probable direction of replication of the molecule is indicated by two light-blue color arrows in Figure 3, and its origin was used as




Figure 3: Mitochondrial genome of *Thecaphora frezii*, outer scale is in base pairs (bp).




Figure 4: Annotation of intron-encoded endonucleases (LAGLIDADG, GIY-YIG) in the mitochondrial genes *cytochrome oxidase subunit 1 (cox1)* and *cytochrome b (cob)* of *Thecaphora frezii*. Introns are represented by light-green color rectangles. Exons and genes are represented as blue arrows. Also encoded in the first intron of *cox1*, and in frame with this gene, is a reverse-transcriptase RNA polymerase. Scales represent number of base pairs (bp).

the start of the nucleotide sequence of the mitogenome. In the *T. frezii* mitogenome, we observed 19 type I introns, 3 type II introns, 6 proteins identified as endonucleases (LAGLIDADG, GIY-YIG, IEND, HEG). Other endonucleases were encoded within the introns of *cox1* and *cob* genes [6 LAGLIDADG, 1 GIY-YIG, and 1 (RT)-RNA-dependent DNA Polymerase (RT_POL)], for clarity these were not included in the map annotation of Figure 3 but are shown separately in Figure 4. The reverse transcriptase (RT_POL) ORF found in *cox1* was encoded in frame with the first exon of *cox1* (overlapping the first 55 aa) and continuing within the first intron. These 895 aa proteins had a reverse transcriptase RVT_1 domain aa400-652 predicted in Pfam (Expected value (E) 1.9×10^{-33}) and a Type II intron maturase at aa680-803 in Pfam (E 3.8×10^{-32}). Also within *cox1*, Pfam predicted LAGLIDADG ORFs in introns 5 (E 1.5×10^{-18}), 8 (E 2.9×10^{-8}), 10 (E 1.4×10^{-4}) and 14 (E 4.5×10^{-12}), and a GIY-YIG (E 2.4×10^{-5}) motif predicted by Pfam within intron 6 (Figure 4). Two open reading frames identified in Pfam as LAGLIDADG endonucleases were found in *cob* gene within introns 1 (E 7.3×10^{-14}) and 3 (E 1.7×10^{-52}), respectively (Figure 4).

cob and *cox1* genes

Primers used to amplify *cob* and *cox1* genes from cDNAs (oligo dT, oligo dT + random hexamers) are listed in Table 1. The sequences of the universal primers T3 and T7 were added as indicated before (Table 1). The presence of long introns in the *cob* gene of *T. frezii* (Figure 4), made its total length 8,478 bp, though its 1,200 bp transcript encoded only 400 aa. The cDNA sequence of the *cob* gene was translated using the yeast-mitochondrial genetic code [36] reading UGA codons 273, 349 and 351 as tryptophane (W) instead of stop codons, underlined in Figure 5. The translated COB protein had molecular weight MW: 45,989.2, isoelectric point (pI): 8.14. This sequence was submitted to GenBank (MH603949).

The total length of *T. frezii cox1* gene was 22,731 bp, however, the cDNA was only 1,586 bp. Two variants of cytochrome oxidase 1 transcripts of *cox1* gene were observed after PCR amplification of *T. frezii* cDNA, sequencing and translation. These were named TF_COX1_v1 and TF_COX1_v2 in Figure 6, submitted to GenBank as MH603947 and MH603948. Both had 528 aa and pI: 6.39, though

1	MRLKSHPIIL	KIANSF F IDS	Q H PYNISYMW	N FGSLL G ACL	VIQILTVFL
51	AMHYTPSVDL	AFISVEHIMR↓	DVNYGWLIRY	LHANTASFFF	IFVVFHIGRG
101	LYYGSYKSPR	TLVWSIGVVI	LILMMAIA F L	GYVL↓ P Y G QMS	LW G ATV↓ITNL
151	LSAIPWIGQD	FVEFVW↓GGFS	VNNATLNRRF	SLHYLLPFVL	AALVAMH L LA
201	LXENGSSNPL	GVSGNTDRLP	FHPY F T F KDL	VTVFVFLVL	SLFVYFMPNA
251	LGHSDNYIPA	NPMQTPASIV	PE↓ W YLLPFYA	ILRSIPSKLI	GVLAMFMSLL
301	ILLAMPILDT	SRIRGSQFRP	FARFAFWTFV	VDFFILMFIG	SHMLLLHM W R
351	W EQQQMRSTF	MVPGSSSSCW	NIEKFLMDIA	LAEITENLK	HLQLHYEQL

Figure 5: Translation of the mitochondrial cytochrome *b* (*cob*) gene (COB) of *Thecaphora frezii*, highlighted in dark red are the amino acids (aa) where mutations have been reported for other fungi in the literature and known to confer resistance to quinone oxidation inhibitor fungicides. Blue arrows indicate the positions where introns are located in the gene. Underlined tryptophane (W) corresponds to UGA codons translated according to yeast mitochondrial genetic code. Amino acids are in groups of 10, and the number on the left correspond to the order of the first amino acid in each row.




Figure 6: Alignment of two versions of cytochrome oxidase subunit 1 (Cox1) translated proteins found in *Thecaphora frezii*. Blue arrows correspond to helix structures, red arrows are strand structures, green arrows point to potential structural variations that may occur in this protein as a result of amino acid substitutions.

they had slightly different molecular weight (58,080.3 and 58,060.2, respectively). Five aa differences were observed for these two molecules between positions aa400 and aa495, predicting protein structural change according to the model of Garnier-Osguthorpe-Robson (Figure 6) [37]. The five amino acid changes were L400F, P451A, L466P, P490A and S495T. In the predicted model, TF_COX1_v2 has a β -sheet instead of turn at aa490-491; a β -sheet instead of a helix at aa462-466; an extended helix at aa449-451 not present in TF_COX1_v1; and an extended turn at aa399-400 in TF_COX1_v2 that is not present in TF_COX1_v1, (Figure 6).

Nuclear-encoded fungicide-target genes

Several iterations of mapping and de novo assemblies were performed searching *T. frezii* sequences for nuclear-encoded genes that are target of fungicides. The *T. frezii* sequences submitted to GenBank corresponded to a 1,648 bp putative succinate dehydrogenase subunit B (*sdhb*) (MH603943), 1379 bp of β -tubulin gene (MH603944), 1,595 bp of cytochrome P450 monooxygenase 51 (*cyp51*) gene (MH603946), and 1,865 bp of ergosterol biosynthesis gene 4 *Erg4* (MH603945). SDHB of the sequenced *T. frezii* presented aa arginine (R) at the position 275, position that corresponds to 272 in SDHB of *B. cinerea* (Pers.) (*Botryotinia fuckeliana*) or 257 in SDHB *U. maydis*, Primers reported here can amplify a 1,379 bp of the β -tubulin gene in *T. frezii* including the region that corresponds to amino acids 198 and 200. In the *T. frezii* isolate sequenced mutations E198A and F200Y normally associated to resistance to benzimidazole fungicides were not observed,

DISCUSSION

Fungicide target genes

Mutations, usually single-nucleotide polymorphisms (SNPs), in

fungal plant-pathogen genes encoding fungicide targets, have been often the cause of developed resistance to strobilurins (QoI), carboxamides (SDHI) and benzimidazole fungicides [21-23,27]. Therefore, monitoring such mutations in field-collected samples has been shown essential to determine fungicide-treatment programs in crops [27,38]. In *T. frezii*, the basic research to discover potential mutations in relation to fungicide resistance has not been possible, because of the lack of genetic information about this pathogen. The genome of *Thecaphora thlaspeos*, a related species that invades Brassicaceae plants, was recently published [39]. This will be a valuable resource to help accelerate research on *T. frezii*. Here we report the complete mitogenome of *T. frezii* and provide tested primer sequences to amplify mitochondrial and nuclear genes that are target of common fungicides used in the peanut crop. Examples of mutations related to fungicide resistance are mentioned for the sole purpose of illustrating their relevance; this is not a complete review of the mutations reported in the literature for fungicide resistance.

Strobilurins, carboxamides and triazoles are the most common fungicides used to control peanut diseases [1, 18-20]. In peanut smut, however, various combinations of the mentioned fungicides have been less than 50% effective even when used at high doses [40,41]. It is known that mutations in genes encoding fungicide targets of pathogenic fungi are often the cause of developed resistance to strobilurins (QoI), carboxamides (SDHI) and triazole fungicides [21-23, 27]. Nevertheless, research on the molecular basis of *T. frezii* fungicide resistance has not been possible due to the lack of genetic information. Here we report the complete mitogenome of *T. frezii*, (accession MH392473), and provide molecular tools to study fungicide target genes such as *sdhb* (MH603943), *cyp51* (MH603946) and β -tubulin

(MH603944). Some of the sequences in this isolate indicated potential resistance to strobilurins and carboxamides.

Cytochrome B: Here we report the complete sequence of DNA and mRNA of the cytochrome B of *T. frezii*, MH603949. Strobilurins fungicides target Cyt B in the mitochondrial complex III, and strobilurins account for approximately 25% of the global market of fungicides [42]. At least eight amino acid substitutions in Cyt B are known to confer resistance to strobilurins in other fungi [21,43]. Thus, the monitoring of mutations in the Cyt B of field samples of fungal pathogens has been essential to modify fungicide-treatment programs in other crops [27,38]. Since the positions of Cyt B mutations reported in other fungi correspond in *T. frezii* COB to different exons in a gene that is 8,478 bp, here we propose the use of cDNA and primers provided in Table 1 as a mean to capture all the potential mutations in this gene. The *T. frezii* isolate sequenced in this study presented the aa phenylalanine (F) at position 17 of the quinone reduction site (Qi) in Cyt B. Substitution I17F has been linked to Diuron and Ilicicilin H resistance in other fungi [43]. Other aa substitutions in Cyt B that normally result in fungicide resistance are F129L, G137R and G143A [21,43] but these were not observed in the *T. frezii* isolate sequenced.

Succinate dehydrogenase b: In most eukaryotes including fungi, all the genes that form the mitochondrial complex II, one being succinate dehydrogenase, are encoded in the nucleus of the cell [44]. The isolate of *T. frezii* sequenced here, presented aa arginine (R) at position 275 of SDHB protein. Presence of aa arginine at the corresponding position of SDHB in *Botrytis cinerea* is known to confer resistance to SDH inhibitors Boscalid and Fluopyram [45]. Thus, it is highly possible that the *T. frezii* isolate sequenced may have resistance to carboxamides.

Cyp51, Erg4 and Nad. Triazole fungicides target the ergosterol biosynthesis pathway by inactivating either the lanosterol 14 β -demethylase *cyp51* gene (member of the P450 family, Erg11 in yeast) [46], or the ergosterol biosynthesis gene 4 (*Erg4*) [47], both nuclear-encoded genes. Mutations in these two genes, *cyp51* gene [24,25] and *Erg4* [47], or in the NADH oxidoreductase subunit of mitochondrial complex I are known to confer resistance to azole fungicides [24]. Here we report partial cDNA sequences of *T. frezii* *cyp51*, *Erg4*, and the *nad* mitochondrial genes of *T. frezii* that form the core of complex I. This information could be used as a starting point to research triazole resistance in *T. frezii*. However, additional mechanisms have recently been described as involved in azole resistance, such as a calcium transporter, transcription activation and a saccharopine dehydrogenase [48].

Beta tubulin: Primers reported here can amplify most of the β -tubulin gene of *T. frezii*, a 1,379 bp, including the coding region corresponding to amino acids 198 and 200. Mutations E198A and F200Y in the β -tubulin gene are known to confer resistance to benzimidazole fungicides in *Monilinia fructicola* and *Botrytis cinerea* [26,27]. The β -tubulin of *T. frezii* had high homology (89%) to *Anthracocystis flocculosa* PF-1 (XM_007882922.1) followed by 87% identity to the β -tubulin gene of *Kalmanozyma brasiliensis* (XM_016438208.1). No evidence of potential fungicide resistance was observed at positions 198 or 200 in the *T. frezii* isolate sequenced.

Mitogenome: endonucleases, gene orientation and heteroplasmy

In the mitochondrial genome of *T. frezii*, genes were oriented in two opposite directions away from the location of GII_MAT (Figure 3). In general, mitochondrial genes can all have the same orientation as for example in *Rhizoctonia solani* (NC_021436) [49], or they can be in opposite orientation as in *Phlebia radiata* Fr [50]. GII_MAT is a Group II transcriptase/maturase (GIIM) required *in vivo* for the splicing of group II introns, GIIM are mobile elements with multiple catalytic activities [51]. We speculate that the location of GII_MAT can be the origin of mitochondrial DNA replication.

BLAST analyses of *T. frezii* sequences showed homology to *Anthracocystis flocculosa* [Traquair et al.] Piatek et al., a fungus in the order Ustilaginales; though *T. frezii* is in the order Urocystidales. This probably occurred because currently there are 595 entries for Ustilaginales for each entry of Urocystidales. The *T. frezii* mitogenome sequenced had 52 genes in addition to intron-encoded homing endonucleases. A 27.5% (34,055 bp) of *T. frezii* mitogenome corresponded to introns, that is the equivalent to twice the size of the entire human mitogenome (16,569 bp) which encodes 37 genes [52]. Fungal mitogenomes vary significantly in size, for example 23,743 bp in the ascomycete *Zasmidium cellare* [53] and 235,849 bp in *Rhizoctonia solani* [49]. Despite having similar number of genes, mitogenome sizes are usually expanded by homing endonucleases. For example, in the 203 Kb mitogenome of *Sclerotinia borealis*, more than half (125 Kb) correspond to introns, most of them containing homing endonucleases (LAGLIDADG and GIY_YIG) [54]. LAGLIDADG endonucleases are encoded within introns type I, type II, archeal introns and inteins to promote lateral transfer [55]. The *T. frezii* mitogenome contained 10 homing endonucleases (LAGLIDADG, GIY-YIG, HEG, IEND), 7 of them encoded in introns of *cob* and *cox1* genes. Whereas the role of these endonucleases is not completely understood; fungal transcriptome analysis of the effect of a new fungicide, JS399-19 on *Fusarium graminearum* showed downregulation of only few proteins, among them was a LAGLIDADG [56]. Thus, it would be interesting to further characterize the role of these endonucleases in *T. frezii* in relation to fungicide response.

In the sequenced isolate of *T. frezii* the total length of cytochrome oxidase I gene (*cox1*) was 22,731 bp due to the presence of 15 introns encoding endonucleases. This gene in *T. frezii* showed heteroplasmy, that is two variants with possible structural differences, within their C-terminus as predicted by Garnier's model [37]. The C-terminal domain of *Cox1* regulates the *Cox1* synthesis by modulating interactions between proteins of the mitochondrial complex IV [57]. Until recently, mitochondrial heteroplasmy in Basidiomycota was considered transient or non-existent, but in 2017, heteroplasmy of *Cox1* was reported for the edible mushroom *Thelephora ganbajun* [58].

We have sequenced and annotated the mitogenome of *T. frezii*, as well as several fungicide target genes outside the mitogenome, these can be used to study the potential fungicide resistance in this pathogen. Amino acids predicted in COB and SDHB proteins indicated potential resistance to strobilurin and carboxamide fungicides. *T. frezii* *cox1* and *cob* genes could be also useful for diagnostic application, as these genes have been successfully used for genetic-barcoding identification in other species [59,60].

AUTHOR CONTRIBUTIONS

Cazon, Conforto, Paredes, Rago: isolation and molecular identification of the fungal pathogen; Lamb, Soave, Buteler: leaders and coordinators in the international agreement for research on peanut smut; Scheffler, Duke, Simpson DNA sequencing; Arias: data analysis and wrote the manuscript, Massa and Sobolev run MAKER for annotation and made editorial comments.

ACKNOWLEDGEMENTS

This work was feasible thanks to the support of USDA-ARS project number 6604-21000-004-00D and 6044-21000-004-00D, National Peanut Research Laboratory, Dawson, Georgia, U.S.A., the collaboration of the Instituto Nacional de Tecnología Agropecuaria (INTA), Cordoba, Argentina, and Fundacion Mani Argentino, Cordoba, Argentina. The authors would like to thank Ms. Valerie Orner at USDA-ARS-NPRL for her technical assistance.

REFERENCES

- Rago AM, Cazon LI, Paredes JA, Molina JPE, Conforto EC, Bisonard EM, et al. Peanut smut: from an emerging disease to an actual threat to argentine peanut production. *Plant Dis*. 2017;101:400-408.
- Marinelli A, March G, Rago AM. El carbon del mani *Thecaphora frezii*, sobre *Arachis hypogaea* L. in VII Congreso de Micología and XVII Jornadas Argentinas de Micología. 1995.
- Fiant S, Alonso C, Fontana T, Spinazze C, Costero D, Agosto G, et al. Caracterización de la producción de mani (*Arachis hypogaea*) Campana 2012/2013. In XXVIII Jornada Nacional de Mani, Cordoba, Argentina. 2013.
- Marinelli A, GJ March, Oddino C. Aspectos biológicos y epidemiológicos del carbon del mani (*Arachis hypogaea* L.) causado por *Thecaphora frezii* Carranza and Lindquist. *Agriscientia*. 2008;1:1-5.
- Gasso AM, Leis R, Marinelli A. Evaluación de incidencia y severidad del carbon de mani (*Thecaphora frezii*) en infecciones artificiales, sobre cultivares comerciales de mani in I Congreso Argentino de Fitopatología. 2008.
- Cazon LI. Detección molecular de *Thecaphora frezii* Carranza and Lindquist en semillas de mani (*Arachis hypogaea* L.). Thesis, Universidad Nacional de Córdoba, Argentina. 2015.
- Oddino C, Marinelli A, March G, Garcia J, Tarditi L, D'Eramo L, et al. Relación entre el potencial inóculo de *Thecaphora frezii*, la intensidad de carbon de mani y el rendimiento del cultivo. In XXV Jornada Nacional de Mani. 2010.
- Paredes JA, Cazon LI, Osella A, Peralta V, Alcalde M, Kearney MI, et al. Relevamiento regional del carbon del mani y estimaciones de pérdidas producidas por la enfermedad. In XXXI Jornada Nacional de Mani. 2016.
- Gunders J. Growers call for ban on Argentinian imports amid peanut smut fears, in ABC Rural 2017: ABC News.
- USDA-APHIS. Plant Protection and Quarantine. 2017.
- Sawe BE. Where are peanuts grown? 2018.
- Chatte L, Ricchetti M. Are mitochondria the Achilles' heel of the kingdom fungi? *Curr Opin Microbiol*. 2014;20:49-54.
- Kuhlbrandt W. Structure and function of mitochondrial membrane protein complexes. *BMC Biol*. 2015;13:89.
- Fujii K, Takakura S. Pyricut (diflufenconazole, UBF-002EC): A new fungicide for ornamental use. *Agrochem Japan*. 1998;72:14-16.
- Avenot HF, Michailides TJ. Progress in understanding molecular mechanisms and evolution of resistance to succinate dehydrogenase inhibiting (SDHI) fungicides in phytopathogenic fungi. *Crop Protection*. 2010;29:643-651.
- Von-Jagow G, Gribble GW, Trumppower BL. Mucidin and strobilurin A are identical and inhibit electron transfer in the cytochrome bc1 complex of the mitochondrial respiratory chain at the same site as myxothiazol. *Biochem*. 1986;25: 775-780.
- National Toxicology Program. Bioassay of triphenyltin hydroxide for possible carcinogenicity. *Natl Cancer Inst Carcinog Tech Rep Ser*. 1978;139:1-99.
- Culbreath AK, Brenneman TB, Kemerait RC, Hammes GG. Effect of the new pyrazole carboxamide fungicide penthiopyrad on late leaf spot and stem rot of peanut. *Pest Manag Sci*. 2009;65:66-73.
- Johnson RC, Cantonwine EG. Post-infection activities of fungicides against *Cercospora arachidicola* of peanut (*Arachis hypogaea*). *Pest Manag Sci*. 2014;70:1202-1206.
- Kemerait RC, Brenneman TB, Culbreath AK. Georgia pest management handbook-commercial edition. In: Horton D, edtr., Special Bulletin. 2016.
- Fernandez-Ortuno D, Tores JA, Vicente AD, Perez-Garcia A. Mechanisms of resistance to QoI fungicides in phytopathogenic fungi. *Int Microbiol*. 2008;11:1-9.
- Gisi U, Sierotzki H, Cook A, McCaffery A. Mechanisms influencing the evolution of resistance to Qo inhibitor fungicides. *Pest Manag Sci*. 2002;58:859-867.
- Sierotzki H, Scalliet G. A review of current knowledge of resistance aspects for the next-generation succinate dehydrogenase inhibitor fungicides. *Phytopathol*. 2013;103:880-887.
- Bowyer P, Mosquera J, Anderson M, Birch M, Bromley M, Denning DW. Identification of novel genes conferring altered azole susceptibility in *Aspergillus fumigatus*. *FEMS Microbiol Lett*. 2012;332:10-19.
- Gisi U, Chin KM, Knapova G, Farber RK, Mohr U, Parisi S, et al. Recent developments in elucidating modes of resistance to phenylamide, DMI and strobilurin fungicides. *Crop Protection*. 2000;19:863-872.
- Fernandez-Ortuno D, Grabke A, Li X, Schnabel G. Independent emergence of resistance to seven chemical classes of fungicides in botrytis cinerea. *Phytopathology*. 2014;105:424-432.
- Luo Y, Ma Z, Michailides TJ. Quantification of allele E198A in beta-tubulin conferring benzimidazole resistance in *Monilinia fructicola* using real-time PCR. *Pest Manag Sci*. 2007;63:1178-1184.
- Conforto C, Cazon I, Fernandez FD, Marinelli A, Oddino C, Rago A, et al., Molecular sequence data of *Thecaphora frezii* affecting peanut crops in Argentina. *European J Plant Pathol*. 2013;137:663-666.
- Conforto C, Cazon I, Fernandez FD, Marinelli A, Oddino C, Rago AM. Molecular detection of *Thecaphora frezii* in peanut (*Arachis hypogaea* L.) seeds. *J Plant Pathol*. 2016;98:327-330.
- Avanzini GV, Perez MA. Análisis de medios de cultivo para la evaluación in-vitro del crecimiento de carbon de mani. 2015.
- Astiz-Gassó MM, Marinelli A. Biología de la germinación de *Thecaphora frezii* in vitro. In XXVIII Jornada Nacional del Mani. 2013.
- Doyle JJ, Doyle JL. Isolation of plant DNA from fresh tissue. *Focus*. 1990;12:13-15.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990;215:403-410.
- Campbell MS, Law MY, Holt C, Stein JC, Moghe GD, Hufnagel DE, et al. MAKER-P: a tool kit for the rapid creation, management and quality control of plant genome annotations. *Plant Physiol*. 2014;164:513-524.

35. Cantarel BL, Korf I, Robb SM, Parra G, Ross E, Moore B, et al. MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome Res.* 2008;18:188-196.
36. Miranda I, Silva R, Santos MA. Evolution of the genetic code in yeasts. *Yeast.* 2006;23:203-213.
37. Garnier J, Osguthorpe DJ, Robson B. Analysis of the accuracy and implications of simple methods for predicting the secondary structure of globular proteins. *J Mol Biol.* 1978;120:97-120.
38. Fontaine S, Remuson F, Tachet LF, Micoud A, Marmeisse R, Melayah D, et al. Monitoring of *Venturia inaequalis* harbouring the QoI resistance G143A mutation in French orchards as revealed by PCR assays. *Pest Manage Sci.* 2009;65:74-81.
39. Courville JK, Frantzeskakis L, Gul S, Haeger N, Kellner R, Hebler N, et al. Smut infection of perennial hosts: the genome and the transcriptome of the Brassicaceae smut fungus *Thecaphora thlaspeos* reveal functionally conserved and novel effectors. *New Phytol.* 2019;222:1474-1492.
40. Paredes JA, Molina E, Cazon LI, Bisonard EM, Rago AM. Alternativas tecnológicas de aplicación de fungicidas para el control de *Thecaphora frezii*. In XXIX Jornada Nacional de Mani. 2014.
41. Paredes JA, Cazon LI, Bisonard EM. Efecto de fungicidas con carboxamidas en el control de *Thecaphora frezii*. in XXX Jornada Nacional de Mani. 2015.
42. China strobilurin fungicides market report 2016 edition. SM Research and Markets, Laura Wood. 2016.
43. Fisher N, Meunier B. Molecular basis of resistance to cytochrome bcl inhibitors. *FEMS Yeast Res.* 2008;8:183-192.
44. Burger G, Lang BF, Reith M, Gray MW. Genes encoding the same three subunits of respiratory complex II are present in the mitochondrial DNA of two phylogenetically distant eukaryotes. *Proc Natl Acad Sci USA.* 1996;93(6):2328-2332.
45. Angelini RMDM, Masiello M, Rotolo C, Pollastro S, Faretra F. Molecular characterisation and detection of resistance to succinate dehydrogenase inhibitor fungicides in *Botryotinia fuckeliana* (*Botrytis cinerea*). *Pest Manag Sci.* 2014;70:1884-1893.
46. Ghannoum MA, Rice LB. Antifungal agents: mode of action, mechanisms of resistance and correlation of these mechanisms with bacterial resistance. *Clin Microbiol Rev.* 1999;12(4):501-517.
47. Lupetti A, Danesi R, Campa M, Tacca MD, Kelly S. Molecular basis of resistance to azole antifungals. *Trends Mol Med.* 2002;8:76-81.
48. Mohd-Assaad N, McDonald BA, Croll D. Multilocus resistance evolution to azole fungicides in fungal plant pathogen populations. *Mol Ecol.* 2016;25:6124-6142.
49. Losada L, Pakala SB, Fedorova ND, Joardar V, Shabalina SA, Hostetler J, et al. Mobile elements and mitochondrial genome expansion in the soil fungus and potato pathogen *Rhizoctonia solani* AG-3. *Fems Microbiol Lett.* 2014;352:165-173.
50. Salavirta H, Oksanen I, Kuuskeri J, Mäkelä M, Laine P, Paulin L, et al. Mitochondrial genome of *Phlebia radiata* is the second largest (156 kbp) among fungi and features signs of genome flexibility and recent recombination events. *PLoS One.* 2014;9(5): e97141.
51. Matsuura M, Saldanha R, Ma H, Wank H, Yang J, Mohr G, et al. A bacterial group II intron encoding reverse transcriptase, maturase and DNA endonuclease activities: biochemical demonstration of maturase activity and insertion of new genetic information within the intron. *Genes Dev.* 1997;11:2910-2924.
52. Anderson S, Bankier AT, Barrell BG, De-Brujin MHL, Coulson AR, Drouin J, et al. Sequence and organization of the human mitochondrial genome. *Nature.* 1981;290:457.
53. Goodwin SB, McCorison CB, Cavaletto JR, Culley DE, LaButti K, Baker SE, et al. The mitochondrial genome of the ethanol-metabolizing, wine cellar mold *Zasmidium cellare* is the smallest for a filamentous ascomycete. *Fungal Biol.* 2016;120:961-974.
54. Mardanov AV, Beletsky AV, Kadnikov VV, Ignatov AN, Ravin NV. The 203 kbp mitochondrial genome of the phytopathogenic fungus *Sclerotinia borealis* reveals multiple invasions of introns and Genomic Duplications. *PLoS One.* 2014;9:e107536.
55. Rosen LE, Morrison HA, Masri S, Brown MJ, Springstubb B, Sussman D, et al. Homing endonuclease I-CreI derivatives with novel DNA target specificities. *Nucleic Acids Res.* 2006;34:4791-4800.
56. Hou Y, Zheng Z, Xu S, Chen C, Zhou M. Proteomic analysis of *Fusarium graminearum* treated by the fungicide JS399-19. *Pestic Biochem Physiol.* 2013;107:86-92.
57. Garcia-Villegas R, Camacho-Villasana Y, Shingu-Vazquez MA, Cabrera-Orefice A, Uribe-Carvajal S, Fox TD, et al. The Cox1 C-terminal domain is a central regulator of cytochrome c oxidase biogenesis in yeast mitochondria. *J Biol Chem.* 2017;292(26):10912-10925.
58. Wang, P, Sha T, Zhang Y, Cao Y, Mi F, Liu C, et al. Frequent heteroplasmy and recombination in the mitochondrial genomes of the basidiomycete mushroom *Thelephora gambajun*. *Sci Rep.* 2017;7:1626.
59. Lin S, Zhang H, Hou Y, Zhuang Y, Miranda L. High-level diversity of dinoflagellates in the natural environment, revealed by assessment of mitochondrial cox1 and cob genes for dinoflagellate DNA barcoding. *Appl Environ Microbiol.* 2009;75:1279-1290.
60. Seifert KA. Progress towards DNA barcoding of fungi. *Mol Ecol Resour.* 2009;9:83-89.