

Purdue University  
**Purdue e-Pubs**

---

Department of Food Science Faculty  
Publications

Department of Food Science

---

9-2018

## Oral Sensations and Secretions

Cordelia Running  
*Purdue University*, [crunning@purdue.edu](mailto:crunning@purdue.edu)

Follow this and additional works at: <https://docs.lib.purdue.edu/foodscipubs>

---

### Recommended Citation

Running, Cordelia, "Oral Sensations and Secretions" (2018). *Department of Food Science Faculty Publications*. Paper 18.  
<https://docs.lib.purdue.edu/foodscipubs/18>

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries.  
Please contact [epubs@purdue.edu](mailto:epubs@purdue.edu) for additional information.

1 This is the author copy of an accepted manuscript, posted to the Purdue University Repository after an  
2 embargo period as permitted by the publisher.

3

4 The published copy can be found at:

5

6

7 [Oral sensations and secretions](#)

8 CA Running

9 Physiology & behavior 193, 234-237

10

11 <https://doi.org/10.1016/j.physbeh.2018.04.011>

12 Oral sensations and secretions

13

14 Cordelia A Running, PhD

15 Department of Nutrition Science, Department of Food Science, Purdue University

16 700 W State St

17 West Lafayette IN, 47907 USA

18 [cunning@purdue.edu](mailto:cunning@purdue.edu)

19

20

21 Abstract

22 Sensations experienced in the mouth influence food choices, both immediately and in the long  
23 term. Such sensations are themselves influenced by experience with flavors, the chemical  
24 environment of the mouth, genetics of receptors for flavors, and individual behavior in the  
25 chewing of food. Gustation, the sense of taste, yields information about nutrients, influences  
26 palatability, and feeds into the human body's preparation to receive those nutrients. Olfaction,  
27 the sense of smell, contributes enormously to defining and identifying food flavors (and is  
28 experienced even after placing food inside the mouth). Another vital component of food flavor is  
29 texture, which contributes to palatability, especially if a food's texture violates a person's  
30 expectations. Next, chemesthesis is the sense of chemically induced irritancy and temperature,  
31 for example spiciness and stinging. All of these sensations are potentially modified by saliva, the  
32 chemical and physical media of the mouth. As a person experiences the culmination of these  
33 oral sensations, modified through an individual's own unique saliva, the flavors in turn influence  
34 both what and how a person eats.

35

36

37

38

39 Keywords: Taste, smell, texture, chemesthesis, saliva, food choice

40

41 1. Introduction

42 Food must be eaten in order to be nutritious (or deleterious). This fundamental fact may be  
43 obvious, but it is also the crux of the dilemmas regarding feeding behavior and health. In the  
44 end, all the healthy food in the world will have absolutely no effect on a person who does not  
45 make the choice to ingest that food. As the most dominant driver of food choice is flavor, e  
46 (IFICF, 2016), the mouth therefore plays a large role in a person's decision to ingest something.  
47 In general, "flavor" is experienced by the brain combining sensory experiences including aroma,  
48 tastes, textures, and perhaps even visual and audible cues from a food (Small, 2012). Not all  
49 fields and researchers agree on how many of these sensory attributes should be included in  
50 strict definitions of "flavor," but for this review the term will be used inclusively of the combination  
51 of sensations that may contribute to an individual's experiences and expectations of a food.  
52 Notably, a number of these potential "flavor" components can be obtained from sensory input  
53 before putting a food in the mouth. For example, expectations about a food may be derived from  
54 appearance, odors are perceived from a distance, and perception of thickness from stirring or  
55 swirling a beverage tracks strongly with in-mouth texture (Christensen & Casper, 1987).  
56 However, the full experience of flavor comes together as the food enters the mouth.  
57 Additionally, the properties (both physical and chemical) of the foods themselves will dictate the  
58 way the food is manipulated by the teeth and tongue. The combination of food sensation and  
59 oral manipulations can also influence the pace of feeding. Some foods take longer to chew than  
60 others. Some foods have dynamic textures which might influence the pace of consumption. But  
61 beyond even those fundamental differences, humans themselves differ in their own personal  
62 oral environments and chewing behaviors, which in turn will influence what they are willing to  
63 eat.

64

65 The purpose of this review is to briefly cover the sensations (gustation, retronasal olfaction,  
66 texture, and chemesthesis/trigeminal sensations) and secretions (saliva) of the oral cavity, and  
67 discuss how these factors may interact with food choice.

68

## 69 2. Gustation and olfaction

70 Gustation is the sense of taste, and will be used in this article to avoid confusion with the verb  
71 “to taste” (i.e., putting something in the mouth to experience/ingest it) or the more common  
72 vernacular meaning of the noun “taste” (referring to flavor in general). While colloquially “taste”  
73 often refers to many aspects of flavor, scientifically the sense of gustation is more limited. In  
74 general, gustation occurs by tastants first dissolving or suspending into saliva. The saliva  
75 passes over taste receptors, which are present on taste cells. The taste cells are organized into  
76 taste buds, which are present throughout much of the oral epithelium including the soft palate  
77 and the esophagus. However, most of the taste buds are found in the fungiform, foliate, and  
78 circumvallate papillae of the tongue (Miller & Bartoshuk, 1991). Tastants bind to the taste  
79 receptors, activating taste cells (either directly or through interaction with neighboring taste  
80 cells) to send a signal through nerves to the brain.

81

82 While scientists tend to agree gustation is limited to only a few qualities, there is little consensus  
83 on the total number of these “primary” gustatory qualities. Sweet, sour, salty, and bitter are  
84 widely accepted as gustatory percepts, but umami/savoriness, oleogustus (fatty acid taste),  
85 starchy taste, and several mineral tastes have also been proposed, with much ambiguity on why  
86 the first four are definitely gustatory sensations while the latter might or might not be. Some  
87 criteria for defining gustatory sensations have, however, been proposed (Mattes, 2011). They  
88 include: the sensation should offer an evolutionary advantage; ligands and receptors should  
89 have been identified; the receptors should activate gustatory specific cells and send signals  
90 along gustatory specific nerves; the sensation should be unique from other gustatory

91 sensations; and the sensation should evoke some kind of physiological or behavioral response.  
92 Still, even among widely accepted tastes, not all fit this list of criteria equally. For example, a  
93 wide variety of chemical structures are all detected as bitter, and the receptors for saltiness and  
94 sourness are still not firmly established. This ambiguity over defining gustation is a critical  
95 reason why the number of gustatory sensations remains debated. For example, the concept of a  
96 gustatory component for fat is not new, dating back at least to the 1500s (Fernel, 1581). Yet,  
97 separating the textural from the gustatory sensation of fat is challenging, not only due to the  
98 physical differences in texture from fat compared to water but also due to the challenges in  
99 distributing fatty molecules in an emulsion with water (Running & Mattes, 2014a, 2014b). The  
100 issue of a “taste” for fat is then further complicated by the apparent unpleasantness of the fatty  
101 acids when used as gustatory stimuli (Running, Craig, & Mattes, 2015; Running, Hayes, &  
102 Ziegler, 2017) compared to the assumed pleasantness of high-fat foods. This observed negative  
103 hedonic experience of fatty acid taste compared to fattiness in general is precisely why a new  
104 term, “oleogustus,” has been proposed to isolate the gustatory experience of fatty acids  
105 (Running et al., 2015). In any case, this particular gustatory sensation is a prime example of  
106 how strict definitions elude us for what is gustation and what is some other oral sensation.

107  
108 Despite the colloquial meaning of the word “taste,” much of a food’s overall flavor actually  
109 comes from odor: specifically, retronasal olfaction. These are the odors that pass through the  
110 back of the mouth into the airway and up into the nasal passages. When nasal passages are  
111 inflamed or otherwise blocked, this movement of air is restricted and results in lack of sensation.  
112 This is why when a person develops a respiratory infection, they can no longer “taste”  
113 anything—in reality, the sense of gustation is intact, but the sense of olfaction is limited. Loss of  
114 the retronasal olfaction reduces the sensation of the food to gustation, texture, and  
115 chemesthesis, and as a result the ability to identify flavors is severely limited.

116

117 The importance of odor on flavor identification is likely because the olfactory system can detect  
118 multitudes of distinct odors, especially compared to the very restricted list of gustatory  
119 sensations described above. To date, over 400 olfactory receptors have been identified, each of  
120 which is expressed on its very own set of individual olfactory neurons (Chess, Simon, Cedar, &  
121 Axel, 1994; Mainland et al., 2014; Zhang & Firestein, 2009). These neurons extend from the  
122 olfactory epithelium, and small piece of tissue in the uppermost part of the nasal passages,  
123 directly into the brain's olfactory bulb, where the signals are processed. When odorants dissolve  
124 into the mucus coating the olfactory epithelium, they stimulate the odor neurons by interacting  
125 with the receptors. The brain interprets the pattern of which neurons were activated, and the  
126 aroma is perceived (Buck & Axel, 1991; Hasin-Brumshtein, Lancet, & Olender, 2009). With the  
127 hundreds of olfactory receptors, and the subsequent plethora of activation combinations,  
128 humans can detect thousands of unique odors.

129

### 130 3. Texture and chemesthesis

131 Texture of foods is derived from physical structure, including the dynamic structural changes  
132 that occur as food interacts with oral surfaces and saliva (Koç, Vinyard, Essick, & Foegeding,  
133 2013). The physical nature of the food stimulates the sense of touch in the mouth, and the  
134 mouth is relatively sensitive to these sensations. Mechanoreceptors on the tongue can have  
135 small receptive fields (around 2.4 mm<sup>2</sup>) and respond to low levels of force (0.15 mN, which is  
136 similar the force exerted by gravity on half a grain of rice) (Trulsson & Essick, 1997). The texture  
137 of foods also directly influences the processes of chewing and swallowing, thus influencing the  
138 time food spends in the mouth as well as overall eating rate. Understanding the dynamics of  
139 food texture in the mouth require consideration of the food's original structure. While gustation  
140 and olfaction often emphasize biochemical reactions, such as receptor-ligand pairs and  
141 inter/intra-cellular trafficking of signaling molecules, the study of food texture requires some  
142 mechanical and engineering perspective in order to model and interpret breakdown of food's



143 physical structures. Inevitably, this physical breakdown of the food in turn influences the other  
144 senses in the mouth, as taste, odor, and chemesthetic compounds are released or re-adhere to  
145 the structures altered by chewing (Dijksterhuis & Piggott, 2000). Importantly, human mouth  
146 behavior may also have a role in the perception of food texture. Recently categories of mouth  
147 behavior have been proposed, which include chewers, crunchers, smooshers, and suckers  
148 (Jeltema, Beckley, & Vahalik, 2015), based on the preferred mouth movements of an individual  
149 and/or the foods that best allow those movements. For examples, “smooshers” seem to prefer  
150 to squeeze their food between the tongue of palate, and this behavior correlates with preference  
151 for more semi-solid foods such as yogurts or oatmeal. However, whether these food  
152 preferences drive the mouth behavior or the mouth behavior drives the food preferences is  
153 unclear. Nevertheless, numerous studies have confirmed that individuals certainly do differ in  
154 their mouth behaviors, including number of chews, shape of the chewing movement, amount of  
155 muscle effort in chewing, chewing rhythm, and more (Brown, Langley, Martin, & MacFie, 1994;  
156 Devezeaux de Lavergne, van de Velde, & Stieger, 2017). Furthermore, many of these  
157 parameters are more consistent within-subject than would be expected for wide differences in  
158 food properties such as hardness or fracturability. For example, individuals who used fewer  
159 chews before swallowing a carrot also tended to have fewer chews before swallowing apples,  
160 pork, salami, shortcake, and toast (Brown et al., 1994). Thus, the chewing behavior appears to  
161 be entrenched or innate in some way that is determined by the individual rather than the food.

162

163 Related to texture perception is chemesthesis. Chemesthesis is the chemical stimulation of  
164 temperature, touch, and irritation, and is also often referred to as “trigeminal” sensation due to  
165 the activity of the trigeminal nerve in carrying these signals from the mouth to the brain. In the  
166 oral cavity, this includes sensations like the spiciness of chilis, cooling of mint, and sting of  
167 carbonation. The trigeminal nerve informs the brain of these chemesthetic signals as well as  
168 physical touch and actual thermal changes. Just as this nerve is shared among these different

169 sensory stimuli, several receptors are also shared. For example, the TRPM8 protein in humans  
170 responds to both cool temperatures and menthol, and TRPV1 response to both capsaicin and  
171 heat (Roper, 2014). Thus, the overlap in words used to describe the sensations, whether  
172 chemical or physical/thermal in nature, makes sense (e.g., coffee and chili peppers can both be  
173 “hot”). Perception of intensity from chemesthesis can vary widely among individuals, but this  
174 variability is best documented for spiciness. Consistently, those who eat more spicy foods and  
175 like spicy foods rate the intensity of spiciness as lower than those who do not eat and do not like  
176 spicy foods (Cowart, 1987; Nolden & Hayes, 2017; Prescott & Stevenson, 1996; Tornwall,  
177 Silventoinen, Kaprio, & Tuorila, 2012). Likely, this association of eating/liking with intensity is a  
178 combination of innate and learned influences (Allen, McGeary, & Hayes, 2014; Byrnes & Hayes,  
179 2013, 2015, 2016; Guimaraes & Jordt, 2007; Tornwall et al., 2012).

180

#### 181 4. Saliva

182 Saliva is the biochemical media of the mouth, as well as a physical lubricant for oral surfaces.  
183 When salivation is impaired, gustation, olfaction, oral touch, chemesthesis, chewing, and  
184 swallowing are also impaired (Mese & Matsuo, 2007; Satoh-Kuriwada et al., 2009). In a single  
185 day, humans may swallow around 0.6-1.5 L of their own saliva (Aliko et al., 2015; Humphrey &  
186 Williamson, 2001), yet only about 0.7-1 mL of saliva is present in the mouth at any given time  
187 (Lagerlof & Dawes, 1984). The amount of saliva in the mouth is increased by stimulation with  
188 tastes and textures, with the strongest stimulations coming from sour taste and chewing (Dawes  
189 & Jenkins, 1964; Proctor & Carpenter, 2014; Watanabe & Dawes, 1988a, 1988b). Odor can also  
190 stimulate saliva, but the effect is generally weaker than taste and texture (Engelen et al., 2003).

191

192 Saliva is not just one fluid. Instead, it is a mixture from several functionally different salivary  
193 glands. “Major” salivary glands include the parotid, submandibular, and sublingual glands and  
194 contribute the larger volume of saliva to the mouth, whereas “minor” glands are distributed as

195 lingual, buccal, palatine, and labial glands and secrete a relatively small volume of saliva  
196 (Humphrey & Williamson, 2001). Importantly, the terms “major” and “minor” refer to the  
197 anatomical size of the glands and thus their volume of secretions, rather than the functional  
198 importance of those secretions. Indeed, many minor glands are crucial to maintaining adequate  
199 protection of oral surfaces (Eliasson & Carlen, 2010; Humphrey & Williamson, 2001). Further,  
200 minor glands in the posterior of the tongue (von Ebner’s glands) secrete directly into the clefts of  
201 the circumvallate and foliate papillae, where the densest population of taste buds in the mouth  
202 are located. Beyond the major and minor glandular distinction, saliva can also be categorized as  
203 serous, mucous, or mixed saliva. Serous saliva is thinner and more watery, while mucous saliva  
204 is thicker and has more gel-like properties. In general, serous saliva appears to be more  
205 involved in solubilizing and processing foods, which mucous saliva is designed more to protect  
206 the oral surfaces (Carpenter, 2013; Eliasson & Carlen, 2010; Humphrey & Williamson, 2001).  
207 The parotid glands (major) and von Ebner’s glands (minor) secrete serous saliva. The other  
208 major glands secrete mucous or mixed saliva, and the other minor glands secrete mucous  
209 saliva.

210

211 While saliva is over 99% water, the proteins and smaller molecules present in saliva significantly  
212 influence the behavior of foods in the mouth. Enzymes like salivary  $\alpha$ -amylase can break down  
213 starch in a matter of seconds, substantially altering texture of food and presumably changing the  
214 time it takes before an individual decides to swallow (Bridges, Smythe, & Reddrick, 2017;  
215 Mandel, Peyrot des Gachons, Plank, Alarcon, & Breslin, 2010). Small molecules in saliva are  
216 released to control ion concentrations in the mouth, such as bicarbonate to neutralize acids  
217 which in turn influences sour taste (Helm et al., 1982; Norris, Noble, & Pangborn, 1984). Beyond  
218 that, proteins in saliva have been proposed to modify astringency (Dinnella, Recchia, Fia,  
219 Bertuccioli, & Monteleone, 2009; Dinnella, Recchia, Vincenzi, Tuorila, & Monteleone, 2010),  
220 saltiness (Stolle et al., 2017), bitterness (Dsamou et al., 2012; Morzel et al., 2014), and

221 fattiness/oleogustus perception (Mounayar, Septier, Chabanet, Feron, & Neyraud, 2013;  
222 Neyraud, Palicki, Schwartz, Nicklaus, & Feron, 2012; Poette et al., 2014; Schmale, Ahlers,  
223 Blaker, Kock, & Spielman, 1993; Schmale, Holtgrevegrez, & Christiansen, 1990; Spielman,  
224 D'Abundo, Field, & Schmale, 1993). Additionally, composition of saliva can cause instability in  
225 emulsions (mixtures of oil and water) leading to different sensory perceptions of those  
226 emulsions among individuals (Dresselhuis, de Hoog, Stuart, Vingerhoeds, & van Aken, 2008;  
227 Vingerhoeds, Blijdenstein, Zoet, & van Aken, 2005). Furthermore, work in rats even indicates  
228 that exposure to bitterness and astringency can change saliva in ways that subsequently alter  
229 the acceptability or intensity of those bitter/astringent compounds (Martin et al., 2018;  
230 Torregrossa et al., 2014). In future years more research will hopefully confirm or clarify these  
231 linkages between diet, flavor, and saliva, and perhaps yield ways in which we could monitor or  
232 alter saliva to improve healthy dietary behaviors.

233

## 234 5. Eating behavior

235 All of these oral factors can, in isolation or combination, influence food choices and eating  
236 behavior. For gustation, sweetness, umami, and saltiness are often thought to enhance the  
237 palatability of food while bitterness, sourness, and oleogustus (the rancid, unpleasant taste of  
238 fatty acids, particularly polyunsaturated fatty acids, not the delicious fatty texture) seem to  
239 reduce palatability. However, these outcomes are not assured. Certainly, sweetness by itself is  
240 accepted even in infants, while sourness and bitterness are rejected (Maone, Mattes,  
241 Bernbaum, & Beauchamp, 1990; Tatzler, Schubert, Timischl, & Simbruner, 1985). Excess  
242 consumption of salt among many cultures would seem to indicate that it is palatable, and the  
243 positive effect of adding monosodium glutamate (prototypical stimulus for umami) to items such  
244 as soups imply this sensation is liked. However, people do not generally drink sugar water, or  
245 salt water, or umami water—instead, we consume foods as mixtures of flavors. Thus, the  
246 context of the food itself is critical for understanding the role of flavor in palatability. While fatty

247 foods are often well-liked, when fat breaks from an emulsion and pools on the top of the food,  
248 the food can be rejected. Further, our own work on oleogustus indicates that the gustatory  
249 sensation from fatty acids is unpalatable. (Running et al., 2015; Running et al., 2017). Similarly,  
250 bitterness in isolation is rated as unpleasant, yet bitter foods such as coffee, chocolate, and tea  
251 have become firmly embedded in many diets. Some of this may be due to post-ingestive  
252 feedback, as these products may have psychoactive (i.e., stimulatory caffeine) or energy  
253 contributions that make them appealing. Associations of these eating consequences with the  
254 flavor of the food can be learned, thus contributing to the wide array of responses of humans to  
255 the sensations experienced in the mouth. Overall, the combination of sensation, saliva, and  
256 experience with flavors influences human food choices.

257

258 In researching these phenomena, the goal is to identify which of these factors are modifiable,  
259 and how, in order to lead to improvements in human diets. With so many potential factors  
260 influencing food choices, there is clearly much room for new work. Combining data and  
261 approaches across these research fields, such as how mouth behavior or movements might  
262 influence dissolution of tastants, or how salivary composition might change over time to alter  
263 chemesthetic ligand activity, will hopefully lead to more targeted understanding of these  
264 phenomena at the individual level. Moreover, a better grasp of which factors are changeable,  
265 and how difficult changes would be to induce, is critical. For example, if a “smooshing” mouth  
266 behavior leads to preference for softer foods, does this in turn lead to excess energy intake  
267 because soft foods are quickly processed in the mouth? More importantly, can we change that  
268 mouth behavior to reduce the excess intake? Is there an ideal life stage to make such  
269 interventions? Could the flavor of the food be modified to help alter the mouth behavior? Will  
270 alterations in the mouth behavior change the secretion of saliva, and will that in turn alter the  
271 flavor experienced? Answers to such questions will be integral as we explore how the oral  
272 environment is more than just the gateway that accepts or rejects food. After all, many of the

273 compounds that are active from an oral sensory perspective, such as sweet sugars, slimy  
274 soluble dietary fiber, or bitter polyphenols, also influence human health. Presumably, our ability  
275 to detect many of these sensations is evolutionarily linked to those health outcomes. As modern  
276 technologies evolve our diets more quickly than we as humans can evolve, understanding the  
277 role of the oral environment in feeding will be paramount to maintaining healthy eating behaviors  
278 and food supplies.

279

## 280 6. Acknowledgements

281 The author would like to acknowledge the Purdue University Ingestive Behavior Research  
282 Center and their conference “The Pace of Life and Feeding: Health Implications” at which this  
283 work was presented.

284

## 285 7. Funding sources

286 This work is supported by the USDA National Institute of Food and Agriculture, Hatch project  
287 1013624.

288

## 289 8. Declaration of interest: none.

290

291

292

293

294

## References

295

- 296 Aliko, A., Wolff, A., Dawes, C., Aframian, D., Proctor, G., Ekström, J., . . . Vissink, A. (2015).  
297 World Workshop on Oral Medicine VI: clinical implications of medication-induced salivary  
298 gland dysfunction. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*,  
299 *120*(2), 185-206. doi:<http://dx.doi.org/10.1016/j.oooo.2014.10.027>  
300 Allen, A. L., McGeary, J. E., & Hayes, J. E. (2014). Polymorphisms in TRPV1 and TAS2Rs  
301 associate with sensations from sampled ethanol. *Alcohol Clin Exp Res*, *38*(10), 2550-  
302 2560. doi:10.1111/acer.12527

303 Bridges, J., Smythe, J., & Reddrick, R. (2017). Impact of salivary enzyme activity on the oral  
304 perception of starch containing foods. *Journal of Texture Studies*, 48(4), 288-293.  
305 doi:10.1111/jtxs.12252

306 Brown, W. E., Langley, K. R., Martin, A., & MacFie, H. J. H. (1994). Characterisation of patterns  
307 of chewing behaviour in human subjects and their influence on texture perception.  
308 *Journal of Texture Studies*, 25(4), 455-468. doi:10.1111/j.1745-4603.1994.tb00774.x

309 Buck, L., & Axel, R. (1991). A novel multigene family may encode odorant receptors: a  
310 molecular basis for odor recognition. *Cell*, 65(1), 175-187.

311 Byrnes, N. K., & Hayes, J. E. (2013). Personality factors predict spicy food liking and intake.  
312 *Food Qual Prefer*, 28(1), 213-221. doi:10.1016/j.foodqual.2012.09.008

313 Byrnes, N. K., & Hayes, J. E. (2015). Gender differences in the influence of personality traits on  
314 spicy food liking and intake. *Food Qual Prefer*, 42, 12-19.  
315 doi:10.1016/j.foodqual.2015.01.002

316 Byrnes, N. K., & Hayes, J. E. (2016). Behavioral measures of risk tasking, sensation seeking  
317 and sensitivity to reward may reflect different motivations for spicy food liking and  
318 consumption. *Appetite*, 103, 411-422.

319 Carpenter, G. H. (2013). The secretion, components, and properties of saliva. *Annu Rev Food*  
320 *Sci Technol*, 4, 267-276. doi:10.1146/annurev-food-030212-182700

321 Chess, A., Simon, I., Cedar, H., & Axel, R. (1994). Allelic inactivation regulates olfactory  
322 receptor gene expression. *Cell*, 78(5), 823-834. doi:http://dx.doi.org/10.1016/S0092-  
323 8674(94)90562-2

324 Christensen, C. M., & Casper, L. M. (1987). Oral and Nonoral Perception of Solution Viscosity.  
325 *Journal of Food Science*, 52(2), 445-447. doi:DOI 10.1111/j.1365-2621.1987.tb06635.x

326 Cowart, B. J. (1987). Oral chemical irritation: Does it reduce perceived taste intensity? *Chemical*  
327 *Senses*, 12(3), 467-479. doi:10.1093/chemse/12.3.467

328 Dawes, C., & Jenkins, G. N. (1964). The effects of different stimuli on the composition of saliva  
329 in man. *J Physiol*, 170, 86-100.

330 Devezeaux de Lavergne, M., van de Velde, F., & Stieger, M. (2017). Bolus matters: the  
331 influence of food oral breakdown on dynamic texture perception. *Food Funct*, 8(2), 464-  
332 480. doi:10.1039/c6fo01005a

333 Dijksterhuis, G. B., & Piggott, J. R. (2000). Dynamic methods of sensory analysis. *Trends in*  
334 *Food Science & Technology*, 11(8), 284-290. doi:https://doi.org/10.1016/S0924-  
335 2244(01)00020-6

336 Dinnella, C., Recchia, A., Fia, G., Bertuccioli, M., & Monteleone, E. (2009). Saliva  
337 characteristics and individual sensitivity to phenolic astringent stimuli. *Chem Senses*,  
338 34(4), 295-304. doi:10.1093/chemse/bjp003

339 Dinnella, C., Recchia, A., Vincenzi, S., Tuorila, H., & Monteleone, E. (2010). Temporary  
340 modification of salivary protein profile and individual responses to repeated phenolic  
341 astringent stimuli. *Chem Senses*, 35(1), 75-85. doi:10.1093/chemse/bjp084

342 Dresselhuis, D. M., de Hoog, E. H. A., Stuart, M. A. C., Vingerhoeds, M. H., & van Aken, G. A.  
343 (2008). The occurrence of in-mouth coalescence of emulsion droplets in relation to  
344 perception of fat. *Food Hydrocolloids*, 22(6), 1170-1183. doi:DOI  
345 10.1016/j.foodhyd.2007.06.013

346 Dsamou, M., Palicki, O., Septier, C., Chabanet, C., Lucchi, G., Ducoroy, P., . . . Morzel, M.  
347 (2012). Salivary protein profiles and sensitivity to the bitter taste of caffeine. *Chem*  
348 *Senses*, 37(1), 87-95. doi: 10.1093/chemse/bjr1070. Epub 2011 Aug 1025.

349 Eliasson, L., & Carlen, A. (2010). An update on minor salivary gland secretions. *Eur J Oral Sci.*,  
350 118(5), 435-442. doi: 410.1111/j.1600-0722.2010.00766.x. Epub 02010 Aug 00724.

351 Engelen, L., de Wijk, R. A., Prinz, J. F., Janssen, A. M., van der Bilt, A., Weenen, H., & Bosman,  
352 F. (2003). A comparison of the effects of added saliva, alpha-amylase and water on  
353 texture perception in semisolids. *Physiol Behav*, 78(4-5), 805-811.

354 Fernel, J. (1581). *Therapeutices universalis seu medendi rationis, libri septem* Frankfurt,  
355 Germany.

356 Guimaraes, M. Z. P., & Jordt, S. E. (2007). TRPA1 : A Sensory Channel of Many Talents. In W.  
357 B. Liedtke & S. Heller (Eds.), *TRP Ion Channel Function in Sensory Transduction and*  
358 *Cellular Signaling Cascades*. Boca Raton (FL).

359 Hasin-Brumshtein, Y., Lancet, D., & Olender, T. (2009). Human olfaction: from genomic  
360 variation to phenotypic diversity. *Trends in Genetics*, 25(4), 178-184.  
361 doi:<http://dx.doi.org/10.1016/j.tig.2009.02.002>

362 Helm, J. F., Dodds, W. J., Hogan, W. J., Soergel, K. H., Egide, M. S., & Wood, C. M. (1982).  
363 Acid Neutralizing Capacity of Human Saliva. *Gastroenterology*, 83, 69-74.  
364 doi:10.1016/S0016-5085(82)80286-2

365 Humphrey, S. P., & Williamson, R. T. (2001). A review of saliva: normal composition, flow, and  
366 function. *J Prosthet Dent*, 85(2), 162-169. doi:10.1067/mpr.2001.113778

367 IFICF. (2016). *2016 Food and Health Survey: Consumer Attitudes Toward Food Safety,*  
368 *Nutrition & Health*. Retrieved from  
369 [http://www.foodinsight.org/sites/default/files/2016\\_executivesummary\\_final\\_web.pdf](http://www.foodinsight.org/sites/default/files/2016_executivesummary_final_web.pdf).

370 Jeltema, M., Beckley, J., & Vahalik, J. (2015). Model for understanding consumer textural food  
371 choice. *Food Sci Nutr*, 3(3), 202-212. doi:10.1002/fsn3.205

372 Koç, H., Vinyard, C. J., Essick, G. K., & Foegeding, E. A. (2013). Food oral processing:  
373 conversion of food structure to textural perception. *Annu Rev Food Sci Technol*, 4, 237-  
374 266. doi:10.1146/annurev-food-030212-182637

375 Lagerlof, F., & Dawes, C. (1984). The volume of saliva in the mouth before and after  
376 swallowing. *J Dent Res*, 63(5), 618-621. doi:10.1177/00220345840630050201

377 Mainland, J. D., Keller, A., Li, Y. R., Zhou, T., Trimmer, C., Snyder, L. L., . . . Matsunami, H.  
378 (2014). The missense of smell: functional variability in the human odorant receptor  
379 repertoire. *Nat Neurosci*, 17(1), 114-120. doi:10.1038/nn.3598

380 Mandel, A. L., Peyrot des Gachons, C., Plank, K. L., Alarcon, S., & Breslin, P. A. (2010).  
381 Individual differences in AMY1 gene copy number, salivary alpha-amylase levels, and  
382 the perception of oral starch. *PLoS One*, 5(10), e13352.  
383 doi:10.1371/journal.pone.0013352

384 Maone, T. R., Mattes, R. D., Bernbaum, J. C., & Beauchamp, G. K. (1990). A New Method for  
385 Delivering a Taste without Fluids to Preterm and Term Infants. *Dev Psychobiol*, 23(2),  
386 179-191.

387 Martin, L. E., Nikonova, L. V., Kay, K., Paedae, A. B., Contreras, R. J., & Torregrossa, A.-M.  
388 (2018). Salivary proteins alter taste-guided behaviors and taste nerve signaling in rat.  
389 *Physiology & Behavior*, 184, 150-161. doi:<https://doi.org/10.1016/j.physbeh.2017.11.021>

390 Mattes, R. D. (2011). Accumulating evidence supports a taste component for free fatty acids in  
391 humans. *Physiol Behav*, 104(4), 624-631. doi:10.1016/j.physbeh.2011.05.002

392 Mese, H., & Matsuo, R. (2007). Salivary secretion, taste and hyposalivation. *J Oral Rehabil.*,  
393 34(10), 711-723.

394 Miller, I. J., & Bartoshuk, L. (1991). Taste Perception, Taste Bud Distribution, and Spatial  
395 Relationships. In T. V. Getchell, R. L. Doty, L. M. Bartoshuk, & J. B. Snow (Eds.), *Smell*  
396 *and Taste in Health and Disease* (pp. 205-234). New York: Raven Press.

397 Morzel, M., Chabanet, C., Schwartz, C., Lucchi, G., Ducoroy, P., & Nicklaus, S. (2014). Salivary  
398 protein profiles are linked to bitter taste acceptance in infants. *Eur J Pediatr*, 173(5), 575-  
399 582. doi:10.1007/s00431-013-2216-z

400 Mounayar, R., Septier, C., Chabanet, C., Feron, G., & Neyraud, E. (2013). Oral fat sensitivity in  
401 humans: Links to saliva composition before and after stimulation by oleic acid.  
402 *Chemosens Percept*, 6(3), 118-126. doi:10.1007/s12078-013-9152-1



403 Neyraud, E., Palicki, O., Schwartz, C., Nicklaus, S., & Feron, G. (2012). Variability of human  
404 saliva composition: Possible relationships with fat perception and liking. *Arch Oral Biol*,  
405 57(5), 556-566. doi:DOI 10.1016/j.archoralbio.2011.09.016

406 Nolden, A. A., & Hayes, J. E. (2017). Perceptual and affective responses to sampled capsaicin  
407 differ by reported intake. *Food Qual Prefer*, 55, 26-34.  
408 doi:10.1016/j.foodqual.2016.08.003

409 Norris, M. B., Noble, A. C., & Pangborn, R. M. (1984). Human saliva and taste responses to  
410 acids varying in anions, titratable acidity, and pH. *Physiol Behav*, 32(2), 237-244.

411 Poette, J., Mekoué, J., Neyraud, E., Berdeaux, O., Renault, A., Guichard, E., . . . Feron, G.  
412 (2014). Fat sensitivity in humans: Oleic acid detection threshold is linked to saliva  
413 composition and oral volume. *Flavour Frag J*, 29(1), 39-49.

414 Prescott, J., & Stevenson, R. J. (1996). Psychophysical responses to single and multiple  
415 presentations of the oral irritant zingerone: relationship to frequency of chili consumption.  
416 *Physiol Behav*, 60(2), 617-624.

417 Proctor, G. B., & Carpenter, G. H. (2014). Salivary secretion: mechanism and neural regulation.  
418 *Monogr Oral Sci*, 24:, 14-29.

419 Roper, S. D. (2014). TRPs in taste and chemesthesis. *Handb Exp Pharmacol*, 223, 827-871.  
420 doi:10.1007/978-3-319-05161-1\_5

421 Running, C. A., Craig, B. A., & Mattes, R. D. (2015). Oleogustus: The unique taste of fat. *Chem*  
422 *Senses*, 40(7), 507-516.

423 Running, C. A., Hayes, J. E., & Ziegler, G. R. (2017). Degree of free fatty acid saturation  
424 influences chocolate rejection in human assessors. *Chem Senses*, 42(2), 161-166.  
425 doi:10.1093/chemse/bjw116

426 Running, C. A., & Mattes, R. D. (2014a). Different oral sensitivities to and sensations of short-,  
427 medium-, and long-chain fatty acids in humans. *Am J Physiol-Gastr L*, 307(3), G381-  
428 G389. doi:10.1152/ajpgi.00181.2014

429 Running, C. A., & Mattes, R. D. (2014b). Humans are more sensitive to the taste of linoleic and  
430  $\alpha$ -linolenic than oleic acid. *Am J Physiol-Gastr L*, 308(5), G442-G449.  
431 doi:10.1152/ajpgi.00394.2014

432 Satoh-Kuriwada, S., Shoji, N., Kawai, M., Uneyama, H., Kaneta, N., & Sasano, T. (2009).  
433 Hyposalivation strongly influences hypogeusia in the elderly. *Journal of Health Science*,  
434 55(5), 689-698. doi:10.1248/jhs.55.689

435 Schmale, H., Ahlers, C., Blaker, M., Kock, K., & Spielman, A. I. (1993). Perireceptor Events in  
436 Taste. In D. Chadwick, J. Marsh, & J. Goode (Eds.), *Ciba Foundation Symposium 179 -*  
437 *Molecular Basis of Smell and Taste Transduction* (Vol. 179, pp. 167-185). Chichester,  
438 UK: John Wiley & Sons, Ltd.

439 Schmale, H., Holtgrevegrez, H., & Christiansen, H. (1990). Possible role for salivary-gland  
440 protein in taste reception indicated by homology to lipophilic-ligand carrier proteins.  
441 *Nature*, 343(6256), 366-369. doi:Doi 10.1038/343366a0

442 Small, D. M. (2012). Flavor is in the brain. *Physiology & Behavior*, 107(4), 540-552.  
443 doi:https://doi.org/10.1016/j.physbeh.2012.04.011

444 Spielman, A. I., D'Abundo, S., Field, R. B., & Schmale, H. (1993). Protein analysis of human von  
445 Ebner saliva and a method for its collection from the foliate papillae. *J Dent Res*, 72(9),  
446 1331-1335.

447 Stolle, T., Grondinger, F., Dunkel, A., Meng, C., Médard, G., Kuster, B., & Hofmann, T. (2017).  
448 Salivary proteome patterns affecting human salt taste sensitivity. *J Agric Food Chem*,  
449 65(42), 9275-9286. doi:10.1021/acs.jafc.7b03862

450 Tatzer, E., Schubert, M. T., Timischl, W., & Simbruner, G. (1985). Discrimination of taste and  
451 preference for sweet in premature babies. *Early Hum Dev*, 12(1), 23-30.

- 452 Tornwall, O., Silventoinen, K., Kaprio, J., & Tuorila, H. (2012). Why do some like it hot? Genetic  
453 and environmental contributions to the pleasantness of oral pungency. *Physiol Behav*,  
454 107(3), 381-389. doi:10.1016/j.physbeh.2012.09.010
- 455 Torregrossa, A. M., Nikonova, L., Bales, M. B., Villalobos Leal, M., Smith, J. C., Contreras, R.  
456 J., & Eckel, L. A. (2014). Induction of salivary proteins modifies measures of both  
457 orosensory and postingestive feedback during exposure to a tannic acid diet. *PLoS One*,  
458 9(8), e105232. doi:10.1371/journal.pone.0105232
- 459 Trulsson, M., & Essick, G. K. (1997). Low-threshold mechanoreceptive afferents in the human  
460 lingual nerve. *J Neurophysiol*, 77(2), 737-748. doi:10.1152/jn.1997.77.2.737
- 461 Vingerhoeds, M. H., Blijdenstein, T. B. J., Zoet, F. D., & van Aken, G. A. (2005). Emulsion  
462 flocculation induced by saliva and mucin. *Food Hydrocolloids*, 19(5), 915-922.  
463 doi:10.1016/j.foodhyd.2004.12.005
- 464 Watanabe, S., & Dawes, C. (1988a). A comparison of the effects of tasting and chewing foods  
465 on the flow rate of whole saliva in man. *Archives of Oral Biology*, 33(10), 761-764.  
466 doi:http://dx.doi.org/10.1016/0003-9969(88)90010-6
- 467 Watanabe, S., & Dawes, C. (1988b). The effects of different foods and concentrations of citric  
468 acid on the flow rate of whole saliva in man. *Arch Oral Biol*, 33(1), 1-5.
- 469 Zhang, X., & Firestein, S. (2009). Genomics of olfactory receptors. *Results Probl Cell Differ*, 47,  
470 25-36. doi:10.1007/400\_2008\_28

471

472