

Neonatal BCG vaccination reduces interferon gamma responsiveness to heterologous pathogens in infants from a randomised controlled trial

Freyne B ^{1,2,3}, Messina NL ^{1,2}, Donath S ^{2,4}, Germano S ¹, Bonnici R ¹, Gardiner K ¹, Casalaz D ⁵, Robins-Browne RM ^{1,6}, Netea MG ^{7,8}, Flanagan KL ^{9,10}, Kollmann T ^{11,12}, Curtis N ^{1,2} and the MIS BAIR group *

¹ Infectious Diseases & Microbiology Group, Murdoch Children's Research Institute, Royal Children's Hospital Melbourne, Parkville, Australia.

² Department of Paediatrics, The University of Melbourne, Parkville, Australia.

³ Institute of Infection and Global Health, University of Liverpool and Malawi-Liverpool Wellcome Trust Research Program, Blantyre, Malawi.

⁴ Clinical Epidemiology & Biostatistics Unit, Murdoch Children's Research Institute, Melbourne, Parkville, Australia.

⁵ Department of Paediatrics, Mercy Hospital for Women, Heidelberg, Australia.

⁶ The Department of Microbiology and Immunology, The University of Melbourne, Parkville, Australia.

⁷ Department of Internal Medicine, Radboud Institute for Molecular Life Sciences, Radboud University Nijmegen Medical Center, The Netherlands.

⁸ Radboud Center for Infectious Diseases, Radboud University Nijmegen Medical Center, The Netherlands.

⁹ University of Tasmania, Launceston, Australia.

¹⁰ Monash University, Clayton, Australia.

¹¹ Department of Experimental Medicine, University of British Columbia, Vancouver, Canada.

¹² Department of Pediatrics, University of British Columbia, Canada.

* MIS BAIR group: Veronica Abruzzo, Katie Allen, Clare Morrison, Anne-Louise Ponsonby,
Peter Vuillermin

Running title (50 characters): BCG alters infant heterologous immune responses

Key words: Bacille Calmette-Guérin, Immunisation, Heterologous, Non-specific effects,
Innate immunity, Infants

Corresponding author:

Prof Nigel Curtis
Department of Paediatrics
University of Melbourne
The Royal Children's Hospital Melbourne
Parkville, VIC 3052 Australia.

Nigel.Curtis@rch.org.au

Phone: +61 3 93456366

Alternate corresponding author

Dr Bridget Freyne,
Malawi-Liverpool Wellcome Trust,
Queen Elizabeth Central Hospital,
PO Box 30096
Chichiri, Blantyre 3

bfreyne@mlw.mw

Phone: +265 995893875

40 word summary

Neonatal BCG vaccination results in a decreased proportion of infants mounting an IFN- γ response to heterologous stimulation at 7 months of age. Both sex and maternal BCG vaccination status influenced the effect of BCG vaccination on heterologous cytokine responses.

Abstract

Background. Bacille Calmette-Guérin (BCG) vaccination has beneficial non-specific (heterologous) effects that protect against non-mycobacterial infections. We have previously reported that BCG vaccination at birth alters *in vitro* cytokine responses to heterologous stimulants in the neonatal period. This study investigated heterologous responses in 167 infants in the same trial seven months after randomisation.

Methods. A whole blood assay was used to interrogate *in vitro* cytokine responses to heterologous stimulants (killed pathogens) and Toll-like receptor (TLR) ligands.

Results. Compared to BCG-naïve infants, BCG-vaccinated infants had increased production of MIG and IFN- γ in response to mycobacterial stimulation and decreased production of IFN- γ in response to heterologous stimulation. Reduced IFN- γ responses to heterologous stimulants and TLR ligands were attributable to a decrease in the proportion of infants who mounted a detectable IFN- γ response. BCG-vaccinated infants also had increased production of MIG and IL-8, and decreased production of IL-10, MIP-1 α and MIP-1 β , the pattern of which varied by stimulant. IL-1Ra responses following TLR1/2 (Pam3CYSK4) stimulation were increased in BCG-vaccinated infants. Both sex and maternal BCG vaccination status influenced the effect of neonatal BCG vaccination.

Conclusions. BCG vaccination leads to changes in IFN- γ responsiveness to heterologous stimulation. BCG-induced changes in other cytokine responses to heterologous stimulation varies by pathogen.

Introduction

Bacille Calmette-Guérin (BCG) vaccination at birth is associated with decreased mortality in infants [1, 2]. This is proposed to be due to heterologous ('non-specific') protection against early-life infections, an effect presumed to result from the immunomodulatory properties of BCG [3, 4]. The beneficial non-specific effects of BCG vaccine persist beyond the neonatal period. BCG vaccination is associated with reduced incidence of lower respiratory tract infections in childhood and sepsis in the first year of life [5, 6] and protection against eczema in infants with an atopic disposition [7]. Maternal BCG, delivery method and gestational age have been reported to modify the non-specific effects of BCG [8, 9].

The mechanisms underlying the beneficial heterologous effects of BCG vaccination remain unclear [10]. We have previously shown that *in vitro* cytokine responses to heterologous stimulants differs between BCG-vaccinated and BCG-naïve neonates [11]. The longer term heterologous immunological effects induced by neonatal BCG are poorly defined but are likely to be different. Maturational changes mean the immune response in infancy is different to that of neonates [12-14]. In addition, a central hypothesis in the literature on the non-specific effects of vaccination is that live vaccines confer protection against all-cause mortality, which is negated by subsequent non-live vaccines in the first six months of life [15]. In the current study, we report the effect of neonatal BCG vaccination on heterologous cytokine responses in infants from a randomised controlled trial at seven months of age.

Methods

Participants were a subset of infants recruited from The Melbourne Infant Study: BCG for the prevention of allergy and infection (MIS BAIR) in which neonates were randomised to vaccination with BCG-Denmark 0.05 mL intradermally or no BCG vaccination (Clinical trials registration NCT01906853). The inclusion and exclusion criteria for MIS BAIR are described elsewhere [16]. MIS BAIR participants were invited to attend a 7-month study visit to provide a blood sample. All participants with a blood sample from December 2014 to April 2016 were recruited to this sub-study. Inclusion criteria for cytokine analysis were: (i) sufficient blood for all stimulants or (ii) participant provided sample in previous study [11]. Exclusion criteria were: (i) fever in the previous 24 hours; (ii) chronic illness (iii) less than two doses of the routine scheduled vaccines; (iv) any vaccination within the last 7 days; and (iv) any blood products since birth (Figure 1).

Venous blood was collected and *in vitro* stimulation assays were done as described previously [17]. Briefly, whole blood, diluted 1:1 with RPMI-1640 medium (GlutaMAX™ Supplement, HEPES, Gibco, Life Technologies), was stimulated at 37°C (5% CO₂: air) for 20 (± 2) hours with RPMI alone (unstimulated control), BCG-Denmark (Serum Statens Institute, Denmark) 75 µg/mL, killed *Mycobacterium tuberculosis* 1.0 x 10⁶ CFU/mL, *Escherichia coli* 1.0 x 10⁶ CFU/mL, *Haemophilus influenzae* type B 1.0 x 10⁶ CFU/mL, *Staphylococcus aureus* 1.0 x 10⁷ CFU/mL, *Streptococcus pyogenes* (GAS) 1.0 x 10⁷ CFU/mL, *Streptococcus pneumoniae* serotype 15C (non-vaccine serotype) 1.0 x 10⁷ CFU/mL, *Listeria monocytogenes* 1.0 x 10⁷ CFU/mL, *Candida albicans* 1.0 x 10⁶ CFU/mL, and the Toll-like receptor (TLR) ligands lipopolysaccharide (LPS; TLR 4 agonist) 100 ng/mL, resiquimod (R848; TLR 7/8) 3.5 µg/mL, peptidoglycan (PEPG; TLR 2/4) 10 µg/mL and (S)-(2,3-bis(palmitoyloxy)-(2-RS)-propyl)-N-palmitoyl-(R)-Cys-(S)-Ser-(S)-

Lys4OH, trihydrochloride (Pam3CSK4; TLR 1/2) 1 µg/mL; all from InvivoGen. When insufficient blood was available for all stimulations, a pre-determined priority order was used. The following cytokines were measured in supernatants using an xMAP Luminex 200 Analyser as described previously: IFN- γ , TNF- α , IL-6, IL-1 β , MIF, IL-8 (CXCL8), MIG (CXCL9), IP-10 (CXCL10), MCP-1 (CCL2), MIP-1 α (CCL3), MIP-1 β (CCL4), IL-10, IL-1Ra [17]. Study staff were blinded to the BCG-vaccination status of participants.

Statistical analysis was done using Stata version 13.1. Cytokine results below the lower limit of detection (LLD) were assigned a value of half the lowest detectable value. There were 14 values above the upper limit of detection (ULD); these were excluded from the analysis. Proportions of detectable values for each cytokine are shown in Supplementary Table 1.

Prior to analysis, all cytokine data were log transformed. Regression analyses, with the log transformed value of the unstimulated cytokine values as a covariate, was used to determine the effect of (i) BCG vaccination, (ii) maternal BCG vaccination status and (iii) infant's sex on cytokine production in response to each stimulant. This approach was used in all regression analyses to account for variability in unstimulated samples and avoids the need for nil subtraction. For cytokine/stimulant pairs with normally distributed data (Supplementary Figure 1), linear regression was used and results presented as geometric mean ratios (GMR) with 95% confidence intervals. For cytokine/stimulant pairs with non-normally distributed data (Supplementary Figure 1), quantile regression was used and results presented as differences in the medians with 95% confidence intervals.

Subgroup analysis (sex and maternal BCG vaccination status) was done when there was statistical evidence of an interaction between BCG vaccination and the subgroup variable, as

determined by interaction analysis (Supplementary Table 4). Given the smaller numbers of participants in sub-group analyses, quantile regression was used, with bootstrapping as required.

Multivariable analysis was done to assess the effect of sex, delivery method, age at infant BCG vaccination (less than or greater than 48 hours of life), maternal BCG vaccination status, age at blood sample and the number of doses of routine vaccinations received.

Multivariable analyses, with the unstimulated values as a covariate, were done using linear or quantile regression as per the corresponding cytokine/stimulant pair in the primary univariate analysis.

For cytokines that showed a dichotomous response, participants were classified according to whether cytokine concentrations in the stimulated supernatants were above ('responders') or below ('non-responders') the LLD. Fisher's exact test was used to compare the proportion of 'responders' between BCG-vaccinated and BCG-naïve participants. To assess the contribution of this observation to the results of the primary analysis, a sensitivity analyses was done for the effect of BCG vaccination when 'non-responders' were excluded.

The study was approved by the research ethics committees of the Royal Children's Hospital (HREC 33025) and the Mercy Hospital for Women (HREC R12/28).

Results

Demographic features of participants were similar between BCG-vaccinated and BCG-naïve participants (Table 1). In unstimulated samples, BCG-vaccinated infants had lower overall cytokine concentrations (particularly IL-10, MIF, IP-10 and IL-8) compared with BCG-naïve

infants (Table 2). In response to all stimulants, there was marked inter-individual variability in cytokine production (Supplementary Figure 1). For IFN- γ , IP-10, IL-1Ra and IL-1 β , there was evidence of dichotomous responses, i.e. participants who did or did not mount a measurable cytokine response following stimulation.

In response to the ‘specific’ mycobacterial antigens BCG and *M. tuberculosis*, BCG-vaccinated infants had higher production of IFN- γ and MIG compared to BCG-naïve infants, and also higher TNF- α and IL-6 following stimulation with BCG (Figure 2, Supplementary Table 1). In response to heterologous stimulation with the Gram-positive bacteria *S. pyogenes* and *S. aureus*, BCG-vaccinated infants had differential production of chemokines including MIF, MIG and MIP-1 β . Chemokine production in response to stimulation with the intracellular pathogens (*L. monocytogenes* and *C. albicans*) was also altered in BCG-vaccinated infants (Figure 2, Supplementary Table 1). In response to stimulation with the Gram-negative bacteria *E. coli* and *H. influenzae*, and the TLR agonist LPS, BCG-vaccinated infants had decreased IFN- γ responses and increased IL-8 response compared to BCG-naïve infants. In response to stimulation with TLR agonists Pam3CYSK4 (TLR1/2) and PEPG (TLR2), BCG-vaccinated infants had increased IL-1Ra production and decreased IFN- γ production respectively (Figure 2, Supplementary Table 1).

There was a dichotomous IFN- γ response to all stimulants aside from R848, with a distinct separation between ‘responders’ and ‘non-responders’. For the ‘specific’ mycobacterial antigens, BCG and *M. tuberculosis*, there was a higher proportion of IFN- γ responders in the BCG-vaccinated group compared to the BCG-naïve group (Figure 3, Supplementary Table 2). Conversely, for the heterologous stimulants, the proportion of IFN- γ responders was significantly lower in BCG-vaccinated infants for all stimulants with the exception of *S.*

pneumoniae (Figure 3, Supplementary Table 2). IP-10, IL-1Ra and IL-1 β responses showed a similar pattern; however, differences in the proportion of responders were not statistically significant between BCG-vaccinated and BCG-naïve infants (Supplementary Table 2).

A sensitivity analysis done after removal of all non-responders showed no significant difference in IFN- γ responses to heterologous stimulants between BCG-vaccinated and BCG-naïve infants, revealing that the observed difference in IFN- γ GMRs was attributable to the change in proportion of ‘responders’ (Supplementary Table 3). In contrast, the difference in the specific IFN- γ response to *in vitro* BCG stimulation between BCG-vaccinated and BCG-naïve infants was still observed.

Infants’ sex also independently influenced cytokine responses. Compared with girls, boys had increased production of both MCP-1 and MIP-1 β in response to almost all stimulants (Supplementary Table 1 & Supplementary Figure 2). Boys also had increased production of IFN- γ in response to stimulation with BCG, *E. coli*, *H. influenzae* and PEPG, and TNF- α in response to most stimulants, particularly *E. coli*, *H. influenzae*, LPS and R848 (Supplementary Table 1 & Supplementary Figure 2). MIG production was also higher in boys following stimulation with *E. coli*. Conversely, boys had decreased production of IL-1Ra in response to *C. albicans* and IL-1 β in response to *L. monocytogenes*, and in response to Pam3CYSK4 (Supplementary Table 1 & Supplementary Figure 2).

A significant interaction between sex and infant BCG vaccination was seen for multiple cytokine/stimulant pairs but most consistently for production of IL-6 and IL-1 β (Supplementary Table 4). Subgroup analyses of the effect of BCG-vaccination in boys and girls separately showed that, compared to BCG-naïve girls, BCG-vaccinated girls had

increased pro-inflammatory cytokine production: (a) IL-6 in response to *E. coli*, *H. influenzae*, LPS and R848 and (b) IL-1 β in response to *S. pneumoniae* (Figure 4, panel a and b), and an increase in the anti-inflammatory cytokine IL-1Ra in response to Pam3CYSK4 (Figure 4, panel c). Overall, compared to BCG-naïve boys, BCG-vaccinated boys had lower cytokine production for all cytokine/stimulant pairs when there was an interaction between infant BCG vaccination and sex. This was statistically significant for IL-1 β /*E. coli*, IFN- γ /Pam3CYSK4 and TNF- α /*C. albicans* responses (Figure 4, panel a-c).

Maternal BCG vaccination status independently influenced cytokine responses to heterologous stimulants and TLR ligands. Compared to infants of mothers who had not had BCG vaccination, infants of mothers who were BCG-vaccinated had increased IFN- γ production in response to stimulation with *M. tuberculosis*, decreased IFN- γ in response to PEPG (TLR2/4); decreased TNF- α in response to BCG, *E. coli*, *H. influenzae*, PEPG (TLR2/4), LPS (TLR4) and R848 (TLR7/8); decreased IL-1Ra in response to *H. influenzae*; decreased MCP-1 in response to *S. pneumoniae* and *S. pyogenes*, decreased IP-10 in response to *L. monocytogenes*, decreased MIP-1 β in response to *E. coli* and *S. pneumoniae* (Supplementary Table 1 and Supplementary Figure 3).

A significant interaction between maternal BCG vaccination and infant BCG vaccination was seen for MIG/*E. coli*, IL-10/BCG and IFN- γ /*S. pneumoniae* cytokine/stimulant pairs (Supplementary Table 4). Separate analyses of the effect of BCG in infants of BCG-vaccinated mothers and infants of BCG-naïve mothers showed that in the former there was lower production of MIG in response to *E. coli* and IFN- γ in response to *S. pneumoniae*, whereas, BCG vaccination had no effect on this response in infants of BCG-naïve mothers

(Figure 4, panel d). Conversely, infants of BCG-vaccinated mothers had higher production of IL-10 in response to stimulation with BCG (Figure 4, panel d).

Multivariable analyses were done to evaluate the potential confounding effect of sex, maternal BCG vaccination status, mode of delivery, age at BCG vaccination, age at blood draw, and number of routine vaccination doses received (Supplementary Table 5). Covariate analysis with these variables did not reveal any meaningful effect on the primary analysis, thus excluding confounding by these variables.

Discussion

We previously reported that BCG vaccination at birth led to decreased production of IL-1ra, IL-6, MCP-1, MIP-1 α and MIP-1 β in response to heterologous stimulants and TLR ligands in the neonatal period (7 days post after randomisation) [17]. This study extends these findings to show that BCG-induced changes in cytokine expression following heterologous stimulation are detectable at 7 months of age, characterised by decreased IFN- γ responsiveness in BCG-vaccinated infants.

Our finding that BCG vaccination was associated with an increase in IFN- γ and pro-inflammatory cytokines (IL-6, IL-1 β and TNF- α) in response to mycobacterial antigens (BCG and *M. tuberculosis*) is consistent with prior studies [13, 14, 18-20]. Furthermore, our finding that two-thirds of BCG-naïve infants produced IFN- γ in response to *in vitro* stimulation with BCG supports our previous finding of a role for NK cells and unconventional T-cells in early IFN- γ responses to BCG [21]. Our main finding that neonatal BCG vaccination is associated with a decrease in the proportion of infants who produce IFN- γ in response to heterologous stimulants has not previously been reported.

Nissen *et al.* reported that, in Danish infants at 12 months of age, BCG vaccination did not lead to differential IFN- γ production either in response to *ex vivo* heterologous bacterial (*E. coli*, *S. pneumoniae*, *C. albicans*) or LPS stimulation [22]. However, in that study approximately half of the BCG-vaccinated participants did not have a specific IFN- γ response to *in vitro* BCG stimulation. Smith *et al.* also assessed the effects of BCG on *in vitro* heterologous cytokine responses in infants, although BCG was given at 6 weeks of age [23]. As in our study, BCG-vaccinated infants had an increase in *M. tuberculosis* lysate-induced IFN- γ response. Although there was no significant difference between BCG-vaccinated and BCG-naïve infants in IFN- γ production following *E. coli*, *S. aureus*, *C. albicans*, LPS or Pam3CYSK4 stimulation, fold-change analysis showed decreased LPS-induced IFN- γ responses in BCG-vaccinated infants, an effect which did not reach statistical significance given the small sample size.

Compared to BCG-naïve infants, BCG-vaccinated infants in our study showed a distinctive pattern of differential chemokine production which varied between biologically-related stimulants. Specifically, in response to mycobacterial, Gram-positive and intracellular heterologous stimulants, chemokine production in BCG-vaccinated infants was characterised by increased production of MIG and a concomitant decrease in MIP-1 α and MIP-1 β . MIG is produced by activated monocytes and is the primary chemoattractant for T-helper type 1 cells, which are central to the IFN- γ -dependent pro-inflammatory response to intracellular bacteria and parasites, and has been recently reported to represent a marker of trained immunity after BCG vaccination [24]. In response to Gram-negative bacteria and LPS, BCG-vaccinated infants had increased IL-8 production. IL-8 is central in the innate response to bacterial infection by promoting neutrophil chemotaxis and macrophage activation. Smith *et*

al reported that, as in our study, BCG-vaccinated infants had significantly higher IL-8 production following stimulation with LPS compared to BCG-naïve infants [23]. It remains to be seen whether variable ‘pathogen-specific’ *in vitro* patterns of innate immune response following BCG-vaccination correlate with clinical protection.

For the majority of heterologous stimulants, IFN- γ responses in our study were characterised by a significantly lower proportion of IFN- γ responders in the BCG-vaccinated group, which was responsible for the lower IFN- γ GMR in these infants compared to BCG-naïve infants. Variable production of IFN- γ between individuals and between populations is well described [13, 20, 25]. This variability in individual IFN- γ responsiveness (which may be genetically or environmentally determined [26, 27]) is a potential modifier of the response to BCG. In a group of BCG-naïve, TST negative, adult volunteers in the Netherlands who received BCG vaccination, ‘non-responders’ were classified as individuals with no IFN- γ response to PPD and expression of CD8⁺ regulatory T-cells which persisted up to one year post-vaccination [28]. Fletcher *et al.* reported differential immune responses following BCG vaccination in infants characterised by differences in monocyte:T-cell ratios and alternatively activated macrophage phenotype (M2) [29]. More recently, Walk *et al* reported two distinct patterns of immune response to human malaria challenge in BCG-vaccinated volunteers. A subgroup of volunteers with strong monocyte and lymphocyte activation were also characterised by increased IFN- γ , granzyme-B production and CRP production [30]. This study is the first to report significant changes in IFN- γ responsiveness in the context of BCG-induced heterologous immunity in infants.

IP-10, IL-1Ra and IL-1 β were the only other cytokines which had a subgroup of non-responders. This is notable in light of the functional relationship of these cytokines to IFN- γ

and their role in the control of inflammation and pathogen persistence following mycobacterial infection [31, 32]. The IL-1 signalling axis is central in mediating the response to BCG vaccination in the setting of both mycobacterial and heterologous infections, and IL-1 β is a crucial component for the induction of trained immunity [24, 33-35]. However, in our study the proportion of IL-1Ra and IL-1 β responders was not significantly different between BCG-vaccinated and BCG-naïve infants. Moreover, sensitivity analysis confirmed that increased IL-1Ra responses following stimulation of TLR 1/2 (Pam3CYSK4) seen in BCG-vaccinated infants was unaffected by removal of ‘non-responders’ from the analysis.

In our previous study, we found that infant sex influenced the effect of BCG vaccination on the production of MIF following heterologous stimulation [17]. In the present study, we observed enhanced *E. coli*, *H. influenzae*, LPS and R848-induced production of the macrophage derived pro-inflammatory cytokines IL-6 and IL-1 β in BCG-vaccinated girls (in contrast to boys). This is interesting in light of the suggestion that beyond the neonatal period, the beneficial effects of BCG vaccination on mortality is stronger in girls [36].

Maternal BCG vaccination status is associated with increased BCG scar formation and decreased all-cause mortality in BCG-vaccinated infants [8]. Mawa *et al.* reported altered cytokine responses to TLR ligands, including LPS, in the cord blood of infants of BCG-vaccinated mothers [37]. Our finding, that infants of BCG-vaccinated mothers had altered cytokine production following heterologous stimulation, independent of infant BCG vaccination status, raises the possibility that maternal BCG vaccination ‘primes’ the response to infant BCG vaccination. Further, an interaction between maternal and infant BCG vaccination on IFN- γ production in response to *S. pneumoniae* was also found in our previous study; however maternal BCG had the opposite effect at this time point [17]. This finding is

particularly intriguing in light of the finding that BCG vaccine protects infants against respiratory infections only in infants whose mothers were also BCG-vaccinated [9, 38].

There is longstanding interest in the variability of BCG vaccine efficacy against tuberculosis infection in different geographic locations. To date, clinically significant heterologous effects of BCG have mainly been documented in African populations [36]. Following BCG vaccination, infant cytokine responses to mycobacteria differ between African and European populations, however it remains uncertain whether this is due to genetic or environmental influences [13, 14, 39]. In this study, the majority of participants were Caucasian precluding subgroup analysis by ethnicity. Exposure to environmental mycobacteria and chronic helminth infection has been associated with altered BCG-induced IFN- γ production in some populations [39]. It is unlikely that exposure to infection was a confounder in our study as proxy measures of exposure to common childhood infections (attendance at childcare and number of children under five years in the household) were similar between groups.

The strengths of this study include its large sample size and the use of samples from participants from a randomised controlled trial. The use of a large number of stimulants and multiplex cytokine analysis enabled a comprehensive interrogation of the immune response. In addition, we used multivariable analysis to identify potential confounders and a highly standardised laboratory protocol to reduce variation. Furthermore, we used the unstimulated sample as a co-variate to control for inter-individual variability in cytokine responses. A sensitivity analysis, excluding the use of unstimulated samples as a co-variate in samples where there was an independent effect on the constitutive expression of a cytokine, showed this did not lead to overcorrection (data not shown). Our study has several limitations which are common to research of this type. Firstly, for some cytokine/stimulant pairs, there were a

lower numbers of participants from whom samples were available. However, this is a large study and our statistical analysis accounted for the varying distributions of individual cytokine/stimulant pair data. Secondly, while we tested for several potential confounders selected due to their biological plausibility and previously published data, it is possible there are others that may have influenced our findings. Thirdly, there was a risk of type 1 error due to multiple comparisons. Statistical correction for multiple comparisons was not done as this was an exploratory analysis that aimed to find consistent patterns in BCG-induced alterations in cytokine responses. In addition, we have shown all analyses in their entirety to aid interpretation of the results in context. Finally, routine vaccines from the Australian immunisation schedule differ to those administered in low and middle-income countries.

In summary, at 7 months of age, following neonatal BCG vaccination, infants had lower IFN- γ responses to heterologous stimulants and TLR ligands compared with BCG-naïve infants. This results from a reduction in the proportion of infants who responded by producing IFN- γ rather than a reduced level of IFN- γ production by all participants. This study is the first to report divergent IFN- γ responses between mycobacterial and heterologous stimulation following BCG-vaccination. BCG-vaccination is associated with variable patterns of chemokine response depending on the stimulating pathogen. There was a significant interaction between both sex and maternal BCG vaccination and neonatal BCG vaccination. Relating the immunological effects of BCG observed in *in vitro* studies with clinical outcome should be a priority for future research.

Table & Figure Legends

Table 1: Demographic features of the study participants. Routine immunisations: Infanrix Hexa™ (hepatitis B, diphtheria-tetanus-pertussis, *H. influenzae* type B, inactivated poliovirus), Prevenar 13™ (13-valent pneumococcal vaccine), rotavirus (Rotarix™ or RotaTeq™).

Figure 1: Flow diagram showing participant inclusion. Table shows number of individual stimulations done for each stimulant.

Table 2: Univariate analysis for the effect of neonatal BCG vaccination versus no BCG vaccination on unstimulated cytokine levels. Results shown as GMR (95% CI) for normally-distributed data and difference in medians (95% CI) from non-parametric quantile regression for data which was not normally distributed.

Figure 2: The effect of neonatal BCG vaccination versus no BCG vaccination on cytokine responses to heterologous stimulants. Significant results $p < 0.05$ are depicted in black. $GMR > 1.0$ indicates cytokine levels were higher in BCG-vaccinated infants compared with BCG-naïve infants. Data which were not normally distributed were analysed using quantile regression and differences in medians (95%CI). As this only applied to IFN- γ , to aid interpretation, all IFN- γ analyses are displayed on stand-alone axes irrespective of the type of analysis done.

Figure 3: Proportion of BCG-vaccinated and BCG-naïve participants with production of IFN- γ in response to heterologous stimulation ('responders') (Fisher's exact test). Data for R848 not included as IFN- γ responses were above the LLD for all participants.

Figure 4: Subgroup analysis for the effect of infant sex and maternal BCG vaccination on *in vitro* cytokine responses. Panels a-c: IL-6 (a), IL-1 β (b), and remaining cytokine/stimulant pairs (c) with a significant interaction between sex and BCG vaccination. Geometric mean ratios (GMR) & 95% confidence limits (CI) are shown for the effect of infant BCG vaccination on (i) all study participants; (ii) in girls; (ii) in boys. Panel d): cytokine/stimulant pairs with a significant interaction between maternal BCG vaccination and BCG vaccination. GMR & 95%CI are shown for the effect of infant BCG vaccination on (i) all study participants; (ii) infants whose mothers were BCG-vaccinated; (iii) infants whose mothers were BCG-naïve. GMR > 1 indicates cytokine production is higher in BCG-vaccinated infants.

Funding

This work was supported by the Australian National Health and Medical Research Council (NHMRC) (grants GNT1051228 and GNT1099680), the University of Melbourne (International Research Scholarship to BF), the European Society of Paediatric Infectious Diseases (Fellowship to BF) and the Clifford Craig Foundation. MGN was supported by a Spinoza grant of the Netherlands Organization for Scientific Research.

Acknowledgements

We thank Judith Spotswood for collection of blood samples, Pedro Ramos for help with processing of blood samples and Casey Goodall for help with data entry. We also thank Dr Laure Pittet for his helpful comments on the manuscript.

Conflict of interest

The authors state that they have no conflict of interest.

References

1. Higgins JP, Soares-Weiser K, Lopez-Lopez JA, et al. Association of BCG, DTP, and measles containing vaccines with childhood mortality: systematic review. *BMJ* **2016**; 355:i5170.
2. Pollard AJ, Finn A, Curtis N. Non-specific effects of vaccines: plausible and potentially important, but implications uncertain. *Archives of disease in childhood* **2017**; 102:1077-81.
3. Aaby P, Roth A, Ravn H, et al. Randomized trial of BCG vaccination at birth to low-birth-weight children: beneficial nonspecific effects in the neonatal period? *J Infect Dis* **2011**; 204:245-52.
4. Biering-Sorensen S, Aaby P, Napirna BM, et al. Small randomized trial among low-birth-weight children receiving bacillus Calmette-Guerin vaccination at first health center contact. *Pediatr Infect Dis J* **2012**; 31:306-8.
5. Hollm-Delgado MG, Stuart EA, Black RE. Acute lower respiratory infection among Bacille Calmette-Guerin (BCG)-vaccinated children. *Pediatrics* **2014**; 133:e73-81.
6. de Castro MJ, Pardo-Seco J, Martinon-Torres F. Nonspecific (Heterologous) Protection of Neonatal BCG Vaccination Against Hospitalization Due to Respiratory Infection and Sepsis. *Clin Infect Dis* **2015**; 60:1611-9.
7. Thostesen LM, Kjaergaard J, Pihl GT, et al. Neonatal BCG vaccination and atopic dermatitis before 13 months of age: A randomized clinical trial. *Allergy* **2018**; 73:498-504.
8. Berendsen MLT, Oland CB, Bles P, et al. Maternal Priming: Bacillus Calmette-Guerin (BCG) Vaccine Scarring in Mothers Enhances the Survival of Their Child With a BCG Vaccine Scar. *J Pediatric Infect Dis Soc* **2019**.
9. Kjaergaard J, Birk NM, Nissen TN, et al. Nonspecific effect of BCG vaccination at birth on early childhood infections: a randomized, clinical multicenter trial. *Pediatr Res* **2016**; 80:681-5.
10. Kandasamy R, Voysey M, McQuaid F, et al. Non-specific immunological effects of selected routine childhood immunisations: systematic review. *BMJ* **2016**; 355:i5225.
11. Freyne B, Donath S, Germano S, et al. Neonatal BCG Vaccination Influences Cytokine Responses to Toll-like Receptor Ligands and Heterologous Antigens. *J Infect Dis* **2018**; 217:1798-808.
12. Kleinnijenhuis J, Quintin J, Preijers F, et al. Long-lasting effects of BCG vaccination on both heterologous Th1/Th17 responses and innate trained immunity. *J Innate Immun* **2014**; 6:152-8.
13. Lalor MK, Floyd S, Gorak-Stolinska P, et al. BCG vaccination induces different cytokine profiles following infant BCG vaccination in the UK and Malawi. *J Infect Dis* **2011**; 204:1075-85.
14. Lalor MK, Smith SG, Floyd S, et al. Complex cytokine profiles induced by BCG vaccination in UK infants. *Vaccine* **2010**; 28:1635-41.
15. Flanagan KL, van Crevel R, Curtis N, Shann F, Levy O. Heterologous ("nonspecific") and sex-differential effects of vaccines: epidemiology, clinical trials, and emerging immunologic mechanisms. *Clin Infect Dis* **2013**; 57:283-9.
16. Messina NL, Gardiner K, Donath S, et al. Study protocol for the Melbourne Infant Study: BCG for Allergy and Infection Reduction (MIS BAIR), a randomised controlled trial to determine the non-specific effects of neonatal BCG vaccination in a low-mortality setting. *BMJ Open* **2019**; 9:e032844. doi: 10.1136/bmjopen-2019-.
17. Freyne B, Donath S, Germano S, et al. Neonatal BCG vaccination influences cytokine responses to Toll-like receptor ligands and heterologous antigens. *J Infect Dis* **2018**.

18. Burl S, Adetifa UJ, Cox M, et al. Delaying bacillus Calmette-Guerin vaccination from birth to 4 1/2 months of age reduces postvaccination Th1 and IL-17 responses but leads to comparable mycobacterial responses at 9 months of age. *J Immunol* **2010**; 185:2620-8.
19. Corral-Fernandez NE, Cortez-Espinosa N, Salgado-Bustamante M, et al. Induction of transcription factors, miRNAs and cytokines involved in T lymphocyte differentiation in BCG-vaccinated subjects. *Mol Immunol* **2016**; 77:44-51.
20. van den Biggelaar AH, Prescott SL, Roponen M, et al. Neonatal innate cytokine responses to BCG controlling T-cell development vary between populations. *J Allergy Clin Immunol* **2009**; 124:544-50, 50.e1-2.
21. Zufferey C, Germano S, Dutta B, Ritz N, Curtis N. The contribution of non-conventional T cells and NK cells in the mycobacterial-specific IFN γ response in Bacille Calmette-Guerin (BCG)-immunized infants. *PLoS One* **2013**; 8:e77334.
22. Nissen TN, Birk NM, Blok BA, et al. Bacillus Calmette-Guerin vaccination at birth and in vitro cytokine responses to non-specific stimulation. A randomized clinical trial. *Eur J Clin Microbiol Infect Dis* **2018**; 37:29-41.
23. Smith SG, Kleinnijenhuis J, Netea MG, Dockrell HM. Whole Blood Profiling of Bacillus Calmette-Guerin-Induced Trained Innate Immunity in Infants Identifies Epidermal Growth Factor, IL-6, Platelet-Derived Growth Factor-AB/BB, and Natural Killer Cell Activation. *Front Immunol* **2017**; 8:644.
24. Joosten SA, van Meijgaarden KE, Arend SM, et al. Mycobacterial growth inhibition is associated with trained innate immunity. *J Clin Invest* **2018**; 128:1837-51.
25. Djuardi Y, Sartono E, Wibowo H, Supali T, Yazdanbakhsh M. A longitudinal study of BCG vaccination in early childhood: the development of innate and adaptive immune responses. *PLoS One* **2010**; 5:e14066.
26. Finan C, Ota MO, Marchant A, Newport MJ. Natural variation in immune responses to neonatal Mycobacterium bovis Bacillus Calmette-Guerin (BCG) Vaccination in a Cohort of Gambian infants. *PLoS One* **2008**; 3:e3485.
27. Newport MJ, Goetghebuer T, Weiss HA, Whittle H, Siegrist CA, Marchant A. Genetic regulation of immune responses to vaccines in early life. *Genes Immun* **2004**; 5:122-9.
28. Boer MC, Prins C, van Meijgaarden KE, van Dissel JT, Ottenhoff TH, Joosten SA. Mycobacterium bovis BCG Vaccination Induces Divergent Proinflammatory or Regulatory T Cell Responses in Adults. *Clin Vaccine Immunol* **2015**; 22:778-88.
29. Fletcher HA, Filali-Mouhim A, Nemes E, et al. Human newborn bacille Calmette-Guerin vaccination and risk of tuberculosis disease: a case-control study. *BMC Med* **2016**; 14:76.
30. Walk J, de Bree LCJ, Graumans W, et al. Outcomes of controlled human malaria infection after BCG vaccination. *Nat Commun* **2019**; 10:874.
31. Bohrer AC, Tocheny C, Assmann M, Ganusov VV, Mayer-Barber KD. Cutting Edge: IL-1R1 Mediates Host Resistance to Mycobacterium tuberculosis by Trans-Protection of Infected Cells. *J Immunol* **2018**; 201:1645-50.
32. Mayer-Barber KD, Andrade BB, Oland SD, et al. Host-directed therapy of tuberculosis based on interleukin-1 and type I interferon crosstalk. *Nature* **2014**; 511:99-103.
33. Arts RJW, Moorlag S, Novakovic B, et al. BCG Vaccination Protects against Experimental Viral Infection in Humans through the Induction of Cytokines Associated with Trained Immunity. *Cell Host Microbe* **2018**; 23:89-100.e5.
34. Koeken V, Verrall AJ, Netea MG, Hill PC, van Crevel R. Trained innate immunity and resistance to Mycobacterium tuberculosis infection. *Clin Microbiol Infect* **2019**.
35. Moorlag S, Roring RJ, Joosten LAB, Netea MG. The role of the interleukin-1 family in trained immunity. *Immunol Rev* **2018**; 281:28-39.

36. Biering-Sorensen S, Jensen KJ, Monterio I, Ravn H, Aaby P, Benn CS. Rapid Protective Effects of Early BCG on Neonatal Mortality Among Low Birth Weight Boys: Observations From Randomized Trials. *J Infect Dis* **2018**; 217:759-66.
37. Mawa PA, Webb EL, Filali-Mouhim A, et al. Maternal BCG scar is associated with increased infant proinflammatory immune responses. *Vaccine* **2017**; 35:273-82.
38. Stensballe LG, Ravn H, Birk NM, et al. BCG Vaccination at Birth and Rate of Hospitalization for Infection Until 15 Months of Age in Danish Children: A Randomized Clinical Multicenter Trial. *Journal of the Pediatric Infectious Diseases Society* **2018**.
39. Weir RE, Black GF, Nazareth B, et al. The influence of previous exposure to environmental mycobacteria on the interferon-gamma response to bacille Calmette-Guerin vaccination in southern England and northern Malawi. *Clin Exp Immunol* **2006**; 146:390-9.