UNIVERSITI TEKNOLOGI MARA

Í

SYNTHESIS AND CHARACTERIZATION OF PALLADIUM(II) AND NICKEL(II) SCHIFF BASE COMPLEXES FOR HOMOGENOUS CATALYTIC AND ANTICANCER STUDIES

SHAHRUL NIZAM BIN AHMAD

Thesis submitted in fulfillment of the requirements for the degree of **Doctor of Philosophy** (Chemistry)

Faculty of Applied Sciences

October 2018

CONFIRMATION BY PANEL OF EXAMINERS

I certify that a Panel of Examiners has met on 20th August 2018 to conduct the final examination of Shahrul Nizam bin Ahmad in his **Doctor of Philosophy** thesis entitled "Synthesis and Characterization of Palladium(II) and Nickel(II) Schiff Base Complexes for Homogenous Catalytic and Anticancer Studies" in accordance with Universiti Teknologi MARA Act 1976 (Akta 173). The Panel of Examiner recommends that the student be awarded the relevant degree. The Panel of Examiners was as follows:

Muhd Fauzi Safian, PhD Associate Professor Faculty of Applied Sciences Universiti Teknologi MARA (Chairman)

Datin Rohaya Ahmad, PhD Professor Faculty of Applied Sciences Universiti Teknologi MARA (Internal Examiner)

Mustaffa Shamsuddin, PhD Professor Faculty of Applied Sciences Universiti Teknologi Malaysia (External Examiner)

Djulia Onggo, PhD Professor Faculty of Applied Sciences Institut Teknologi Bandung (External Examiner)

PROF SR TS DR HAJI ABDUL HADI HAJI NAWAWI Dean Institute of Graduates Studies Universiti Teknologi MARA Date: 22 October 2018

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of Universiti Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of Student	:	Shahrul Nizam bin Ahmad	
Student I.D. No.	:	2014818278	
Programme	:	Doctor of Philosophy (Chemistry)- AS950	
Faculty	:	Applied Sciences	
Thesis Title	:	Synthesis and Characterization of Palladium(II) and Nickel(II) Schiff Base Complexes for Homogenous Catalytic and Anticancer Studies	
Signature of Student	:	8W	

Date : 22 October 2018

ABSTRACT

This thesis reports the synthesis, characterization, catalytic investigation as well as anticancer screening of palladium(II) and nickel(II) Schiff base complexes in homogenous copper-free Sonogashira reaction. Two groups of symmetrical tetradentate ONNO Schiff bases namely L1 and L2 series were successfully prepared from condensation reaction between salicylaldehyde and its derivatives with two different primary amines namely ortho-phenylenediamine and 2,2-dimethyl-1,3-propanediamine, in 2:1 ratio. The synthesized ligands were reacted with palladium(II) acetate and nickel(II) acetate tetrahydrate in 1:1 ratio, vielding palladium(II) and nickel(II) Schiff base complexes. A total of 12 ligands and 17 complexes were successfully isolated and characterized through elemental analysis, melting point, ¹H and ¹³C Nuclear Magnetic Resonance (NMR), Fourier Transform Infrared (FTIR), UV-Visible spectroscopy and molar conductivity and magnetic susceptibility studies. All spectral and physicochemical data supported the proposed chemical structures. Low molar conductivity values suggested non-electrolytic nature of all complexes. The diamagnetism of all complexes suggested their square planar geometry. This was corroborated by single crystal X-ray crystallography which revealed the square planar geometry of PdL2F and PdL2OMe with crystal structures showing the tetradentate dinitrogen dioxygen [ONNO] nature of L1H, L1C and L2F. Coordination to metal centres occur through two phenolic oxygen and two imine nitrogen donor atoms, assenting to the indication through FTIR, NMR and UV-Visible spectroscopy. Palladium(II) complexes of the L2 series were employed to find the optimum reaction conditions for copper-free Sonogashira reaction by varying type of bases, amount of catalysts, and temperature. Copper salt was not used as co-catalyst to avoid the possible homocoupling of diphenylacetylene. All palladium(II) and nickel(II) complexes were employed as catalysts in the reaction after optimization. All palladium(II) complexes were found to be effective as catalysts with 100% conversion of iodobenzene after 3 hours of reaction. Nickel(II) complexes revealed lower catalytic activities than palladium(II) complexes, with fairly moderate efficiency of 91% conversion after 12 hours of reaction. From GC-FID, a single peak of the product, diphenylacetylene, was detected at 11.69 min with no extra peaks, suggesting the absence of homocoupling products. The product of the catalysed reaction was isolated with a yield of 38% and successfully characterized via ¹H and ¹³C Nuclear Magnetic Resonance (NMR) and Fourier Transform Infrared (FTIR) spectroscopy. All compounds were also tested against human colon cancer cell lines (HCT116) where PdL2H showed the highest potency with IC_{50} of 0.60 µg/ml, lower than that of the standard drug used in this study, 5-FU with IC₅₀ of 1.70 μ g/ml. A moderate binding affinity was observed between PdL2H with DNA with a binding constant of $5.715 \times 10^5 \text{ M}^{-1}$. Hypochromism value of 24-33% indicated intercalation mode of interaction with π - π end stacking of the complex with DNA.

TABLE OF CONTENT

		Page
CON	ii	
AUTHOR'S DECLARATION		
ABSTRACT		
ACK	v	
TABLE OF CONTENT		
LIST OF TABLES		xii
LIST	OF FIGURES	xiv
LIST	OF SYMBOLS	xviii
LIST	OF ABBREVIATIONS	XX
CHA	TER ONE: INTRODUCTION	1
1.1	1	
	1.1.1 Schiff Bases	1
	1.1.1.1 Formation of Schiff base	2
	1.1.1.2 Mechanism of Schiff base formation	3
	1.1.2 Palladium	3
	1.1.3 Nickel	4
	1.1.4 Catalytic Studies	5
	1.1.4.1 Sonogashira Coupling Reaction	5
	1.1.5 Anticancer Studies	6
	1.1.5.1 Colorectal Cancer	6
	1.1.5.2 Metal Complexes as Anticancer Drugs	6
1.2	Problem Statement	6
1.3	Objectives	8
1.4	Scope and Limitation	8
1.5	Significance of Study	9
CHA	TER TWO: LITERATURE REVIEW	10
2.1	Schiff Base and Metal Complexes	10