



Short Note

2-Hydroxy-*N'*-(4-Fluorobenzoyl)Benzohydrazide

Harry Santosa, Tegar Achsendo Yuniarta *, Dini Kesuma and Galih Satrio Putra

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Surabaya, Jalan Raya Kali Rungkut, Surabaya 60293, Indonesia; harry_s@staff.ubaya.ac.id (H.S.); dinikesuma@gmail.com (D.K.); galih_satrio_putra@yahoo.co.id (G.S.P.)

* Correspondence: tegar.achsendo@staff.ubaya.ac.id

Received: 7 November 2019; Accepted: 18 December 2019; Published: 24 December 2019



Abstract: 2-Hydroxy-*N'*-(4-fluorobenzoyl)benzohydrazide was synthesized in two steps using methyl salicylate as the starting material. The reaction took place via microwave-aided hydrazinolysis, followed by acylation using 4-fluorobenzoyl chloride at low temperature to yield the target compound.

Keywords: benzoylsalicylhydrazide; methyl salicylate; microwave-assisted synthesis; nucleophilic substitution


1. Introduction

Salicylate-containing folk medicine has been used to treat pain and reduce fever since two millennia ago [1]. However, it was not until the identification and characterization of its chemical constituent in the 19th century that this class of compounds transformed into modern therapeutic use for various medical indications and for cosmetic usage on a global scale [2,3]. In order to improve its biological activity, several attempts to modify the structure has been done with varying degrees of success [4]. Some of derivatives even reveals another possible biological mode of action, such as salicylhydrazide which is known to possess inhibitory activity of HIV-integrase 1 [5–7]. Based on this notion, synthesis of 4-fluorobenzoyl derivative of salicylhydrazide with methyl salicylate as starting material was performed in this study. This ester is contained abundantly in *Gaultheria* sp. which can be found in several regions of Indonesia [8,9]. The reaction was performed according to previous research [10] and the target compound (2-hydroxy-*N'*-(4-fluorobenzoyl)benzohydrazide) was characterized using a combination of IR, NMR, and MS methods to establish its chemical structure.

2. Results and Discussion


Target compound 2-hydroxy-*N'*-(4-fluorobenzoyl)benzohydrazide (3) was synthesized in two-steps reaction (Scheme 1). Initially methyl salicylate was hydrazinolysed with the aid of microwave irradiation for three minutes. Hydrazinolysis of esters has been long used as a mean to produce hydrazide [11], although this reaction has a drawback in time efficiency, especially for unreactive esters [12,13]. However, recent developments showed that microwave-assisted synthesis has reduced the reaction time significantly while slightly improving product yield for not only esters [10,14,15], but also carboxylic acid containing starting materials [16,17]. After simple work-up procedure and recrystallization using ethanol, salicylhydrazide (2) was obtained with decent yield of 61.1%. This result provided further evidence of efficiency in the synthesis of benzohydrazide using microwave irradiation compared to the more conventional method. For example, it took approximately 6 hours of heating under reflux to complete the synthesis of various benzohydrazide from their respective methyl or ethyl ester of benzoic acid, with the overall yield ranging from 40 to 67% [12]. Another method which employed the addition of coupling agent under milder temperature showed an improvement in the reaction rate by three times for methyl benzoate, giving yield of 74% [13]. However, microwave-aided

synthesis has significantly speeded up the reaction to the extent where it is completed in less than 20 min. This was proven by previous researches [10,14,15] and this study, all of which used methyl salicylate as starting material. Furthermore it can be observed that the increase in microwave power employed in reaction correlates with the increase in reaction rate.




Scheme 1. Synthesis of 2-hydroxy-*N'*-(4-fluorobenzoyl)benzohydrazide.

The following step was acylation of **2** with 4-fluorobenzoyl chloride. This reaction generally yields 1,2-diacylhydrazine spontaneously under various organic solvents [18,19]. However, care must be taken with acyl chloride group since it tends to produce disubstituted hydrazine as major product. Therefore, low temperature was implemented (0–5 °C) to suppress the reactivity of the functional group, ensuring 1,2-diacylhydrazine as the sole product [11]. Both reactions took place via nucleophilic substitution (Schemes 2 and 3). After 30 min, similar work-up procedure was performed to obtain **3** with a yield of 50%. Overall, the yield obtained from this two-steps reaction was 30.55%. The structure of **3** was verified by IR, NMR, and MS data.



Scheme 2. Reaction mechanism of synthesis of salicylhydrazide (**2**).




Scheme 3. Reaction mechanism of synthesis of 2-hydroxy-*N'*-(4-fluorobenzoyl)benzohydrazide (**3**).

Infrared spectrum showed strong absorption peak of two amide carbonyls at around 1635 and 1660 cm^{-1} , and the -NH peak at 3317 cm^{-1} (Supplementary Figure S1). This finding corresponds the characteristic band of dibenzoylhydrazine compound [20], despite the presence of only one single absorption peak of -NH due to overlapping with -OH broad peak at the same region (3300–3400 cm^{-1}) [21]. ^{13}C -NMR indicated the presence of two carbonyls at 166 and 168 ppm (Supplementary Figure S2). In addition, it is also observed the doublet splitting patterns of all carbon peaks of *p*-fluorobenzoyl moiety. This is due to the carbon-fluorine coupling of the carbon ipso to the fluorine atom up to the para position. Calculation of their coupling constant give the value of 252.5, 20.2, 10.1, and 3.0 Hz for $J_{\text{C1-F}}$, $J_{\text{C2-F}}$, $J_{\text{C3-F}}$, and $J_{\text{C4-F}}$ respectively (Supplementary Figures S2, S12, and S13). These are approximately comparable to the data published for compounds

bearing p-fluorocarbonyl moiety such as p-fluorobenzaldehyde [22] and p-fluoroacetophenone [23]. Meanwhile, $^1\text{H-NMR}$ spectrum results showed the existence of two peaks of -NH indicating the formation of hydrazide group. It is found that these peaks were observed clearly (10.66 and 10.69 ppm) under temperature slightly above the freezing point of the solvent ($\text{DMSO-}d_6$), which in this case was $22\text{ }^\circ\text{C}$. On the other hand, proton spectrum analysis under room temperature only showed a broad singlet peak due to rapid exchange phenomenon (Supplementary Figures S3 and S4). The amine and hydroxy proton assignment was concluded with the aid of Double Pulsed Field Gradient Spin Echo-NOE (DPFGSE-NOE) analysis. It is shown that proton peak at 10.66 ppm belongs to the -NH group vicinal to salicylate carbonyl group, while the other -NH proton has the chemical shift of 10.69 ppm (Supplementary Figure S5–S7). Furthermore, various 2D-NMR methods ($^1\text{H-}^1\text{H}$ COSY, $^1\text{H-}^{13}\text{C}$ HMQC, and HMBC) were carried out in order to verify the hydrogen and carbon peak assignment, thus determining the chemical structure of target molecule (Table 1 and Supplementary Figures S8–S10). Ultimately, mass spectrum data indicated the presence of the target compound which possesses m/z value of 275.0848 ($[\text{M} - \text{H}]^+$) (Supplementary Figure S11).

Table 1. Cross-peaks in COSY, DPGSE-NOE, HMBC, and HMQC of compound (3).



No *	HMQC		COSY/ DPFGSE-NOE	HMBC
	H-NMR	C-NMR		
1	7.89 (dd, $J = 7.9, 1.7$ Hz, 1 H)	128.9	H2	C3; C5; C7
2	6.93 (m, 1 H)	119.4	H1; H3	C1; C4; C6
3	7.43 (ddd, $J = 8.3, 7.2,$ 1.7 Hz, 1 H)	134.7	H2; H4	C1; C5
4	6.95 (m, 1 H)	117.9	H3	C2; C5; C6
5	-	159.8	-	-
6	-	115.2	-	-
7	-	168.2(C = O)	-	-
8	10.66(NH)	-	H1	-
9	10.69(NH)	-	H16/H12	-
10	-	165.0 (C = O)	-	-
11	-	129.3 (d, $J = 3.0$ Hz, 1C)	-	-
12	7.97 (m, 1 H)	130.8 (d, $J = 10.1$ Hz, 1C)	H13	C10; C14; C16
13	7.34 (m, 1 H)	116.1 (d, $J = 20.2$ Hz, 1C)	H12	C11; C15
14	-	164.8 (d, $J = 252.5$ Hz, 1C)	-	-
15	7.34 (m, 1 H)	116.1 (d, $J = 20.2$ Hz, 1C)	H16	C11; C13
16	7.97 (m, 1 H)	130.8 (d, $J = 10.1$ Hz, 1C)	H15	C10; C12; C14
17	11.88(OH)	-	H4	-

* Numbering of the compound only for the purpose of structure determination and does not represent IUPAC nomenclature.

In summary, it is found that 2-hydroxy-*N'*-(4-fluorobenzoyl)benzohydrazide can be synthesized using two steps reaction in a time efficient manner.

3. Materials and Methods

3.1. Instruments

All reagents and solvents were purchased from Sigma Aldrich and Merck (Singapore), aside from methyl salicylate which was obtained from local chemical supplier with the purity level of technical grade. Infrared spectra were measured using JASCO FT/IR-4200, provided by Research Laboratory, Faculty of Pharmacy, University of Surabaya. NMR spectra measurements were conducted using JEOL ECS-400 spectrophotometer (400 MHz for proton, 101 MHz for carbon) in Institute of Tropical Disease, Airlangga University. NMR data are reported as follows (chemical shift (δ , in ppm), multiplicity, coupling constant (J , in Hz), integration). $^1\text{H-NMR}$ and DPGSE-NOE spectra were recorded in the controlled temperature of 22 °C, while the others were analyzed under room temperature. HRMS-TOF spectra were recorded in Padjadjaran University. Commercial microwave (LG MH6548FR) was employed for hydrazinolysis step. Thin Layer Chromatography was used to monitor the completion of reaction (Type 60 F254, Merck). Detection of the spots was carried out using three different eluent combinations (Acetone-Ethyl Acetate 2:1; Acetone-Chloroform 2:1; Chloroform-Ethyl Acetate 1:1) and visualized under 254 nm UV lamp. Melting point determination was performed using Sybron-Thermolyne-MP12615. Synthesis procedure was performed in the Laboratory of Organic Chemistry, Faculty of Pharmacy, University of Surabaya.

3.2. Synthesis

The synthesis was performed according to previously published work [10]. 20 mmol of methyl salicylate (2.6 mL) (**1**) was mixed with 80 mmol of hydrazine hydrate 80% (3.2 mL) and irradiated in microwave (360 W). Reaction completion was monitored using TLC. After 3 min the reaction was deemed complete. The mixture was then washed with distilled water and the separated solid was collected by filtration. Recrystallization with ethanol 96% was performed to obtain salicylhydrazide (**2**) with the yield of 61.1%. The following step was acylation of salicylhydrazide (1.3 g) (**2**) with equimolar amount of 4-fluorobenzoyl chloride (1.0 mL) (8.5 mmol), pre-mixed in THF. The reaction was conditioned at low temperature (0–5 °C) in an ice bath. Reaction was monitored using TLC until completion after 30 min. The mixture was then washed with cold ethanol and filtered in vacuo, followed by recrystallization with ethanol 96% to obtain the target compound 2-hydroxy-*N'*-(4-fluorobenzoyl)benzohydrazide (**3**).

2-Hydroxy-N'-(4-fluorobenzoyl)benzohydrazide (**3**). 1.16 g (Overall yield: 30.55%; three replications); white crystalline; m.p.: 229–230 °C; IR: (KBr pellet) 3441 (OH), 3317 (NH), 1660, (C=O, amide), 1635 (C=O, amide) cm^{-1} ; $^1\text{H-NMR}$: (400 MHz, $\text{DMSO-}d_6$) δ 11.81 (s, 1H), 10.67 (s, 1H), 8.00–7.93 (m, 2H), 7.89 (dd, $J = 7.9, 1.7$ Hz, 1H), 7.43 (ddd, $J = 8.3, 7.2, 1.7$ Hz, 1H), 7.38–7.31 (m, 2H), 6.97–6.89 (m, 2H); $^{13}\text{C-NMR}$: (101 MHz, $\text{DMSO-}d_6$) δ 168.2, 165.0, 164.8 (d, $J = 252.5$ Hz, 1C), 159.8, 134.7, 130.8 (d, $J = 10.1$ Hz, 1C), 129.3 (d, $J = 3.0$ Hz, 1C), 128.9, 119.6, 117.9, 116.1 (d, $J = 20.2$ Hz, 1C), 115.2; $^1\text{H-}^1\text{H COSY}$, HMQC, and HMBC as presented in Table 1; MS-TOF (ESI): m/z calcd. $[\text{M} - \text{H}]^+$: 275.0832, found: 275.0848; Spectroscopic data are provided in Supplementary Materials.

Supplementary Materials: The following are available online, Figure S1: FT-IR spectrum Figure S2: $^{13}\text{C-NMR}$ spectrum, Figure S3: $^1\text{H-NMR}$ spectrum (400 MHz, 22 °C, $\text{DMSO-}d_6$), Figure S4: $^1\text{H-NMR}$ spectrum (400 MHz, 25 °C, $\text{DMSO-}d_6$), Figures S5–S7: DPGSE-NOE spectrum, Figure S8: COSY spectrum, Figure S9: HMQC spectrum, Figure S10: HMBC spectrum, Figure S11: MS spectrum, Figures S12 and S13: $^{13}\text{C-NMR}$ spectrum of p-fluorobenzoyl moiety

Author Contributions: The research was designed by H.S., D.K., and G.S.P. Funding was acquired by H.S. and T.A.Y. Synthesis was carried out by H.S. Spectroscopic data recording and analysis was performed by T.A.Y., D.K., and G.S.P. The manuscript was written by T.A.Y. and G.S.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Lembaga Penelitian dan Pengabdian Masyarakat University of Surabaya.

Acknowledgments: The authors would like to thank Institute of Tropical Disease (Airlangga University) and Padjadjaran University for the help in NMR and MS data acquisition.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lévesque, H.; Lafont, O. L'aspirine à travers les siècles: Rappels historiques. *Rev. Médecine Interne* **2000**, *21*, 8–17.
2. Wu, K.K. Salicylates and their spectrum of activity. *Anti-Inflamm. Anti-Allergy Agents Med. Chem.* **2007**, *6*, 278–292. [[CrossRef](#)]
3. Cosmetic Ingredient Review Expert Panel. Safety assessment of salicylic acid, butyloctyl salicylate, calcium salicylate, C12-15 alkyl salicylate, capryloyl salicylic acid, hexyldodecyl salicylate, isocetyl salicylate, isodecyl salicylate, magnesium salicylate, MEA-salicylate, ethylhexyl salicylate, potassium salicylate, methyl salicylate, myristyl salicylate, sodium salicylate, TEA-salicylate, and tridecyl salicylate. *Int. J. Toxicol.* **2003**, *22*, 1–108.
4. Ekinci, D.; Şentürk, M.; Küfevrioglu, Ö.İ. Salicylic acid derivatives: Synthesis, features, and usage as therapeutic tools. *Expert Opin. Ther. Pat.* **2011**, *21*, 1831–1841. [[CrossRef](#)] [[PubMed](#)]
5. Neamati, N.; Hong, H.; Owen, J.M.; Sunder, S.; Winslow, H.E.; Christensen, J.L.; Zhao, H.; Burke, T.R.; Milne, G.W.A.; Pommier, Y. Salicylhydrazine-containing inhibitors of HIV-1 integrase: Implication for a selective chelation in the integrase active site. *J. Med. Chem.* **1998**, *41*, 3202–3209. [[CrossRef](#)]
6. Neamati, N.; Lin, Z.; Karki, R.G.; Orr, A.; Cowansage, K.; Strumberg, D.; Pais, G.C.G.; Voigt, J.H.; Nicklaus, M.C.; Winslow, H.E.; et al. Metal-dependent inhibition of HIV-1 integrase. *J. Med. Chem.* **2002**, *45*, 5661–5670. [[CrossRef](#)]
7. Al-Mawsawi, L.Q.; Dayam, R.; Taheri, L.; Witvrouw, M.; Debyser, Z.; Neamati, N. Discovery of novel non-cytotoxic salicylhydrazide containing HIV-1 integrase inhibitors. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 6472–6475. [[CrossRef](#)]
8. Liu, W.R.; Qiao, W.L.; Liu, Z.Z.; Wang, X.H.; Jiang, R.; Li, S.Y.; Shi, R.B.; She, G.M. Gaultheria: Phytochemical and pharmacological characteristics. *Molecules* **2013**, *18*, 12071–12108. [[CrossRef](#)]
9. Sadili, A.; Kartawinata, K.; Kartonegoro, A.; Soedjito, H.; Sumadijaya, A. Floristic composition and structure of subalpine summit habitat of Mt. Gede-Pangrango Complex, Cibodas Biosphere Reserve, West Java, Indonesia. *Reinwardtia* **2009**, *12*, 393–404.
10. Santosa, H.; Putra, G.S.; Yuniarta, T.A.; Budiati, T. Synthesis and molecular docking studies of N'-benzoylsalicylhydrazide derivatives as antituberculosis through InhA enzyme inhibition. *Indones. J. Pharm.* **2018**, *29*, 198–205. [[CrossRef](#)]
11. Smith, P.A.S. *Organic Reactions*; Adams, R., Ed.; Wiley: New York, NY, USA, 1946; Chapter 9; Volume 3, pp. 366–369.
12. Yale, H.L.; Losee, K.; Martins, J.; Holsing, M.; Perry, F.M.; Bernstein, J. Chemotherapy of experimental tuberculosis. VIII. The synthesis of acid hydrazides, their derivatives and related compounds. *J. Am. Chem. Soc.* **1933**, *75*, 1933–1942. [[CrossRef](#)]
13. Smith, R.F.; Bates, A.C.; Battisti, A.J.; Byrnes, P.G.; Mroz, C.T.; Smearing, T.J.; Albrecht, F.X. Reactions of hydrazines with esters and carboxylic acids. *J. Org. Chem.* **1968**, *33*, 851–855. [[CrossRef](#)]
14. Peng, Y.; Song, G. Simultaneous microwave and ultrasound irradiation: A rapid synthesis of hydrazides. *Green Chem.* **2001**, *3*, 302–304. [[CrossRef](#)]
15. Jain, A.K.; Gupta, P.K.; Ganesan, K.; Pande, A.; Malhotra, R.C. Rapid solvent-free synthesis of aromatic hydrazides under microwave irradiation. *Def. Sci. J.* **2007**, *57*, 267–270. [[CrossRef](#)]
16. Peng, Y.; Song, G. An efficient microwave-assisted one-pot conversion of carboxylic acids into hydrazides. *J. Chem. Res.* **2003**, *12*, 768–769. [[CrossRef](#)]
17. Saha, A.; Kumar, R.; Kumar, R.; Devakumar, C. Development and assessment of green synthesis of hydrazides. *Indian J. Chem.* **2010**, *49B*, 526–531.
18. Paulsen, H.; Stoye, D. *The Chemistry of Amides*; Zabicky, J., Ed.; Interscience: New York, NY, USA, 1970; Chapter 10; pp. 555–557.

19. Khattab, S.N. Synthesis biological activity of novel amino acid-(*N'*-benzoyl) hydrazide and amino acid-(*N'*-nicotiny)l hydrazide derivatives. *Molecules* **2005**, *10*, 1218–1228. [[CrossRef](#)]
20. Mashima, M. Infrared absorption spectra of hydrazides. I. Hydrazides of aromatic acids. *Bull. Chem. Soc. Jpn.* **1962**, *35*, 332–337. [[CrossRef](#)]
21. El-Baradie, H.Y.F.; Khattab, M.A.; Issa, R.M.; Maghrabi, J.Y. Substituent effects on the spectral behaviour and acid-base properties of arylidene derivatives of salicylic hydrazide. *J. Chem. Technol. Biotechnol.* **1983**, *33A*, 123–129.
22. Weigert, F.J.; Roberts, J.D. ¹³C nuclear magnetic resonance spectroscopy. Determination of carbon-fluorine couplings. *J. Am. Chem. Soc.* **1971**, *93*, 2361–2369.
23. Muller, N.; Carr, T.D. Carbon-13 splittings in fluorine nuclear magnetic resonance spectra. *J. Phys. Chem.* **1963**, *67*, 112–115. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Search:

[Advanced](#)

Journals / Molbank / Volume 2020

[Submit to Molbank](#)[Review for Molbank](#)

Journal Menu

- [Molbank Home](#)
- [Aims & Scope](#)
- [Editorial Board](#)
- [Instructions for Authors](#)
- [Special Issues](#)
- [Sections & Collections](#)
- [Article Processing Charge](#)

Molbank, Volume 2020 (2020)

- [Issue 1 \(March 2020\)](#)



Editorial Board

- [Editorial Board](#)
- [Natural Products Section](#)
- [Organic Synthesis Section](#)
- [Structure Determination Section](#)

Editor

Prof. Dr. René T. Boéré [E-Mail](#) [Website](#)

Editor-in-Chief

Department of Chemistry and Biochemistry, University of Lethbridge, 4401 University Drive W., Lethbridge, Alberta T1K3M4, Canada

Interests: organoelement chemistry; sulfur-nitrogen compounds; heterocycles; amidines; heteroamidines; free-radicals; electrochemistry; supramolecular chemistry; structural methods; crystallography; electron paramagnetic resonance spectroscopy



Previous Editor

Prof. Dr. Norbert Haider * [E-Mail](#) [Website](#)

Former Editor-in-Chief

Department of Pharmaceutical Chemistry, University of Vienna, Althanstrasse 14, 1090 Vienna, Austria
Tel. +438642204677

Interests: medicinal heterocyclic chemistry; pyridazines; carbazoles; nitrogen hetarenes

* former Editor-in-Chief from 2007 up to 30 September 2019; The Editorial Board of *Molecules* also governs *Molbank*. Former Editor-in-Chief: up to 2001 Dr. Luc Patiny; in 2002 Prof. Dr. Bruce A. Hathaway; in 2003 Reto Mueller (organo-chemistry.org)

[Special Issues and Collections in MDPI journals](#)



Editorial Board Members (23)

Prof. Dr. Rodrigo Abonia [E-Mail](#) [Website](#)

Department of Chemistry, Universidad del Valle, Calle 13 No 100-00, A. A. 25380, Cali, Colombia

Interests: organic synthesis, methodology and spectroscopic analysis; heterocyclic chemistry; natural products; asymmetric synthesis; nitrogenated compounds of pharmaceutical interest; multicomponent reactions

Dr. R. Alan Aitken [E-Mail](#) [Website](#)

School of Chemistry, University of St Andrews North Haugh, St Andrews Fife, KY16 9ST, UK
Tel. (01334) 463865; Fax: (01334) 463865

Interests: synthesis; synthetic use of flash vacuum pyrolysis; heterocyclic chemistry; reactive intermediates; organophosphorus; organosulfur; heavier main group chemistry



Prof. Dr. Fawaz Aldabbagh [E-Mail](#) [Website](#)

Department of Pharmacy, School of Life Sciences, Pharmacy & Chemistry, Kingston University, Penrhyn Road, Kingston upon Thames, KT1 2EE, UK

Interests: free radical organic and polymer chemistry; heterocyclic and medicinal chemistry

Special Issues and Collections in MDPI journals:

Special Issue in *Molecules*: [Free Radicals in Organic Synthesis](#)

Special Issue in *Molecules*: [Recent Advances in Nitrogen-Containing Aromatic Heterocycles](#)

Special Issue in *Molbank*: [Heterocycle Reactions](#)

Special Issue in *Molecules*: [Synthetic Heterocyclic Chemistry](#)



Prof. Dr. Ian R. Baxendale [E-Mail](#) [Website](#)

Department of Chemistry, Durham University, University Science Laboratories, South Road, Durham DH1 3LE, UK

Interests: organic synthesis and methodology; automation; chemical reactors; flow chemistry; enabling technologies; immobilised reagents and scavengers; heterocyclic chemistry



Dr. Jose Berna [E-Mail](#) [Website](#)

Department of Organic Chemistry, Faculty of Chemistry, University of Murcia, 30100 Murcia, Spain

Interests: self-assembly; mechanically interlocked molecules (rotaxanes and catenanes); hydrogen bond; template synthesis; molecular recognition; supramolecular chemistry

Prof. Dr. Fang-Rong Chang [E-Mail](#) [Website](#)

Graduate Institute of Natural Products, College of Pharmacy, Kaohsiung Medical University, No. 100, Shih-Chuan 1st Road, Kaohsiung, 80708, Taiwan

Interests: natural products chemistry; medicinal chemistry; transgenic plant (arabidopsis) reporter assay; epigenetic modulation for microbial secondary metabolites; functional food; ethnopharmacology

Prof. Dr. Gregory K. Friestad [E-Mail](#) [Website](#)

Department of Chemistry, University of Iowa, Iowa City, IA 52242 USA

Tel. 1-319-335-1364; Fax: +1 319 335 1270

Interests: asymmetric synthesis methodology; free radical reactions; organometallic reagents; natural product synthesis; asymmetric catalysis

Prof. Dr. Bartolo Gabriele [E-Mail](#) [Website1](#) [Website2](#)

Laboratory of Industrial and Synthetic Organic Chemistry (LISOC), Department of Chemistry and Chemical Technologies, University of Calabria, Via Pietro Bucci, 12/C, 87036 – Arcavacata di Rende (Cosenza), Italy

Tel. +390984492815; Fax: +390984492044

Interests: new syntheses of high value added molecules through catalytic assembly of simple units; innovative syntheses of heterocyclic molecules of pharmaceutical, agrochemical, or applicative interest; carbonylation chemistry; use of non-conventional solvents in organic synthesis; synthesis and semi-synthesis of bioactive compounds of pharmaceutical or agrochemical interest; synthesis of new materials for advanced applications; extraction, characterization, and evaluation of the biological activity of bioactive principles from natural matrices

Special Issues and Collections in MDPI journals:

Special Issue in *Molecules*: [Cascade Catalysis](#)

Special Issue in *Molecules*: [Palladium-Catalyzed Organic Synthesis: Commemorative Issue in Honor of Professor Richard Frederick Heck \(1931–2015\)](#)

Special Issue in *Molbank*: [Metal-Catalyzed Synthesis](#)

Special Issue in *Molecules*: [Organic Synthesis via Transition Metal-Catalysis](#)



Prof. Dr. Panayiotis A. Koutentis [E-Mail](#) [Website](#)

Department of Chemistry, University of Cyprus, P. O. Box 20537, 1678 Nicosia, Cyprus
Tel. 0035722892783

Interests: heterocyclic chemistry; sulfur-nitrogen heterocycles; synthetic methods; azaaocenes; zwitterionic aocenes; stable organic radicals; biologically active heterocycles; isothiazoles; 1,2,3-dithiazoles; 1,2,6-thiadiazines; 1,2,4-benzotriazines

Special Issues and Collections in MDPI journals:

Special Issue in *Molecules*: [Design and Synthesis of Novel Conjugated and Non Conjugated Small Molecules](#)

Special Issue in *Molecules*: [Sulfur-Nitrogen Heteroaromatics](#)

Special Issue in *Molbank*: [Heteroatom Rich Organic Heterocycles](#)

Special Issue in *Molecules*: [Non-Natural Multi-Heteroatom Heterocycles: New Chemical Space](#)



Prof. Dr. Conrad Kunick [E-Mail](#) [Website](#)

Institut für Medizinische und Pharmazeutische Chemie, Technische Universität Braunschweig, Beethovenstraße 55, 38106 Braunschweig, Germany

Interests: medicinal chemistry; pharmaceutical chemistry; synthesis of heterocycles; protein kinase inhibitors

Dr. Raffaella Mancuso [E-Mail](#)

Laboratory of Industrial and Synthetic Organic Chemistry (LISOC), Department of Chemistry and Chemical Technologies, University of Calabria, Via Pietro Bucci 12/C, 87036 Arcavacata di Rende (CS), Italy

Interests: innovative syntheses of high value molecules through catalytic process; new syntheses of heterocyclic compounds of pharmaceutical interest; carbonylation catalyzed chemistry; application of unconventional solvents in advanced organic synthesis; synthesis of novel materials for advanced applications

Special Issues and Collections in MDPI journals:

Special Issue in *Catalysts*: [Catalytic Carbonylation Reactions](#)



Prof. Dr. Ge Meng [E-Mail](#) [Website](#)

Chemistry Department, Fudan University, Shanghai 200433, China
Fax: +86 029 82655424

Interests: medicinal heterocyclic chemistry; indoles; indazoles; pyridines; thiazoles; thiadiazoles; pyrroles; computer aided drug design; molecular probe



Prof. Dr. Hideto Miyabe [E-Mail](#) [Website](#)

School of Pharmacy, Hyogo University of Health Sciences, 1-3-6 Minatojima, Chuo-ku, Kobe 650-8530, Japan

Interests: organic synthesis and methodology; radical reactions; organocatalysis; asymmetric catalysis; photochemistry; aryne chemistry

Special Issues and Collections in MDPI journals:

Special Issue in *Molbank*: [Molecules from Radical Reactions](#)



Prof. Dr. Luke R. Odell [E-Mail](#) [Website](#)

Organic Pharmaceutical Chemistry, Department of Medicinal Chemistry, BMC, Uppsala University, Box 574, SE-751 23 Uppsala, Sweden
Tel. +4618-471 4297

Interests: heterocyclic chemistry; multicomponent reactions; catalysis; medicinal chemistry; synthetic methodology

Special Issues and Collections in MDPI journals:

Special Issue in *Molbank*: [Molecules from Multicomponent Reactions](#)



Prof. Dr. Thierry Ollevier [E-Mail](#) [Website](#)

Department of Chemistry, Pavillon Alexandre-Vachon, 1045, Avenue de la Médecine, Université Laval, Québec, G1V 0A6, Canada

Tel. +1 418 656 5034; Fax: +1 418 656 7916

Interests: synthetic methodology; asymmetric catalysis; Lewis acids

Dr. Luc Patiny [E-Mail](#) [Website](#)

Director of Chemical Information, Institute of Chemical Sciences and Engineering (ISIC), Ecole Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland

Interests: chemoinformatics; chemical information; infrastructure for chemical research; online processing and data mining

Special Issues and Collections in MDPI journals:

Special Issue in *Challenges: Contest: Open Source Cheminformatic Tool / 9th Workshop in Chemical Information*

Prof. Dr. Vincenzo Piccialli [E-Mail](#) [Website](#)

Department of Chemical Sciences, Università degli Studi di Napoli Federico II, Via Cintia 21, 80126 Napoli, Italy

Interests: organic and medicinal chemistry; organic synthesis; catalytic oxidative processes; marine natural products; nucleosides chemistry

Special Issues and Collections in MDPI journals:

Special Issue in *Molecules: New Studies on the Synthesis of Biologically Active Products*



Prof. Dr. Oleg A. Rakitin [E-Mail](#) [Website](#)

1. N. D. Zelinsky Institute of Organic Chemistry Russian Academy of Sciences, 47 Leninsky Prospekt, 119991 Moscow, Russia

2. Nanotechnology Education and Research Center, South Ural State University, 454080 Chelyabinsk, Russia

Tel. +7-499-1355327

Interests: heterocyclic chemistry; sulfur-nitrogen heterocycles; selenium heterocycles; synthetic methods; biologically active compounds; organic sensitizers for DSSCs and OLEDs; sulfur monochloride

Special Issues and Collections in MDPI journals:

Special Issue in *Molecules: Polysulfur- and Sulfur-Nitrogen Heterocycles*



Dr. Bernd Schneider [E-Mail](#) [Website](#)

Max Planck Institute for Chemical Ecology, Hans-Knöll-Str. 8, Beutenberg Campus, 07745 Jena, Germany

Interests: natural products chemistry; chemical ecology; plant natural products; NMR of small molecules

Prof. Dr. Jonathan W. Steed [E-Mail](#) [Website](#)

Department of Chemistry, Durham University, University Science Laboratories, South Road, Durham DH1 3LE, UK

Fax: +44 191 384 4737

Interests: supramolecular chemistry; gels; molecular materials and the molecular solid state; pharmaceutical solid forms; platinum group transition metal chemistry

Prof. Dr. Ping-Jyun Sung [E-Mail](#) [Website](#)

1. National Museum of Marine Biology and Aquarium, Pingtung 944, Taiwan
2. Graduate Institute of Marine Biology, National Dong Hwa University, Pingtung 944, Taiwan
3. Department of Marine Biotechnology and Resources, National Sun Yat-sen University, Kaohsiung 804, Taiwan
4. Chinese Medicine Research and Development Center, China Medical University Hospital, Taichung 404, Taiwan
5. Graduate Institute of Natural Products, Kaohsiung Medical University, Kaohsiung 807, Taiwan

Tel. 886-8-8825037; Fax: 886-8-8825087

Interests: marine natural products; marine chemical ecology; bioactive substances from cultured marine invertebrates

Special Issues and Collections in MDPI journals:

Special Issue in *Marine Drugs*: [Development and Application of Herbal Medicine from Marine Origin](#)

Dr. Hidenori Tanaka [E-Mail](#) [Website1](#) [Website2](#)

Center for Highly Advanced Integration of Nano and Life Sciences (G-CHAIN), Gifu University, 1-1 Yanagido, Gifu, Gifu 501-1193, Japan

Special Issues and Collections in MDPI journals:

Special Issue in *Molecules*: [Carbohydrate Chemistry](#)

Special Issue in *Molbank*: [Organic Synthesis of Carbohydrates](#)



Prof. Dr. Kristof Van Hecke [E-Mail](#) [Website](#)

Universiteit Gent, Ghent, Belgium

Tel. +32(0)9264 48 70

Interests: structures; X-ray diffraction; bioinorganic chemistry; metal complexes; coordination chemistry


Special Issues and Collections in MDPI journals:

Special Issue in *Molecules*: [Metal Coordination Complexes for Biomedical Applications](#)



Table of Contents

Molbank, Volume 2020, Issue 1 (March 2020) – 5 articles

- Issues are regarded as officially published after their release is announced to the [table of contents alert mailing list](#).
- You may [sign up for e-mail alerts](#) to receive table of contents of newly released issues.
- PDF is the official format for papers published in both, html and pdf forms. To view the papers in pdf format, click on the "PDF Full-text" link, and use the free [Adobe Reader](#)  to open them.

Order results

Publication Date

Result details

Normal

Show export options 

Open Access Short Note



2-Hydroxy-*N'*-(4-Fluorobenzoyl)Benzohydrazide

by  Harry Santosa ,  Tegar Achsendo Yuniarta ,  Dini Kesuma and  Galih Satrio Putra

Molbank 2020, 2020(1), M1103; <https://doi.org/10.3390/M1103> - 24 Dec 2019

Abstract

2-Hydroxy-*N'*-(4-fluorobenzoyl)benzohydrazide was synthesized in two steps using methyl salicylate as the starting material. The reaction took place via microwave-aided hydrazinolysis, followed by acylation using 4-fluorobenzoyl chloride at low temperature to yield the target compound. [Full article](#)
(This article belongs to the Section [Organic Synthesis](#))

Open Access Short Note



Bicyclo[4.2.0]octa-1,3,5-trien-3-yl-dimethyl((*E*)-styryl)-silane

by  Konstantin S. Levchenko ,  Konstantin A. Chudov ,  Dmitri Yu. Demin ,  Pavel S. Shmelin and  Evgeny P. Grebennikov

Molbank 2020, 2020(1), M1102; <https://doi.org/10.3390/M1102> - 23 Dec 2019

Abstract

Bicyclo[4.2.0]octa-1,3,5-trien-3-yl-dimethyl-((*E*)-styryl)-silane was synthesized via three stage synthesis starting from benzocyclobutene and (2-bromo-vinyl)-benzene. The structure of the product was determined using ¹H- and ¹³C-NMR and HRMS. [Full article](#)

Open Access Short Note



Bis(*N'*-(3-chlorobenzoyl)isonicotinohydrazide)Iron(III) Complex

by  Richa Mardianingrum ,  Susanti and  Ruswanto Ruswanto

Molbank 2020, 2020(1), M1101; <https://doi.org/10.3390/M1101> - 20 Dec 2019

Abstract The bis(*N'*-(3-chlorobenzoyl)isonicotinohydrazide)iron(III) complex was synthesised from *N'*-(3-chlorobenzoyl)isonicotinohydrazide and iron(III) metal by reflux in an ethanol solution. The title compound was characterised by Fourier-transform infrared spectroscopy (FTIR) spectroscopy, differential thermal analysis/thermogravimetric analysis (DTA/TGA) and UV-visible spectroscopy. The results indicate that coordination [...] [Read more.](#)

(This article belongs to the Section [Organic Synthesis](#))

Open Access Communication



Synthesis of N-Heterocyclic Analogues of 28-O-Methyl Betulinate, and Their Antibacterial and Antifungal Properties

by Elvira R. Shakurova and Lyudmila V. Parfenova

Molbank 2020, 2020(1), M1100; <https://doi.org/10.3390/M1100> - 19 Dec 2019

Abstract The paper presents the results on the one-pot pyridine quaternization using betulinic 28-O-methyl ester (1) and Tempo⁺Br₃⁻ cation followed by reduction of the resulting salt (2) to 1,2,5,6-tetrahydropyridine derivative (3). The [...] [Read more](#).

(This article belongs to the Section [Natural Products](#))

[► Show Figures](#)

Open Access Short Note



N-(2-Hydroxy-1,1-dimethylethyl)-3-methylbenzamide

by Hamad H. Al Mamari and Yousuf Al Lawati

Molbank 2020, 2020(1), M1099; <https://doi.org/10.3390/M1099> - 19 Dec 2019

Abstract The title compound, N-(2-hydroxy-1,1-dimethylethyl)-3-methylbenzamide was synthesized by reacting 3-methylbenzoyl chloride or 3-methylbenzoic acid with 2-amino-2-methyl-1-propanol. In the present report, the synthesized target compound was fully characterized by various spectroscopic methods (¹H NMR, ¹³C NMR, IR, GC-MS), its composition confirmed [...] [Read more](#).

(This article belongs to the Special Issue [Molecules from Side Reactions](#))

[Show export options](#)