



Host Range and Symbiotic Effectiveness of N₂O Reducing *Bradyrhizobium* Strains

Kedir Woliy^{1*}, Tulu Degefu² and Åsa Frostegård^{1*}

¹ Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Ås, Norway, ² International Crops Research Institute for the Semi-Arid Tropics, Addis Ababa, Ethiopia

OPEN ACCESS

Edited by:

David Richardson,
University of East Anglia,
United Kingdom

Reviewed by:

Jose María Vinardell,
University of Seville, Spain
Petri Penttinen,
Sichuan Agricultural University, China

*Correspondence:

Kedir Woliy
kedir.wolijillo@nmbu.no
Åsa Frostegård
asa.frostegard@nmbu.no

Specialty section:

This article was submitted to
Microbial Physiology and Metabolism,
a section of the journal
Frontiers in Microbiology

Received: 19 August 2019

Accepted: 12 November 2019

Published: 29 November 2019

Citation:

Wolij K, Degefu T and
Frostegård Å (2019) Host Range
and Symbiotic Effectiveness of N₂O
Reducing *Bradyrhizobium* Strains.
Front. Microbiol. 10:2746.
doi: 10.3389/fmicb.2019.02746

Emissions of the potent greenhouse gas N₂O is one of the environmental problems associated with intensive use of synthetic N fertilizers, and novel N₂O mitigation strategies are needed to minimize fertilizer applications and N₂O release without affecting agricultural efficiencies. Increased incorporation of legume crops in agricultural practices offers a sustainable alternative. Legumes, in their symbiosis with nitrogen fixing bacteria, rhizobia, reduce the need for fertilizers and also respond to the need for increased production of plant-based proteins. Not all combinations of rhizobia and legumes result in efficient nitrogen fixation, and legume crops therefore often need to be inoculated with compatible rhizobial strains. Recent research has demonstrated that some rhizobia are also very efficient N₂O reducers. Several nutritionally and economically important legumes form root nodules in symbiosis with bacteria belonging to *Bradyrhizobium*. Here, the host-ranges of fourteen N₂O reducing *Bradyrhizobium* strains were tested on six legume hosts; cowpea, groundnut, mung bean, haricot bean, soybean, and alfalfa. The plants were grown for 35 days in pots in sterile sand supplemented with N-free nutrient solution. Cowpea was the most promiscuous host nodulated by all test strains, followed by groundnut (11 strains) and mungbean (4 strains). Three test strains were able to nodulate all these three legumes, while none nodulated the other three hosts. For cowpea, five strains increased the shoot dry weight and ten strains the shoot nitrogen content (pairwise comparison; $p < 0.05$). For groundnut the corresponding results were three and nine strains. The symbiotic effectiveness for the different strains ranged from 45 to 98% in cowpea and 34 to 95% in groundnut, relative to fertilized controls. The N₂O reduction capacity of detached nodules from cowpea plants inoculated with one of these strains confirmed active N₂O reduction inside the nodules. When released from senescent nodules such strains are expected to also act as sinks for N₂O produced by denitrifying organisms in the soil microbial community. Our strategy to search among known N₂O-reducing *Bradyrhizobium* strains for their N₂-fixation effectiveness successfully identified several strains which can potentially be used for the production of legume inoculants with the dual capacities of efficacious N₂-fixation and N₂O reduction.

Keywords: *Bradyrhizobium*, denitrification, symbiotic nitrogen fixation, nitrous oxide, nitrous oxide reductase

INTRODUCTION

Synthetic fertilizers are the major sources of fixed N in most agricultural systems and have contributed to boosting crop productivity since the beginning of the green revolution (Kox and Jetten, 2015). Excessive use of fertilizer, however, comes with environmental risks and has created problems such as pollution of water bodies and emission of nitrous oxide (N₂O), a potent greenhouse gas and destructor of the stratospheric ozone layer (Ravishankara et al., 2009; Montzka et al., 2011; Reay et al., 2012; Butterbach-Bahl et al., 2013). Agriculture is a major source of N₂O, and it has been estimated that more than 60% of the N₂O emitted into the atmosphere is derived from N fertilized agricultural soils, mainly through the microbial processes of denitrification and nitrification (Thomson et al., 2012). Biological N₂ fixation is an environmentally friendly alternative to synthetic fertilizers (Jensen and Hauggaard-Nielsen, 2003). Legumes in symbiosis with N₂ fixing bacteria (rhizobia) account for the major part of the biological N₂ fixation, with rates in different agricultural systems being about 115 kg N ha⁻¹ year⁻¹ for crop legumes and 110–227 for pasture and fodder legumes (Herridge et al., 2008). In comparison, other endophytic and free-living bacteria fixed <5 to 25 kg N ha⁻¹ year⁻¹. Thus, the incorporation of legumes in pastures and in crop rotation practices, or intercropping legumes with other plants, will lessen the need for synthetic N fertilizers (Dakora et al., 1987; Peoples et al., 2009; Jensen et al., 2012) and thereby contribute in the battle against N₂O emissions.

The establishment of the legume-rhizobium symbiosis is based on a highly specialized signaling system, leading to the development of root nodules in the plant cortex. In most cases, rhizobia gain entrance to the plant via an infection thread formed in the root hairs, but in some plants, for example groundnut, the rhizobia instead enter through wounds in the root. The rhizobia then enter the cytoplasm of plant cells in the nodule where they, surrounded by plant-derived membranes, will differentiate into bacteroids. These are specialized bacterial cells that fix N₂ into NH₃, which is used by the plant for the synthesis of amino acids and proteins (Poole et al., 2018). Rhizobia vary in the type of hosts they nodulate. Some strains are highly specific, nodulating only a limited number of hosts, while others are promiscuous with a wide host range (Pueppke and Broughton, 1999). Moreover, some combinations of plant–rhizobia are efficient, others not, and the nitrogen fixation effectiveness may vary even between closely related plant cultivars/bacterial strains. Since soils may lack matching and/or effective rhizobia for certain legumes, especially when newly introduced to an area (Rodríguez-Echeverría et al., 2012), seeds of many legume crops need to be inoculated with compatible rhizobia. Strain screening and selection is an important step when developing new inoculants that are better specialized for selected crops and agricultural regions, aiming to optimize yields and plant nitrogen content.

Another feature of some rhizobia is their ability to denitrify, i.e., use nitrogen oxides as electron acceptors for respiration under anoxic conditions. Rhizobia with a complete

denitrification pathway carry genes coding for the periplasmic nitrate reductase NapAB, the copper (Cu) containing nitrite reductase NirK, a cytochrome c-dependent nitric oxide reductase NorCB and a clade I NosZ (Bedmar et al., 2005; Sameshima-Saito et al., 2006; Torres et al., 2014; Mania et al., 2019). Genes for the membrane-bound NarG appear to be lacking (Mania et al., 2019), while a cd1-type nitrite reductase NirS has been reported in a few strains (Sánchez and Minamisawa, 2018). Denitrification has hitherto only been investigated in a limited number of rhizobia, and it is therefore not known how widespread this metabolism is in this group of organisms. Incomplete denitrification pathways lacking one or more of the reduction steps have been reported for strains of *Ensifer*, *Mesorhizobium*, and *Bradyrhizobium* (Falk et al., 2010; Bueno et al., 2015; Mania et al., 2019). Depending on if they carry and express the gene *nosZ*, coding for NosZ, different strains of rhizobia may serve either as sources or sinks for N₂O emission. Thus, some rhizobia may have the potential to both enhance crop production and mitigate N₂O emission, serving to counteract both problems of agricultural productivity and environmental risks associated with the use of synthetic N fertilizers. Among the N₂O reducing rhizobia with such potentials are members of the genus *Bradyrhizobium*, of which many strains can form effective nodules with a wide range of economically important legume plants such as cowpea, peanut, soybean, and mungbean. Reports from greenhouse and field experiments demonstrate that inoculation of soybean with N₂O-reducing strains of *Bradyrhizobium* can mitigate N₂O emissions (Hénault and Revellin, 2011; Itakura et al., 2013; Akiyama et al., 2016). These findings are promising, and call for investigations of a wider range of rhizobia to identify N₂O reducing strains that can be further investigated for their suitability as inoculants for different legumes.

In a recent study, we screened 39 *Bradyrhizobium* strains, isolated from legume trees and herbal crops in Ethiopia, and characterized them with respect to their taxonomy and phylogeny (Wolde-Meskel et al., 2004b, 2005). Among them, we found 20 *Bradyrhizobium* strains capable of denitrifying NO₃⁻ to N₂ (Mania et al., 2019). The phylogeny of these strains had been analyzed through multilocus sequence analyses (MLSA), which clustered them into seven genospecies (Degefu et al., 2017). The study by Mania et al. (2019) showed that 18 of the complete denitrifiers belonged to three different genospecies within the *Bradyrhizobium japonicum* superclade, while the other two belonged to the *Bradyrhizobium elkanii* superclade (genospecies defined by Degefu et al., 2017). When supplied with both NO₃⁻ and N₂O, these strains all showed a strong preference for reduction of N₂O over NO₃⁻, probably because the electron transport pathway to NosZ competes very efficiently for electrons compared to the pathway to NapAB (Mania et al., 2019). The strong capacity to reduce N₂O was characteristic for all the complete denitrifying bradyrhizobia tested.

Aiming to find bradyrhizobia with the dual capacity of efficacious N₂-fixation and N₂O reduction, we here investigated the host ranges of 14 of these strongly N₂O reducing

Bradyrhizobium strains (Mania et al., 2019), all clustered in genospecies I (Degefu et al., 2017), against six legume hosts: cowpea (*Vigna unguiculata*), groundnut (*Arachis hypogea*), mung bean (*Vigna radiata*), soybean (*Glycine max*), haricot bean (*Phaseolus vulgaris*), and alfalfa (*Medicago sativa*). We also determined their symbiotic N₂ fixation effectiveness in cowpea (12 strains) and groundnut (11 strains) and compared the symbiotic performance of six of the best performing strains with cowpea seeds collected from three locations in Ethiopia (Hawassa, Ziway, and Bale). Furthermore, we investigated the N₂O reduction capacity of detached cowpea nodules formed by two different *Bradyrhizobium* strains, AC29c and AC70c. A major outcome of our study was that several of the N₂O reducing *Bradyrhizobium* test strains fixed N₂ effectively with different legume hosts. Such organisms are promising candidates for sustainable legume crop production and may simultaneously serve as a sink for N₂O emission from the legume rhizosphere.

MATERIALS AND METHODS

Bradyrhizobium Strains and Growth Conditions

Fourteen of the 39 *Bradyrhizobium* strains previously isolated from the tree and crop legumes growing in Ethiopia (Wolde-Meskel et al., 2004b) and capable of N₂O reduction (Mania et al., 2019) were used (Table 1). Each strain was grown at 28°C for 6 days on yeast mannitol agar (YMA). A single colony was then aseptically picked and cultured to a cell density of $\approx 10^9$ cells ml⁻¹

in a test tube containing 10 ml yeast mannitol broth (YMB) at 28°C. The YMA/YMB medium was as described in Mania et al. (2019).

Plant Growth, Host Range, and Symbiotic Effectiveness

The experiments were performed at Hawassa University, Ethiopia, in a polyhouse (a type of “shadehouse” with polyethylene roof and walls and fences to prevent animals from entering), at the temperature and humidity of the surrounding environment. We used six grain legume hosts: cowpea (local landrace, Hawassa, Ethiopia), groundnut (local landrace, Gofa, Ethiopia), mung bean (local landrace, Gofa), soybean (variety Ethio-Yugoslavia), haricot bean (variety Hawassa Dume) and alfalfa (variety 1086). Seeds were surface sterilized by immersion in 70% ethanol for 1 min, then transferred to a 3% sodium hypochlorite solution for 1 min followed by rinsing them in six changes of sterile water. Sterility was verified by streaking 2–3 seeds from each germination plate onto YMB agar. Sterilized seeds were imbibed in water for 1 h and subsequently incubated for germination at 28°C on sterile petri dishes lined with moist tissue paper for 3–4 days, or until the emergence of the radicle. Plants were grown in modified plastic cups with two separate parts: the lower part providing a reservoir for nutrient solution, and the upper part filled with washed and sterilized river sand, with a centrally positioned cotton wick extending through a hole out to the reservoir used as a rooting medium (Yates et al., 2016). Two germinated seeds were transferred to each plastic pot and thinned to one after successful seedling establishment. The experiments were carried out

TABLE 1 | Host range of N₂O reducing *Bradyrhizobium* strains tested on six legume crops.

Strains*	Host of isolation	Legume hosts tested					
		Cowpea ^a	Groundnut ^b	Mungbean ^c	Soybean ^d	Haricot bean ^e	Alfalfa ^f
AC70c	<i>Phaseolus vulgaris</i>	✓	✓	✓	–	–	–
AC79a	<i>Erythrina brucei</i>	✓	–	–	–	–	–
AC79b1	<i>Erythrina brucei</i>	✓	–	–	–	–	–
AC79b3	<i>Erythrina brucei</i>	✓	✓	–	–	–	–
AC79c2	<i>Erythrina brucei</i>	✓	✓	–	–	–	–
AC86b2	<i>Cajanus cajan</i>	✓	✓	–	–	–	–
AC86d2	<i>Cajanus cajan</i>	✓	✓	–	–	–	–
AC87h	<i>Milletia ferruginea</i>	✓	✓	–	–	–	–
AC87j1	<i>Milletia ferruginea</i>	✓	✓	–	–	–	–
AC87j2	<i>Milletia ferruginea</i>	✓	✓	–	–	–	–
AC92d	<i>Acacia gummifera</i>	✓	✓	–	–	–	–
AC101b	<i>Acacia saligna</i>	✓	✓	✓	–	–	–
AC101c	<i>Acacia saligna</i>	✓	✓	✓	–	–	–
AC101e	<i>Acacia saligna</i>	✓	–	✓	–	–	–

The plants were harvested 35 days after inoculation, and host ranges were assessed by scoring the presence of nodules on the roots. *The phylogeny of 39 *Bradyrhizobium* strains isolated from legume plants in Ethiopia (Wolde-Meskel et al., 2004b) was determined through multilocus sequence analyses (MLSA) which clustered the strains into seven genospecies (Degefu et al., 2017). Here we tested the host range of 14 *Bradyrhizobium* strains in genospecies I, identified as “strong N₂O reducers” (Mania et al., 2019). ^aLocal landrace Hawassa; ^blocal landrace from Gofa; ^clocal landrace from Gofa; ^dvariety Ethio Yugoslavia; ^evariety Hawassa Dume; ^fvariety 1086.

with three replicates using a randomized complete block design. The seedlings in each pot were inoculated with 1 ml (approximately 10^9 cells ml⁻¹) of rhizobial culture. Non-inoculated seedlings, either supplied with mineral nitrogen (as 0.05 g L⁻¹ KNO₃ weekly) or grown without nitrogen, were used as positive and negative controls, respectively. The seedlings inoculated with rhizobia were supplied with sterile, quarter-strength of Jensen's modified N-free nutrient solution twice per week and with sterilized, distilled water as necessary (Somasegaran and Hoben, 1994).

Host ranges were assessed by scoring the number of root nodules after carefully uprooting the whole plant 35 days after inoculation. Symbiotic effectiveness of the strains was determined by measuring the nodule number (NN), nodule dry weight (NDW), and shoot dry weight (SDW) as described by Somasegaran and Hoben (1994). Plant shoot samples were oven-dried at 70°C for 48 h to determine the SDW, then ground to a fine powder to determine their nitrogen content using a LECO Truspec carbon, hydrogen and nitrogen analyzer (St. Joseph, MI, United States). The shoot nitrogen content is given as SN (%) of the sample dry weight according to the Dumas method (Nelson and Sommers, 1996). The symbiotic effectiveness (SE) was estimated as the percentage of SDW of the plants supplied with N (+N control) (Yates et al., 2016). Shoot nitrogen was estimated as percent N₂ fixed from the atmosphere (Ndfa %): $[(SN_{\text{inoculated}} - SN_{\text{uninoculated}})/SN_{\text{inoculated}}] \times 100$. Since the plants were grown on sterile sand medium not supplied with an external source of nitrogen, the N accumulated in the plant shoots was assumed to be derived from the atmosphere through symbiotic N₂ fixation. The shoot nitrogen content of the uninoculated plants (SN_{uninoculated}) was considered to be equal to the N content of the seeds (Rennie, 1984). Average values \pm standard deviations ($n = 3$) are reported if not otherwise stated. Comparisons of the symbiotic effectiveness in the different treatments were done using analysis of variance (ANOVA) and Tukey's HD test. Correlations between the different parameters for symbiotic effectiveness were determined by Pearson's correlation coefficient.

Symbiotic Effectiveness of the Best Performing Strains in Cowpea Plants

After the first round of experiments, the six *Bradyrhizobium* strains that performed best with the cowpea landraces collected from Hawassa (located about 175 miles south of the Ethiopian capital Addis Ababa) were tested against two more cowpea landraces collected from Ziway (100 miles south of Addis Ababa) and Bale (300 miles south-east of Addis Ababa). Procedures for determining the N₂ fixation effectiveness and shoot N were as described previously.

N₂O Reduction Capacity by Intact Legume Nodules

We also investigated the capacity of bradyrhizobia to reduce N₂O when living inside nodules. This experiment was performed in a temperature-controlled greenhouse facility at the Norwegian

University of Life Sciences, Norway. To obtain nodules, cowpea seeds were sterilized and germinated as described above. Seedlings were planted in pots with vermiculite as rooting medium and inoculated with either the *Bradyrhizobium* strain AC29c, which is incapable of reducing N₂O, or with the N₂O-reducing *Bradyrhizobium* strain AC70c (Mania et al., 2019). Plants were uprooted 40 days after inoculation and nodules were cut from the roots and immediately washed with sterile distilled water. Equal numbers of healthy-looking nodules of approximately similar size were placed in three 120 ml sample vials. For each vial, nodules from three plants were pooled to obtain measurable N₂O reduction rates, thus nine plants were used in total. The dry weight of the nodules in each vial was approximately 18.9 mg. The vials were sealed with gas-tight butyl rubber septa and flushed with Helium (He) gas for 180 s to remove other gases in the vials. After releasing the He overpressure, 5% O₂ and about 400 ppm N₂O were injected into the headspace. The N₂O production/reduction capacity of the nodules was monitored using a robotic incubation system (Molstad et al., 2007).

RESULTS

Host Range of *Bradyrhizobium* Strains Capable of N₂O Reduction

Among the 14 tested strains, all formed pink, N₂ fixing nodules on cowpea, while 11 strains nodulated groundnut and 4 strains nodulated mungbean. We identified three strains (AC70c, AC101b, and AC101c) that were able to nodulate all these three crops. None of the strains nodulated soybean, haricot bean or alfalfa. Uninoculated controls did not form any nodule in any of the treatments. A complete list of the strains, their host of isolation and the cross-inoculation results is given in **Table 1**.

Symbiotic Effectiveness

Twelve of the fourteen strains that nodulated cowpea "Hawassa" and all eleven that nodulated groundnut were further tested for their N₂-fixation effectiveness. The treatments resulted in significant differences in SDW and SN (%) for both plants ($p < 0.001$). Results of multiple pairwise comparisons using Tukey's HD test at $\alpha = 0.05$ showed that the SDW of cowpea plants nodulated by five of the twelve N₂O reducing *Bradyrhizobium* strains was significantly higher than the SDW of the uninoculated -N controls (three strains at $p < 0.001$ and two strains at $p < 0.05$) (**Table 2**). There were large variations in SDW between replicates within the treatments with a coefficient of variation (CV) ranging between 4.0 and 56.6%. The SDW ranged from 280 ± 30 (standard deviation, $n = 3$) mg plant⁻¹ to 570 ± 30 mg plant⁻¹ with the corresponding SE of 45 and 91%, respectively. Similarly, groundnut plants nodulated by three out of the eleven N₂O reducing *Bradyrhizobium* strains had significantly higher SDW (one strain at $p < 0.001$ and two strains at $p < 0.05$) than uninoculated -N controls (**Table 4**). The SDW of groundnut plants ranged from 328 ± 90 to 919 ± 110 mg plant⁻¹ with

TABLE 2 | Symbiotic performance of 12 N₂O reducing *Bradyrhizobium* strains inoculated with cowpea was determined based on nodule number (NN), nodule dry weight (NDW), shoot dry weight (SDW), symbiotic effectiveness (SE, calculated as % relative to SDW of +N controls), shoot nitrogen content (SN, given as % of SDW), and shoot nitrogen derived from the atmosphere (Ndfa, calculated as % SN relative to -N controls).

Treatment	NN	NDW (mg plant ⁻¹)	SDW (mg plant ⁻¹)	SE (%)	SN (%)	Ndfa (%)
AC70c	44 ± 11 ^a	93 ± 12 ^a	570 ± 44 ^{ab}	91	4.07 ± 0.37 ^{abc}	76
AC101b	59 ± 20 ^a	90 ± 26 ^a	557 ± 116 ^{ab}	89	4.24 ± 0.15 ^{ab}	77
AC92d	31 ± 12 ^a	107 ± 55 ^a	500 ± 144 ^{abc}	80	3.34 ± 0.68 ^{abcd}	70
AC79c2	45 ± 13 ^a	100 ± 00 ^a	443 ± 29 ^{abc}	71	2.99 ± 0.81 ^{bcd}	66
AC101c	57 ± 37 ^a	103 ± 12 ^a	430 ± 56 ^{abc}	67	2.62 ± 0.82 ^{cd}	60
AC101e	29 ± 15 ^a	70 ± 10 ^a	393 ± 59 ^{abcd}	63	3.81 ± 0.07 ^{abc}	75
AC86d2	37 ± 4 ^a	83 ± 21 ^a	367 ± 32 ^{bcd}	59	3.08 ± 0.21 ^{abcd}	69
AC87H	40 ± 9 ^a	83 ± 51 ^a	363 ± 115 ^{bcd}	58	2.23 ± 0.81 ^{de}	52
AC79b3	46 ± 19 ^a	60 ± 36 ^a	323 ± 183 ^{bcd}	52	3.45 ± 0.54 ^{abcd}	72
AC86b2	27 ± 7 ^a	70 ± 30 ^a	297 ± 40 ^{cd}	47	2.91 ± 0.10 ^{bcd}	67
AC87j2	40 ± 17 ^a	63 ± 15 ^a	290 ± 69 ^{cd}	46	2.54 ± 0.54 ^{cd}	61
AC87j1	30 ± 13 ^a	50 ± 17 ^a	280 ± 35 ^{cd}	45	2.10 ± 0.67 ^{de}	50
+N control	na	na	627 ± 25 ^a	na	4.57 ± 0.00 ^a	na
-N control	na	na	147 ± 31 ^d	na	0.97 ± 0.02 ^e	na
Average	40 ± 17	81 ± 30	399 ± 147	64 ± 20	3.07 ± 1.03	66 ± 12
F statistics	1.16 ^{ns}	1.23 ^{ns}	7.3 ^{***}	3.69 ^{**}	10.45 ^{***}	2.55 [*]

Average values are given ($n = 3$) ± standard deviation. +N control: fertilized uninoculated; -N: non-fertilized uninoculated. Average values in the same column, followed by the same letter, are not significantly different at the 5% probability level by Tukey's test. na, not applicable; ns, non-significant; *significant at 0.05; **significant at 0.01; ***significant at 0.001.

SE of 34 and 95%, respectively (Table 4). There were also large variations in SDW between replicates of groundnut plants, with CV ranging between 5.4 and 47.9%. The highest SDW scores, and thus SE% values, were achieved for both cowpea and groundnut by inoculation with strain AC70c. For these combinations the SE values were 91 and 95%, respectively. This exceeded the symbiotic effectiveness significantly ($p < 0.05$) of three cowpea-nodulating strains and two groundnut-nodulating strains. For cowpea, strains AC101b and AC92d were also highly effective with $SE \geq 80\%$ of the +N control. In total, 9 strains out of 12 that nodulated cowpea and 9 strains out of 11 that nodulated groundnut were rated from moderately to highly effective, with SDW exceeding 50% of the +N controls.

The cowpea plants inoculated with our N₂O reducing *Bradyrhizobium* strains (11 out of 12) accumulated significantly higher amounts of N in their shoots ($p < 0.001$) than the uninoculated control (Table 2). All cowpea plants inoculated with these strains were able to derive most of their shoot nitrogen from the atmosphere (Ndfa $\geq 50\%$) (Table 2). Similarly, groundnut plants inoculated with 9 out of 11 strains accumulated significantly higher levels of N in their shoots ($p < 0.001$) than the uninoculated control plants (Table 4). Unlike cowpea plants, most of the nodulated groundnut plants derived <50% of their shoot N from the atmosphere through symbiotic nitrogen fixation (Table 4).

The number of nodules (NN) produced by cowpea was 40 ± 17 plant⁻¹, while the average NN for groundnut was 20 ± 13 (Tables 2, 4). There was no significant difference in NN between treatments of the same plant species ($p > 0.05$). However, NDW for groundnut differed significantly ($p < 0.05$)

between treatments with different bacterial strains, ranging between 21 ± 18 and 111 ± 44 mg plant⁻¹. For cowpea, the NDW ranged between 50 ± 17 to 107 ± 55 mg plant⁻¹, but the difference between the treatments was not statistically significant.

The Pearson correlation coefficients of the different parameters of symbiotic effectiveness in cowpea and groundnut are presented in Tables 3, 5, respectively. Nodule number for cowpea was significantly correlated with NDW ($r = 0.38$, $p < 0.05$) and SDW ($r = 0.51$, $p < 0.01$). In groundnut, NN showed a significant positive correlation only with NDW ($r = 0.66$, $p < 0.001$). NDW showed a significant positive correlation with SDW both in cowpea ($r = 0.65$, $p < 0.001$) and groundnut ($r = 0.48$, $p < 0.01$). Neither NN nor NDW showed any significant correlation with SN (%) in both legumes ($p > 0.05$). However, SDW showed a significant positive correlation with SN (%) both in cowpea ($r = 0.74$, $p < 0.001$) and in groundnut (0.53 , $p < 0.01$).

TABLE 3 | Pearson correlation coefficient among parameters of symbiotic performance of cowpea inoculated with 12 N₂O reducing *Bradyrhizobium* strains (Table 2).

	NN	NDW	SDW	SE (%)	SN (%)	Ndfa (%)
NN	1.00					
NDW	0.381 [*]	1.00				
SDW	0.510 ^{**}	0.648 ^{***}	1.00			
SE (%)	0.510 ^{**}	0.648 ^{***}	1.00 ^{***}	1.00		
SN (%)	0.305 ^{ns}	0.147 ^{ns}	0.736 ^{***}	0.548 ^{***}	1.00	
Ndfa (%)	0.327 ^{ns}	0.235 ^{ns}	0.456 ^{**}	0.456 ^{**}	0.936 ^{***}	1.00

ns, non-significant; *significant at 0.05; **significant at 0.01; ***significant at 0.001.

TABLE 4 | Symbiotic performance of 11 N₂O reducing *Bradyrhizobium* strains inoculated with groundnut was determined based on nodule number (NN), nodule dry weight (NDW), shoot dry weight (SDW), symbiotic effectiveness (SE, calculated as% relative to SDW of +N controls), shoot nitrogen content (SN, given as% of SDW), and shoot nitrogen derived from the atmosphere (Ndfa, calculated as% SN relative to –N controls).

Treatment	NN	NDW (mg plant ⁻¹)	SDW (mg plant ⁻¹)	SE (%)	SN (%)	Ndfa (%)
AC70c	24 ± 16 ^a	83 ± 41 ^a	919 ± 185 ^a	95	1.9 ± 0.21 ^{bc}	41
AC79b3	19 ± 10 ^a	56 ± 9 ^a	733 ± 207 ^{ab}	76	1.8 ± 0.30 ^{bc}	38
AC86b2	30 ± 4 ^a	111 ± 44 ^a	723 ± 244 ^{ab}	75	1.9 ± 0.04 ^{bc}	44
AC87j1	28 ± 15 ^a	89 ± 7 ^a	691 ± 165 ^{abc}	72	1.6 ± 0.23 ^{bc}	30
AC79c2	14 ± 8 ^a	52 ± 20 ^a	664 ± 60 ^{abc}	69	1.9 ± 0.30 ^{bc}	43
AC92d	16 ± 12 ^a	56 ± 9 ^a	651 ± 85 ^{abc}	66	1.8 ± 0.14 ^{bc}	39
AC101c	23 ± 18 ^a	87 ± 65 ^a	580 ± 134 ^{abc}	60	1.9 ± 0.10 ^{bc}	43
AC101b	10 ± 8 ^a	21 ± 18 ^a	576 ± 222 ^{abc}	60	2.0 ± 0.33 ^{bc}	44
AC86d2	28 ± 22 ^a	57 ± 12 ^a	564 ± 88 ^{abc}	59	2.4 ± 0.05 ^b	54
AC87H	15 ± 12 ^a	37 ± 33 ^a	423 ± 49 ^{bc}	44	1.7 ± 0.23 ^{cd}	35
AC87j2	10 ± 3 ^a	24 ± 21 ^a	328 ± 157 ^{bc}	34	1.8 ± 0.27 ^{cd}	38
+N control	na	na	965 ± 52 ^a	na	3.1 ± 0.00 ^a	na
–N control	na	na	277 ± 17 ^c	na	1.1 ± 0.12 ^d	na
Mean	20 ± 13	61 ± 38	623 ± 231	65 ± 21	1.91 ± 0.47	41 ± 9
F statistics	0.93 ^{ns}	2.55 [*]	5.67 ^{***}	2.99 [*]	14.42 ^{***}	2.09 ^{ns}

Average values are given ($n = 3$) ± standard deviation. +N control: fertilized uninoculated; –N: non-fertilized uninoculated. Average values in the same column, followed by the same letter, are not significantly different at the 5% probability level by Tukey's test. na, not applicable; ns, non-significant; * significant at 0.05; ***significant at 0.001.

Relative Effectiveness of the Best Performing Strains in Three Cowpea Landraces

The six *Bradyrhizobium* strains that showed the best symbiotic performance with the cowpea landrace CPH (AC70c, AC101b, AC79c2, AC101c, AC101e, and AC92d) were also tested with landraces Cowpea Bale (CPB) and Cowpea Ziway (CPZ) (Table 6). The SDW and SN (%) values were higher to much higher than the –N control, but due to the variation between replicates, this was not statistically significant in most cases. Thus, unlike for CPH, no significant increase in SDW was found for CPZ and CPB after inoculation with the different strains ($p > 0.05$), except for strain AC79c2 with CPB ($p < 0.05$). Significant increases in SN (%) were, however, seen for CPZ inoculated with five of the strains ($p < 0.001$), and for CPB inoculated with two of the strains ($p < 0.05$). No significant differences were found when we compare the different strains with respect to their symbiotic performance within the same cowpea landrace. Comparison between individual strains, on the other hand, revealed differences in symbiotic effectiveness with the three landraces. Strain AC70c showed the best performance with CPH, resulting in the highest SDW recorded among the inoculated plants ($570 ± 44$ mg plant⁻¹), while strain 101b performed best with CPZ, a combination that resulted in the highest SE (95% of the SDW of the +N control). Strain AC79c2 performed best with CPB. The shoot N content (given as % N of the SDW) also varied between different bacterial-plant combinations. The highest SN (%) recorded was in CPH inoculated with AC101b and AC70c, with values of $4.24 ± 0.15$ and $4.07 ± 0.37\%$, respectively. All the CPH and CPZ plants derived most of their SN (%) from the atmosphere through symbiotic nitrogen fixation (Ndfa > 50%). Contrary to this, three

strain-CPB combinations derived less than 50% of their shoot nitrogen from the atmosphere (Ndfa < 50%).

N₂O Reduction by Legume Nodules

To evaluate whether bradyrhizobia inside nodules may contribute to the reduction of N₂O, intact nodules collected from cowpea plants were placed in airtight vials containing He mixed with 5% O₂ and 4–5 μmol N₂O–N, and gas concentrations (O₂, N₂O, and N₂) were monitored during 100 h (Figure 1). Nodules from plants inoculated with *Bradyrhizobium* strain AC70c, previously shown to reduce N₂O when grown in batch culture (Mania et al., 2019), reduced exogenously supplied N₂O to N₂ at a rate of 6 μmol N g⁻¹ nodules simultaneously with O₂ consumption until all N₂O was reduced to N₂ (Figure 1). In contrast, nodules of cowpea plants inoculated with strain AC29c, classified as a phenotype lacking N₂O reduction capacity (Mania et al., 2019) failed to reduce exogenously supplied N₂O to N₂. Cowpea nodules containing strain AC70c reduced exogenously

TABLE 5 | Pearson correlation coefficient among parameters of symbiotic performance of groundnut inoculated with 11 N₂O reducing *Bradyrhizobium* strains (Table 4).

	NN	NDW	SDW	SE (%)	SN (%)	Ndfa (%)
NN	1.00					
NDW	0.657 ^{***}	1.00				
SDW	0.254 ^{ns}	0.484 ^{**}	1.00			
SE (%)	0.254 ^{ns}	0.484 ^{**}	1.00	1.00		
SN (%)	0.229 ^{ns}	0.042 ^{ns}	0.528 ^{**}	0.079 ^{ns}	1.00	
Ndfa (%)	0.204 ^{ns}	0.045 ^{ns}	0.111 ^{ns}	0.111 ^{ns}	0.984 ^{***}	1.00

ns, non-significant; **significant at 0.01; ***significant at 0.001.

TABLE 6 | Relative effectiveness of the six best performing *Bradyrhizobium* strains inoculated with two additional cowpea landraces Cowpea Ziway (CPZ) and Cowpea Bale (CPB), compared with Cowpea Hawassa (CPH) from first round of experiments.

Treatment	SDW (mg plant ⁻¹)			SE (%)			SN (%)			Ndfa (%)		
	CPH	CPZ	CPB	CPH	CPZ	CPB	CPH	CPZ	CPB	CPH	CPZ	CPB
AC70c	570 ± 44 ^{ab}	350 ± 200 ^{ab}	425 ± 35 ^{ab}	91	73	77	4.07 ± 0.37 ^{ab}	3.52 ± 0.61 ^a	2.61 ± 0.02 ^{ab}	76	71	50
AC101b	557 ± 116 ^{ab}	457 ± 202 ^{ab}	460 ± 28 ^{ab}	89	95	84	4.24 ± 0.15 ^{ab}	3.18 ± 0.44 ^a	3.78 ± 0.30 ^a	77	68	66
AC92d	500 ± 144 ^{ab}	347 ± 159 ^{ab}	495 ± 35 ^{ab}	80	72	90	3.34 ± 0.68 ^{a-c}	3.37 ± 0.57 ^a	3.01 ± 0.88 ^{ab}	70	70	55
AC79c2	443 ± 29 ^{ab}	300 ± 139 ^{ab}	540 ± 57 ^a	71	63	98	3.00 ± 0.81 ^{b-c}	3.23 ± 0.57 ^a	2.41 ± 1.2 ^{ab}	66	68	38
AC101c	430 ± 57 ^{ab}	357 ± 109 ^{ab}	450 ± 156 ^{ab}	69	74	82	2.62 ± 0.82 ^c	2.74 ± 0.69 ^a	2.01 ± 0.38 ^{ab}	60	62	34
AC101e	393 ± 59 ^b	307 ± 75 ^{ab}	490 ± 14 ^{ab}	63	64	89	3.81 ± 0.07 ^{a-c}	2.59 ± 0.98 ^{ab}	2.38 ± 0.06 ^{ab}	75	56	45
+N control	627 ± 25 ^a	483 ± 104 ^a	550 ± 42 ^a	na	na	na	4.57 ± 0.00 ^a	3.90 ± 0.00 ^a	4.01 ± 0.00 ^a	na	na	na
-N control	147 ± 31 ^c	83 ± 6 ^b	260 ± 57 ^b	na	na	na	0.97 ± 0.02 ^d	1.01 ± 0.00 ^b	1.30 ± 0.00 ^b	na	na	na
Average	458 ± 155	335 ± 164	458 ± 101	77 ± 15	73 ± 29	87 ± 11	3.33 ± 0.24	2.94 ± 0.20	2.67 ± 0.24	70 ± 9	65 ± 10	48 ± 15
F statistics	11.77 ^{***}	2.29 ^{ns}	3.69 [*]	2.13 ^{ns}	0.39 ^{ns}	0.64 ^{ns}	16.54 ^{***}	7.15 ^{***}	5.27 [*]	2.34 ^{ns}	1.02 ^{ns}	1.22 ^{ns}

Shoot dry weight (SDW), symbiotic effectiveness (SE, calculated as % relative to SDW of +N controls), shoot nitrogen content (SN, given as % of SDW) and shoot nitrogen derived from the atmosphere (Ndfa, calculated as % SN relative to -N controls). Average values are given (n = 3) ± standard deviation. +N control: fertilized uninoculated; -N: non-fertilized uninoculated. Average values in the same column, followed by the same letter, are not significantly different at the 5% probability level by Tukey's test. na, not applicable; ns, non-significant; * significant at 0.05; *** significant at 0.001. The means and levels of significance for each parameter were re-calculated for CPH relative to +N and -N controls.




FIGURE 1 | N₂O reduction by detached nodules obtained from cowpea plants that had been inoculated with either the N₂O reducing *Bradyrhizobium* strain AC70c, or with the non-N₂O reducing *Bradyrhizobium* strain AC29c. The nodules (approximately 18.9 mg dry weight of nodules per vial) were incubated in vials containing He mixed with 5% O₂ and 4–5 μmol N₂O–N (corresponding to about 400–500 ppm N₂O). n = 3; bars indicate standard deviation.

supplied N₂O to N₂ within 40 h of incubation, unlike those containing strain AC29c which did not produce N₂. The slight decrease in N₂O in flasks containing nodules with strain AC29c is due to sampling loss.

DISCUSSION

Emissions of N₂O are steadily increasing, and the International Panel of Climate Change foresees continued increases during the rest of the present century unless effective measures are undertaken (Rogelj et al., 2018). In accordance, Davidson and Kanter (2014) pointed out the need for novel mitigation options beyond “good agronomic practice” such as avoiding excessive use of fertilizers (Thomson et al., 2012). One such option could be to increase the production of nutrient-rich legumes, since this would reduce the needs for fertilizers and at the same time provide nutritious food and feed for people and livestock. Inoculation of legume crops with N₂O reducing rhizobia, as suggested by, e.g., Hénault and Revellin (2011) and Itakura et al. (2013), would further enforce the N₂O mitigation. The potential of this is emphasized by the recent findings that a range of taxonomically diverse bradyrhizobia exhibit very efficient N₂O

reduction, showing a strong preference for N₂O over NO₃⁻ as electron acceptor caused by the N₂O reduction pathway being a powerful competitor for electrons compared to that of the periplasmic nitrate reductase Nap (Mania et al., 2019). This is likely to be a general trait common to all organisms, including rhizobia, that carry N₂O reductase as well as Nap but lack the membrane-bound nitrate reductase Nar. Such organisms have the potential to act as sinks for N₂O, produced in soil both by themselves and by other microorganisms.

Rhizobia vary in their compatibility as well as in their N₂ fixation effectiveness (Musiyiwa et al., 2005). Consequently, an increased, yet sustainable, production of legume crops will require a wider variety of rhizobial inoculants, selected for providing the highest possible N₂-fixation efficacy in combination with different types of legume crops, and at the same time carrying the capacity to reduce N₂O. In a recent investigation of a wide diversity of *Bradyrhizobium* strains, we found that approximately 50% were able of N₂O reduction (Mania et al., 2019), and in another collection of strains isolated mostly from nodules of peanut, only about one-third reduced N₂O (Å. Frostegård, pers comm). The search for promising strains should focus on performing the time-consuming and laborious cross-inoculation tests on strains that are selected based on their ability to reduce N₂O.

The 14 *Bradyrhizobium* strains studied here, all known to have a strong capacity for N₂O reduction, nodulated at least one of the legume species, and some showed efficacious N₂-fixation capacity (Tables 1, 2, 4). The results showed, however, clear compatibility differences between different combinations of test strain and the legume host. Both legume hosts and microsymbionts varied regarding their levels of promiscuity. Cowpea was the most promiscuous host and was nodulated by all the test strains, followed by peanut (11 strains) and mung bean (4 strains), while none of the strains nodulated soybean, haricot bean and alfalfa. This is in agreement with earlier reports showing that cowpea, groundnut and mungbean belong to the same cross-inoculation group (Somasegaran and Hoben, 1994) which can be nodulated by several *Bradyrhizobium* strains and are reported as promiscuous hosts (Peoples et al., 1989; Steenkamp et al., 2008; Pule-Meulenberg et al., 2010; Jaiswal et al., 2017; Ndungu et al., 2018). On the microsymbiont side, the three strains AC70c, AC101b, and AC101c that nodulated cowpea, groundnut, and mung bean may, therefore, be regarded as promiscuous.

Rhizobial strains with a wide host range and proven N₂ fixation effectiveness are often preferred for the development of inoculants suitable for different legume crops (Drew et al., 2012). Except for strain AC70c, which was isolated from trap host species *P. vulgaris*, other test strains included in this study were isolated from tree legumes growing in Ethiopia (Wolde-Meskel et al., 2004b). This indicates that rhizobia either isolated from wild legumes or trapped from soil can be domesticated and used as inoculants for legume crops as reported earlier (Wange, 1989; Zhang et al., 1991). This potential can be further exploited as Ethiopia hosts a wide diversity of *Bradyrhizobium* strains in its soil (Aserse et al., 2012; Degefu et al., 2017, 2018) that can be used

as inoculants for grain legumes. Our test strains failed to nodulate soybean, haricot bean, and alfalfa. This could be due to the specificity of these hosts to a certain group of rhizobia (Peoples et al., 1989). Reports show that alfalfa and haricot bean form a symbiotic association with fast-growing rhizobial strains belonging to genus *Ensifer* and *Rhizobium*, respectively, but not with bradyrhizobia (Somasegaran and Hoben, 1994). The inability of strain AC70c to nodulate haricot bean, its trap host (Wolde-Meskel et al., 2004a), could also be due to the specificity of the host genotype to a particular strain (Abi-Ghanem et al., 2011). We speculate that failure of our *Bradyrhizobium* test strains to nodulate soybean, a common host for *B. japonicum* strains, might be due to incompatibility as soybean is an exotic crop introduced to Ethiopia in the 1950s (Aserse et al., 2012). Classification of soybean under the cross-inoculation group different from cowpea and groundnut (Somasegaran and Hoben, 1994) further supports this.

Nodulation alone might not always result in an effective symbiosis that can fulfill the host's N requirement (Terpolilli et al., 2008), necessitating careful screening and selection for elite rhizobial symbionts that can form effective N₂ fixing nodules with the host legumes. The symbiotic effectiveness (SE) in our study ranged between 45 and 98% in the different cowpea landraces and between 34 and 95% in groundnut. Most of the test strains that nodulated both legume crops, however, were rated as moderately to highly effective with SDW exceeding 50% of the N fertilized controls, indicating their potential suitability as inoculants for cowpea and groundnut provided that they are competitive with other rhizobia in different soil environments. Inspection of the phylogenetic tree constructed based on the symbiosis-related gene *nodA* shows that most of our test strains are closely related to *Bradyrhizobium* strain CB756, a widely used inoculant of cowpea and peanut (Degefu et al., 2017), supporting their suitability as inoculants for the tested legumes. This should, however, be followed by field evaluation of the strain's ability to compete with indigenous rhizobia for nodulation and adaptation to the conditions in the soil (moisture content, temperature, acidity, and others) that might limit the efficiency and survival of the inoculant (O'Hara et al., 2002). In cases where the soil harbors high numbers of less effective indigenous rhizobia, the selected strain should be both competitive for nodule occupancy and have a high N fixation effectiveness with the selected legume crop, to ensure the success of inoculation (Toro, 1996).

Among the most effective strains in the present study was strain AC70c that occupied its own distinct branch designated as C on the *nodA* phylogenetic tree (Degefu et al., 2017). It was the best performing strain both with cowpea and groundnut, with SE values of 98 and 95%, respectively. Differences in nodulation and symbiotic effectiveness of rhizobial strains can also be extended to genotypes of the same host legume species (Pazdernik et al., 1997; Abi-Ghanem et al., 2011; Yates et al., 2016). Our analysis of the relative effectiveness of the six best-performing strains on the three cowpea landraces (Table 6) also demonstrated such variations, confirming the need to evaluate the host range and

symbiotic effectiveness of rhizobial strains with different cultivars of the same legume species.

Our results also showed that all the cowpea and groundnut plants inoculated by the test strains accumulated more N from the atmosphere through symbiotic N₂ fixation than uninoculated non-fertilized controls. Cowpea plants derived most of their shoot N from the atmosphere through symbiotic N₂ fixation. The low Ndfa in groundnut (<50% in most strain-plant combinations) might be due to the somewhat short duration of the experiment (35 days). Growth up to 7 weeks was recommended for large-seeded legumes such as groundnut before assessing nodulation and effectiveness of the inoculants (Yates et al., 2016).

Soybean nodules containing *Bradyrhizobium* strains have been reported to reduce N₂O (Hirayama et al., 2011; Hénault and Revellin, 2011; Inaba et al., 2012). In accordance, our results showed that detached nodules from cowpea plants, inoculated with the N₂O reducing strain AC70c (Mania et al., 2019), efficiently reduced exogenously provided N₂O. This strain expressed N₂O reductase *in vivo*, indicating the presence of bacterial cells with active anaerobic respiration in the root nodules. Yet, it is likely that the most important N₂O reduction activity will be performed by N₂O reductase expressing bradyrhizobia when they are released from senescent nodules into the soil where they can act as sinks for N₂O produced by denitrifying organisms in the microbial community (Akiyama et al., 2016).

REFERENCES

- Abi-Ghanem, R., Carpenter-Boggs, L., and Smith, J. L. (2011). Cultivar effects on nitrogen fixation in peas and lentils. *Biol. Fertil. Soils* 47, 115–120. doi: 10.1007/s00374-010-0492-6
- Akiyama, H., Hoshino, Y. T., Itakura, M., and Shimomura, Y. (2016). Mitigation of soil N₂O emission by inoculation with a mixed culture of indigenous *Bradyrhizobium diazoefficiens*. *Nat. Publ. Gr.* 6:32869. doi: 10.1038/srep32869
- Aserse, A. A., Räsänen, L. A., Aseffa, F., Hailemariam, A., and Lindström, K. (2012). Phylogenetically diverse groups of *Bradyrhizobium* isolated from nodules of *Crotalaria* spp., *Indigofera* spp., *Erythrina brucei* and *Glycine max* growing in Ethiopia. *Mol. Phylogenet. Evol.* 65, 595–609. doi: 10.1016/j.ympev.2012.07.008
- Bedmar, E. J., Robles, E. F., and Delgado, M. J. (2005). The complete denitrification pathway of the symbiotic, nitrogen-fixing bacterium *Bradyrhizobium japonicum*. *Biochem. Soc. Trans.* 33, 141–144. doi: 10.1042/bst0330141
- Bueno, E., Mania, D., Frostegård, Å., Bedmar, E. J., Bakken, L. R., and Delgado, M. J. (2015). Anoxic growth of ensifer meliloti 1021 by N₂O-reduction, a potential mitigation strategy. *Front. Microbiol.* 6:537. doi: 10.3389/fmicb.2015.00537
- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S. (2013). Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philos. Trans. R. Soc. B Biol. Sci.* 368:20130122. doi: 10.1098/rstb.2013.0122
- Dakora, F. D., Aboyinga, R. A., Mahama, Y., and Apaseku, J. (1987). Assessment of N₂ fixation in groundnut (*Arachis hypogaea* L.) and cowpea (*Vigna unguiculata* L.). *Mircen J. Appl. Microbiol. Biotechnol.* 3, 389–399. doi: 10.1007/BF00935697
- Davidson, E. A., and Kanter, D. (2014). Inventories and scenarios of nitrous oxide emissions. *Environ. Res. Lett.* 9:105012. doi: 10.1088/1748-9326/9/10/105012
- Degefu, T., Wolde-Meskel, E., and Rasche, F. (2018). Genetic diversity and symbiotic effectiveness of *Bradyrhizobium* strains nodulating selected annual grain legumes growing in Ethiopia. *Int. J. Syst. Evol. Microbiol.* 68, 449–460. doi: 10.1099/ijsem.0.002486
- Degefu, T., Wolde-meskel, E., Woliy, K., and Frostegård, Å. (2017). Phylogenetically diverse groups of *Bradyrhizobium* isolated from nodules of tree and annual legume species growing in Ethiopia. *Syst. Appl. Microbiol.* 40, 205–214. doi: 10.1016/j.syapm.2017.04.001
- Drew, E., Herridge, D., Ballard, R., Hara, G. O., Deaker, R., Denton, M., et al. (2012). *Inoculating Legumes: A practical guide*. Canberra: Grains Research and Development Corporation.
- Falk, S., Liu, B., and Braker, G. (2010). Isolation, genetic and functional characterization of novel soil *nirK*-type denitrifiers. *Syst. Appl. Microbiol.* 33, 337–347. doi: 10.1016/j.syapm.2010.06.004
- Hénault, C., and Revellin, C. (2011). Inoculants of leguminous crops for mitigating soil emissions of the greenhouse gas nitrous oxide. *Plant Soil* 346, 289–296. doi: 10.1007/s11104-011-0820-0
- Herridge, D. F., Peoples, M. B., and Boddey, R. M. (2008). Global inputs of biological nitrogen fixation in agricultural systems. *Plant Soil* 311, 1–18. doi: 10.1007/s11104-008-9668-3
- Hirayama, J., Eda, S., Mitsui, H., and Minamisawa, K. (2011). Nitrate-dependent N₂O emission from intact soybean nodules via denitrification by *Bradyrhizobium japonicum* bacteroids. *Appl. Environ. Microbiol.* 77, 8787–8790. doi: 10.1128/AEM.06262-11
- Inaba, S., Ikenishi, F., Itakura, M., Kikuchi, M., Eda, S., Chiba, N., et al. (2012). N₂O emission from degraded soybean nodules depends on denitrification by *Bradyrhizobium japonicum* and other microbes in the rhizosphere. *Microbes Environ.* 27, 470–476. doi: 10.1264/jisme.2.me12100
- Itakura, M., Uchida, Y., Akiyama, H., Hoshino, Y. T., Shimomura, Y., Morimoto, S., et al. (2013). Mitigation of nitrous oxide emissions from soils by *Bradyrhizobium japonicum* inoculation. *Nat. Clim. Chang.* 3, 208–212. doi: 10.1038/nclimate1734
- Jaiswal, S. K., Msimbira, L. A., and Dakora, F. D. (2017). Phylogenetically diverse group of native bacterial symbionts isolated from root nodules of groundnut (*Arachis hypogaea* L.) in South Africa. *Syst. Appl. Microbiol.* 40, 215–226. doi: 10.1016/j.syapm.2017.02.002
- Jensen, E. S., and Hauggaard-Nielsen, H. (2003). How can increased use of biological N₂ fixation in agriculture benefit the environment? *Plant Soil* 252, 177–186. doi: 10.1023/a:1024189029226

DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding authors.

AUTHOR CONTRIBUTIONS

KW, TD, and ÅF planned the experiments, analyzed the results, and wrote the manuscript. KW and TD performed the experiments.

FUNDING

KW was supported by a grant from the Norwegian State Educational Loan Fund.

- Jensen, E. S., Peoples, M. B., Boddey, R. M., Gresshoff, P. M., Henrik, H. N., Alves, B. J. R., et al. (2012). Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries. *Rev. Agron. Sustain. Dev.* 32, 329–364. doi: 10.1007/s13593-011-0056-7
- Kox, M. A. R., and Jetten, M. S. M. (2015). “The nitrogen cycle,” in *Principles of Plant-Microbe Interactions: Microbes for Sustainable Agriculture*, ed B. Lugtenberg, (Berlin: Springer), 205–214.
- Mania, D., Woliy, K., Degefu, T., and Frostegård, Å. (2019). A common mechanism for efficient N₂O reduction in diverse isolates of nodule-forming bradyrhizobia. *Environ. Microbiol.* (in press). doi: 10.1111/1462-2920.14731
- Molstad, L., Dörsch, P., and Bakken, L. R. (2007). Robotized incubation system for monitoring gases (O₂, NO, N₂O N₂) in denitrifying cultures. *J. Microbiol. Methods* 71, 202–211. doi: 10.1016/j.mimet.2007.08.011
- Montzka, S. A., Dlugokencky, E. J., and Butler, J. H. (2011). Non-CO₂ greenhouse gases and climate change. *Nature* 476, 43–50. doi: 10.1038/nature10322
- Musiyiwa, K., Mpepereki, S., and Giller, K. E. (2005). Symbiotic effectiveness and host ranges of indigenous rhizobia nodulating promiscuous soybean varieties in Zimbabwean soils. *Soil Biol. Biochem.* 37, 1169–1176. doi: 10.1016/j.soilbio.2004.12.004
- Ndungu, S. M., Messmer, M. M., Ziegler, D., Gamper, H. A., Mézáros, É, Thuita, M., et al. (2018). Cowpea (*Vigna unguiculata* L. Walp) hosts several widespread bradyrhizobial root nodule symbionts across contrasting agro-ecological production areas in Kenya. *Agric. Ecosyst. Environ.* 261, 161–171. doi: 10.1016/j.agee.2017.12.014
- Nelson, D. W., and Sommers, L. E. (1996). “Total carbon, organic carbon and organic matter,” in *Methods of Soil Analysis. Part 3: Chemical Methods*, ed D. L. Sparks (Madison, WI: Soil Science Society of America), 961–1010.
- O’Hara, G., Yates, R., and Howieson, J. (2002). “Selection strains of root nodule bacteria to improve inoculant performance and increase legume productivity in stressful environments,” in *Inoculants and Nitrogen Fixation of Legumes in Viet*, ed D. Herridge, (Canberra: ACIAR Proc), 75–80.
- Pazdernik, D. L., Vance, C. P., Sadowsky, M. J., Graham, P. H., and Orf, J. H. (1997). A host-controlled, serogroup-specific, ineffective-iodulation system in the *Bradyrhizobium-Soybean (Glycine max)* symbiosis. *Mol. Plant Microbe Interact* 10, 994–1001. doi: 10.1094/mpmi.1997.10.8.994
- Peoples, M. B., Brockwell, J., Herridge, D. F., Rochester, I. J., Alves, B. J. R., Urquiaga, S., et al. (2009). The contributions of nitrogen-fixing crop legumes to the productivity of agricultural systems. *Symbiosis* 48, 1–17. doi: 10.1007/bf03179980
- Peoples, M. B., Faizah, A. W., Rerkasem, B., and Herridge, D. F. (1989). Methods for evaluating nitrogen fixation by nodulated legumes in the field. *Aust. Cent. Int. Agric. Res.* 11:76.
- Poole, P., Ramachandran, V., and Terpolilli, J. (2018). Rhizobia: from saprophytes to endosymbionts. *Nat. Rev. Microbiol.* 16, 291–303. doi: 10.1038/nrmicro.2017.171
- Pueppke, S. G., and Broughton, W. J. (1999). *Rhizobium* sp. Strain NGR234 and *R. fredii* USDA257 share exceptionally broad, nested host ranges. *Mol. Plant-Microbe Interact.* 12, 293–318. doi: 10.1094/mpmi.1999.12.4.293
- Pule-Meulenber, F., Belane, A. K., Krasova-wade, T., and Dakora, F. D. (2010). Symbiotic functioning and bradyrhizobial biodiversity of cowpea (*Vigna unguiculata* L. Walp.) in Africa. *BMC Microbiol.* 10:89. doi: 10.1186/1471-2180-10-89
- Ravishankara, A. R., Daniel, J. S., and Portmann, R. W. (2009). Nitrous oxide (N₂O): The dominant ozone-depleting substance emitted in the 21st century. *Science* 326, 123–125. doi: 10.1126/science.1176985
- Reay, D. S., Davidson, E. A., Smith, K. A., Smith, P., Melillo, J. M., Dentener, F., et al. (2012). Global agriculture and nitrous oxide emissions. *Nat. Clim. Chang.* 2, 410–416.
- Rennie, R. J. (1984). Comparison of N balance and ¹⁵N Isotope dilution to quantify N₂ fixation in field-grown Legumes 1. *Agron. J.* 76, 785–790.
- Rodríguez-Echeverría, S., Fajardo, S., Ruiz-Díez, B., and Fernández-Pascual, M. (2012). Differential effectiveness of novel and old legume-rhizobia mutualisms: implications for invasion by exotic legumes. *Oecologia* 170, 253–261. doi: 10.1007/s00442-012-2299-7
- Rogelj, J., Shindell, D., Jiang, K., Fifita, S., Forste, P., Ginzburg, V., et al. (2018). “Mitigation pathways compatible with 1.5°C in the context of sustainable development,” in *Global Warming of 1.5°C. An IPCC Special Report on the Impacts of 537 Global warming of 1.5°C above Pre-Industrial Levels and Related Global Greenhouse Gas 538 Emission Pathways, in the Context of Strengthening the Global Response to the Threat of 539 Climate Change, Sustainable Development, and Efforts to Eradicate Poverty*, eds V. P. Masson-Delmotte, H.-O. Zhai, D. Pörtner, J. Roberts, P. Skea, R. Shukla, et al. (Genève: Intergovernmental Panel on Climate Change).
- Sameshima-Saito, R., Chiba, K., Hirayama, J., Itakura, M., Mitsui, H., Eda, S., et al. (2006). Symbiotic *Bradyrhizobium japonicum* reduces N₂O surrounding the soybean root system via nitrous oxide reductase. *Appl. Environ. Microbiol.* 72, 2526–2532. doi: 10.1128/aem.72.4.2526-2532.2006
- Sánchez, C., and Minamisawa, K. (2018). Redundant roles of *Bradyrhizobium oligotrophicum* Cu-type (NirK) and cd1-type (NirS) nitrite reductase genes under denitrifying conditions. *FEMS Microbiol. Lett.* 365, 1–7. doi: 10.1093/femsle/fny015
- Somasegaran, P., and Hoben, H. J. (1994). *Handbook for Rhizobia Methods in Legume-Rhizobium Technology*. Berlin: Springer Science & Business Media.
- Steenkamp, E. T., Stepkowski, T., Przymusiak, A., Botha, W. J., and Law, I. J. (2008). Cowpea and peanut in southern Africa are nodulated by diverse *Bradyrhizobium* strains harboring nodulation genes that belong to the large pantropical clade common in Africa. *Mol. Phylogenet. Evol.* 48, 1131–1144. doi: 10.1016/j.ympev.2008.04.032
- Terpolilli, J. J., O’Hara, G. W., Tiwari, R. P., Dilworth, M. J., and Howieson, J. G. (2008). The model legume *Medicago truncatula* A17 is poorly matched for N₂ fixation with the sequenced microsymbiont *Sinorhizobium meliloti* 1021. *New Phytol.* 179, 62–66. doi: 10.1111/j.1469-8137.2008.02464.x
- Thomson, A. J., Giannopoulos, G., Pretty, J., Baggs, E. M., and Richardson, D. J. (2012). Biological sources and sinks of nitrous oxide and strategies to mitigate emissions. *Philos. Trans. R. Soc. B Biol. Sci.* 367, 1157–1168. doi: 10.1098/rstb.2011.0415
- Toro, N. (1996). Nodulation competitiveness in the *Rhizobium*-legume symbiosis. *World J. Microbiol. Biotechnol.* 12, 157–162. doi: 10.1007/BF00364680
- Torres, M. J., Rubia, M. I., De La Peña, T. C., Pueyo, J. J., Bedmar, E. J., and Delgado, M. J. (2014). Genetic basis for denitrification in *Ensifer meliloti*. *BMC Microbiol.* 14:142. doi: 10.1186/1471-2180-14-142
- Wange, S. S. (1989). Response of groundnut (*Arachis hypogaea* L.) to inoculation with *Rhizobium* strains isolated from wild arboreal legumes. *MIRCEN J. Appl. Microbiol. Biotechnol.* 5, 135–141. doi: 10.1007/bf01741836
- Wolde-Meskel, E., Terefework, Z., Frostegård, Å, and Lindström, K. (2005). Genetic diversity and phylogeny of rhizobia isolated from agroforestry legume species in southern Ethiopia. *Int. J. Syst. Evol. Microbiol.* 55, 1439–1452. doi: 10.1099/ijs.0.63534-0
- Wolde-Meskel, E., Terefework, Z., Lindström, K., and Frostegård, Å (2004a). Metabolic and genomic diversity of rhizobia isolated from field standing native and exotic woody legumes in southern Ethiopia. *Syst. Appl. Microbiol.* 27, 603–611. doi: 10.1078/0723202041748145
- Wolde-Meskel, E., Terefework, Z., Lindström, K., and Frostegård, Å (2004b). Rhizobia nodulating African *Acacia* spp. and *Sesbania sesban* trees in southern Ethiopian soils are metabolically and genomically diverse. *Soil Biol. Biochem.* 36, 2013–2025. doi: 10.1016/j.soilbio.2004.05.020
- Yates, R. J., Howieson, J. G., Hungria, M., Bala, A., O’Hara, G. W., and Terpolilli, J. (2016). “Authentication of rhizobia and assessment of the legume symbiosis in controlled plant growth systems,” in *Working with rhizobia*, eds J. G. Howieson, and M. J. Dilworth, (Canberra: Australian Centre for International Agricultural Research), 73–108.
- Zhang, X., Harper, R., Karsisto, M., and Lindström, K. (1991). Diversity of *Rhizobium* bacteria isolated from the root nodules of leguminous trees. *Int. J. Syst. Evol. Microbiol.* 41, 104–113. doi: 10.1099/00207713-41-1-104

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The handling editor declared a past co-authorship with one of the authors, ÅF.

Copyright © 2019 Woliy, Degefu and Frostegård. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.