

Original Article

Cite this article: Selinheimo S, Vasankari T, Jokela M, Kanervisto M, Pirkola S, Suvisaari J, Paunio T (2018). The association of psychological factors and healthcare use with the discrepancy between subjective and objective respiratory-health complaints in the general population. *Psychological Medicine* **49**, 121–131. <https://doi.org/10.1017/S0033291718000582>

Received: 11 July 2017

Revised: 25 November 2017

Accepted: 13 February 2018

First published online: 20 March 2018

Key words:

Alexithymia; health anxiety; healthcare use; illness perception; population-based study; respiratory symptoms; sense of coherence; subjective health complaints

Author for correspondence:

Sanna Selinheimo, E-mail: sanna.selinheimo@ttl.fi or Tiina Paunio, E-mail: tiina.paunio@helsinki.fi

The association of psychological factors and healthcare use with the discrepancy between subjective and objective respiratory-health complaints in the general population

Sanna Selinheimo^{1,2,3}, Tuula Vasankari^{4,5}, Markus Jokela³, Merja Kanervisto⁶, Sami Pirkola⁷, Jaana Suvisaari² and Tiina Paunio^{1,2,8}

¹Finnish Institute of Occupational Health, Helsinki, Finland; ²Department of Health, National Institute for Health and Welfare, Helsinki, Finland; ³Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Helsinki, Finland; ⁴Finnish Lung Health Association, Helsinki, Finland; ⁵Department of Pulmonary Diseases and Clinical Allergology, University of Turku, Turku, Finland; ⁶Independent Science Researcher Ylöjärvi, Finland; ⁷Health Sciences, and University Hospital Psychiatric Department, University of Tampere, Tampere, Finland and ⁸Department of Psychiatry, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

Abstract

Background. We examined the prevalence of self-perceived respiratory symptoms (SRS) in the absence of any objective findings of respiratory pathology, and the association of such prevalence with psychological factors and healthcare use in the general population.

Methods. The study was conducted among a nationally representative sample of Finnish adults (BRIF8901). Respiratory functioning was measured by a spirometry test. Structured questionnaires were used to measure SRS, physician visits and psychological factors of alexithymia, sense of coherence, illness worry and common mental disorders. Individuals with a diagnosed respiratory disease or a severe psychiatric disorder, determined in a diagnostic interview, were excluded, giving a sample comprising 4544 participants.

Results. Twenty-six per cent of the general population and 36% of those with no diagnosed severe psychiatric disorder or respiratory disease experienced SRS despite a normal spirometry result. Psychological factors were associated with SRS ($0.0001 < p < 0.032$), and on the number of physician visit explaining 42.7% of the difference in visits between individuals with and without SRS, respectively. Illness worry was associated most strongly with SRS [odds ratio (OR) 1.29, 95% confidence interval (CI) 1.19–1.41, $p < 0.0001$] and higher numbers of physician visits (OR 1.35, CI 1.32–1.38, $p < 0.00001$), even after several adjustments.

Conclusions. Respiratory symptoms without objective findings are common in the general population. The study results underline the role of psychological factors in the reporting of respiratory symptoms and the associated medical burden, thereby indicating the functional nature of the symptomatology.

Introduction

Somatic symptoms are highly prevalent in the general population (Petrie *et al.* 2014; Eliassen *et al.* 2016), and up to two-thirds of them remain unexplained (Steinbrecher *et al.* 2011). Estimations of perceived but medically unexplained health complaints[†] vary widely, between 4% and 30% in the population and between 10% and as much as 50% in primary-care consultations (Rief *et al.* 2001; Kroenke *et al.* 2002; Kirmayer *et al.* 2004; Loenggaard *et al.* 2015; van der Sluijs *et al.* 2015). Of these, respiratory symptomatology with an unclear background varies from approximately 5% to 40% of the cases (Reid *et al.* 2001; Rief *et al.* 2001; Nimnuan *et al.* 2001). The highly variable prevalence estimates underline the need for a more comprehensive understanding of recurring subjective respiratory-health complaints, in which psychological factors are suggested to play a prominent part (Lehrer *et al.* 2002).

Individuals differ in their attributional styles with regard to somatic symptoms (Robbins & Kirmayer, 1991; Barsky *et al.* 2001b; Goodwin & Engstrom, 2002; Petrie & Weinman, 2012). This is related to how accurately both healthy people and patients with respiratory diseases perceive their respiratory symptomatology, and it predicts the outcome of illness (Lehrer *et al.* 2002; Cohen *et al.* 2003; Hynninen *et al.* 2005; Nici *et al.* 2006; Janssens *et al.* 2011). Psychological factors such as a pervasive tendency to experience negative emotions (negative affectivity) have proved to be influential in terms of distorting symptom-perception processes related to respiratory diseases (Lehrer *et al.* 2002). Failure to perceive symptoms results in problems of self-management and the overutilisation of health services (Frostholm *et al.* 2005;

© Cambridge University Press 2018

CAMBRIDGE
UNIVERSITY PRESS

[†]The notes appear after the main text.

Serrano *et al.* 2006; Rhee *et al.* 2011). Depression is a risk factor for incident asthma (Brunner *et al.* 2014) and its severity (Eisner *et al.* 2005), and respiratory symptoms are at least as strongly associated with common psychiatric disorders as they are with objective lung-function tests (Katon *et al.* 2007). Thus, the involvement of psychological and psychiatric factors in symptoms of respiratory diseases is established, but no general population samples have been used to examine whether these factors are related to the perceived respiratory-symptom burden in populations with normal lung functioning.

Studies involving selected patient populations have shown that it is essential to identify the determinants that influence an individual's ability to cope with and adjust to chronic conditions, and thereby to enhance understanding of the prognosis of the disease (Stewart & Yuen, 2011; Disler *et al.* 2012). Antonovsky's theory of a sense of coherence (SOC) (Antonovsky, 1987) conceptualises such an ability: a high SOC is related to the numbers and types of adaptive health-promoting coping strategies and a better health status in the general population as well as in different populations of chronically ill patients (Suominen *et al.* 2001; Pallant & Lae, 2002; Eriksson & Lindström, 2005, 2006; Simonsson *et al.* 2008). There is also evidence that a low SOC increases the risk of acquiring new diseases (Poppius *et al.* 1999; Agardh *et al.* 2003), but how it relates to the respiratory-symptom burden remains to be studied.

Alexithymia is also connected to an individual's ability to adjust to chronic conditions, being linked to having difficulty interpreting and expressing somatic sensations related to emotional arousal (Taylor *et al.* 1999; De Gucht & Heiser, 2003; Waller & Scheidt, 2006; Mattila *et al.* 2008). This could result in the less effective regulation of somatic conditions and, consequently, play a role in increased subjective health complaints. It has been shown in clinical samples, for example, that alexithymia complicates the optimal management of asthma due to the failure to perceive dyspnoea (Feldman *et al.* 2002; Vazquez *et al.* 2010; Baiardini *et al.* 2011), resulting in a larger number of healthcare contacts compared with non-alexithymic patients (Serrano *et al.* 2006). According to clinical trials, alexithymia may be detrimental to the perception of respiratory symptoms, but there has been little population-level research based on the objective measurement of pulmonary functioning.

Health anxiety is defined as the fear of having an illness, based on the misinterpretation of general somatic symptoms. It has been shown to indicate attentional bias towards illness-related cues (Owens *et al.* 2004), which results in the increased use of health services (Barsky *et al.* 2001a; Frosthalm *et al.* 2005; Bleichhardt & Hiller, 2007; Sunderland *et al.* 2013). Health anxiety, measured in terms of self-reported illness worry, is relatively common among patients in primary and secondary care (Robbins & Kirmayer, 1996; Clarke *et al.* 2008), including those with respiratory complaints (Tyrer *et al.* 2011). However, its influence on help-seeking behaviour is only partially recognised in the planning of medical treatment for somatic symptoms (Henningesen *et al.* 2007). A variety of somatic symptoms linked to illness worries appear to associate with a poor prognosis and high healthcare utilisation in a way that cannot be solely attributed to concurrent psychiatric disorders or somatic diseases, for example (Barsky *et al.* 2001a; Creed & Barsky, 2004). The research thus far has focused on the prevalence of the diagnostic concept of health anxiety and its correlation with the burden of somatic symptoms. The association of dispositional illness worry with objectively measured lung functioning and health complaints nevertheless remains to be explored in the context of respiratory medicine.

Further population-level research is needed to assess the association of a poor SOC, a high level of alexithymia and illness worry with subjective respiratory-health complaints. In the present study, we first determined the prevalence of self-reported respiratory symptoms with no objective signs of impaired lung functioning in the general population. Second, we examined the extent to which a SOC, alexithymia and illness worry associated with subjective respiratory-health complaints, when objective respiratory status was taken into account. Third, given that a high number of somatic symptoms is associated with an impaired health status and high healthcare use in a way that cannot be attributed to concurrent mood disorders or medical illness (Tomenson *et al.* 2013), we sought to determine whether respiratory-health complaints are related to increased healthcare use in the general population, and if so whether psychological factors strengthen the association.

Materials and methods

Design

Sample

The Health 2000 Survey (BRIF8901) is a nationally representative survey of Finnish adults aged 30 years and over conducted by the National Institute for Health and Welfare. The data were collected in Finland in 2000 and 2001 (see Aromaa & Koskinen (2004) for a detailed description of the sampling procedure). The base sample comprised 8028 subjects, of whom 6005 participated in a clinical health examination focusing on lung functioning, and in a structured interview covering respiratory symptomatology and common depressive, anxiety and alcohol-use disorders, in line with the Munich version of the Composite International Diagnostic Interview (CIDI) (WHO, 1993, 1997). The sample analysed for this study included subjects with complete information on lung functioning ($N = 4626$) and information on self-reported respiratory symptoms ($N = 4544$). Three exclusion criteria were applied: (1) to subjects with any psychotic disorder ($N = 249$) as described in Perala *et al.* (2007); (2) to one participant who was diagnosed after the study sample was modified; and (3) to those who either reported a diagnosed respiratory disease (asthma, COPD, chronic bronchitis, other) or on whom there was missing information on these items. The mean age of the 4544 participants in this sample was 51 (s.d. = 14.13), with a range of 30–94, and 54% of them were female (Table 1). Figure 1 outlines the study flow.

Measures

Lung functioning and symptoms

Spirometry is the most efficient tool for diagnosing bronchial obstruction. A Vitalograph bellow spirometer (Vitalograph Ltd. Buckingham, UK) model was used in this survey. The lower limit of normal (LLN) for the FEV₁/FVC ratio [the proportion of their vital capacity that people can expire: the forced expiratory volume in 1 s (FEV₁) and the forced vital capacity (FVC) of 6 s duration] were used as airflow-obstruction criteria in the study. The LLN is defined as the value that identifies the lower fifth percentile of a population. It is calculated by subtracting 1.645 times the standard deviation from the mean, that is, the expected value, and it is age-corrected: a value below -1.645 predicts bronchial obstruction (Quanjer *et al.* 2012). Acceptable spirometry results were obtained from 2087 males (57% of the base sample) and

Table 1. Baseline characteristics of the study population and comparison between included and excluded participants

		Group A LLN + NS		Group B LLN + S		Group C LLN–		Groups A–C		Excluded	
		N	%	N	%	N	%	N	%	N	%
Gender	Male	1358 ^B	49	649	40	80 ^B	56	2087	46	1550	44
	Female	1407	51	986 ^{AC}	60	64	44	2457	54	1934	56
AGE	30–44	1261 ^{BC}	46	434	27	39	27	1734 ^E	38	935	27
	45–54	769	28	409	25	41	28	1219 ^E	27	701	20
	55–64	405	15	332 ^A	20	23	16	760 ^E	17	522	15
	65+	330	12	460 ^A	28	41 ^A	28	831	18	1326 ^I	38
Education	Basic	790	29	763 ^A	47	75 ^A	52	1628	36	1473 ^I	52
	Secondary	1004 ^B	36	494	30	39	27	1537 ^E	34	741	26
	Higher	971 ^{BC}	35	377	23	30	21	1378 ^E	30	601	21
Marital status	Married	1752 ^{BC}	63	955	58	76	53	2783	61	1309	46
	Cohabit	362 ^B	13	153	9	19	13	534	12	237	8
	Divorce or separation	230	8	161	10	17	12	408	9	328 ^I	12
	Widow	125	5	204 ^A	12	16 ^A	11	345	8	547 ^I	19
	Unmarried	296	11	161	10	16	11	473	10	421 ^I	15
Main activity	Full-time	1848 ^{BC}	67	712	44	65	45	2625 ^E	58	925	33
	Part-time	115	4	86	5	4	3	205	5	108	4
	Student	35	1	17	1	2	1	54	1	31 ^I	1
	Retired	507	18	652 ^A	40	56 ^A	39	1215	27	1504	53
	Unemployed, laid off	183	7	125	8	15	10	323	7	208	7
	Homemaker	66	2	33	2	2	1	101	2	49	2
	Other	11		10	1	0	0	21		17	1
Type of pension	Disability or other early retirement	95	19	132	20	10	18	237	20	313	21
	Retirement (old-age)	363	72	478	73	43	77	884	73	1102	73
	Unemployment	39 ^B	8	27	4	2	4	68 ^E	6	40	3
	Other	10	2	15	2	1	2	26	2	48	3
BMI	<18,49	18	1	8		2 ^{AB}	1	28	1	46 ^I	1
	18,5–25,99	1459 ^B	53	597	37	92	64	2148 ^E	47	1256	36
	26–29,99	839 ^C	30	535 ^C	33	29	20	1403 ^E	31	750	22
	>30	449	16	495 ^{AC}	30	21	15	965	21	1432 ^I	41
Smoking	Daily	571	21	385	24	62 ^{AB}	43	1018	22	562	22
	Occasionally	142 ^B	5	53	3	5	3	200 ^E	4	69	3
	Former 1–12mths ago	57	2	27	2	3	2	87	2	34	1
	Former over 12mths ago	532	19	327	20	28	19	887	20	522	21
	Never	1463	53	841 ^C	52	46	32	2350	52	1328	53
CIDI: Alcohol	Yes	242	9	166	11	17	12	425	10	220 ^I	13
	No	2430	91	1382	89	120	88	3932 ^E	90	1428	87
CIDI: Mood	Yes	129	5	148 ^A	10	11	8	288	7	133 ^I	8
	No	2543 ^B	95	1400	90	126	92	4069 ^E	93	1515	92
CIDI: Anxiety	Yes	116	4	118 ^A	8	5	4	239	5	154 ^I	9

(Continued)

Table 1. (Continued.)

		Group A LLN + NS		Group B LLN + S		Group C LLN-		Groups A-C		Excluded	
		N	N %	N	N %	N	N %	N	N %	N	N %
CIDI: Sum	No	2556 ^B	96	1430	92	132	96	4118 ^E	95	1494	91
	Yes	413	15	348 ^A	22	29	21	790	18	387 ^I	23
	No	2259 ^B	85	1200	78	108	79	3567 ^E	82	1261	77
Physician visits during 12 months	0	846 ^B	31	346	21	42	29	1234	27	734	27
	1	530 ^B	19	263	16	17	12	810	18	452	16
	2	404	15	261	16	27	19	692 ^E	15	348	13
	3	310	11	196	12	19	13	525	12	294	11
	4-5	353	13	286 ^A	18	22	15	661	15	374	14
	6-50	318	12	280 ^A	17	16	11	614	14	558 ^I	20

LLN + NS, group includes participants whose spirometry lower limits of normal (LLN) ratio FEV₁/FVC is normal (≥ -1.645) and who do not report respiratory symptoms; LLN + S, participants whose LLN is normal (≥ -1.645) and who report respiratory symptoms; LLN-, participants whose LLN is low (< -1.645). Excluded were participants ($N=3484$) with diagnosed respiratory disease ($N=1391$), psychosis ($N=250$), or information on self-reported respiratory symptoms ($N=1027$) or primary outcomes missing ($N=816$). BMI, body mass index; CIDI Alcohol, alcohol-related disorders by Mental Health Composite International Diagnostic Interview (CIDI); CIDI Mood, mood disorders; CIDI Anxiety, anxiety disorders; CIDI sum, the number of participants with one or more mental-health diagnoses. Column proportions are compared by Z-tests, the results are based on two-sided tests with Bonferroni correction, $p < 0.05$. For each significant pair, the key of the category with the smaller column proportion appears in the category with the larger column proportion; comparisons between included and excluded participant groups are based on Pearson's χ^2 tests, $p < 0.05$.


2457 females (56% of the base sample) included in this study who had not been diagnosed with a respiratory disease.

Self-reported respiratory symptoms were assessed in line with WHO-recommended sets of relevant questions (Fletcher *et al.* 1959; Rose & Blackburn, 1968): 'Do you cough or bring up phlegm on most days?' 'Do you become short of breath when you are hurrying on ground level or walking up a slight hill?' 'Do you feel breathless when walking with people of the same age on the level, or do you have to stop for breath when walking at your own pace on the level?' The subjects gave dichotomous

(yes-no) responses to each question, and if they answered 'yes' to one or more, they were classified as having self-reported respiratory symptoms. (online Supplementary 1: complete symptom questionnaire).

For the categorical analyses, we divided the participants into three groups based on the spirometry values and the self-reported respiratory symptoms. Group A (LLN + NS) includes those with a spirometry LLN FEV₁/FVC ratio ≥ -1.645 who did not report respiratory symptoms ($N=2765$); group B (LLN + S) includes those with a spirometry LLN ≥ -1.645 who reported respiratory

Fig. 1. The flow chart of the study population: LLN + NS = group includes participants whose spirometry lower limits of normal (LLN) ratio FEV₁/FVC is normal (≥ -1.645) and who do not report respiratory symptoms; LLN + S = group includes participants whose LLN is normal (≥ -1.645) and who report respiratory symptoms; LLN- = group includes participants whose spirometry LLN was low (< -1.645). *Information of spirometry missing $N=734$, **information of self-reported respiratory symptoms missing $N=82$. Self-reported respiratory symptoms without signs of bronchial obstruction (group B) were present in 36% of the general population when patients with psychosis and those diagnosed with respiratory diseases were excluded ($N=4544$). In all of the Health 2000 sample including those with no or missing information on respiratory symptoms, clinical health examinations, respiratory diseases and psychosis ($N=6185$), the prevalence of self-reported respiratory symptoms without signs of bronchial obstruction was 26%.



symptoms ($N = 1635$); group C (LLN-) includes those with a LLN spirometry ratio FEV_1/FVC of below -1.645 ($N = 144$) (Fig. 1).

Sense of coherence: SOC scale

SOC was assessed on 12 items of the Finnish SOC-13 scale compiled from the original SOC-29 scale (Antonovsky, 1987; Bernabe *et al.* 2009). Item 9 was not included in the final questionnaire of the Health 2000 Survey. The subjects gave their responses on seven-point semantic-differential scales with two opposite anchoring phrases (1 = very often and 7 = very seldom or never, sum-score range 12–84; a high score indicates a strong SOC). The scores for the five negatively keyed items (4, 5, 6, 16 and 25 from the original scale) were reversed. The Cronbach's α for the total scale was 0.85 in this study population. The psychometric properties of the SOC have proved to be good, and the measurements have been validated both in normal populations and in several clinical populations (Eriksson & Lindström, 2005).

Alexithymia: Toronto Alexithymia Scale-20

Alexithymia was assessed on the Finnish version of the Toronto Alexithymia Scale-20 (TAS-20) (Bagby *et al.* 1994a; Joukamaa *et al.* 2001). The subjects indicated their responses on five-point semantic-differential scales with two opposite anchoring phrases (1 = strongly disagree and 5 = strongly agree). The scores for the five negatively keyed items (4, 5, 10, 18 and 19) were reversed. The score is the sum score of responses to all 20 items, and a high score indicates alexithymic characteristics. A Cronbach's α coefficient of 0.85 was obtained for TAS-20 in a study based on the Health 2000 sample (Mattila *et al.* 2008). The psychometric properties of the Finnish version of TAS-20 have proved to be good (Bagby *et al.* 1994a, b; Joukamaa *et al.* 2001), and this measurement has been validated in the general population and in several patient populations (Joukamaa *et al.* 2001; Feldman *et al.* 2002; Waller & Scheidt, 2006; Vazquez *et al.* 2010; Baiardini *et al.* 2011).

Health anxiety: the Whiteley Index

The Whiteley Index (WI) is an instrument that is widely used for measuring health anxiety in the form of hypochondriac worries, and beliefs and convictions about illness, for example. In this study, we used a seven-item measure rated on a five-point Likert scale (Whiteley-7, range 7–35) (Fink *et al.* 1999), adapted from the original 20-item WI (Pilowsky, 1967). A high score indicates a tendency towards health anxiety. The scale has proved to be reliable and to have high internal consistency (Speckens *et al.* 1996). In this study, the Cronbach's α coefficient was 0.77 for the seven-item scale.

To make these three scales comparable, we standardised the sum scores in the analysis to a mean of zero and a standard deviation of one.

Demographic variables

We categorised age into four classes for the descriptive data analysis (30–44, 45–54, 55–64 and 65–100 years), and used it as a standardised value in the main analysis. Marital status was broken down into five categories: single, married, cohabiting, divorced/separated and widowed. The information related to socioeconomic position concerned education and main activity. The education variable comprised three levels (basic, secondary and higher education), and main activity was divided into full-time employed, part-time employed, unemployed or laid off, retired,

homemaker, student and other. Among the retired, the alternative grounds for the retirement were old age, disability, unemployment and other. Table 1 shows the demographics of the participants.

Other covariates

In further analyses, we used the standardised body mass index (BMI), spirometry LLN FEV_1/FVC ratio, and the maximum hand-grip strength of the dominating hand to measure physical condition, and C-reactive protein (CRP) as a measure of acute inflammation served as a continuous explanatory variable. The categorical variables were gender and smoking, the latter categorised as non-smoker or former smoker (stopped over 12 months previously) and current smoker (daily, occasionally or stopped in the preceding 12 months). The most common diagnosis for depressive, anxiety and alcohol-use disorders obtained from the CIDI interview was used as a categorical explanatory variable (three groups separately and one sum score: no diagnosis–diagnosis) (Pirkola *et al.* 2005).

Statistical analysis

First, Z-testing was used to compare the proportional differences in the descriptive data between the three groups (A: LLN + NS; B: LLN + S; C: LLN-). Bonferroni adjustment was used to account for multiple testing, the results being based on two-sided tests with $p < 0.05$. Second, analyses of covariance were conducted to compare the groups in terms of psychological factors (TAS, SOC and WI), controlled for gender and age. The results are presented as standardised values, including the confidence intervals (CIs) of the standard error of the means.

Third, binary logistic regression was used to estimate the influence of psychological factors on the reporting of respiratory symptoms (no symptoms–symptoms) (model 1). The results are presented as odds ratios (ORs), including 95% CIs and p values. Given their potential clinical significance, the analyses were adjusted for gender, age, spirometry LLN, CIDI diagnosis, smoking, BMI, grip strength and CRP (model 2). CIDI diagnoses were tested separately in the model, but because the results remained the same, only the sum score is shown. The dependent variables were self-reported respiratory symptoms (0 = no symptoms, 1 = symptoms). Fourth, a separate binary logistic regression analysis was conducted, with self-reported symptoms as the dependent variable and in separate models SOC, WI and TAS-20, with gender, age and spirometry LLN values as explanatory variables. The interactions between the psychological factors and gender, age and LLN were added to the models separately to estimate the possible interactions between the factors and the covariates (3×3 models). Both CIDI diagnosis and smoking status were controlled for in the analyses together (nine models) and separately (nine models). Fifth, Poisson regression was run to predict the number of physician visits during the previous 12 months as a dependent variable based on gender, age and study group (model 1), psychological factors (model 2) and other markers (model 3) as independent variables. Again, CIDI diagnoses were adjusted for separately in the model, but because the association did not change, only the sum score is shown. The results are presented with OR including 95% CIs and p values, and cover the subjects who had no missing values in the variables. IBM-SPSS 22.0 for Windows (SPSS Illinois, Chicago, Illinois, USA) software was used for the statistical analyses.

Results

Participant characteristics and the presence of respiratory symptoms in the population in relation to objective findings

Our primary study sample comprised 4544 individuals from the population-based Health 2000 Study, and excluded self-reported respiratory disease diagnosed by a medical doctor as well as participants with psychosis. Thirty-six per cent of the subjects experienced frequent respiratory symptoms with no objective signs of bronchial obstruction (LLN+S group). Three per cent had a bronchial obstruction (LLN-), and the majority (61%) had no respiratory symptoms or objective signs of bronchial obstruction (LLN+NS) (Fig. 1). The prevalence of self-reported respiratory symptoms without signs of bronchial obstruction in the whole Health 2000 sample (with available information on respiratory symptoms, spirometry, respiratory diseases and severe psychiatric disorders; $N=6185$) was 26%. Females were significantly over-represented in the LLN+S group compared with the other two groups, whereas LLN+NS was the youngest group. Being married, highly educated and in full-time work were significantly more frequent in the LLN+NS compared with the other two groups. Of the health-related factors, severe overweight was significantly more frequent in the LLN+S group than in the other two, whereas daily smoking was significantly more infrequent in the LLN+NS group. Compared with the members of the LLN+NS group, those in the LLN+S group had more depression and anxiety-disorder diagnoses, and reported significantly more frequent physician visits. Compared with the excluded participants, being 64 years old or younger, married, highly educated and in full-time work were significantly more frequent in the included participants, who also had fewer overall CIDI diagnoses. Physician visits were also more frequent among the excluded participants, which closer analysis showed was attributable to the presence of respiratory diseases (data not shown) (Table 1).

Psychological factors related to respiratory symptoms

There were significant differences on the WI [$F_{(2, 4118)} = 64.11$, $p < 0.0001$], the TAS-20 [$F_{(2, 4974)} = 37.29$, $p < 0.0001$] and the

SOC scale [$F_{(2, 4151)} = 40.86$, $p < 0.0001$]: the LLN+NS group reported higher SOC, fewer alexithymic characteristics and less health anxiety than the other groups (Fig. 2).

A high SOC was associated with a decrease in the odds of self-reported respiratory symptoms, and scoring high on the WI or TAS-20 was associated with an increase in self-reported symptoms even following adjustment for age, gender and the spirometry LLN FEV% value (Table 2, model 1). Adjusted for all the covariates, health anxiety was related to an increase in self-reported symptoms even when the covariates were accounted for (OR = 1.29, $p < 0.0001$). High alexithymia (OR = 1.11, $p = 0.032$) and a low SOC (OR = 0.89, $p = 0.015$) were significantly associated with an increase in respiratory symptoms, although to a lesser extent.

There were no interactions linking gender, age or spirometry LLN FEV% with alexithymia or SOC ($p < 0.05$) when predicting subjective respiratory-health complaints as a dependent variable. The WI interacted significantly with the spirometry LLN FEV% value (OR = 1.10, 95% CI 1.02–1.20, $p = 0.017$) even following adjustment for CIDI diagnoses, smoking or both. Online Supplementary Figs S3 and S4 show the interaction for males and females separately.

Healthcare usage

Members of the LLN+S study group reported 1.25 (95% CI 1.15–1.35, $p < 0.0001$) times more physician visits than those in the reference group LLN+NS (Table 3, model 1). The difference between the LLN+S and LLN+NS groups attenuated by 42.7% (from 1.25 to 1.10) following adjustment for psychological factors. The direction of the difference between the groups remained the same when the rest of the variables were added in model 3. However, no statistically significant result for SOC emerged in model 3.

Discussion

Subjective respiratory-health complaints without objective signs of impaired respiratory function were very common in this national sample of Finnish adults, with a prevalence estimate of 26% in the general population and 36% excluding cases of respiratory or severe psychiatric disease. Subjective respiratory symptoms




Fig. 2. Means and standard errors (s.e.) standardised (Z) measurements between the groups. Age and gender were controlled for in the analysis. The LLN+NS group includes participants who do not report respiratory symptoms with a normal spirometry results [lower-than-normal (LLN) spirometry ratio FEV₁/FVC ≥ -1.645]; the LLN+S group includes participants who do report respiratory symptoms with a normal spirometry results (LLN ratio FEV₁/FVC ≥ -1.645); the LLN- group includes participants with low spirometry (LLN ratio FEV₁/FVC < -1.645).

Table 2. Predictors of self-reported respiratory symptoms: the results obtained from the binary logistic regression

	Model 1 (N = 3784)			Model 2 (N = 3617)		
	OR	95% CI	p	OR	95% CI	p
Gender (female)	0.71	(0.61, 0.82)	0.000	0.92	(0.71, 1.19)	0.514
Age ^z	1.63	(1.50, 1.74)	0.000	1.51	(1.37, 1.66)	0.000
Spiro LLN% ^z	0.87	(0.81, 0.95)	0.001	0.88	(0.81, 0.96)	0.003
SOC ^z	0.85	(0.78, 0.93)	0.000	0.89	(0.81, 0.98)	0.015
WI ^z	1.32	(1.22, 1.43)	0.000	1.29	(1.19, 1.41)	0.000
TAS ^z	1.13	(1.04, 1.24)	0.007	1.11	(1.01, 1.22)	0.032
CIDI diagnosis				1.37	(1.13, 1.67)	0.002
Smoking				1.52	(1.28, 1.80)	0.000
BMI ^z				1.40	(1.29, 1.51)	0.000
GRIP ^z				0.78	(0.68, 0.90)	0.001
CRP ^z				1.13	(1.03, 1.24)	0.007

Spiro LLN %, spirometry lower limits of normal (LLN) ratio FEV₁/FVC; SOC, sense of coherence; WI, Whiteley Index; TAS 20, Toronto Alexithymia Scale; CIDI, categorical predictor according to the Mental Health Composite International Diagnostic Interview diagnosis, no diagnosis as a reference category; Smoking, categorical predictor based on self-reported smoking, never or no smoking for more than 12 months as the reference category; BMI, body mass index; GRIP, maximum hand-grip strength; CRP, C-reactive protein; predictors with ^z are used as standardised values. In model 1, R² = .084 (Hosmer and Lemeshow), 0.139 (Nagelkerke); in model 2, R² = .277 (Hosmer and Lemeshow), 0.180 (Nagelkerke).

were significantly related to the measures of SOC, alexithymia and illness worry, indicating that these specific psychological traits may enhance the perceived burden of the symptoms. The individuals concerned were more likely to visit a physician, and their psychological profile explained 42.7% of this association.

These results are in line with those of previous studies showing that perceived symptomatology has an influence on healthcare utilisation (Kolk *et al.* 2002; Burton, 2003; Tomenson *et al.* 2013), with a focus on the respiratory system. We found that the difference in the numbers of physician visits between the participants with no

Table 3. Predictors of physician visits during the previous 12 months: the results obtained from the Poisson regression

	Model 1 (N = 4536)			Model 2 (N = 3780)			Model 3 (N = 3613)		
	OR	95% CI	p	OR	95% CI	p	OR	95% CI	p
Female	1.37	(1.27, 1.48)	0.000	1.34	(1.29, 1.40)	0.000	1.27	(1.18, 1.36)	0.000
Male	1.00			1.00			1.00		
LLN–	1.05	(0.86, 1.28)	0.627	0.98	(0.86, 1.11)	0.725	0.98	(0.86, 1.11)	0.735
LLN + S	1.25	(1.15, 1.35)	0.000	1.10	(1.06, 1.15)	0.000	1.08	(1.03, 1.13)	0.001
LLN + NS	1.00			1.00			1.00		
Age ^z	1.01	(0.98, 1.05)	0.497	0.98	(0.95, 1.00)	0.032	0.97	(0.94, 0.99)	0.0012
TAS ^z				0.92	(0.90, 0.95)	0.000	0.93	(0.90, 0.95)	0.000
WI ^z				1.36	(1.34, 1.39)	0.000	1.35	(1.32, 1.38)	0.000
SOC ^z				0.97	(0.95, 1.00)	0.020	1.00	(0.97, 1.03)	0.933
CIDI diagnosis = yes							1.29	(1.22, 1.34)	0.000
CIDI diagnosis = no							1.00		
Smoking = yes							0.98	(0.94, 1.03)	0.444
Smoking = no							1.00		
GRIP ^z							0.94	(0.90, 0.97)	0.001
CRP ^z							1.02	(1.00, 1.04)	0.041
BMI ^z							1.04	(1.03, 1.07)	0.000

The LLN + S group includes participants with lower spirometry limits than normal (LLN) ratio FEV₁/FVC ≥ -1.645 and who report respiratory symptoms; the LLN + NS group includes participants whose spirometry LLN ≥ -1.645 and who do not report respiratory symptoms; the LLN– group includes participants whose spirometry LLN < -1.645 ; SOC, sense of coherence; WI, Whiteley Index; TAS 20, Toronto Alexithymia Scale; CIDI, categorical predictor according to the Mental Health Composite International Diagnostic Interview diagnosis; Smoking, categorical predictor based on self-reported smoking; no, never or no smoking for more than 12 months; yes, Current smoker or stopped during the previous 12 months; BMI, body mass index; GRIP, maximum hand-grip strength; CRP, C-reactive protein; predictors with ^z are used as standardised values. Model 1 deviance = 1 521 657 (df = 4531, $\chi^2 = 3.36$) and BIC = 2 493 716; in model 2, deviance = 1 166 556 (df = 3772, $\chi^2 = 3.09$) and BIC = 1 976 686; and in model 3, deviance = 1 098 372 (df = 3600, $\chi^2 = 3.05$) and BIC = 1 880 254.

signs of impaired lung functioning but with or without respiratory symptoms diminished following adjustment for psychological factors. In fact, SOC, alexithymia and illness worry even contributed independently to perceived symptoms when indicators of poor physical-health and mental-health diagnoses were accounted for. These findings suggesting the presence of a functional component in the symptomatology reflect the extent of various psychological variables linked to subjective health complaints. Our results are compatible with those reported in earlier experimental studies on symptom perception in patients with respiratory disease. It has been shown, for example, that a distressing context reduces the accuracy of symptom perception among people who are prone to negative affectivity: they rely more on environmental cues than on their own physiological responses in such situations (Bogaerts *et al.* 2005; Janssens *et al.* 2011). Thus, the presence of an underlying functional component appears to affect somatic perception, even on the population level, and could strongly affect overall subjective health complaints.

A higher SOC is considered to have health-promoting effects against somatic complaints (Eriksson & Lindström, 2005; Gallagher *et al.* 2008; Stewart & Yuen, 2011). Somewhat surprisingly, even though the predictive validity of SOC has proved to be good in several studies, its predictive power remained modest across the analyses. Kivimäki *et al.* (2000) found that a low SOC predicted significantly more adverse health problems than a high SOC, and concluded that the low rather than the high end of the SOC continuum was relevant in terms of health maintenance. Given these results, it would be worth evaluating the discriminant validity between high and low SOC scores.

Our findings imply that alexithymic characteristics are risk factors for the prevalence of somatic symptoms without a clear organic explanation (Deary *et al.* 1997; Bailey & Henry, 2007; Mattila *et al.* 2008), although the association between alexithymia and increased physician visits was modest. It has been suggested that the constitutive dimensions of alexithymia relate differently to self-perceived symptoms, and that separate factors have greater discriminant validity than general alexithymia measured on the TAS-20 (Lundh & Simonsson-Sarnecki, 2001; De Gucht & Heiser, 2003). Further analyses should evaluate the discriminant validity of its factor structure (Kooiman *et al.* 2004) in comparison with an SOC continuum.

The results on illness worry merit closer attention in that its predictive power remained the most stable across the analyses independently of a large body of adjustments. As a diagnostic concept, high health anxiety reflects the risk of high somatic-symptom counts, and predicts increased healthcare utilisation (Barsky *et al.* 2001a; Jyväsjärvi *et al.* 2001). It has also been found to have prognostic power in the course of hypochondriasis (olde Hartman *et al.* 2009). Thus, one may well argue that illness worry as a phenomenon only occurs simultaneously with poor self-perceived health, which is shown to associate with numerous somatic symptoms (Eliassen *et al.* 2016). Frosthalm *et al.* (2007) even showed that illness perceptions were more strongly associated with mental-health problems than poor physical health. We used several covariates to counter these arguments, including facets of poor physical and mental health. The fact that our results remained significant after these adjustments implies that patients' perceived health complaints reflect psychological vulnerability related to health behaviour, and not only disease severity from a biomedical perspective (Frosthalm *et al.* 2005). This is also supported by the interaction we found between illness worry and the spirometry results: high illness worry increased the probability of self-reported symptoms

regardless of the spirometry results, whereas the association with symptoms was stronger at low levels of illness worry. Thus, high illness worry seems to have an impact on self-reported symptoms, and should therefore be targeted in treatment planning [see also (Petrie & Weinman, 2012)].

Psychiatric disorders are associated with somatic symptoms without a clear medical explanation, and depression, anxiety and somatisation have been shown largely to overlap (Barsky & Borus, 1999; Henningsen *et al.* 2003; Haug *et al.* 2004; Löwe *et al.* 2008). In some cases, such as chest tightness among patients with asthma, psychiatric disorders may explain more of the variance in disease-specific symptoms than the specific disease (Lehrer *et al.* 2002; Barsky *et al.* 2005; Katon *et al.* 2007). We also found that depression and anxiety disorders were more prevalent in the self-reported symptoms than in the no-self-reported symptoms study group. Psychological distress is also strongly associated with the prevalence of somatic symptoms (Simon *et al.* 1996; Katon & Walker, 1998), and even after adjustment for psychiatric disorders, our results indicate that psychological distress should be considered separately from a psychiatric diagnosis in symptom evaluations. It is further suggested in previous studies that measurements of SOC, alexithymia and illness worry could facilitate recognition of psychological distress and non-adaptive health behaviour (Speckens *et al.* 1996; Joukamaa *et al.* 2001; Eriksson & Lindström, 2005). Thus, our findings support the need to further investigate the pragmatic value to clinicians, in terms of clinical standards or cut-off scores for diagnostic evaluation, of assessing precisely these factors as markers of non-adaptive health behaviour.

Our study has several strengths, including the large sample size and the generalisability to populations with several data sources (self-reported measures, interviewer-based diagnostic interviews and objective laboratory data). We adjusted for smoking, as well as for the spirometry value, to control for possible non-diagnosed respiratory disorders that might affect perceived respiratory-health complaints. It has been shown that attributing somatic symptoms to lifestyle factors such as smoking behaviour or alcohol consumption decreases the probability of reporting self-perceived health complaints (Nimnuan *et al.* 2001). Thus, we also adjusted for indicators of poor physical condition and acute inflammation, which have been associated with worsened somatic conditions including respiratory symptomatology. Finally, we adjusted for psychiatric disorders, which are comorbid with a high number of somatic symptoms and somatisation (Creed & Barsky, 2004; Löwe *et al.* 2008). The results were consistent across the analyses even following adjustment for these factors.

The study has its limitations. First, when we compared the study groups and the excluded participants in terms of demographics, we noted some differences. Hence, the findings are likely to be particularly valid for a population under 65 years old with no respiratory diseases, but not necessarily generalisable to elderly populations. Symptoms were identified via validated questionnaires focusing on current and recent (during the previous 12 months) symptoms, whereas information on respiratory diseases was retrospective and self-reported. Thus, although we controlled for diagnosed chronic respiratory diseases, the study population may have included individuals with rare forms of the disease that spirometry cannot recognise, and who might have been misclassified as healthy. This could reduce the value of the information on the presence/absence of excluded respiratory diseases, even though we controlled for poor physical health-related factors affecting perceived respiratory symptomatology. Moreover,

comorbid medical disorders explain high somatic-symptom counts only to a limited extent (Kroenke & Spitzer, 1998), and symptoms that are considered 'disease-specific' are almost as prevalent in patients without the specific disease (Michael *et al.* 2005). In adjusting for psychiatric diagnoses, we also controlled for comorbidities that have been shown to frequently co-occur with proneness to somatisation (Löwe *et al.* 2008).

Another potential limitation of our study is that the Health 2000 data may slightly underestimate the true frequency of physician visits, again especially in older age groups: it has been shown in previous studies that a long recall period may cause underestimation of self-reported yearly healthcare utilisation (Bhandari & Wagner, 2006; Short *et al.* 2009). Moreover, although the psychological factors included in this study are reportedly stable traits (Eriksson & Lindström, 2005; Hiirola *et al.* 2017), we could not control for dispositional personality factors such as negative affectivity. This might correlate with our measurements. Indeed, it has been shown to moderate the number of subjective health complaints even in the absence of medical problems (Watson & Pennebaker, 1989; Bogaerts *et al.* 2005, Bogaerts *et al.* 2010) and to influence the progression of the illness (Suominen *et al.* 2001; Hynninen *et al.* 2005): this may limit the generalisability of our results. A longitudinal study design would allow researchers to evaluate the extent to which psychological factors also influence the incidence of a new disease, as previous studies imply (Poppius *et al.* 1999; Agardh *et al.* 2003; olde Hartman *et al.* 2009).

In sum, our results clearly show that self-perceived respiratory-health complaints are highly prevalent in the general population, and they are not attributable to common mood, anxiety, substance-use or psychotic and respiratory disorders. The findings from our objective clinical data clearly emphasise the independent contribution of psychological factors to such complaints, and their associations with on the medical burden, independently of psychiatric and physical comorbidities. The subjective symptoms are important in themselves in that they constitute the basis of compliance with medical treatment, regardless of the diagnostic information they may hold (Barsky *et al.* 2001b; Petrie & Weinman, 2012). According to our results, intrinsic psychological vulnerability may independently explain a large proportion of the tendency to suffer from somatic distress. Thus, one would need to optimise the treatment of medically unexplained respiratory complaints by facilitating individual coping with symptoms as an integral part of individual symptom management, as recommended for the management of respiratory diseases (Nici *et al.* 2006; Yorke *et al.* 2007). The multifactorial background of the symptom experience should be considered an essential aspect in the diagnosis and treatment of respiratory symptoms.

Note

¹ Because of a lack of consensus concerning the definition of diffuse physical symptoms of uncertain aetiology, medically unexplained, functional somatic symptoms have become the more common term for subjective health complaints. However, given that our article focuses on the subjective faculties behind self-perceived health, we use the expression 'subjective health complaints' throughout.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291718000582>

Acknowledgements. SS is grateful to the Ida Montini Foundation and the Foundation of Non-Institutional Care, which funded the work. MJ is funded by the Academy of Finland (grant 311578).

Conflicts of interest. None.

Ethical approval. The approval of the Ethics Committee of Epidemiology and Public Health in the Hospital District of Helsinki and Uusimaa was obtained for the Health 2000 Study (31.005.2000).

References

- Agardh EE, Ahlbom A, Andersson T, Efendic S, Grill V, Hallqvist J, Norman A and Östenson C-G (2003) Work stress and low sense of coherence is associated with type 2 diabetes in middle-aged Swedish women. *Diabetes Care* **26**, 719–724.
- Antonovsky A (1987) *Unraveling the Mystery of Health: How People Manage Stress and Stay Well*. San Francisco: Jossey-Bass.
- Aromaa A and Koskinen S (eds.) (2004). *Health and Functional Capacity in Finland. Baseline Results of the Health 2000 Health Examination Survey*. Helsinki: Publications of the National Public Health Institute = Kansanterveyslaitos.
- Bagby RM, Parker JD and Taylor GJ (1994a) The twenty-item Toronto Alexithymia Scale-I. Item selection and cross-validation of the factor structure. *Journal of Psychosomatic Research* **38**, 23–32.
- Bagby RM, Taylor GJ and Parker JD (1994b) The twenty-item Toronto Alexithymia Scale-II. Convergent, discriminant, and concurrent validity. *Journal of Psychosomatic Research* **38**, 33–40.
- Baiardini I, Braidò F, Ferraioli G, Menoni S, Bruzzone M, Conte ME, Gani F, Ridolo E, Scordamaglia A and Canonica GW (2011) Pitfalls in respiratory allergy management: alexithymia and its impact on patient-reported outcomes. *Journal of Asthma* **48**, 25–32.
- Bailey PE and Henry JD (2007) Alexithymia, somatization and negative affect in a community sample. *Psychiatry Research* **150**, 13–20.
- Barsky AJ and Borus JF (1999) Functional somatic syndromes. *Annals of Internal Medicine* **130**, 910–921.
- Barsky AJ, Ettner SL, Horsky J and Bates DW (2001a) Resource utilization of patients with hypochondriacal health anxiety and somatization. *Medical Care* **39**, 705–715.
- Barsky AJ, Orav EJ and Bates DW (2005) Somatization increases medical utilization and costs independent of psychiatric and medical comorbidity. *Archives General Psychiatry* **62**, 903–910.
- Barsky AJ, Peekna HM and Borus JF (2001b) Somatic symptom reporting in women and men. *Journal of General Internal Medicine* **16**, 266–275.
- Bernabe E, Tsakos G, Watt RG, Suominen-Taipale AL, Uutela A, Vahtera J and Kivimäki M (2009) Structure of the sense of coherence scale in a nationally representative sample: the Finnish Health 2000 survey. *Quality of Life Research* **18**, 629–636.
- Bhandari A and Wagner T (2006) Self-reported utilization of health care services: improving measurement and accuracy. *Medical Care Research and Review* **63**, 217–235.
- Bleichhardt G and Hiller W (2007) Hypochondriasis and health anxiety in the German population. *British Journal of Health Psychology* **12**, 511–523.
- Bogaerts K, Notebaert K, Van Diest I, Devriese S, De Peuter S and Van den Bergh O (2005) Accuracy of respiratory symptom perception in different affective contexts. *Journal of Psychosomatic Research* **58**, 537–543.
- Bogaerts K, Van Eylen L, Li W, Bresseleers J, Van Diest I, De Peuter S, Stans L, Decramer M and Van den Bergh O (2010) Distorted symptom perception in patients with medically unexplained symptoms. *Journal of Abnormal Psychology* **119**, 226.
- Brunner WM, Schreiner PJ, Sood A and Jacobs Jr. DR (2014) Depression and risk of incident asthma in adults. The CARDIA study. *American Journal of Respiratory and Critical Care Medicine* **189**, 1044–1051.
- Burton C (2003) Beyond somatisation: a review of the understanding and treatment of medically unexplained physical symptoms (MUPS). *The British Journal of General Practice* **53**, 231–239.
- Clarke DM, Piterman L, Byrne CJ and Austin DW (2008) Somatic symptoms, hypochondriasis and psychological distress: a study of somatisation in Australian general practice. *Medical Journal of Australia* **189**, 560.
- Cohen S, Doyle WJ, Turner RB, Alper CM and Skoner DP (2003) Emotional style and susceptibility to the common cold. *Psychosomatic Medicine* **65**, 652–657.

- Creed F and Barsky A** (2004) A systematic review of the epidemiology of somatisation disorder and hypochondriasis. *Journal of Psychosomatic Research* **56**, 391–408.
- Deary IJ, Scott S and Wilson JA** (1997) Neuroticism, alexithymia and medically unexplained symptoms. *Personality and Individual Differences* **22**, 551–564.
- De Gucht V and Heiser W** (2003) Alexithymia and somatisation: a quantitative review of the literature. *Journal of Psychosomatic Research* **54**, 425–434.
- Disler R, Gallagher R and Davidson P** (2012) Factors influencing self-management in chronic obstructive pulmonary disease: an integrative review. *International Journal of Nursing Studies* **49**, 230–242.
- Eisner MD, Katz PP, Lactao G and Iribarren C** (2005) Impact of depressive symptoms on adult asthma outcomes. *Annals of Allergy, Asthma & Immunology* **94**, 566–574.
- Eliassen M, Kreiner S, Ebstrup JF, Poulsen CH, Lau CJ, Skovbjerg S, Fink PK and Jørgensen T** (2016) Somatic symptoms: prevalence, co-occurrence and associations with self-perceived health and limitations due to physical health – a Danish population-based study. *PLoS ONE* **11**, e0150664.
- Eriksson M and Lindström B** (2005) Validity of Antonovsky's sense of coherence scale: a systematic review. *Journal of Epidemiology and Community Health* **59**, 460–466.
- Eriksson M and Lindström B** (2006) Antonovsky's sense of coherence scale and the relation with health: a systematic review. *Journal of Epidemiology and Community Health* **60**, 376–381.
- Feldman JM, Lehrer PM and Hochron SM** (2002) The predictive value of the Toronto Alexithymia Scale among patients with asthma. *Journal of Psychosomatic Research* **53**, 1049–1052.
- Fink P, Ewald H, Jensen J, Sørensen L, Engberg M, Holm M and Munk-Jørgensen P** (1999) Screening for somatization and hypochondriasis in primary care and neurological in-patients: a seven-item scale for hypochondriasis and somatization. *Journal of Psychosomatic Research* **46**, 261–273.
- Fletcher CM, Elmes PC, Fairbairn AS and Wood CH** (1959) The significance of respiratory symptoms and the diagnosis of chronic bronchitis in a working population. *British Medical Journal* **2**, 257–266.
- Frostholm L, Fink P, Christensen KS, Toft T, Oernboel E, Olesen F and Weinman J** (2005) The patients' illness perceptions and the use of primary health care. *Psychosomatic Medicine* **67**, 997–1005.
- Frostholm L, Oernboel E, Christensen KS, Toft T, Olesen F, Weinman J and Fink P** (2007). Do illness perceptions predict health outcomes in primary care patients? A 2-year follow-up study. *Journal of Psychosomatic Research* **62**, 129–138.
- Gallagher R, Donoghue J, Chenoweth L and Stein-Parbury J** (2008) Self-management in older patients with chronic illness. *International Journal of Nursing Practice* **14**, 373–382.
- Goodwin R and Engstrom G** (2002) Personality and the perception of health in the general population. *Psychological Medicine* **32**, 325–332.
- Haug TT, Mykletun A and Dahl AA** (2004) The association between anxiety, depression, and somatic symptoms in a large population: the HUNT-II study. *Psychosomatic Medicine* **66**, 845–851.
- Henningsen P, Zimmermann T and Sattel H** (2003) Medically unexplained physical symptoms, anxiety, and depression: a meta-analytic review. *Psychosomatic Medicine* **65**, 528–533.
- Henningsen P, Zipfel S and Herzog W** (2007) Management of functional somatic syndromes. *The Lancet* **369**, 946–955.
- Hiirola A, Pirkola S, Karukivi M, Markkula N, Bagby R, Joukamaa M, Jula A, Kronholm E, Saarijärvi S and Salminen J** (2017) An evaluation of the absolute and relative stability of alexithymia over 11 years in a Finnish general population. *Journal of Psychosomatic Research* **95**, 81–87.
- Hynninen KMJ, Breivte MH, Wiborg AB, Pallesen S and Nordhus IH** (2005) Psychological characteristics of patients with chronic obstructive pulmonary disease: a review. *Journal of Psychosomatic Research* **59**, 429–443.
- Janssens T, Verleden G, De Peuter S, Petersen S and Van den Bergh O** (2011) The influence of fear of symptoms and perceived control on asthma symptom perception. *Journal of Psychosomatic Research* **71**, 154–159.
- Joukamaa M, Miettunen J, Kokkonen P, Koskinen M, Julkunen J, Kauhanen J, Jokelainen J, Veijola J, Laksy K and Jarvelin MR** (2001) Psychometric properties of the Finnish 20-item Toronto Alexithymia Scale. *Nordic Journal of Psychiatry* **55**, 123–127.
- Jyväsjärvi S, Joukamaa M, Väisänen E, Larivaara P, Kivelä S-L and Keinänen-Kiukaanniemi S** (2001) Somatizing frequent attenders in primary health care. *Journal of Psychosomatic Research* **50**, 185–192.
- Katon W, Lin EH and Kroenke K** (2007) The association of depression and anxiety with medical symptom burden in patients with chronic medical illness. *General Hospital Psychiatry* **29**, 147–155.
- Katon WJ and Walker EA** (1998) Medically unexplained symptoms in primary care. *The Journal of Clinical Psychiatry* **59**(Suppl 20), 15–21.
- Kirmayer LJ, Groleau D, Looper KJ and Dao MD** (2004) Explaining medically unexplained symptoms. *The Canadian Journal of Psychiatry* **49**, 663–672.
- Kivimäki M, Feldt T, Vahtera J and Nurmi J-E** (2000) Sense of coherence and health: evidence from two cross-lagged longitudinal samples. *Social Science & Medicine* **50**, 583–597.
- Kolk A, Hanewald G, Schagen S and van Wijk CG** (2002) Predicting medically unexplained physical symptoms and health care utilization: a symptom-perception approach. *Journal of Psychosomatic Research* **52**, 35–44.
- Kooiman CG, Bolk JH, Rooijmans HG and Trijsburg RW** (2004) Alexithymia does not predict the persistence of medically unexplained physical symptoms. *Psychosomatic Medicine* **66**, 224–232.
- Kroenke K and Spitzer RL** (1998) Gender differences in the reporting of physical and somatoform symptoms. *Psychosomatic Medicine* **60**, 150–155.
- Kroenke K, Spitzer RL and Williams JB** (2002) The PHQ-15: validity of a new measure for evaluating the severity of somatic symptoms. *Psychosomatic Medicine* **64**, 258–266.
- Lehrer P, Feldman J, Giardino N, Song H-S and Schmalting K** (2002) Psychological aspects of asthma. *Journal of Consulting and Clinical Psychology* **70**, 691.
- Loegaard K, Bjorner JB, Fink PK, Burr H and Rugulies R** (2015) Medically unexplained symptoms and the risk of loss of labor market participation – a prospective study in the Danish population. *BioMedCentral Public Health* **15**, 844.
- Löwe B, Spitzer RL, Williams JB, Mussell M, Schellberg D and Kroenke K** (2008) Depression, anxiety and somatization in primary care: syndrome overlap and functional impairment. *General Hospital Psychiatry* **30**, 191–199.
- Lundh LG and Simonsson-Sarnecki M** (2001) Alexithymia, emotion, and somatic complaints. *Journal of Personality* **69**, 483–510.
- Mattila AK, Kronholm E, Jula A, Salminen JK, Koivisto A-M, Mielonen R-L and Joukamaa M** (2008) Alexithymia and somatization in general population. *Psychosomatic Medicine* **70**, 716–722.
- Michael CS, Callahan CM, Counsell SR, Westmoreland GR, Stump TE and Kroenke K** (2005) Physical symptoms as a predictor of health care use and mortality among older adults. *The American Journal of Medicine* **118**, 301–306.
- Nici L, Donner C, Wouters E, Zuwallack R, Ambrosino N, Bourbeau J, Carone M, Celli B, Engelen M and Fahy B** (2006) American thoracic society/European respiratory society statement on pulmonary rehabilitation. *American Journal of Respiratory and Critical Care Medicine* **173**, 1390–1413.
- Nimmun C, Hotopf M and Wessely S** (2001) Medically unexplained symptoms: an epidemiological study in seven specialities. *Journal of Psychosomatic Research* **51**, 361–367.
- olde Hartman TC, Borghuis MS, Lucassen PL, van de Laar FA, Speckens AE and van Weel C** (2009) Medically unexplained symptoms, somatisation disorder and hypochondriasis: course and prognosis. A systematic review. *Journal of Psychosomatic Research* **66**, 363–377.
- Owens KM, Asmundson GJ, Hadjistavropoulos T and Owens TJ** (2004) Attentional bias toward illness threat in individuals with elevated health anxiety. *Cognitive Therapy and Research* **28**, 57–66.
- Pallant JF and Lae L** (2002) Sense of coherence, well-being, coping and personality factors: further evaluation of the sense of coherence scale. *Personality and Individual Differences* **33**, 39–48.
- Perala J, Suvisaari J, Saarni SI, Kuoppasalmi K, Isometsa E, Pirkola S and Partonen T, Tuulio-Henriksson A, Hintikka J, Kiesepa T, Harkanen T, Koskinen S and Lonnqvist J** (2007) Lifetime prevalence of psychotic and bipolar I disorders in a general population. *Archives of General Psychiatry* **64**, 19–28.
- Petrie KJ, Faasse K, Crichton F and Grey A** (2014) How common are symptoms? Evidence from a New Zealand national telephone survey. *British Medical Journal Open* **4**, e005374.

- Petrie KJ and Weinman J (2012) Patients' perceptions of their illness: the dynamo of volition in health care. *Current Directions in Psychological Science* **21**, 60–65.
- Pilowsky I (1967) Dimensions of hypochondriasis. *The British Journal of Psychiatry* **113**, 89–93.
- Pirkola SP, Isometsä E, Suvisaari J, Aro H, Joukamaa M, Poikolainen K, Koskinen S, Aromaa A and Lönnqvist JK (2005) DSM-IV mood-, anxiety- and alcohol use disorders and their comorbidity in the Finnish general population. *Social Psychiatry and Psychiatric Epidemiology* **40**, 1–10.
- Poppius E, Tenkanen L, Kalimo R and Heinsalmi P (1999) The sense of coherence, occupation and the risk of coronary heart disease in the Helsinki Heart Study. *Social Science & Medicine* **49**, 109–120.
- Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, Enright PL, Hankinson JL, Ip MS and Zheng J (2012) Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *European Respiratory Journal* **40**, 1324–1343.
- Reid S, Wessely S, Crayford T and Hotopf M (2001) Medically unexplained symptoms in frequent attenders of secondary health care: retrospective cohort study. *British Medical Journal* **322**, 767.
- Rhee H, Belyea MJ and Halterman JS (2011) Adolescents' perception of asthma symptoms and health care utilization. *Journal of Pediatric Health Care* **25**, 105–113.
- Rief W, Hessel A and Braehler E (2001) Somatization symptoms and hypochondriacal features in the general population. *Psychosomatic Medicine* **63**, 595–602.
- Robbins JM and Kirmayer LJ (1991) Attributions of common somatic symptoms. *Psychological Medicine* **21**, 1029–1045.
- Robbins JM and Kirmayer LJ (1996) Transient and persistent hypochondriacal worry in primary care. *Psychological Medicine* **26**, 575–589.
- Rose GA and Blackburn H (1968) *Cardiovascular Survey Methods*. Geneva: World Health Organization.
- Serrano J, Plaza V, Sureda B, De Pablo J, Picado C, Bardagi S, Lamela J, Sanchis J and Group SHRAR (2006) Alexithymia: a relevant psychological variable in near-fatal asthma. *European Respiratory Journal* **28**, 296–302.
- Short ME, Goetzel RZ, Pei X, Tabrizi MJ, Ozminkowski RJ, Gibson TB, DeJoy DM and Wilson MG (2009) How accurate are self-reports? An analysis of self-reported healthcare utilization and absence when compared to administrative data. *Journal of Occupational and Environmental Medicine/American College of Occupational and Environmental Medicine* **51**, 786.
- Simon G, Gater R, Kisely S and Piccinelli M (1996) Somatic symptoms of distress: an international primary care study. *Psychosomatic Medicine* **58**, 481–488.
- Simonsson B, Nilsson KW, Leppert J and Diwan VK (2008) Psychosomatic complaints and sense of coherence among adolescents in a county in Sweden: a cross-sectional school survey. *BioPsychoSocial Medicine* **2**, 1.
- Speckens AE, Spinhoven P, Sloekers PP, Bolk JH and van Hemert AM (1996) A validation study of the Whitely Index, the Illness Attitude Scales, and the Somatosensory Amplification Scale in general medical and general practice patients. *Journal of Psychosomatic Research* **40**, 95–104.
- Steinbrecher N, Koerber S, Frieser D and Hiller W (2011) The prevalence of medically unexplained symptoms in primary care. *Psychosomatics* **52**, 263–271.
- Stewart DE and Yuen T (2011) A systematic review of resilience in the physically ill. *Psychosomatics* **52**, 199–209.
- Sunderland M, Newby JM and Andrews G (2013) Health anxiety in Australia: prevalence, comorbidity, disability and service use. *The British Journal of Psychiatry* **202**, 56–61.
- Suominen S, Helenius H, Blomberg H, Uutela A and Koskenvuo M (2001) Sense of coherence as a predictor of subjective state of health: results of 4 years of follow-up of adults. *Journal of Psychosomatic Research* **50**, 77–86.
- Taylor GJ, Bagby RM and Parker JD (1999) *Disorders of Affect Regulation: Alexithymia in Medical and Psychiatric Illness*. Cambridge: Cambridge University Press.
- Tomenson B, Essau C, Jacobi F, Ladwig KH, Leiknes KA, Lieb R, Meinschmidt G, McBeth J, Rosmalen J and Rief W (2013) Total somatic symptom score as a predictor of health outcome in somatic symptom disorders. *The British Journal of Psychiatry* **203**, 373–380.
- Tyrer P, Cooper S, Crawford M, Dupont S, Green J, Murphy D, Salkovskis P, Smith G, Wang D and Bhogal S (2011) Prevalence of health anxiety problems in medical clinics. *Journal of Psychosomatic Research* **71**, 392–394.
- van der Sluijs J, Ten Have M, Rijnders C, van Marwijk H, de Graaf R and van der Feltz-Cornelis C (2015) Medically unexplained and explained physical symptoms in the general population: association with prevalent and incident mental disorders. *PLoS ONE* **10**, e0123274.
- Vazquez I, Sáñez E, González-Freire B, Romero-Frais E, Blanco-Aparicio M and Vereá-Hernando H (2010) The role of alexithymia in quality of life and health care use in asthma. *Journal of Asthma* **47**, 797–804.
- Waller E and Scheidt CE (2006) Somatoform disorders as disorders of affect regulation: a development perspective. *International Review of Psychiatry* **18**, 13–24.
- Watson D and Pennebaker JW (1989) Health complaints, stress, and distress: exploring the central role of negative affectivity. *Psychological Review* **96**, 234.
- World Health Organization (1993) *Composite International Diagnostic Interview (CIDI), Version 1.1*. Geneva: World Health Organization.
- World Health Organization (1997) *Composite International Diagnostic Interview (CIDI), Version 2.1*. Geneva: World Health Organization.
- Yorke J, Fleming SL and Shuldham C (2007) Psychological interventions for adults with asthma: a systematic review. *Respiratory Medicine* **101**, 1–14.