

Article

Classifying Evolution Algebras of Dimensions Two and Three

M. Eugenia Celorrio ^{1,†}  and M. Victoria Velasco ^{2,*,†} 

¹ Department of Mathematics and Statistics, Lancaster University, Lancaster LA1 4YF, UK; m.celorrioramirez@lancaster.ac.uk

² Departamento de Análisis Matemático, Facultad de Ciencias, Universidad de Granada, 18071 Granada, Spain

* Correspondence: vvelasco@ugr.es; Tel.: +34-958-241-000

† These authors contributed equally to this work.

Received: 17 October 2019; Accepted: 10 December 2019; Published: 13 December 2019



Abstract: We classified evolution algebras of dimensions two and three. Evolution algebras of dimensions three were classified recently obtaining 116 non-isomorphic types of algebras. Herein, with a new approach, we classify these algebras into 14 non-isomorphic types of algebra, so that this new classification is easier to deal with.

Keywords: evolution algebra; evolution operator; genetic algebra

1. Introduction

Mathematics and biology are intimately tied, and genetic algebras are an example of this link, as these are algebras with biological meanings. In this paper we are going to work with evolution algebras, kinds of genetic algebra introduced by Tian in [1] in 2008 that are used to model non-Mendelian genetics laws, although this is not their only application. In fact, they are strongly connected with group theory, Markov processes, the theory of knots, dynamic systems and graph theory. Due to the versatility of these algebras, the amount of literature studying them has grown immensely since 2008. In [2], the authors studied the relationship between evolution algebras and the spaces of functions defined by the Gibbs measure of a graph, which led into direct applications in biology, physics and mathematics itself. In works such as [3–10] they studied purely mathematical notions, such as nilpotency and solvency of evolution algebras, as well as the interpretation of these mathematical notions, relating, for example, the nilpotency of an element to gametes that go extinct after some generations. Chains of evolution algebras were studied in [11–14]. These are dynamic systems where the state of each system can be seen as an evolution algebra. Some derivatives of evolution algebras were studied in [1,15,16].

An important topic is the classification of evolution algebras of a given dimension up to isomorphism. There are several papers related with classification of evolution algebras, such as [17–24]. The first classification of evolution algebras of dimension two was given in [6], and some years later, in [17] (as part of a doctoral thesis [25]) another classification of these evolution algebras was provided, together with a classification of three dimensional evolution algebras into 116 non-isomorphic types.

Classifying a class of algebras consists of determining a classification criterion; i.e., defining different types of these algebras such that these different types are non-isomorphic to each other and such that each algebra belongs to exactly one of these types. In an intuitive way, we are constructing a cupboard with different drawers, in a such a way that each of the elements we are classifying belongs to one (and only one) of those drawers, but you can have more than one thing in each one. Nevertheless, if we change the shape of the drawers, the final cupboard will look completely different, even if it contains the same objects. As mentioned before, in [17] evolution algebras of

dimension three were classified into 116 types of non-isomorphic evolution algebras. In this paper, we classify three-dimensional evolution algebras into 14 non-isomorphic types (Theorem 11). To do so, our classifying criteria are based on distinguishing whether these algebras are degenerate or not and whether they are reducible or not. In the case of irreducible, non-degenerate algebras we differentiate three situations: when they have a basic ideal of dimension one and none of dimension two, when they have a two-dimensional basic ideal and they do not have a one dimensional basic ideal and when they have no basic ideals. According to the same criteria, we also obtain a classification of two-dimensional evolution algebras (Theorem 3). This shall be helpful for the classification of reducible three-dimensional evolution algebras. Since we reduce the study of evolution algebras of dimension three to 14 non-isomorphic types, this classification is much more practical than the classification provided in [17].

Note that every weighted digraph with three nodes is associated to an evolution algebra with dimension three in a one-to-one way. So, with this classification for evolution algebras we also have a classification of weighted digraphs with three vertices, particularly of discrete Markov processes with a state space of size three, as Markov processes are particular cases of evolution algebras. As a matter of fact, Markov processes are evolution algebras whose structure matrix is stochastic.

2. Basic Background

As the problem addressed in this paper is that of classifying evolution algebras of dimension two and three, in what follows we shall consider only evolution algebras of finite dimensions. Also, the algebras considered are defined over a field \mathbb{K} (where $\mathbb{K} = \mathbb{R}$ or \mathbb{C}).

We recall that an **evolution algebra** is an algebra A that has a natural basis $B = \{e_1, \dots, e_n\}$, which is a basis of A such that $e_i e_j = 0$ if $i \neq j$. For a fixed natural basis B , the square of each element can be written as $e_i^2 = \sum_{k=1}^n w_{ki} e_k$, and so, we can define the **structure matrix of A relative to B** in the following way

$$M_B(A) = \begin{pmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{n1} & \dots & w_{nn} \end{pmatrix},$$

where the i th column is given by the coefficients of e_i^2 with respect to B . When the basis B is clear we shall refer to this matrix as the structure matrix of A , without any further specification to B . This matrix determines the **product** of A . Indeed, given $a = \sum_{k=1}^n \alpha_k e_k$ and $b = \sum_{k=1}^n \beta_k e_k$ elements of A , it follows that

$$ab = \sum_{k=1}^n \gamma_k e_k, \text{ where}$$

$$\begin{pmatrix} \gamma_1 \\ \vdots \\ \gamma_n \end{pmatrix} = \begin{pmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{n1} & \dots & w_{nn} \end{pmatrix} \begin{pmatrix} \alpha_1 \beta_1 \\ \vdots \\ \alpha_n \beta_n \end{pmatrix}.$$

The next definition shall be useful to understand whether the natural basis is essentially unique or not.

Definition 1. Let A be an evolution algebra with dimension $n \in \mathbb{N}$, and let B and \tilde{B} be two natural basis of A . We say that B and \tilde{B} are **related** if $B = \{e_1, \dots, e_n\}$ and $\tilde{B} = \{\alpha_1 e_{\tau(1)}, \dots, \alpha_n e_{\tau(n)}\}$, where τ is a permutation of the set $\{1, \dots, n\}$ and $\alpha_1, \dots, \alpha_n$ are non-zero scalars.

In the next result, we shall see that, if the structure matrix of an evolution algebra A relative to one natural basis has non-zero determinant, then the natural basis is “essentially unique.”

Proposition 1. Let A be an evolution algebra. Let $B = \{e_1, \dots, e_n\}$ be a natural basis of A and $M_B(A)$ the corresponding structure matrix of A relative to B . Suppose that $\det(M_B(A)) \neq 0$. Then any other natural basis \tilde{B} of A is related to B , and moreover, $\det(M_{\tilde{B}}(A)) \neq 0$.

Proof. Since $A^2 = \text{lin}\{e_1^2, \dots, e_n^2\}$, the condition $\det(M_B(A)) \neq 0$ is equivalent to $\dim A^2 = n$. Thus, if $\tilde{B} = \{u_1, \dots, u_n\}$, since $A^2 = \text{lin}\{u_1^2, \dots, u_n^2\}$ we obtain that $\det(M_{\tilde{B}}(A)) \neq 0$, as $\{u_1^2, \dots, u_n^2\}$ must also be linearly independent. Now, consider $\alpha_{ij} \in \mathbb{K}$ for all $i, j \in \{1, \dots, n\}$ such that

$$\begin{aligned} u_1 &= \alpha_{11}e_1 + \dots + \alpha_{n1}e_n \\ &\vdots \\ u_n &= \alpha_{1n}e_1 + \dots + \alpha_{nn}e_n. \end{aligned}$$

Consider $\Lambda = \{\alpha_{ki} \neq 0 : 1 \leq k, i, \leq n\}$, which has a maximum of $n \times n$ elements. Fix i in $\{1, \dots, n\}$. There must exist $k \in \{1, \dots, n\}$ such that $\alpha_{ki} \neq 0$; otherwise, we would have $u_i = 0$ (a contradiction). Thus, $|\Lambda| \geq n$. But as $u_i u_j = 0$ for all $j \neq i$ we have that $\sum_{k=1}^n \alpha_{ki} \alpha_{kj} e_k^2 = 0$ and as e_1^2, \dots, e_n^2 are linearly independent; then, $\alpha_{ki} \alpha_{kj} = 0$ whenever $i \neq j$. Thus, it must be $\alpha_{kj} = 0$ for all $j \neq i$ as $\alpha_{ki} \neq 0$. Rephrasing what we just obtained, for each $1 \leq k \leq n$ there is at most one sub-index i such that $\alpha_{ki} \neq 0$. Then, $|\Lambda| \leq n$, which immediately implies that $|\Lambda| = n$. But as said before, each u_i is non-zero, so we must have that there exists a unique $k \in \{1, \dots, n\}$ such that $u_i = \alpha_{ki} e_k$, and the result is clear. \square


We shall now explain how to assign a graph to each evolution algebra. At first, it might depend on the natural basis selected, although we shall see that in some situations this graph is again “essentially unique.” For a discussion about this topic see [3].

Definition 2. Let $B = \{e_i : i \in \Lambda\}$ be a natural basis of an evolution algebra A , and $M_B = (w_{ij})$ its structure matrix. The **graph associated to A with respect to B** is the graph E_A^B whose set of vertices is B and the adjacency matrix is $M_B^1(A)$. The **simplified graph associated to A with respect to B** , is defined as the associated graph but without taking into account the weights; i.e., just considering whether there is a link between two vertices or not. Again, whenever the basis is clear, we shall speak about the graph (respectively simplified graph) associated to A , without any further specification.


Example 1. Let A be an evolution algebra with dimension 3, and consider a natural base $B = \{e_1, e_2, e_3\}$ being

$$\begin{aligned} e_1^2 &= e_1 - 2e_2 + 3e_3, \\ e_2^2 &= -5e_2 + e_3, \\ e_3^2 &= -2e_1. \end{aligned}$$

Then, the structure matrix is given by $M_B(A) = \begin{pmatrix} 1 & 0 & -2 \\ -2 & -5 & 0 \\ 3 & 1 & 0 \end{pmatrix}$, while the associated graph is



When we ignore the weights w_{ij} of the arrows of the graph we obtain the corresponding simplified graph:



We shall see now how the structure matrix changes when we multiply the elements of the natural basis by non-zero scalars, and how it affects the associated graph.

Proposition 2. Let A be an evolution algebra, $B = \{e_1, \dots, e_n\}$ a natural basis, and $M_B(A) = \begin{pmatrix} w_{11} & \dots & w_{1n} \\ \vdots & \ddots & \vdots \\ w_{n1} & \dots & w_{nn} \end{pmatrix}$ the structure matrix of A relative to B . If $\alpha_1, \dots, \alpha_n$ are non-zero scalars, then $\tilde{B} = \{\alpha_1 e_1, \dots, \alpha_n e_n\}$ is a natural basis of A and its structure matrix is given by

$$M_{\tilde{B}}(A) = \begin{pmatrix} \alpha_1 w_{11} & \frac{\alpha_2^2}{\alpha_1} w_{12} & \dots & \frac{\alpha_n^2}{\alpha_1} w_{1n} \\ \frac{\alpha_1^2}{\alpha_2} w_{21} & \alpha_2 w_{22} & & \frac{\alpha_n^2}{\alpha_2} w_{2n} \\ \vdots & & & \vdots \\ \frac{\alpha_1^2}{\alpha_n} w_{n1} & \frac{\alpha_2^2}{\alpha_n} w_{n2} & \dots & \alpha_n w_{nn} \end{pmatrix}.$$

Thus, the corresponding simplified graphs of A with respect to B and \tilde{B} coincide.

Proof. Taking into account that

$$(\alpha_i e_i)^2 = \alpha_i^2 e_i^2 = \sum_{k=1}^n \alpha_i^2 w_{ki} e_k = \sum_{k=1}^n \frac{\alpha_i^2}{\alpha_k} w_{ki} (\alpha_k e_k),$$

the conclusion is obtained straightforwardly. \square

Corollary 1. Let A be an evolution algebra and let B and B' be related natural basis. Then, the simplified graphs of A associated to B and B' respectively coincide, up to relabelling of the vertices.

Proof. The result follows from the above result together with the following fact: if $B = \{e_1, \dots, e_n\}$ is a natural basis and τ is a permutation of the set $\{1, \dots, n\}$, then $B' = \{e_{\tau(1)}, \dots, e_{\tau(n)}\}$ is another natural basis whose associated graph coincides with the graph associated to B up to relabelling of the vertices if needed. \square

Corollary 2. Let A_1 and A_2 be two evolution algebras with finite dimensions. Let B be a natural basis of A_1 such that $|M_B(A_1)| \neq 0$, and let $\theta : A_1 \rightarrow A_2$ be an algebra isomorphism. Then:

- (i) $\theta(B)$ defines a natural basis of A_2 such that $|M_{\theta(B)}(A)| \neq 0$.
- (ii) Every two natural basis of A_2 (respectively of A_1) are related.

Proof. As θ is an algebra isomorphism, it is clear that $\theta(B)$ is a natural basis of A_2 . Let $B = \{e_1, \dots, e_n\}$. As $\{e_1^2, \dots, e_n^2\}$ are linearly independent and $\theta(e_i^2) = \theta(e_i)^2$, we conclude that $\theta(e_i^2)$ are also linearly independent, and so $|M_{\theta(B)}(A)| \neq 0$. Then, as a consequence of Proposition 1, we can conclude that any two natural bases of A_2 (respectively of A_1) are related. \square

We see that whenever the structure matrix of an evolution algebra A associated to a natural basis B has non-zero determinant, then the basis as well as the associated graph are essentially unique (in

fact, all the natural bases are related and the simplified associated graph is unique). This leads to a result which is useful to proving that two evolution algebras are non-isomorphic.

Corollary 3. *Let A_1 and A_2 be two evolution algebras with dimension $n \in \mathbb{N}$. Let B_1 and B_2 be natural bases of A_1 and A_2 respectively. If $|M_{B_1}(A_1)| \neq 0$ and the simplified graph of A_1 associated to B_1 does not coincide with the simplified graph of A_2 associated to B_2 (up to relabelling the vertices), then A_1 and A_2 are non-isomorphic.*

Proof. If there exists an isomorphism $\theta : A_1 \rightarrow A_2$, then $\theta^{-1}(B_2)$ defines a natural basis on A_1 which is related to B_1 by Proposition 1. Thus, by Corollary 1, the simplified graphs with respect to these natural bases coincide, and therefore, the result follows. \square

Definition 3. *An evolution algebra A is **non-degenerate** if*

$$An(A) = \{b \in A : ab = 0 \forall a \in A\} = 0,$$

*and we say it is **degenerate** if $An(A) \neq 0$. As proven in ([3], Proposition 2.28), the latter is equivalent to the fact that any natural basis of A contains elements with zero square.*

Definition 4. *Let A be an algebra. An **ideal** of A is a linear subspace I such that $AI \subseteq I$ and $IA \subseteq I$ (note that for commutative algebras $AI = IA$). This means that, the quotient linear space A/I is an algebra with the canonical product $(a + I)(b + I) = ab + I$, for $a, b \in I$.*

*We say that an ideal I is **basic** if there exists $B = \{e_1, \dots, e_n\}$ a natural basis of A such that $I = \text{lin}\{e_{j_1}, \dots, e_{j_k}\}$ where $\{e_{j_1}, \dots, e_{j_k}\} \subseteq B$. This means that the ideal I is an evolution subalgebra provided with a natural basis that can be extended to a natural basis of A .*

Note that if B_I is a natural basis of I that is contained in B_A a natural basis of A , and if B'_I is another natural basis of I , then $B'_I \cup (B_A \setminus B_I)$ is a natural basis of A containing B'_I .

In [17] basic ideals are called evolution ideals with the extension property; meanwhile, in [10] they are simply called ideals. Note that the image of a basic ideal by a homomorphism is a basic ideal. For a discussion about the above notions with explanatory examples, see [3].

Definition 5. *An evolution algebra A is **reducible** if there exists non-zero proper ideals I and J such that $A = I \oplus J$. If A is not reducible then we say that it is **irreducible**.*

The following result was proven in [3], where the problem of the reducibility of an evolution algebra was deeply studied.

Theorem 1. *Let A be a non-degenerate evolution algebra and $B = \{e_1, \dots, e_n\}$ a natural basis of A . Then, A is reducible if and only if for some reorganisation B' of B the structure matrix $M_{B'}$ is of the type*

$$M_{B'} := \begin{pmatrix} W_{m \times m} & 0_{m \times (n-m)} \\ 0_{(n-m) \times m} & Y_{(n-m) \times (n-m)} \end{pmatrix},$$

with $m \in \mathbb{N}$, $m < n$ and $W_{m \times m}, Y_{(n-m) \times (n-m)}$ matrices with entries in \mathbb{K} .

In this case, we have that $A = I \oplus J$ where I and J are the ideals given by $I = \text{lin}\{e_i : i = 1, \dots, m\}$ and $J = \text{lin}\{e_i : i = m + 1, \dots, n\}$.

Note that the ideals I and J given in the above theorem are basic ideals, so that A is reducible as a direct sum of ideals if and only if A is reducible as a direct sum of basic ideals.

Before starting the study of the two-dimensional and three-dimensional evolution algebras, we need to clarify the notation we shall be using. From now on, a non-zero entry of a matrix shall be denoted by $*$. Note that two symbols $*$ in the same matrix may represent different (non-zero) values.

3. Classification of Two-Dimensional Evolution Algebras

In this section, we make a classification of two-dimensional evolution algebras taking into account the following properties: whether they are irreducible or not and whether they are degenerate or not.

As we can see in the following result, the degenerate case is easy to study.

Theorem 2. *Let A be a two-dimensional degenerate evolution algebra. Then:*

(i) *A is irreducible if and only if there exists a natural basis B of the evolution algebra A such that the structure matrix is $M_B(A) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$;*

(ii) *A is reducible if and only if there exists a natural basis B of A such that either $M_B(A) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ or $M_B(A) = \begin{pmatrix} 0 & w \\ 0 & * \end{pmatrix}$,*

where $*$ denotes a non-zero scalar and $w \in \mathbb{K}$.

Proof. Let $B = \{e_1, e_2\}$ be a natural basis of the evolution algebra A and denote by π_i the canonical projection of A on the subspace $\mathbb{K}e_i$, for $i = 1, 2$. As A is degenerate, we know that $e_i^2 = 0$ for some $i \in \{1, 2\}$ (according to [3], Proposition 2.28).

Case 1. $e_1^2 = e_2^2 = 0$. Then $A = I \oplus J$ where $I = \mathbb{K}e_1$ and $J = \mathbb{K}e_2$ and

$$M_B(A) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

Case 2. One of the elements of the natural basis has a zero square while the other has a non-zero square. We suppose that $e_1^2 = 0$ and $e_2^2 \neq 0$ (which is not restrictive). Hence, $e_2^2 = w_{12}e_1 + w_{22}e_2$ where $w_{12}, w_{22} \in \mathbb{K}$ with at least one of those scalars being non-zero.

Case 2.1. Suppose that $w_{22} = 0$. Then $e_2^2 = w_{12}e_1$ with $w_{12} \neq 0$, so A is irreducible. In fact, assume the contradiction that I and J are non-zero proper ideals, such that $A = I \oplus J$. Then, we can suppose that $\pi_2(I) \neq 0$ (which is not restrictive, as it cannot be $\pi_2(I) = \pi_2(J) = \{0\}$). It follows that $e_2^2 \in I$, and so, $e_1 = \frac{1}{w_{12}}e_2^2 \in I$. Thus, $I = A$, a contradiction, as the decomposition $A = I \oplus J$ is non-trivial.

Consequently, if $M_B(A) = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}$, then A is irreducible.

Case 2.2. Suppose that $w_{22} \neq 0$. Then, as $e_1^2 = 0$, the structure matrix is $M_B(A) = \begin{pmatrix} 0 & w \\ 0 & * \end{pmatrix}$ with $w = w_{12}$ and $e_2^2 = w_{12}e_1 + w_{22}e_2$. Then, $A = I \oplus J$ with $I = \mathbb{K}e_1$ and $J = \mathbb{K}e_2$. \square

In order to study the non-degenerate case, the following corollary is useful.

Lemma 1. *Let A be an evolutionary algebra, $B = \{e_1, e_2\}$ a natural basis and $M_B(A) = \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{pmatrix}$ the corresponding structure matrix. Then the structure matrix of A related to the natural basis $B' = \{e_2, e_1\}$ is given by $M_{B'}(A) = \begin{pmatrix} w_{22} & w_{21} \\ w_{12} & w_{11} \end{pmatrix}$.*

Proof. If $B' = \{v_1, v_2\}$ where $v_1 = e_2$, and $v_2 = e_1$, is clear that $v_1^2 = w_{22}v_1 + w_{12}v_2$ and $v_2^2 = w_{21}v_1 + w_{11}v_2$ as $e_2^2 = w_{12}e_1 + w_{22}e_2$ and $e_1^2 = w_{21}e_2 + w_{11}e_1$. \square

Corollary 4. Let A be a non-degenerate evolution algebra, $B = \{e_1, e_2\}$ a natural basis and $M_B(A) = \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{pmatrix}$ the corresponding structure matrix. Then A is reducible if and only if $w_{12} = w_{21} = 0$, in which case $M_B(A) = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$.

Proof. Since the only possible reordination of B is $B' = \{e_2, e_1\}$ and the corresponding structure matrix is $M_{B'}(A) = \begin{pmatrix} w_{22} & w_{21} \\ w_{12} & w_{11} \end{pmatrix}$ from the above lemma, the result follows from Theorem 1, jointly with the fact that the columns of $M_B(A)$ cannot be zero because A is non-degenerate. \square

Theorem 3. Let A be a non-degenerate evolution algebra, with $\dim A = 2$. Then:

- (i) A is reducible if and only if there exists a natural basis B of A such that $M_B(A) = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$;
- (ii) A is irreducible if and only if there exists a natural basis B of A such that either $M_B(A) = \begin{pmatrix} 0 & * \\ * & w \end{pmatrix}$ or $M_B(A) = \begin{pmatrix} w & \tilde{w} \\ * & * \end{pmatrix}$, with $w, \tilde{w} \in \mathbb{K}$.

Proof. As A is a non-degenerate evolution algebra, by Corollary 4, we obtain that A is reducible if and only if the structure matrix of A respect any natural basis B is of the type

$$\begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}.$$

Let us suppose now that A is irreducible and let $B = \{e_1, e_2\}$ be a natural basis. Then, by Corollary 4, and keeping in mind that A is non-degenerate, it is clear that $M_B(A)$ must be in one of the following cases:

$$\begin{pmatrix} * \\ * \end{pmatrix}, \begin{pmatrix} * \\ * \end{pmatrix}.$$

Moreover, as A is non-degenerate, we have that the structure matrix must be in one of the following cases:

$$M_1 = \begin{pmatrix} * & * \\ * & * \end{pmatrix}, M_2 = \begin{pmatrix} * & * \\ * & * \end{pmatrix}, M_3 = \begin{pmatrix} * & * \\ * & * \end{pmatrix}.$$

We can see that both M_2 and M_3 lead to the same evolution algebra. In fact, by switching if needed the elements of the basis (see Corollary 4) we obtain $\begin{pmatrix} w & \tilde{w} \\ * & * \end{pmatrix}$. The case M_1 produces the following situations:

$$\begin{pmatrix} 0 & * \\ * & w \end{pmatrix} \text{ and } \begin{pmatrix} * & * \\ * & w \end{pmatrix}.$$

If in the situation $\begin{pmatrix} * & * \\ * & w \end{pmatrix}$ we switch the elements of the basis then we get $\begin{pmatrix} w & * \\ * & * \end{pmatrix}$, a matrix of the type $\begin{pmatrix} w & \tilde{w} \\ * & * \end{pmatrix}$ as desired. \square

We can gather all this information in the following theorem, where we shall show, additionally, that the evolution algebras that we have found are not isomorphic.

Theorem 4. Let A be a two-dimensional evolution algebra. Then, A is one of the following non-isomorphic ones, where $r, s \in \mathbb{K} \setminus \{0\}$ and $w, \tilde{w} \in \mathbb{K}$.

- $A_1 = \text{lin}\{e_1, e_2\}$ is such that $e_1^2 = e_2^2 = 0$ and $e_1e_2 = 0$.
- $A_2 = \text{lin}\{e_1, e_2\}$ is such that $e_1^2 = 0, e_2^2 = we_1 + re_2$ and $e_1e_2 = 0$.
- $A_3 = \text{lin}\{e_1, e_2\}$ is such that $e_1^2 = re_1$ and $e_2^2 = se_2$ and $e_1e_2 = 0$.
- $A_4 = \text{lin}\{e_1, e_2\}$ is such that $e_1^2 = 0$ and $e_2^2 = re_1$ and $e_1e_2 = 0$.
- $A_5 = \text{lin}\{e_1, e_2\}$ is such that $e_1^2 = re_2$ and $e_2^2 = se_1 + we_2$ and $e_1e_2 = 0$.
- $A_6 = \text{lin}\{e_1, e_2\}$ is such that $e_1^2 = we_1 + re_2, e_2^2 = \tilde{w}e_1 + se_2$ with $e_1e_2 = 0$.

As a matter of fact, we have the following classification:

	Degenerate	Non-Degenerate
Reducible	A_1, A_2	A_3
Irreducible	A_4	A_5, A_6

Proof. Let us consider the matrices

$$M_1 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, M_2 = \begin{pmatrix} 0 & w \\ 0 & * \end{pmatrix}, M_3 = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix},$$

$$M_4 = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}, M_5 = \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix}, M_6 = \begin{pmatrix} w & \tilde{w} \\ * & * \end{pmatrix}.$$

The evolution algebras A_i described above have a natural basis B_i whose corresponding structure matrix is M_i with $i = 1, 2, 3, 4, 5, 6$. By Theorem 2 we know that whenever A is degenerate, then A is reducible if and only if there exists a natural basis such that its corresponding structure matrix is like either M_1 or M_2 . Meanwhile, A is irreducible if and only if A has a natural basis which structure matrix is like M_4 . If A is non-degenerate, A is reducible if and only if there exists a basis for which structure matrix is of the type M_3 , as it is shown in Theorem 3. Moreover, by this last result, if A is non-degenerate, then A is irreducible if the structure matrix for some natural basis is either $M = \begin{pmatrix} 0 & * \\ * & w \end{pmatrix}$ or $\tilde{M} = \begin{pmatrix} w & \tilde{w} \\ * & * \end{pmatrix}$. If the structure matrix of A is of the type of \tilde{M} , then it is within M_6 , whereas M can be identified with M_5 if $w = 0$ or with M_6 if $w \neq 0$.

Now, we just need to prove that these evolution algebras are non-isomorphic. In order to do so, we need to take into account that the properties of being degenerate and being irreducible are maintained by algebra isomorphisms. Thus, A_3 and A_4 cannot be isomorphic to any of the others. To verify that A_1 and A_2 are non-isomorphic, we just need to realise that A_1 is a zero-product evolution algebra while A_2 is not. To check that A_5 and A_6 are not isomorphic, note that if $|M_6| = 0$, then the conclusion follows from Corollary 2 as $|M_5| \neq 0$. Otherwise, we have $|M_5| \neq 0$ and $|M_6| \neq 0$, but then the simplified associated graphs do not coincide because the corresponding to M_6 has a loop and the given by M_5 does not have any loop, and Corollary 3 applies. \square

4. Classification of Three-Dimensional Evolution Algebras

In order to start with this classification, we need to see the different structure matrices that we can obtain by just reordering the elements of a natural basis of a three-dimensional evolution algebra. As seen before, if $B = \{e_1, e_2, e_3\}$ is a natural basis of an evolution algebra, A , and σ is a permutation of the set $\{1, 2, 3\}$, then $\{e_{\sigma(1)}, e_{\sigma(2)}, e_{\sigma(3)}\}$ is a natural basis of A . We shall describe below the structure matrix associated to each possible reorganisation of a natural basis B .

Lemma 2. Let A be an evolution algebra and $B = \{e_1, e_2, e_3\}$ a natural basis. Let σ be a permutation of the set $\{1, 2, 3\}$. If the structure matrix of A relative to B is given by $M_B(A)$, then the corresponding structure matrix of A relative to $B_\sigma = \{e_{\sigma(1)}, e_{\sigma(2)}, e_{\sigma(3)}\}$ is $M_{B_\sigma}(A)$ where

$$M_B(A) = \begin{pmatrix} w_{11} & w_{12} & w_{13} \\ w_{21} & w_{22} & w_{23} \\ w_{31} & w_{32} & w_{33} \end{pmatrix} \text{ and } M_{B_\sigma}(A) = \begin{pmatrix} w_{\sigma(1)\sigma(1)} & w_{\sigma(1)\sigma(2)} & w_{\sigma(1)\sigma(3)} \\ w_{\sigma(2)\sigma(1)} & w_{\sigma(2)\sigma(2)} & w_{\sigma(2)\sigma(3)} \\ w_{\sigma(3)\sigma(1)} & w_{\sigma(3)\sigma(2)} & w_{\sigma(3)\sigma(3)} \end{pmatrix}.$$

Proof. Straightforward. \square

We can easily check that, for all permutation σ of the set $\{1, 2, 3\}$ the corresponding associated graphs to $M_{B_\sigma}(A)$ are identical up to relabelling of the vertices (and coincide with the one associated to $M_B(A)$), as seen in Section 2.

From Theorem 1 (see also Remark 5.7 in [3]) we deduce the following result:

Corollary 5. Let A be a non-degenerate evolution algebra and $B = \{e_1, e_2, e_3\}$ a natural basis. Then, A is reducible if and only if $M_B(A)$ is within one of the following types of matrices:

$$M_1 := \begin{pmatrix} w_{11} & 0 & 0 \\ 0 & w_{22} & w_{23} \\ 0 & w_{32} & w_{33} \end{pmatrix}, M_2 := \begin{pmatrix} w_{11} & 0 & w_{13} \\ 0 & w_{22} & 0 \\ w_{31} & 0 & w_{33} \end{pmatrix}, M_3 := \begin{pmatrix} w_{11} & w_{12} & 0 \\ w_{21} & w_{22} & 0 \\ 0 & 0 & w_{33} \end{pmatrix},$$

with $w_{ij} \in \mathbb{K}$, $i, j = 1, 2, 3$.

Proof. By the Theorem 1 we have that A is reducible if and only if the structure matrix of some reordenation of B is diagonal by blocks (that is like either M_1 or M_3). Taking into account the Lemma 2, the result follows. \square

Proposition 3. A three-dimensional evolution algebra A has a two-dimensional basic ideal if and only if there exists a natural basis B with respect to which the structure matrix $M_B(A)$ is of one of the following types of matrices:

$$M_1 = \begin{pmatrix} & & \\ 0 & 0 & \end{pmatrix}; M_2 = \begin{pmatrix} & & \\ 0 & 0 & \end{pmatrix} M_3 = \begin{pmatrix} 0 & 0 & \\ & & \end{pmatrix}. \tag{1}$$

Proof. If $B = \{e_1, e_2, e_3\}$ and if I is a basic ideal of dimension 2 associated to B then, either $I = I_1 := \text{lin}\{e_1, e_2\}$ or $I = I_2 := \text{lin}\{e_1, e_3\}$ or $I = I_3 := \text{lin}\{e_2, e_3\}$. The result follows from the fact that I_i is an ideal of A if and only $M_B(A)$ is like M_i , for $i = 1, 2, 3$, respectively. \square

Finally, we shall see the relationship between two structure matrices $M_B(A)$ and $M_{\tilde{B}}(A)$ of an evolution algebra A associated to two different natural bases B and \tilde{B} . This can be seen in [1], but we shall also show the proof for completeness.

Proposition 4. Let A be an evolution algebra; consider $B = \{e_1, e_2, e_3\}$ and $\tilde{B} = \{\tilde{e}_1, \tilde{e}_2, \tilde{e}_3\}$ two natural bases of A ; and let $M_B(A)$ and $M_{\tilde{B}}(A)$ be the corresponding structure matrices. If $\tilde{e}_i = p_{1i}e_1 + p_{2i}e_2 + p_{3i}e_3$ for $i = 1, 2, 3$, then

$$P = \begin{pmatrix} p_{11} & p_{12} & p_{13} \\ p_{21} & p_{22} & p_{23} \\ p_{31} & p_{32} & p_{33} \end{pmatrix}$$

is a nonsingular matrix such that $PM_{\tilde{B}}(A) = M_B(A)P^{[2]}$, where

$$P^{[2]} = \begin{pmatrix} p_{11}^2 & p_{12}^2 & p_{13}^2 \\ p_{21}^2 & p_{22}^2 & p_{23}^2 \\ p_{31}^2 & p_{32}^2 & p_{33}^2 \end{pmatrix}.$$

Proof. Consider the structure matrices associated, respectively, to both basis:

$$M_B(A) = \begin{pmatrix} \omega_{11} & \omega_{12} & \omega_{13} \\ \omega_{21} & \omega_{22} & \omega_{23} \\ \omega_{31} & \omega_{32} & \omega_{33} \end{pmatrix} \text{ and } M_{\tilde{B}}(A) = \begin{pmatrix} \tilde{\omega}_{11} & \tilde{\omega}_{12} & \tilde{\omega}_{13} \\ \tilde{\omega}_{21} & \tilde{\omega}_{22} & \tilde{\omega}_{23} \\ \tilde{\omega}_{31} & \tilde{\omega}_{32} & \tilde{\omega}_{33} \end{pmatrix}.$$

If $\tilde{e}_i = p_{1i}e_1 + p_{2i}e_2 + p_{3i}e_3$ for $p_{1i}, p_{2i}, p_{3i} \in \mathbb{K}$ and $i \in \{1, 2, 3\}$ then we have, $\tilde{e}_i^2 = q_{1i}e_1 + q_{2i}e_2 + q_{3i}e_3$ where

$$\begin{pmatrix} q_{1i} \\ q_{2i} \\ q_{3i} \end{pmatrix} = \begin{pmatrix} \omega_{11} & \omega_{12} & \omega_{13} \\ \omega_{21} & \omega_{22} & \omega_{23} \\ \omega_{31} & \omega_{32} & \omega_{33} \end{pmatrix} \begin{pmatrix} p_{1i}^2 \\ p_{2i}^2 \\ p_{3i}^2 \end{pmatrix}.$$

But also $\tilde{e}_i^2 = \tilde{\omega}_{1i}\tilde{e}_1 + \tilde{\omega}_{2i}\tilde{e}_2 + \tilde{\omega}_{3i}\tilde{e}_3$ and so

$$\begin{pmatrix} p_{11} & p_{12} & p_{13} \\ p_{21} & p_{22} & p_{23} \\ p_{31} & p_{32} & p_{33} \end{pmatrix} \begin{pmatrix} \tilde{\omega}_{1i} \\ \tilde{\omega}_{2i} \\ \tilde{\omega}_{3i} \end{pmatrix} = \begin{pmatrix} q_{1i} \\ q_{2i} \\ q_{3i} \end{pmatrix}.$$

Hence $PM_{\tilde{B}}(A) = M_B(A)P^{[2]}$ as desired. \square

4.1. The Non-Degenerate Case

According to Theorem 1, if an evolution algebra A has no basic proper ideals then A is irreducible. In particular, if $\dim A = 3$ and A has no basic ideals of dimension two, then A is irreducible. In the following result we shall characterize this fact for a particular type of evolution algebra. Recall that two $*$ symbols in the same matrix do not necessarily have the same non-zero value.

Lemma 3. *Let A be an evolution algebra with a natural basis $B = \{e_1, e_2, e_3\}$ with respect to which the structure matrix is like*

$$M_B(A) = \begin{pmatrix} * & * & w \\ 0 & * & * \\ 0 & * & * \end{pmatrix}$$

with $|M_B(A)| = 0$. Then, A has a basic ideal with dimension 2 if and only if $M_B(A) = \begin{pmatrix} * & a & \lambda a \\ 0 & b & \lambda b \\ 0 & c & \lambda c \end{pmatrix}$ with

$a, b, c, \lambda \in \mathbb{K} \setminus \{0\}$ and $\lambda \neq -\frac{b^2}{c^2}$.

Proof. Suppose that I is a basic ideal of A with $\dim I = 2$. Since it cannot be $I = \mathbb{K}e_1$ (as the ideal I has dimension 2), we obtain that $e_2^2 = ae_1 + be_2 + ce_3$ belongs to I . In fact, either $\pi_2(I) \neq 0$ or $\pi_3(I) \neq 0$. Thus, if $u \in I$ is such that $\pi_2(u) \neq 0$, then $e_2u = \mu e_2^2 \in I$ for some $\mu \neq 0$, and hence, $e_2^2 \in I$. Similarly if $\pi_3(u) \neq 0$ then $e_3^2 \in I$ and it follows that $e_2^2 \in I$ as $e_2e_3^2 \in I$. Consequently $e_1^2 = \frac{1}{a}e_1e_2^2 \in I$. Therefore, $e_1 \in I$ and $I = \text{lin}\{e_1, be_2 + ce_3\}$. Since I is a basic ideal, there exists $v = \alpha e_1 + \beta e_2 + \gamma e_3$ such that $\tilde{B} = \{e_1, be_2 + ce_3, v\}$ is a natural basis of A . From $ve_1 = 0$ we obtain that $\alpha = 0$, and hence, $v = \beta e_2 + \gamma e_3$ where $|\beta| + |\gamma| \neq 0$. Since,

$$(be_2 + ce_3)v = (be_2 + ce_3)(\beta e_2 + \gamma e_3) = b\beta e_2^2 + c\gamma e_3^2 = 0,$$

it follows that $\beta \neq 0$ and $\gamma \neq 0$ simultaneously, as $e_2^2 \neq 0$ and $e_3^2 \neq 0$. Hence, we obtain that e_2^2 and e_3^2 are proportional and non-zero. In fact, $e_3^2 = \lambda e_2^2$ with $\lambda = -\frac{b\beta}{c\gamma}$. Moreover, as $be_2 + ce_3$ and $v = \beta e_2 + \gamma e_3$ are linearly independent we obtain that $b\gamma - c\beta \neq 0$. Thus, $\frac{\beta}{\gamma} \neq \frac{b}{c}$ and so $\lambda \neq -\frac{b^2}{c^2}$.

Conversely, if $e_3^2 = \lambda e_2^2$ with $\lambda \neq -\frac{b^2}{c^2}$, then it follows that $B = \{e_1, be_2 + ce_3, e_2 - \frac{b}{c\lambda}e_3\}$ is a natural basis of A and $I := \text{lin}\{e_1, be_2 + ce_3\}$ is a proper two-dimensional basic ideal. \square

Theorem 5. *Let A be a three-dimensional, irreducible, non-degenerate evolution algebra. Then, A has a one-dimensional basic ideal and has no two-dimensional basic ideals if and only if A has a natural basis B such that the structure matrix $M_B(A)$ is within the following types (non-isomorphic each other), where $*$ denotes a non-zero scalar and $w, \tilde{w} \in \mathbb{K}$:*

- (i) $M_1 = \begin{pmatrix} * & * & 0 \\ 0 & w & * \\ 0 & * & \tilde{w} \end{pmatrix}$ with $|M_1| \neq 0$.
- (ii) $M_2 = \begin{pmatrix} * & * & * \\ 0 & w & * \\ 0 & * & \tilde{w} \end{pmatrix}$ with $|M_2| \neq 0$.
- (iii) $M_3 = \begin{pmatrix} * & * & w \\ 0 & * & * \\ 0 & * & * \end{pmatrix}$ with $|M_3| = 0$ and either $M_3 = \begin{pmatrix} * & a & -\frac{b^2}{c^2}a \\ 0 & b & -\frac{b^2}{c^2}b \\ 0 & c & -\frac{b^2}{c^2}c \end{pmatrix}$ or M_3 having no proportional columns.

Proof. For the sufficient condition, suppose that A has a natural basis of type M_1, M_2 , or M_3 . Then A has a basic ideal with dimension 1 (namely $I = \mathbb{K}e_1$). Moreover, A does not have a basic ideal with dimension 2. In the case of M_1 and M_2 this last assertion follows from the fact that all the natural basis of A are related by Proposition 1, and from Lemma 2 none of the related natural basis of M_1 or M_2 are of the type

$$M_a = \begin{pmatrix} * & * \\ 0 & 0 \end{pmatrix}, M_b = \begin{pmatrix} * & * \\ 0 & 0 \end{pmatrix}, M_c = \begin{pmatrix} * & * \\ 0 & 0 \end{pmatrix}, \tag{2}$$

which together with Proposition 3 shows that A does not have any two-dimensional basic ideals. In the case of M_3 , it follows from Lemma 3 that A does not have any two-dimensional basic ideals.

For the necessary condition, suppose that A has a basic ideal of dimension one and does not have any two-dimensional basic ideals. Then, it is not restrictive to assume that $I = \mathbb{K}e_1$ is a one-dimensional basic ideal. On the other hand, by Proposition 3, we have that $M_B(A)$ is not in any of the situations of (2) (otherwise A has a two-dimensional basic ideal). As A is non-degenerate and $I = \mathbb{K}e_1$ is a basic ideal we have that

$$\begin{pmatrix} * \\ 0 \\ 0 \end{pmatrix}.$$

In order to not be in the cases of (2), we must have

$$\begin{pmatrix} * & & \\ 0 & * & \\ 0 & * & \end{pmatrix}.$$

Still, this matrix could be of the type M_c in (2). Consequently we have the two following possibilities:

Case 1. $\begin{pmatrix} * & 0 \\ 0 & * \\ 0 & * \end{pmatrix}$. Again because of (2) we must have $\begin{pmatrix} * & * & 0 \\ 0 & & * \\ 0 & * & \end{pmatrix}$. Therefore, $M_B(A)$ is of type M_1 .

Case 2. $\begin{pmatrix} * & * \\ 0 & * \\ 0 & * \end{pmatrix}$. Here we have either $\begin{pmatrix} * & * & * \\ 0 & & * \\ 0 & * & \end{pmatrix}$ or $\begin{pmatrix} * & 0 & * \\ 0 & & * \\ 0 & * & \end{pmatrix}$.

Case 2.1. $M_B(A) = \begin{pmatrix} * & * & * \\ 0 & & * \\ 0 & * & \end{pmatrix}$. We consider the following situations:

Case 2.1.1. $M_B(A) = \begin{pmatrix} * & * & * \\ 0 & & * \\ 0 & * & \end{pmatrix}$ with $|M_B(A)| \neq 0$ and we are within the type M_2 .

Case 2.1.2. $M_B(A) = \begin{pmatrix} * & * & * \\ 0 & & * \\ 0 & * & \end{pmatrix} = \begin{pmatrix} * & * & * \\ 0 & w & * \\ 0 & * & \tilde{w} \end{pmatrix}$ with $|M_B(A)| = 0$. Therefore we have that

$\left| \begin{pmatrix} w & * \\ * & \tilde{w} \end{pmatrix} \right| = 0$ so $w \neq 0$ and $\tilde{w} \neq 0$. Consequently, $M_B(A) = \begin{pmatrix} * & * & * \\ 0 & * & * \\ 0 & * & * \end{pmatrix}$ with $|M_B(A)| = 0$,

which is included in $M = \begin{pmatrix} * & * & w \\ 0 & * & * \\ 0 & * & * \end{pmatrix}$ with $|M| = 0$. By Lemma 3 either $M_B(A) = \begin{pmatrix} * & * & w \\ 0 & * & * \\ 0 & * & * \end{pmatrix}$

with no proportional columns or, $M_B(A) = \begin{pmatrix} * & a & -\frac{b^2}{c^2}a \\ 0 & b & -\frac{b^2}{c^2}b \\ 0 & c & -\frac{b^2}{c^2}c \end{pmatrix}$ otherwise.

Finally, note that these three types of algebra are not isomorphic. In fact, if A_i is an evolution algebra with a structure matrix of the type M_i respectively, for $i = 1, 2, 3$, then obviously A_i with $i = 1, 2$ is not isomorphic to A_3 because $|M_3| = 0$ and $|M_i| \neq 0$ for $i = 1, 2$. Also, from Corollary 2 all the natural basis of A_1 are related, and from Lemma 2 we know that A_1 does not have a related structure matrix of the type M_2 . Therefore, A_1 and A_2 are not isomorphic. \square

Theorem 6. Let A be an irreducible, three-dimensional, non-degenerate evolution algebra. Then, A has a basic ideal of dimension two if and only if there exists a natural basis B such that the structure matrix associated $M_B(A)$ is within the following type, where $*$ denotes a non-zero scalar and $\alpha, \beta, \gamma \in \mathbb{K}$:

$$M_B(A) = \begin{pmatrix} & \alpha & \gamma \\ \beta & & * \\ 0 & 0 & \end{pmatrix}, \text{ with } |\alpha| + |\beta| + |\gamma| \neq 0, \text{ and no zero columns.}$$

Proof. As A has a two-dimensional basic ideal, there is a natural basis B such that the structure matrix is

$$M_B(A) = \begin{pmatrix} & & \\ & & \\ 0 & 0 & \end{pmatrix}.$$

Also, by Theorem 1, we have that the two first entries of the third column cannot be zero simultaneously (otherwise A is reducible). We have then two possibilities:

$$\begin{pmatrix} & * \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} & \\ 0 & 0 \end{pmatrix}. \tag{3}$$

In addition, we cannot have the following cases

$$\begin{pmatrix} 0 & \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & \end{pmatrix}$$

as in the first one $A = I \oplus J$ with $I = \mathbb{K}e_2$ and $J = \text{lin}\{e_1, e_3\}$, and in the second one $A = I \oplus J$ with $I = \mathbb{K}e_1$ and $J = \text{lin}\{e_2, e_3\}$. Hence, applying this to (3) we have either

$$\begin{pmatrix} \beta & * \\ \alpha & \gamma \\ 0 & 0 \end{pmatrix} \text{ or } \begin{pmatrix} \beta & \gamma \\ \alpha & * \\ 0 & 0 \end{pmatrix} \tag{4}$$

with $|\alpha| + |\beta| + |\gamma| \neq 0$. Nevertheless, if $B = \{e_1, e_2, e_3\}$ is such that $M_B(A) = \begin{pmatrix} \beta & * \\ \alpha & \gamma \\ 0 & 0 \end{pmatrix}$ and we consider $\tilde{B} = \{e_2, e_1, e_3\}$, then the structure matrix relative to the new natural basis is $M_{\tilde{B}}(A) = \begin{pmatrix} \alpha & \gamma \\ \beta & * \\ 0 & 0 \end{pmatrix}$. So in both cases we arrive (after reorganisation of the basis if needed) at a structure

matrix of the type $\begin{pmatrix} \alpha & \gamma \\ \beta & * \\ 0 & 0 \end{pmatrix}$ with $|\alpha| + |\beta| + |\gamma| \neq 0$ and non-zero columns.

Reciprocally, if A is an evolution algebra and B a natural basis of A such that the structure matrix associated to it is $M_B(A) = \begin{pmatrix} \alpha & \gamma \\ \beta & * \\ 0 & 0 \end{pmatrix}$ without zero columns and with $|\alpha| + |\beta| + |\gamma| \neq 0$, then $I = \text{lin}\{e_1, e_2\}$, is a basic ideal of A and by Corollary 5 it is clear that A is irreducible. \square

Lemma 4. Let A be a non-degenerate evolution algebra such that $\dim A^2 = 2$. Then, A^2 is a basic ideal if and only if there exists a natural basis $B = \{e_1, e_2, e_3\}$ such that

$$M_B(A) = \begin{pmatrix} \alpha_1 & \beta_1 & \gamma_1 \\ \alpha_2 & \beta_2 & \gamma_2 \\ 0 & 0 & 0 \end{pmatrix}, \tag{5}$$

with non zero columns and $\alpha_i, \beta_i, \gamma_i \in \mathbb{K}$, for $i = 1, 2$.

Proof. If A^2 is a proper basic ideal with $\dim A^2 = 2$, then there exists a natural basis $B = \{e_1, e_2, e_3\}$ such that $A^2 = \text{lin}\{e_1, e_2\}$. Thus, $e_3x = 0$ for every $x \in A^2$. As B is a natural basis, there exist $\alpha_i, \beta_i, \gamma_i \in \mathbb{K}$ for $i = 1, 2, 3$ such that

$$\begin{aligned} e_1^2 &= \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \\ e_2^2 &= \beta_1 e_1 + \beta_2 e_2 + \beta_3 e_3, \\ e_3^2 &= \gamma_1 e_1 + \gamma_2 e_2 + \gamma_3 e_3. \end{aligned}$$

Equivalently, the structure matrix of A relative to B is

$$M_B(A) = \begin{pmatrix} \alpha_1 & \beta_1 & \gamma_1 \\ \alpha_2 & \beta_2 & \gamma_2 \\ \alpha_3 & \beta_3 & \gamma_3 \end{pmatrix}.$$

Since $e_3e_i^2 = 0$ for $i = 1, 2, 3$ and $e_3^2 \neq 0$ as A is non-degenerate, $\alpha_3 = \beta_3 = \gamma_3 = 0$. Thus,

$$M_B(A) = \begin{pmatrix} \alpha_1 & \beta_1 & \gamma_1 \\ \alpha_2 & \beta_2 & \gamma_2 \\ 0 & 0 & 0 \end{pmatrix}$$

and $A^2 = \text{lin}\{e_1^2, e_2^2, e_3^2\}$ is such that $\dim A^2 = 2$. Clearly, all columns are non-zero, as A is a non-degenerate algebra.

Conversely, suppose there exists a natural basis $B = \{e_1, e_2, e_3\}$ such that the structure matrix of A is given by (5). Then $A^2 = \text{lin}\{e_1, e_2\}$ is a basic ideal with dimension 2, as desired. \square

Lemma 5. *Let A be an evolution algebra and let $B = \{e_1, e_2, e_3\}$ be a natural basis such that the structure matrix of A relative to B is of the type*

$$M_B(A) = \begin{pmatrix} & * & \\ \alpha & & * \\ \beta & \gamma & \end{pmatrix} \text{ with } |\beta| + |\alpha\gamma| \neq 0.$$

If $\dim A^2 = 1$ then A has proper basic ideals.

Proof. If $\beta = 0$ then $\alpha \neq 0$ and $\gamma \neq 0$, so $\dim A^2 = 2$, a contradiction. Thus, $\beta \neq 0$, and hence, since the columns of $M_B(A)$ are proportional, as $\dim A^2 = 1$, we have

$$M_B(A) = \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix} = \begin{pmatrix} a & \lambda a & \mu a \\ b & \lambda b & \mu b \\ c & \lambda c & \mu c \end{pmatrix},$$

with $a, b, c, \lambda, \mu \in \mathbb{K} \setminus \{0\}$. We shall split the proof in two cases:

Case 1. $\lambda b^2 + \mu c^2 \neq 0$. Then, $B = \{e_1, be_2 + ce_3, e_2 - \frac{b\lambda}{c\mu}e_3\}$ is a natural basis of A and $I_1 = \text{lin}\{e_1, be_2 + ce_3\}$ is a basic ideal of A .

Case 2. $\lambda b^2 + \mu c^2 = 0$. Then, $\lambda = -\frac{\mu c^2}{b^2}$, so $e_2^2 = -\frac{\mu c^2}{b^2}e_1^2$. Thus,

$$M_B(A) = M_1 = \begin{pmatrix} * & * & * \\ * & * & * \\ * & * & * \end{pmatrix} = \begin{pmatrix} a & -\frac{\mu c^2}{b^2}a & \mu a \\ b & -\frac{\mu c^2}{b^2}b & \mu b \\ c & -\frac{\mu c^2}{b^2}c & \mu c \end{pmatrix}.$$

Let us consider $\tilde{B} = \{\tilde{e}_1, \tilde{e}_2, \tilde{e}_3\} = \{e_1, \frac{b}{c}e_2, e_3\}$. Then, by applying Proposition 2 with $\alpha_1 = 1$, $\alpha_2 = \frac{b}{c}$, and $\alpha_3 = 1$ we get that $\alpha_2^2 \frac{c^2}{b^2} = 1$ and

$$M_{\tilde{B}}(A) = \begin{pmatrix} \alpha_1 a & -\frac{\alpha_2^2}{\alpha_1} \frac{\mu c^2}{b^2} a & \frac{\alpha_3^2}{\alpha_1} \mu a \\ \frac{\alpha_2}{\alpha_1} b & -\alpha_2 \frac{\mu c^2}{b^2} b & \frac{\alpha_3}{\alpha_2} \mu b \\ \frac{\alpha_3}{\alpha_1} c & -\frac{\alpha_2^2}{\alpha_3} \frac{\mu c^2}{b^2} c & \alpha_3 \mu c \end{pmatrix} = \begin{pmatrix} a & -\mu a & \mu a \\ c & -\mu c & \mu c \\ c & -\mu c & \mu c \end{pmatrix}.$$

We shall consider two different cases again.

Case 2.1. $a^2 + \mu c^2 \neq 0$. Then, $B = \{\tilde{e}_2, a\tilde{e}_1 + c\tilde{e}_3, c\tilde{e}_1 - \frac{a}{\mu}\tilde{e}_3\}$ is a natural basis and $I_{e_2} = \text{lin}\{\tilde{e}_2, a\tilde{e}_1 + c\tilde{e}_3\}$ is a basic ideal of dimension two.

Case 2.2. $a^2 + \mu c^2 = 0$. Then,

$$M_{\tilde{B}}(A) = \begin{pmatrix} a & -\mu a & \mu a \\ c & -\mu c & \mu c \\ c & -\mu c & \mu c \end{pmatrix} = \begin{pmatrix} a & \frac{a^2}{c^2}a & -\frac{a^2}{c^2}a \\ c & \frac{a^2}{c^2}c & -\frac{a^2}{c^2}c \\ c & \frac{a^2}{c^2}c & -\frac{a^2}{c^2}c \end{pmatrix} = \begin{pmatrix} a & \frac{a^3}{c^2} & -\frac{a^3}{c^2} \\ c & \frac{a^2}{c} & -\frac{a^2}{c} \\ c & \frac{a^2}{c} & -\frac{a^2}{c} \end{pmatrix}.$$

Let us consider now $\tilde{B} = \{\tilde{e}_1, \tilde{e}_2, \tilde{e}_3\} = \{\alpha_1 \tilde{e}_1, \alpha_2 \tilde{e}_2, \alpha_3 \tilde{e}_3\} = \{\frac{1}{a} \tilde{e}_1, -\frac{c}{a^2} \tilde{e}_2, -\frac{c}{a^2} \tilde{e}_3\}$. By applying Proposition 2 with $\alpha_1 = \frac{1}{a}$, $\alpha_2 = -\frac{c}{a^2}$, and $\alpha_3 = \frac{c}{a^2}$ we have,

$$M_{\hat{B}}(A) = \begin{pmatrix} \alpha_1 a & \frac{\alpha_2^2 a^3}{\alpha_1 c^2} & -\frac{\alpha_3^2 a^3}{\alpha_1 c^2} \\ \frac{\alpha_1^2}{\alpha_2} c & \alpha_2 \frac{a^2}{c} & -\frac{\alpha_3^2 a^2}{\alpha_2 c} \\ \frac{\alpha_1^2}{\alpha_3} c & \frac{\alpha_2^2 a^2}{\alpha_3 c} & -\alpha_3 \frac{a^2}{c} \end{pmatrix} = \begin{pmatrix} 1 & 1 & -1 \\ -1 & -1 & 1 \\ 1 & 1 & -1 \end{pmatrix}.$$

Thus, $B = \{\hat{e}_1^2, u, v\}$, where $\hat{e}_1^2 = \hat{e}_1 - \hat{e}_2 + \hat{e}_3$; $u = \hat{e}_1 + \hat{e}_2$; $v = \hat{e}_1 - \hat{e}_2 + 2\hat{e}_3$ is a natural basis of A and A^2 is the ideal generated by \hat{e}_1^2 . Note that A^2 is a basic proper one-dimensional ideal. \square

From Theorem 1 we can deduce that whenever a non-degenerate evolution algebra A has no basic ideals of dimension one or two, then A is irreducible. We shall obtain a necessary and sufficient condition for this property.

Lemma 6. *Let A be a non-degenerate three-dimensional evolution algebra with no proper basic ideals, and let $M_B(A)$ be the structure matrix of A with respect to a natural basis B . Then $M_B(A)$ cannot be within any of the following types of matrices*

$$M_1 = \begin{pmatrix} & \\ 0 & \\ 0 & \end{pmatrix}; M_2 = \begin{pmatrix} 0 & \\ & \\ 0 & \end{pmatrix}; M_3 = \begin{pmatrix} & 0 \\ & 0 \\ & \end{pmatrix}$$

$$M_4 = \begin{pmatrix} & \\ & \\ 0 & 0 \end{pmatrix}; M_5 = \begin{pmatrix} & \\ 0 & 0 \\ & \end{pmatrix}; M_6 = \begin{pmatrix} & 0 & 0 \\ & & \\ & & \end{pmatrix}.$$

Proof. If $M_B(A) = M_i$ with $i = 1, 2, 3$ then $\mathbb{K}e_i$ is a basic ideal of dimension 1 (with $i = 1, 2, 3$ respectively). Similarly, by Proposition 3, the structure matrix of A has to be different from M_i for $i = 4, 5, 6$, as otherwise A has a basic ideal of dimension two (namely, $I_4 = \text{lin}\{e_1, e_2\}$, $I_5 = \text{lin}\{e_1, e_3\}$ and $I_6 = \text{lin}\{e_2, e_3\}$ respectively). \square

Theorem 7. *Let A be a non-degenerate three-dimensional evolution algebra with no proper basic ideals. Then, A has a natural basis $B = \{e_1, e_2, e_3\}$ such that*

$$M_B(A) = \begin{pmatrix} & * & \\ \alpha & & * \\ \beta & \gamma & \end{pmatrix} \text{ with } |\beta| + |\alpha\gamma| \neq 0. \tag{6}$$

Moreover, B can be reordered in a way such that either

$$M_B(A) = M_1 := \begin{pmatrix} & * & \\ & & * \\ * & & \end{pmatrix} \text{ or } M_B(A) = M_2 := \begin{pmatrix} & * & 0 \\ * & & * \\ 0 & * & \end{pmatrix}.$$

Proof. First, we shall see that whenever $M_B(A)$ is like (6), then B can be reordered such that either $M_B(A) = M_1$ or $M_B(A) = M_2$. Indeed, if $\beta \neq 0$ then $M_B(A) = M_1$. Otherwise, $\beta = 0$ and so

$$M_B(A) = \begin{pmatrix} & * & \\ * & & * \\ 0 & * & \end{pmatrix}.$$

Then either $M_B(A) = M_2 = \begin{pmatrix} & * & 0 \\ * & & * \\ 0 & * & \end{pmatrix}$ or $M_B(A) = \begin{pmatrix} & * & * \\ * & & * \\ 0 & * & \end{pmatrix}$. But the latter case is gathered in $M_B(A) = \begin{pmatrix} & * & \\ * & & * \\ & * & \end{pmatrix}$. If we consider the reordering $B' = \{e_1, e_3, e_2\}$ then, by Lemma 2, we have

$$M_B(A) = \begin{pmatrix} & * & \\ * & & * \\ & * & \end{pmatrix} \equiv M_{B'}(A) = \begin{pmatrix} & * & \\ & & * \\ * & & \end{pmatrix} = M_1, \tag{7}$$

which proves the claim.

We shall prove now that whenever A is non-degenerate and has no proper basic ideals, then A has a natural basis $B = \{e_1, e_2, e_3\}$ such that

$$M_B(A) = \begin{pmatrix} & * & \\ \alpha & & * \\ \beta & \gamma & \end{pmatrix} \text{ with } |\beta| + |\alpha\gamma| \neq 0. \tag{8}$$

We shall split the proof into the following two cases: $\beta = 0$ and $\beta \neq 0$ (Cases 1 and 2).

Case 1. $M_B(A) = \begin{pmatrix} & * & \\ & & * \\ 0 & & \end{pmatrix}$. By Lemma 6 we must have $\begin{pmatrix} * & \\ 0 & * \end{pmatrix}$. Again by Lemma 6 the choices are the following:

Case 1.1 $\begin{pmatrix} & * & \\ * & & * \\ 0 & * & \end{pmatrix}$ which is a particular case of $\begin{pmatrix} & * & \\ * & & * \\ & * & \end{pmatrix}$ which is equivalent to M_1 by (7).

Case 1.2 $\begin{pmatrix} & & 0 \\ * & & * \\ 0 & * & \end{pmatrix}$. Hence, again from Lemma 6, we have $\begin{pmatrix} & * & 0 \\ * & & * \\ 0 & * & \end{pmatrix}$, so we arrive to M_2 .

Case 2. $M_B(A) = \begin{pmatrix} & & \\ & & \\ * & & \end{pmatrix}$. Here we consider the following situations:

Case 2.1 $\begin{pmatrix} & & 0 \\ * & & \\ & & \end{pmatrix}$. By Lemma 6 we have that $\begin{pmatrix} & * & 0 \\ & & * \\ * & & \end{pmatrix}$, but this is a particular case of M_1 .

Case 2.2 $\begin{pmatrix} & & * \\ & & \\ * & & \end{pmatrix}$. We have the following possibilities:

$$\begin{pmatrix} & * & \\ * & & \\ * & & \end{pmatrix} \text{ and } \begin{pmatrix} & * & \\ 0 & & \\ * & & \end{pmatrix}.$$

Case 2.2.1 $\begin{pmatrix} & * \\ * & \\ * & \end{pmatrix}$. Since by Lemma 6 we cannot have $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ then we get the cases

$$\begin{pmatrix} & * \\ * & \\ * & * \end{pmatrix} \text{ and } \begin{pmatrix} & * & * \\ * & \\ * & \end{pmatrix}.$$

Case 2.2.1.1 $\begin{pmatrix} & * \\ * & \\ * & * \end{pmatrix}$. This is a particular case of $\begin{pmatrix} & * \\ * & \\ * & * \end{pmatrix}$ that is equivalent to M_1 by (7).

Case 2.2.1.2 $\begin{pmatrix} & * & * \\ * & \\ * & \end{pmatrix}$. Here we have either $\begin{pmatrix} & * & * \\ * & \\ * & * \end{pmatrix}$ or $\begin{pmatrix} & * & * \\ * & \\ * & 0 \end{pmatrix}$. The matrix

$\begin{pmatrix} & * & * \\ * & \\ * & * \end{pmatrix}$ is gathered in $\begin{pmatrix} & * \\ * & \\ * & \end{pmatrix}$. For the case $\begin{pmatrix} & * & * \\ * & \\ * & 0 \end{pmatrix}$ the choices are

$$\begin{pmatrix} & * & * \\ * & 0 \\ * & 0 \end{pmatrix} \text{ and } \begin{pmatrix} & * & * \\ * & \\ * & 0 \end{pmatrix}.$$

The first one is equivalent to $\begin{pmatrix} & * & 0 \\ * & * \\ 0 & * \end{pmatrix}$ for $B' = \{e_2, e_1, e_3\}$, and the second one is contained

in the case $\begin{pmatrix} & * & \\ * & * \\ * & \end{pmatrix}$ and hence in $\begin{pmatrix} & * \\ * & \\ * & \end{pmatrix}$ by (7).

Case 2.2.2. $\begin{pmatrix} & * \\ 0 & \\ * & \end{pmatrix}$. From Lemma 6, we arrive to $\begin{pmatrix} & * \\ 0 & * \\ * & \end{pmatrix}$. Now the choices are either

$\begin{pmatrix} & * & * \\ 0 & * \\ * & \end{pmatrix}$ or $\begin{pmatrix} & 0 & * \\ 0 & * \\ * & \end{pmatrix}$. The first case is contained in $\begin{pmatrix} & * \\ * & \\ * & \end{pmatrix}$ for $B' = \{e_1, e_3, e_2\}$.

For the second one, by Lemma 6, we obtain $\begin{pmatrix} & 0 & * \\ 0 & * \\ * & * \end{pmatrix}$ which is equivalent to $\begin{pmatrix} & * & 0 \\ * & * \\ 0 & * \end{pmatrix}$ by

considering $B' = \{e_1, e_3, e_2\}$. \square

Corollary 6. Let A be a non-degenerate three-dimensional evolution algebra. Then the following assertions are equivalent:

- (i) A has no basic proper ideals;
- (ii) A has a natural basis B with respect to which,

$$M_B(A) = \begin{pmatrix} & * \\ \alpha & * \\ \beta & \gamma \end{pmatrix} \text{ with } |\beta| + |\alpha\gamma| \neq 0, \tag{9}$$

and either $\dim A^2 = 3$ or $\dim A^2 = 2$, and there does not exist a nonsingular matrix P such that $PM_{\bar{B}}(A) = M_B(A)P^{[2]}$ where $M_{\bar{B}}(A)$ is a matrix of the type

$$M_{\tilde{B}}(A) = \begin{pmatrix} \alpha_1 & \beta_1 & \gamma_1 \\ \alpha_2 & \beta_2 & \gamma_2 \\ 0 & 0 & 0 \end{pmatrix}.$$

Proof. (i) \implies (ii) Let us suppose that A has no basic proper ideals. Then, by Theorem 7 there is a natural basis B such that its structure matrix is within the type (9). Also, by Lemma 5 we know that if $\dim A^2 = 1$, A has basic ideals. If $\dim A^2 = 3$ then A has no proper ideals. Indeed, as β and γ cannot be zero simultaneously, it follows that every non-zero ideal I contains A^2 . If $\dim A^2 = 2$ the conclusion follows from Lemma 4 joint with Proposition 4 (In this last case, the entries of third row of $M_B(A)$ cannot be zero simultaneously).

(ii) \implies (i) Whenever A has a structure matrix of the type (9), then any ideal contains A^2 . Hence, if $\dim A^2 = 3$ then the conclusion is clear, and if $\dim A^2 = 2$, then by Lemma 4, A cannot have a proper ideal. \square

The following result is nothing but Corollary 6 in the particular case that $\dim A^2 = 2$: keep in mind Proposition 4 and the fact that the associated graph to a natural basis $B = \{e_1, e_2, e_3\}$ does not have a source in e_i if and only if the i th row of the structure matrix associated to B is a zero row.

Corollary 7. *Let A be a non-degenerate evolution algebra with $\dim A^2 = 2$. Then A has no proper basic ideals if and only if A has a natural basis respect to which the structure matrix is of the type (9) and all the natural bases of A have an associated graph with no source vertices (this is a graph such that every vertex has some incoming edge).*

We shall study now when a non-degenerate evolution algebra A is reducible. In this case, $A = I \oplus J$ where $\dim I = 1$ and it has no zero product (otherwise A is degenerate) and $\dim J = 2$ with J a non-degenerate two-dimensional basic ideal. Therefore, by Theorem 4, J has a natural basis $\{e_2, e_3\}$ with respect to which the structure matrix is of the type $M_{J_1} = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}$, $M_{J_2} = \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix}$ or $M_{J_3} = \begin{pmatrix} w & \tilde{w} \\ * & * \end{pmatrix}$, being that these types are non-isomorphic. Thus, A has a natural basis $B = \{e_1, e_2, e_3\}$ with respect to which $M_B(A)$ is within the following types:

$$M_1 = \begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ 0 & 0 & * \end{pmatrix}, M_2 = \begin{pmatrix} * & 0 & 0 \\ 0 & 0 & * \\ 0 & * & 0 \end{pmatrix}, M_3 = \begin{pmatrix} * & 0 & 0 \\ 0 & w & \tilde{w} \\ 0 & * & * \end{pmatrix}.$$

We shall see now that these algebras are non-isomorphic. In order to do so, the following Lemma shall be useful.

Lemma 7. *Let A be a three-dimensional evolution algebra such that $A = I \oplus J$ with $\dim I = 1$ and $\dim J = 2$. If J is irreducible, then the decomposition of A is unique.*

Proof. Let us suppose that $A = I \oplus J = \hat{I} \oplus \hat{J}$ where $\dim I = \dim \hat{I} = 1$. Then, $J \simeq A/I = (\hat{I} \oplus \hat{J})/I \simeq \hat{I}/I \oplus \hat{J}/I$. As J cannot be decomposed we have that $\hat{I}/I = 0$; note that $\dim \hat{I}/I \leq 1$. Then, $I = \hat{I}$ as $\dim I = \dim \hat{I} = 1$, so $J \subseteq \hat{J}$ or equivalently $J = \hat{J}$, as they have the same dimension. \square

Theorem 8. *Let A be a three-dimensional reducible and non-degenerate evolution algebra. Then, A has a natural basis B such that the structure matrix associated to it, $M_B(A)$, is within the following non-isomorphic types:*

$$M_1 = \begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ 0 & 0 & * \end{pmatrix}, M_2 = \begin{pmatrix} * & 0 & 0 \\ 0 & 0 & * \\ 0 & * & 0 \end{pmatrix}, M_3 = \begin{pmatrix} * & 0 & 0 \\ 0 & w & \tilde{w} \\ 0 & * & * \end{pmatrix},$$

with $w, \tilde{w} \in \mathbb{K}$.

Proof. As said above, by Theorem 4 there exists a natural base B such that $M_B(A)$ has the form of M_1, M_2 or M_3 . Let us denote by A_i the algebra given by the structure matrix M_i with $i = 1, 2, 3$. By the above Lemma it follows that the decomposition of A_2, A_3 is unique, and so A_i are non-isomorphic, for $i = 2, 3$, as J_i are not isomorphic either. But similarly, neither A_2 nor A_3 can be isomorphic to A_1 , as J_1 is reducible and J_2, J_3 are irreducible. \square

4.2. The Degenerate Case

The following result describes whether an evolution algebra is reducible according with the number of elements in the natural basis having zero square.

Theorem 9. Let A be a degenerate evolution algebra and let $B = \{e_1, e_2, e_3\}$ be a natural basis of A .

- (i) Suppose that $e_1^2 = e_2^2 = e_3^2 = 0$. Then, A has zero product and is reducible.
- (ii) Suppose that $e_1^2 = e_2^2 = 0$ and $e_3^2 \neq 0$. Then, A is reducible.
- (iii) Suppose that $e_1^2 = 0, e_2^2 \neq 0$ and $e_3^2 \neq 0$. Then we have one of the following situations

(a) e_2^2 and e_3^2 are linearly dependent, and so $M_B(A) = \begin{pmatrix} 0 & \alpha & t\alpha \\ 0 & \beta & t\beta \\ 0 & \gamma & t\gamma \end{pmatrix}$. Then:

(a.1) A is reducible if and only if $|\beta| + |\gamma| \neq 0$.

(a.2) A is irreducible if and only if $M_B(A) = \begin{pmatrix} 0 & * & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$.

(b) e_2^2 and e_3^2 are linearly independent. Then:

(b.1) A is reducible if and only if $M_B(A) = \begin{pmatrix} 0 & w_{12} & w_{13} \\ 0 & w_{22} & w_{23} \\ 0 & w_{32} & w_{33} \end{pmatrix}$ with $w_{22}w_{33} - w_{32}w_{23} \neq 0$.

(b.2) A is irreducible if and only if $M_B(A) = \begin{pmatrix} 0 & w & \tilde{w} \\ 0 & \alpha & \alpha t \\ 0 & \beta & \beta t \end{pmatrix}$ with $\tilde{w} \neq wt$ and $|\alpha| + |\beta| \neq 0$.

Proof. (i) It is clear that if A has a zero product then A is reducible. Indeed, $A = I \oplus J$ where $I = \text{lin}\{e_1, e_2\}$ and $J = \mathbb{K}e_3$.

(ii) Suppose that $e_1^2 = e_2^2 = 0$ and $e_3^2 \neq 0$. Consider $e_3^2 = w_{13}e_1 + w_{23}e_2 + w_{33}e_3$. We have the following possibilities:

Case (ii)(1) $w_{33} \neq 0$. Then, $A = I \oplus J$ where $I = \text{lin}\{e_1, e_2\}$ and $J = \mathbb{K}e_3^2$.

Case (ii)(2) $w_{33} = 0$. We shall consider the following situations.

Case (ii)(2.1) $w_{23} \neq 0$. Then $A = I \oplus J$ being $I = \text{lin}\{e_3, e_3^2\} = \text{lin}\{e_3, w_{13}e_1 + w_{23}e_2\}$ and $J = \mathbb{K}e_1$.

Case (ii)(2.2) $w_{23} = 0$, which immediately implies $w_{13} \neq 0$. Then $A = I \oplus J$ with $I = \text{lin}\{e_1, e_3\}$ and $J = \mathbb{K}e_2$.

(iii) Let assume that $e_1^2 = 0, e_2^2 \neq 0$ and $e_3^2 \neq 0$. We shall split the proof of this assertion in two cases (a) and (b).

Case (iii)(a) e_2^2 and e_3^2 are linearly dependent. Hence, there exists $t \in \mathbb{K} \setminus \{0\}$ such that $te_2^2 = e_3^2 \neq 0$. We claim that whenever A is reducible then one of the ideals is $\mathbb{K}e_1$. To prove the claim suppose

that $A = I \oplus J$. Since it cannot be $\pi_2(I) = \pi_2(J) = 0$ (as $\pi_2(A) \neq 0$), it is not restrictive to assume that $\pi_2(I) \neq 0$. But $\pi_2(I) \neq 0$ implies that $\pi_2(J) = 0$, as otherwise $e_2^2 \in I \cap J$. Also, whenever $\pi_2(I) \neq 0$ then $\pi_3(I) \neq 0$. Indeed, if $\pi_3(I) = 0$, then $\pi_3(J) \neq 0$ (otherwise $\pi_3(A) = 0$). Thus, $e_2^2 = te_3^2 \in I \cap J = \{0\}$, a contradiction. Therefore $\pi_2(I) \neq 0$ and $\pi_3(I) \neq 0$ while $\pi_2(J) = \pi_3(J) = 0$ (as $I \cap J = \{0\}$ and $e_2^2 = te_3^2 \neq 0$). Thus, we obtain $J = \mathbb{K}e_1$, as desired to prove the claim.

We shall consider two different situations:

Case (iii)(a.1) $e_2^2 = te_3^2 \in \mathbb{K}e_1$, with $t \neq 0$. Then, from the former claim it follows that A is not reducible. Indeed, let us assume that $A = I \oplus J$. Then, $J = \mathbb{K}e_1$ and $\pi_2(I) \neq 0$ as shown above. Hence, $e_2^2 = te_3^2 \in I \cap J$, and thus, $e_1 \in I \cap J \neq \{0\}$, a contradiction. Thus, whenever the structure matrix of A is of the type

$$M_B(A) = \begin{pmatrix} 0 & * & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

the evolution algebra is irreducible.

Case (iii)(a.2) $e_3^2 = te_2^2 \notin \mathbb{K}e_1$, with $t \neq 0$. Then $e_2^2 = \alpha e_1 + \beta e_2 + \gamma e_3$ with $|\beta| + |\gamma| \neq 0$. We consider the following possibilities:

Case (iii)(a.2.1) $\alpha = 0$. Then $A = I \oplus J$ with $I = \text{lin}\{e_2, e_3\}$ and $J = \mathbb{K}e_1$.

Case (iii)(a.2.2) $\alpha \neq 0$. Then, as $|\beta| + |\gamma| \neq 0$, we have the following possibilities:

Case (iii)(a.2.2.1) $\beta \neq 0$. Then $A = I \oplus J$ with $I = \text{lin}\{e_2^2, e_3\} = \text{lin}\{\alpha e_1 + \beta e_2, e_3\}$ and $J = \mathbb{K}e_1$.

Case (iii)(a.2.2.2) $\beta = 0$. Then, $\gamma \neq 0$ and so $A = I \oplus J$ where $J = \mathbb{K}e_1$ and $I = \text{lin}\{e_2, e_2^2\} = \text{lin}\{\alpha e_1 + \gamma e_3, e_2\}$.

Case (iii)(b) e_2^2 and e_3^2 are linearly independent. We claim that A is reducible if and only if $e_1 \notin \text{lin}\{e_2^2, e_3^2\}$. To prove the claim, suppose that $e_1 \in \text{lin}\{e_2^2, e_3^2\}$. Then, A is reducible since $A = I \oplus J$ with $I = \text{lin}\{e_2^2, e_3^2\}$ and $J = \mathbb{K}e_1$. In this case, the structure matrix is

$$M_B(A) = \begin{pmatrix} 0 & w_{12} & w_{13} \\ 0 & w_{22} & w_{23} \\ 0 & w_{32} & w_{33} \end{pmatrix}$$

with $w_{22}w_{33} - w_{32}w_{23} \neq 0$. To finish the proof of the claim, suppose now that $e_1 \in \text{lin}\{e_2^2, e_3^2\}$ and let us show that A is irreducible. Assume towards contradiction that $A = I \oplus J$ is a non-trivial decomposition and suppose that $\pi_2(I) \neq 0$ while $\pi_2(J) = 0$, which is not restrictive. Let $e_2^2 = \sum_{i=1}^3 w_{i2}e_i$

and $e_3^2 = \sum_{i=1}^3 w_{i3}e_i$. From the fact that $e_1 \in \text{lin}\{e_2^2, e_3^2\}$ it follows that

$$\left| \begin{pmatrix} w_{22} & w_{23} \\ w_{32} & w_{33} \end{pmatrix} \right| = 0.$$

Therefore, there exists $t \in \mathbb{K}$ such that

$$\begin{pmatrix} w_{22} & w_{23} \\ w_{32} & w_{33} \end{pmatrix} = \begin{pmatrix} w_{22} & tw_{22} \\ w_{32} & tw_{32} \end{pmatrix}.$$

We are going to distinguish between two cases:

Case (iii)(b.1) $w_{32} \neq 0$. Then, since $e_2^2 \in I$ (because $\pi_2(I) \neq 0$) we deduce that $e_3^2 \in I$, and hence, $e_1 \in I$ as $e_1 \in \text{lin}\{e_2^2, e_3^2\}$. Also, we have that $\pi_3(J) = 0$ (otherwise $e_3^2 \in I \cap J$) and similarly $\pi_2(J) = 0$. Thus, $J = \mathbb{K}e_1$ and $e_1 \in I \cap J$, a contradiction.

Case (iii)(b.2) $w_{32} = 0$. Then:

Case (iii)(b.2.1) If $\pi_3(I) \neq 0$ then, since $\pi_2(I) \neq 0$, we have that $I = \text{lin}\{e_2^2, e_3^2\}$, and hence, $e_1 \in I$, and, as in the case (b.1), we get $e_1 \in I \cap J$, a contradiction.

Case (iii)(b.2.2) Suppose $\pi_3(I) = 0$, and recall that $\pi_2(I) \neq 0$. Consequently, $\pi_3(J) \neq 0$ and $\pi_2(J) = 0$. It follows that $w_{23} = w_{32} = 0$ as $e_2^2 = \sum_{i=1}^3 w_{i2}e_i \in I$ with $\pi_3(I) = 0$ and $e_3^2 = \sum_{i=1}^3 w_{i3}e_i \in J$ with $\pi_2(J) = 0$. Therefore

$$\begin{pmatrix} w_{22} & w_{23} \\ w_{32} & w_{33} \end{pmatrix} = \begin{pmatrix} w_{22} & tw_{22} \\ w_{32} & tw_{32} \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

which contradicts the fact that e_2^2 and e_3^2 are linearly independent.

Hence, we conclude that if e_2^2 and e_3^2 are linearly independent, then A is irreducible if and only if $e_1 \in \text{lin}\{e_2^2, e_3^2\}$; i.e., if and only if $\begin{pmatrix} w_{22} \\ w_{23} \end{pmatrix}$ and $\begin{pmatrix} w_{32} \\ w_{33} \end{pmatrix}$ are proportional but $\begin{pmatrix} w_{21} \\ w_{22} \\ w_{23} \end{pmatrix}$ and $\begin{pmatrix} w_{31} \\ w_{32} \\ w_{33} \end{pmatrix}$ are not as e_2^2 and e_3^2 are linearly independent. This is equivalent to

$$M_B(A) = \begin{pmatrix} 0 & w & \tilde{w} \\ 0 & \alpha & \alpha t \\ 0 & \beta & \beta t \end{pmatrix}$$

with $\tilde{w} \neq wt$, and $|\alpha| + |\beta| \neq 0$. The rest is clear. \square

Theorem 10. Let A be a three-dimensional degenerate evolution algebra. Then we have the following:

(i) A is reducible if and only if there exists a natural basis B of A whose structure matrix is within the following types:

(a) $M_B(A) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$

(b) $M_B(A) = \begin{pmatrix} 0 & 0 & \alpha \\ 0 & 0 & \beta \\ 0 & 0 & \gamma \end{pmatrix}$ with $|\alpha| + |\beta| + |\gamma| \neq 0, \alpha, \beta, \gamma \in \mathbb{K}.$

(c) $M_B(A) = \begin{pmatrix} 0 & \alpha & t\alpha \\ 0 & \beta & t\beta \\ 0 & \gamma & t\gamma \end{pmatrix}$ with $|\beta| + |\gamma| \neq 0$ and $t \neq 0, \alpha, \beta, \gamma \in \mathbb{K}.$

(d) $M_B(A) = \begin{pmatrix} 0 & w & \tilde{w} \\ 0 & \alpha & \beta \\ 0 & \gamma & \delta \end{pmatrix}$ with $w, \tilde{w}, \alpha, \beta, \gamma, \delta \in \mathbb{K}$ and $\alpha\delta - \gamma\beta \neq 0.$

(ii) A is reducible if and only if there exists a natural basis B of A whose structure matrix is within the following types:

(e) $M_B(A) = \begin{pmatrix} 0 & * & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$

(f) $M_B(A) = \begin{pmatrix} 0 & w & \tilde{w} \\ 0 & \alpha & \alpha t \\ 0 & \beta & \beta t \end{pmatrix}$ with $\tilde{w} \neq wt$ and $|\alpha| + |\beta| \neq 0.$

The former type of evolution algebras are non-isomorphic to each other.

Proof. We obtained this classification because of Theorem 9. Let us check that they are non-isomorphic. First of all, none of the algebras in the group (i) can be isomorphic with any of the algebras in the group (ii), as those in the former group are reducible while those in the latter group are not. In the group (i), the algebra (a) is clearly non-isomorphic to any of the others, as its product is zero. On the other hand, (b) cannot be isomorphic to (c) or (d) because its annihilator has two-dimensional, while (c) and (d) and have a one-dimensional annihilator. To verify that (c) is not isomorphic with (d), we just point out that in (c), $\dim A^2 = 1$, while in (d), $\dim A^2 = 2$. In the group (ii), finally, (e) cannot be isomorphic to (f), as in (e) we have $\dim A^2 = 1$; meanwhile, in (f) we have $\dim A^2 = 2$. \square

4.3. The Main Result

To sum up, we gather Theorems 5 and 6, Corollary 6, Theorems 8 and 10 in the theorem below, showing that when we classify three-dimensional evolution algebras according to their degeneracy and their reducibility we obtain 14 non-isomorphic types of evolution algebras.

Theorem 11. *Let A be an evolution algebra with $\dim A = 3$ and let us consider $t, a, b, c \in \mathbb{K} \setminus \{0\}$ and $\alpha, \beta, \gamma, \delta, w, \tilde{w} \in \mathbb{K}$. Then:*

(i) *Suppose that A is degenerate and reducible. Then, there exists a natural basis B such that the structure matrix of A relative to B is like M_1, M_2, M_3 or M_4 , where:*

- $M_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix};$
- $M_2 = \begin{pmatrix} 0 & 0 & \alpha \\ 0 & 0 & \beta \\ 0 & 0 & \gamma \end{pmatrix}$ with $|\alpha| + |\beta| + |\gamma| \neq 0;$
- $M_3 = \begin{pmatrix} 0 & \alpha & t\alpha \\ 0 & \beta & t\beta \\ 0 & \gamma & t\gamma \end{pmatrix}$ with $|\beta| + |\gamma| \neq 0;$
- $M_4 = \begin{pmatrix} 0 & w & \tilde{w} \\ 0 & \alpha & \beta \\ 0 & \gamma & \delta \end{pmatrix}$ with $\alpha\delta - \gamma\beta \neq 0.$

(ii) *Suppose that A is degenerate and irreducible. Then, there exists a natural basis B of A whose structure matrix is like M_5 or M_6 , where:*

- $M_5 = \begin{pmatrix} 0 & * & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix};$
- $M_6 = \begin{pmatrix} 0 & w & \tilde{w} \\ 0 & \alpha & \alpha t \\ 0 & \beta & \beta t \end{pmatrix}$ with $\tilde{w} \neq wt$ and $|\alpha| + |\beta| \neq 0.$

(iii) *Suppose that A is non-degenerate and reducible. Then, there exists a natural basis B of A which structure matrix is like M_7, M_8 or M_9 , where:*

- $M_7 = \begin{pmatrix} * & 0 & 0 \\ 0 & * & 0 \\ 0 & 0 & * \end{pmatrix};$
- $M_8 = \begin{pmatrix} * & 0 & 0 \\ 0 & 0 & * \\ 0 & * & 0 \end{pmatrix};$

- $M_9 = \begin{pmatrix} * & 0 & 0 \\ 0 & w & \tilde{w} \\ 0 & * & * \end{pmatrix}.$

(iv) Suppose that A is non-degenerate and irreducible. Then, there exists a natural basis B of A such that the structure matrix associated to it is like $M_{10}, M_{11}, M_{12}, M_{13}$ or M_{14} , where:

- $M_{10} = \begin{pmatrix} * & * & 0 \\ 0 & w & * \\ 0 & * & \tilde{w} \end{pmatrix}$ with $|M_{10}| \neq 0$;

- $M_{11} = \begin{pmatrix} * & * & * \\ 0 & w & * \\ 0 & * & \tilde{w} \end{pmatrix}$ with $|M_{11}| \neq 0$;

- $M_{12} = \begin{pmatrix} * & * & w \\ 0 & * & * \\ 0 & * & * \end{pmatrix}$ with $|M_{12}| = 0$ and either $M_{12} = \begin{pmatrix} * & a & -\frac{b^2}{c^2}a \\ 0 & b & -\frac{b^2}{c^2}b \\ 0 & c & -\frac{b^2}{c^2}c \end{pmatrix}$ or M_{12} has no proportional columns;

- $M_{13} = \begin{pmatrix} & \alpha & \gamma \\ \beta & & * \\ 0 & 0 & \end{pmatrix}$ with $|\alpha| + |\beta| + |\gamma| \neq 0$, and no zero columns;

- $M_{14} = \begin{pmatrix} & * & \\ \alpha & & * \\ \beta & \gamma & \end{pmatrix}$ with $|\beta| + |\alpha\gamma| \neq 0$, range of M_{14} greater than 1, and such that it does

not exist a nonsingular matrix P such that $PX = M_{14}P^{[2]}$, where $X = \begin{pmatrix} \alpha_1 & \beta_1 & \gamma_1 \\ \alpha_2 & \beta_2 & \gamma_2 \\ 0 & 0 & 0 \end{pmatrix}.$

Moreover, if A has a basic ideal of dimension 1 and has no basic ideals with dimension 2, then $M_B(A)$ is given by either M_{10}, M_{11} or M_{12} . If A has basic ideals of dimension 2 then $M_B(A)$ is like M_{13} and if A has no proper basic ideals then $M_B(A)$ is like M_{14} .

In fact, for $1 \leq i \leq 14$, denote by A_i an evolutionary algebra having a natural basis of the type M_i described above. Then, these algebras are not isomorphic and we obtain the following classification of evolution algebras with three dimensions:

	Degenerate	Non-Degenerate
Reducible	A_1, A_2, A_3, A_4	A_7, A_8, A_9
Irreducible	A_5, A_6	$A_{10}, A_{11}, A_{12}, A_{13}, A_{14}$

Therefore, we have obtained 14 non-isomorphic types of evolution algebras of dimension 3.

This means that an algebra of the type A_i is not isomorphic to an algebra of the type A_j whenever $i \neq j, 1 \leq i, j \leq 14$. Nevertheless, we found several non-isomorphic evolution algebras that belong to the same type. As a matter of fact, by considering the 116 types of non-isomorphic three-dimensional evolution algebras described in [17], we have reclassified them into the 14 different types A_i described above. It is easy to check when one of the algebras stated in [17] belongs to one of the types obtained in this paper by just considering the properties:

- (a) Being reducible or not;
- (b) Being degenerate or not,
- (c) Having a basic ideals of dimension 1 and no basic ideals of dimension 2;
- (d) Having a basic ideal of dimension 2;
- (e) Having no proper basic ideals.

Author Contributions: All authors contributed equally to this manuscript.

Funding: This work was partially supported by the Spanish Project MTM2016-76327-C3-2-P (AEI\FEDER, UE).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Tian, J.P. *Evolution Algebras and Their Applications*; Lecture Notes in Math. 1921; Springer: Berlin, Germany, 2008.
2. Rozikov, U.A.; Tian, J.P. Evolution algebras generated by Gibbs measures. *Lobachevskii J. Math.* **2011**, *32*, 270–277. [[CrossRef](#)]
3. Cabrera Casado, Y.; Siles Molina, M.; Velasco, M.V. Evolution algebras of arbitrary dimension and their decompositions. *Linear Algebra Appl.* **2016**, *495*, 122–162. [[CrossRef](#)]
4. Camacho, L.M.; Gómez, J.R.; Omirov, B.A.; Turdibaev, R.M. Some properties of evolution algebras. *Bull. Korean Math. Soc.* **2013**, *50*, 1481–1494. [[CrossRef](#)]
5. Casas, J.M.; Ladra, M.; Rozikov, U.A. On nilpotent index and dibaricity of evolution algebras. *Linear Algebra Appl.* **2013**, *43*, 90–105. [[CrossRef](#)]
6. Casas, J.M.; Ladra, M.; Rozikov, U.A. On evolution algebras. *Algebra Colloq.* **2014**, *21*, 331–342. [[CrossRef](#)]
7. Dzhumadil'daev, A.; Omirov, B.A.; Rozikov, U.A. On a class of evolution algebras of “chicken” population. *Internat. J. Math.* **2014**, *25*, 1450073. [[CrossRef](#)]
8. Labra, A.; Ladra, M.; Rozikov, U.A. An evolution algebra in population genetics. *Linear Algebra Appl.* **2014**, *45*, 348–362. [[CrossRef](#)]
9. Rozikov, U.A.; Murodov, S.N. Dynamics of two-dimensional evolution algebras. *Lobachevskii J. Math.* **2013**, *3*, 344–358. [[CrossRef](#)]
10. Tian, J.P.; Vojtechovsky, P. Mathematical concepts of evolution algebras in non-Mendelian genetics. *Quasigroups Relat. Syst.* **2006**, *14*, 111–122.
11. Casas, J.M.; Ladra, M.; Rozikov, U.A. A chain of evolution algebras. *Linear Algebra Appl.* **2011**, *435*, 852–870. [[CrossRef](#)]
12. Murodov, S.N. Classification dynamics of two-dimensional chains of evolution algebras. *Int. J. Math.* **2014**, *24*, 140012. [[CrossRef](#)]
13. Omirov, B.A.; Rozikov, U.A.; Tulenbayev, K.M. On real chains of evolution algebras. *Linear Multilinear A* **2015**, *63*, 586–600. [[CrossRef](#)]
14. Rozikov, U.A.; Murodov, S.N. Chain of evolution algebras of “chicken” population. *Linear Algebra Appl.* **2014**, *450*, 186–201. [[CrossRef](#)]
15. Camacho, L.M.; Gómez, J.R.; Omirov, B.A.; Turdibaev, R.M. The derivations of some evolution algebras. *Linear Multilinear A* **2013**, *6*, 309–322. [[CrossRef](#)]
16. Ladra, M.; Rozikov, U.A. Evolution algebra of a bisexual population. *J. Algebra* **2013**, *378*, 153–172. [[CrossRef](#)]
17. Cabrera, Y.; Siles Molina, M.; Velasco, M.V. Classification of three-dimensional evolution algebras. *Linear Algebra Appl.* **2017**, *524*, 68–108. [[CrossRef](#)]
18. Ahmed, H.; Bekbaev, U.; Rakhimov, I. Classification of 2-dimensional evolution algebras, their groups of automorphisms and derivation algebras. *arXiv* **2017**, arXiv:1702.08616.
19. Cabrera Casado, Y.; Kanuni, M.; Siles Molina, M. Classification of four dimensional perfect non-simple evolution algebras. *arXiv* **2018**, arXiv:1801.03856.
20. Camacho, L.M.; Khudoyberdiyev, A.K.; Omirov, B.A. On the property of subalgebras of evolution algebras. *Algebras Represent. Theor.* **2019**, *22*, 281–296. [[CrossRef](#)]
21. Falcón Ganfornina, O.; Falcón, R.; Valdés, J. Classifications of evolution algebras over finite fields. In Proceedings of the EACA's III International School on Computer Algebra and Its Applications, Sevilla, Spain, 18–21 January 2016.
22. Hegazi, A.S.; Abdelwahab, H. Five-dimensional nilpotent evolution algebras. *arXiv* **2015**, arXiv:1508.07442.
23. Imomkulov, A. Classification of a family of three dimensional real evolution algebras. *arXiv* **2019**, arXiv:1812.03669.

24. Murodov, S.N. Classification of two-dimensional real evolution algebras and dynamics of some two-dimensional chains of evolution algebras. *arXiv* **2013**, arXiv:1305.6416.
25. Cabrera Casado, Y. Evolution Algebras. Ph.D. Thesis, Universidad de Málaga, Málaga, Spain, 2017.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).