Environmental Research Letters

TOPICAL REVIEW • OPEN ACCESS

Human-carnivore relations: conflicts, tolerance and coexistence in the American West

To cite this article: Mónica Expósito-Granados et al 2019 Environ. Res. Lett. 14 123005

View the article online for updates and enhancements.

Environmental Research Letters

TOPICAL REVIEW CrossMark

OPEN ACCESS

RECEIVED 12 July 2019

REVISED 14 October 2019

ACCEPTED FOR PUBLICATION 5 November 2019

PUBLISHED 6 December 2019

Original content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Human-carnivore relations: conflicts, tolerance and coexistence in the American West

Mónica Expósito-Granados^{1,14}, Antonio J Castro^{2,3}, Jorge Lozano⁴, Jose A Aznar-Sanchez¹, Neil H Carter⁵, Juan M Requena-Mullor⁶, Aurelio F Malo^{7,8}, Agnieszka Olszańska⁹, Zebensui Morales-Reyes¹⁰, Marcos Moleón¹¹, José A Sánchez-Zapata¹⁰, Ainara Cortés-Avizanda¹², Joern Fischer¹³ and Berta Martín-López^{13,14}

- Department of Economics and Business, Research Centre CIAIMBITAL and CAESCG, University of Almería, La Cañada de San Urbano, E-04120, Almería, Spain
- Department of Biology and Geology, Social-Ecological Research Laboratory, Research Centre CAESCG, University of Almería, La Cañada de San Urbano, E-04120, Almería, Spain
- Department of Biological Sciences, Idaho State University, 921 South 8th Avenue, Pocatello, ID, 83209, United States of America
- Department of Biodiversity, Ecology and Evolution, Complutense University of Madrid, C/José Antonio Novais 12, E-28040, Madrid, Spain
- School for Environment and Sustainability, University of Michigan 440 Church Street Ann Arbor, MI 48109, United States of America Department of Biological Sciences, Boise State University, 1910 University Drive, ID-83725, Boise, United States of America
- Department of Life Sciences, GloCEE Global Change Ecology and Evolution Research Group, University of Alcalá, E-28805, Madrid, Spain
- Department of Life Sciences, Imperial College London, Ascot, United Kingdom
- Institute of Nature Conservation Polish Academy of Sciences, al. Adama Mickiewicza 33, 31-120 Krakow, Poland
- 10 Department of Applied Biology, Miguel Hernández University of Elche, Avda. de la Universidad, s/n, E-03202, Elche, Alicante, Spain
- ¹¹ Department of Zoology, University of Granada, Avda. de Fuente Nueva s/n, E-18071, Granada, Spain
- ¹² Animal Demography and Ecology Unit (GEDA), IMEDEA (CSIC)-UIB, C/. Miquel Marqués 21, E-07190, Esporles, Illes Balears, Spain 13
 - Leuphana University of Lüneburg, Faculty of Sustainability, Universitätsallee 1, D-21335, Lüneburg, Germany
- 14 Authors to whom any correspondence should be addressed.

E-mail: moexposit@ual.es and martinlo@leuphana.de

Keywords: ecosystem services, human-wildlife interactions, multi-use landscapes, shared landscapes, socio-ecological systems Supplementary material for this article is available online

Abstract

Carnivore and humans live in proximity due to carnivore recovery efforts and ongoing human encroachment into carnivore habitats globally. The American West is a region that uniquely exemplifies these human-carnivore dynamics, however, it is unclear how the research community here integrates social and ecological factors to examine human-carnivore relations. Therefore, strategies promoting human-carnivore coexistence are urgently needed. We conducted a systematic review on human-carnivore relations in the American West covering studies between 2000 and 2018. We first characterized human-carnivore relations across states of the American West. Second, we analyzed similarities and dissimilarities across states in terms of coexistence, tolerance, number of ecosystem services and conflicts mentioned in literature. Third, we used Bayesian modeling to quantify the effect of social and ecological factors influencing the scientific interest on coexistence, tolerance, ecosystem services and conflicts. Results revealed some underlying biases in humancarnivore relations research. Colorado and Montana were the states where the highest proportion of studies were conducted with bears and wolves the most studied species. Non-lethal management was the most common strategy to mitigate conflicts. Overall, conflicts with carnivores were much more frequently mentioned than benefits. We found similarities among Arizona, California, Utah, and New Mexico according to how coexistence, tolerance, services and conflicts are addressed in literature. We identified percentage of federal/private land, carnivore family, social actors, and management actions, as factors explaining how coexistence, tolerance, conflicts and services are addressed in literature. We provide a roadmap to foster tolerance towards carnivores and successful coexistence strategies in the American West based on four main domains, (1) the dual role of carnivores as providers of both

beneficial and detrimental contributions to people, (2) social-ecological factors underpinning the provision of beneficial and detrimental contributions, (3) the inclusion of diverse actors, and (4) cross-state collaborative management.

Introduction

Carnivore and human populations are often in proximity to each other, due to carnivore recovery efforts in some areas (e.g. North America, Europe) and ongoing human encroachment into carnivore habitats globally. Increasing proximity prompts more frequent interactions between carnivores and humans. These interactions are often viewed through the lens of 'conflict' or 'risk' to human communities, such as livestock depredation, impacts on abundances of game species, and threats to human safety (e.g. Treves and Karanth 2003, Treves et al 2004, Inskip and Zimmermann 2009, Dickman 2010, Miller 2015, van Eeden et al 2018, Lozano et al 2019). However, carnivores can also benefit humans by the provision of ecosystem services such as the mitigation of diseases (Harris and Dunn 2010), carcass removal (Moleón et al 2014, O'Bryan et al 2018) and opportunities provided for ecotourism (Willemen et al 2015, Arbieu et al 2017). Carnivores and humans are therefore considered parts of integrated social-ecological systems, whereby the mutual wellbeing is inextricably linked (Carter et al 2014, Darimont et al 2018, Dressel et al 2018, Lischka et al 2018, Lozano et al 2019).

The American West is an evocative and unique region that exemplifies dynamic human-carnivore relations (e.g. Kellert et al 1996, Young et al 2015, Bruskotter et al 2017, Slagle et al 2017, Jones et al 2019). In this region there are areas with intact carnivore guilds and wilderness; yet rapid human development and polarizing debate about carnivore management threaten the future of these animals and can hinder effective policy-making (Bangs and Shivik 2001, Linnell et al 2001, Bruskotter 2013, Bradley et al 2015, Smith et al 2016). In particular, interaction between livestock and carnivores generates intense controversy in the American West (van Eeden et al 2018), where publically-owned grazing land is ubiquitous and livestock production is an important economic sector (Sarchet 2005). Legal hunting has also generated intense debate among actors about the ways to solve human-carnivore conflicts and conserve carnivores (Treves and Naughton-Treves 2005, Treves 2009). Strategies that promote coexistence between humans and carnivores in multi-use landscapes are therefore urgently needed in order to balance the goals of nature preservation and livelihood protection in the American West.

Using the definition by Carter and Linnell (2016), we characterize coexistence as the 'dynamic but sustainable state in which humans and carnivores coadapt to living in shared landscapes where human relations with carnivores are governed by effective institutions that ensure long-term carnivore population persistence, social legitimacy, and tolerable levels of risk and damage.' Then, we use coexistence as an umbrella concept that encompasses 'tolerance', which could be defined (based on Bruskotter and Wilson 2014) as the 'human acceptance' of the risks and damages caused by carnivores, a necessary condition to achieve a permanent coexistence (Carter and Linnell 2016).

Given these definitions, many ecological and social factors, heterogeneous in both space and time, can facilitate or limit human-carnivore coexistence (Frank et al 2019). Among the ecological factors, the species involved (Kansky et al 2014) and the ecosystems that they inhabit, are likely relevant in determining coexistence. Also, several social factors might foster or constrain coexistence, such as the type of actors involved, gender, or education level (e.g. Morzillo et al 2007, 2010, Agarwala et al 2010, Smith et al 2014, Lute et al 2016), as well as the attributes of the governance systems (Borrini-Feyerabend and Hill 2015). However, it is unclear the extent to which the research community integrates social and ecological factors to examine human-carnivore coexistence in the American West. Furthermore, studies tend to emphasize conflicts with carnivores and rarely assess the variety of services they provide to people (Lozano et al 2019). Yet, acknowledging and understanding the multiple ecosystem services (i.e. benefits) and disservices (i.e. risks and damages) carnivores provide to people allow for a more comprehensive, and defensible, evaluation of the trade-offs of coexisting with these species (e.g. Ripple et al 2014, Braczkowski et al 2018, Morales-Reves et al 2018). Finally, recent research suggests the need of reconnecting people with nature (Folke et al 2011, Ives et al 2017). The concept of human-nature connectedness integrates different relationships between social and natural systems (Ives et al 2017, 2018). In this context, experiential (e.g. recreational activities in nature), emotional (e.g. affective response to nature), and cognitive (e.g. knowledge, beliefs and attitudes) connections are important for human well-being and play a useful role in fostering conservation and tolerance of carnivores. Therefore, identifying and clarifying knowledge gaps in coexistence research in the American West can shed light on where the field has been, where it stands, and where it might go in the future.

Here, we conducted a systematic review of the literature on human-carnivore relations in the American West published between 2000 and 2018. To do so, we first characterized human-carnivore relations across states of the American West according to ecological (e.g. species, biomes) and social factors (e.g. actors and management). Second, we analyzed the similarities and dissimilarities across states of the American West in terms of coexistence and tolerance (mentioned or evaluated in the reviewed literature), the number of ecosystem services and conflicts mentioned. Third, we quantified the effect of social and ecological factors that influence the mention of coexistence, tolerance, ecosystem services and conflicts in the scientific literature across states of the American West. Through this analysis, we reveal the underlying biases in humancarnivore relations research in the American West and outline a road map for advancing the theory and practice of human-carnivore coexistence in this important region.

Methods


We searched for articles indexed by the Scopus database following guidelines of Pullin and Stewart (2006). We based the search on the systematic review conducted by Lozano et al (2019) and their final 2000–2016 database¹. We used the same search query to include articles published until 2018 and only carried out in the American West. The search string included four main elements: (1) human-carnivore relations, (2) ecosystem services, (3) conflicts and (4) the taxonomic groups of terrestrial carnivores (see appendix A for the full search string). We used different terms referring to conflicts (i.e. 'conflict"' OR 'damage*' OR 'impair*' OR 'harm*'), ecosystem services (i.e. 'ecosystem service"' OR 'ecosystem good"' OR 'environmental service*') and human-carnivore relations (i.e. 'human-wildlife' OR 'human-carnivore*' OR 'human-felid*' OR 'human-canid*'), since these can represent negative, positive or neutral relations with carnivores, respectively. The search was applied to the fields title, abstract and keywords (see Lozano et al 2019), and the final number of selected articles for in-depth review was 71 (see appendix B for detailed methods of the review process).

For all the articles reviewed (see appendix C), we registered the ecological and social factors considered. Regarding ecological variables, we included: (1) biome type (based on MA (Millennium Ecosystem Assessment) 2005), (2) carnivore family and (3) carnivore species. Regarding social factors, we considered (1) type of social actor (based on Lozano *et al* 2019), and (2) type of management action (according to Inskip and Zimmerman (2009) and Lozano *et al* 2019). We also coded different variables representing human-carnivore relations: (1) whether human-carnivore 'coexistence' was mentioned (based on Carter and Linnell 2016), (2) whether 'tolerance' or 'acceptance' of carnivores were mentioned or evaluated in the article (based on Gore *et al* 2006, Bruskotter and

Wilson 2014 and Kansky *et al* 2014), (3) type of ecosystem services mentioned (based on MA (Millennium Ecosystem Assessment) 2005) and (4) type of humancarnivore conflict considered (based on Peterson *et al* 2010 and Lozano *et al* 2019). For a detailed description of the variables included, see appendix D.

Firstly, we performed a descriptive analysis to present the state of knowledge regarding human-carnivore relations research in the American West. Secondly, similarities regarding human-carnivore relations research across states of the American West were analyzed by nonparametric multidimensional scaling (NMDS) using the 'vegan' R package (Oksanen et al 2019). 17 out of 71 articles were conducted simultaneously in several states of the American West. Then, to obtain the different variables across states, the information was disaggregated and assigned to each of the states separately, so information provided by one article can be assigned to different states (N = 120). The states of the American West were arranged on a Cartesian axis based on the terms of coexistence, tolerance toward carnivores (mentioned or evaluated), number of ecosystem services and number of conflicts reported by articles in each state. A shorter distance between states would mean greater similarity in the way scientists approach research on human-carnivore relations. We used the Mahalanobis distance, which takes into account the potential correlation between the variables used in the ordination. In addition, we fitted the number of types of human-nature connection (according to Ives et al 2017) mentioned in articles using penalized splines (Oksanen et al 2019). Two types of human-nature connection were fitted: experiential and emotional; we excluded cognitive connections as this category was not mentioned in the articles considered. We used Kruskal's stress (Kruskal 1964) to check for the goodness of fit of the NMDS. Kruskal's stress measures the agreement in the rank order of the inter-state distances observed and those predicted from the similarities. According to Clarke's (1993) guidelines for stress values, values lower than 0.3 indicate that the arrangement reached is better than one obtained randomly.

Finally, we used a Bayesian modeling approach to quantify the effect of ecological and social variables influencing the number of (1) articles mentioning human-carnivore coexistence, (2) articles mentioning or evaluating tolerance toward carnivores, (3) ecosystem services mentioned, and (4) conflicts reported. All variables were aggregated by states of the American West (see above). We estimated the effect of the following predictor variables: percentage of federal and private land (average of percentage of 2000, 2010 and 2015; obtained from Vincent *et al* 2017), number of articles according to carnivore family, number of social actors mentioned (i.e. three types included: local, non-local, and manager/academia), number of management actions mentioned (i.e. three types



included: non-lethal actions, community development programs, and lethal control interventions), number of ecosystem services mentioned (i.e. two types included: regulating and cultural; we exclude provisioning services because these were only mentioned twice in one article), number of conflicts reported (i.e. four types included: damage to human food, damage to human property, damage to human safety and human-human conflicts; we excluded damage to biodiversity as it was only mentioned in one article). The number of articles across states was also included in the models as a covariate. All the variables were standardized to have a mean of zero and standard deviation of 1. We built spatial models assuming a Besag-York-Mollie specification (Besag et al 1991) using the R-integrated nested Laplace approximation (INLA) package (Rue et al 2009) (see appendix E for details and models parameterization). INLA is a computationally efficient method for fitting Bayesian models while accounting for spatial dependence of residuals (i.e. model residuals at nearby states are not independent) (Blangiardo and Cameletti 2015). Altogether, we built separate models for each of the following response variables: the number of mentions of human-carnivore coexistence, tolerance towards carnivores, ecosystem services, and conflicts (i.e. 56 models in total). Each of the models included one intercept, one predictor variable listed above (a different one each model), the number of articles conducted in each state as a covariate and one spatially structured term. All the analyses were performed using R software version 3.6.0 (R Core Team 2019).

Results

State-of-the-science in human-carnivore relations

The number of published articles since 2000 concerning human-carnivore relations in the American West has rapidly increased, with a peak in 2018 with 14 articles (figure 1(a)). The largest proportion of research was conducted in Colorado (29.6% of articles), followed by Montana (26.8% of articles), Wyoming (22.5% of articles), and Washington (18.3% of articles), whereas Arizona and New Mexico received relatively less attention (figure 1(b)). Most articles were carried out in mountain areas (59.2%) and temperate forests (50.7%), whereas Mediterranean ecosystems (7.0%) and polar environments (2.8%) were scarcely represented in articles (figure 1(c)).

Families most studied were Ursidae (63.4% of articles) followed by Canidae (wolves and coyotes; 31%) and Felidae (cougars and bobcats; 14.1%). Most articles only included one species (88.7% of articles). The American black bear (Ursus americanus; 39.4% of articles) was the most frequently studied followed by brown bear (U. arctos; 28.2%), grey wolf (Canis lupus; 21.1%), cougar (Puma concolor; 14.1%) and coyotes (C. latrans; 11.3%). Only 2.8% of articles focused on other carnivores such as polar bears (U. maritimus) and bobcats (Lynx rufus). Seven out of 71 articles (9.9%) dealt with reintroduced carnivores (mainly wolves). Interestingly, there was no article focusing on small or medium-sized carnivores. Among the three most studied carnivores, American black bear was most frequently studied in Colorado (22.6%) and California (13.2%), brown bear in Alaska (30.8%) and




Figure 2. Number of articles that mentioned (a) types of social actors involved, and (b) management actions implemented for dealing with human-carnivore conflicts.

Wyoming (30.8%) and grey wolf in Montana (28.6%) and Idaho (20%) (figure 1(d)).

Coexistence and tolerance toward carnivores in the American West

Coexistence was mentioned in 25.4% of articles, whereas human tolerance toward carnivores was mentioned or evaluated in 43.7% of articles. Coexistence was more frequently mentioned in articles conducted in Colorado (22.6%) and Montana (19.4%), while tolerance was mainly mentioned or evaluated in Montana (22%), Colorado (15%) and Idaho (13%).

Social actors

Sixty-four articles (90.1%) mentioned any type of social actor involved in the study of human-carnivore relations. Local actors were the most frequently mentioned social actor type (70.4% of articles), particularly rural (31.0% of articles), urban residents (28.2%) and hunters (28.2%), while subsistence and commercial farmers were mentioned in 22.5% and 14.1% of articles, respectively. Thirty-tree articles (46.5%) mentioned non-local actors, specifically 36.6% of articles referred to general public and 19.7% of articles to tourists. In addition, actors from academia and managers were mentioned in 48 articles out of 71 (67.6%), environmental managers were mentioned in 66.2 % of articles, while decision makers in governments were considered in 12.7% of articles, NGOs/conservationists in 9.9%, and researchers were mentioned less frequently (4.2%) (figure 2(a)).

Human-carnivore management actions


Management recommendations towards carnivores were mentioned in 90.1% of articles. Non-lethal measures were the most mentioned management action to alleviate human-carnivore conflicts (66.2% of articles); 31% of articles mentioned the use of deterrents and barriers (e.g. specialized electric fencing, lights and loud noises), 26.8% of articles reported zoning (i.e. separating livestock grazing from carnivores' habitat) as a management action. Translocation was mentioned in 19.7% of articles, and aversive conditioning and verification of attacks were both mentioned in 16.9% of the articles, whilst livestock guarding (11.3%) and husbandry techniques (11.3%) were less frequently mentioned (figure 2(b)).

Management actions that target community development were mentioned in 47.9% of the articles. In particular, 39.4% of articles mentioned actions related to education programs while economic incentives (12.7% of articles) and co-management (11.3% of articles) were less mentioned. Finally, 33 articles out of 71 (46.5%) mentioned lethal management actions, specifically individual removal (39.4% of articles) and hunting permit regulations (28.2% of articles) were reported (figure 2(b)). Despite this, only 20 articles out of 71 (28.2%) tested the effectiveness of different management practices in reducing damages occurrence (i.e. garbage disposal, deterrent and aversive techniques, or effectiveness of education programs).

Ecosystem services and human-carnivore conflicts

More than half of articles focused only on conflicts (62.0% of articles), followed by articles mentioning both services and conflicts (36.6%). Only 1.4% of articles exclusively mentioned services of carnivores to society. Regarding the ecosystem services provided by carnivores, 35.2% of articles mentioned cultural services, mainly sport hunting (25.4% of articles), while 7% of the articles referred to regulating services, such as important roles of apex predators (5.6% of articles), and 1.4% of articles mentioned provisioning services (i.e. fur and food) (figure 3(a)). Most articles mentioning cultural services were conducted in Wyoming (15%) and Washington (15%). Regulating services were more frequently mentioned in Alaska (28.6%) than in other states, and this state was the only one that mentioned provisioning services. New Mexico (2.5%) and Utah (2.5%) scarcely mentioned any ecosystem services (in both states regulating services were mentioned once) while Arizona did not mention any ecosystem services (figure 3(b)).

Seventy out of 71 articles mentioned conflicts, 64.8% of articles mentioned damages to human food



resources (mainly predation on livestock and poultry), 50.7% of articles mentioned damage to human safety, 36.6% of articles mentioned damage to human property and human-human conflicts were considered by 16.9% of articles (figure 3(c)). Damage to biodiversity was only mentioned in one article (Ziegltrum and Nolte 2000). Regarding conflicts per state, damage to food were more frequently mentioned in articles carried out in Montana (18.4%), Wyoming (14.9%) and Colorado (14.9%). In addition, damage to human property (e.g. damage to trash containers, or noisy activities) was mainly mentioned in Montana (27.8%) and Colorado (22.2%), damage to human safety was mostly frequently mentioned in Colorado (24.1%), and human-human conflicts were more frequently mentioned in Washington (27.8%) (figure 3(d)).


Measuring similarity across states

The states of the American West were optimally arranged according to coexistence mentioned, tolerance mentioned or evaluated, number of ecosystem services and conflicts mentioned by articles (Kruskal's stress 0.16 < 0.3). The spatial configuration of the states reached by the NMDS analysis (figure 4) suggests that there are different human-carnivore relations in the American West, with some states sharing the composition of these relations, and others having a more dissimilar one. For example, we found that Arizona, California, Utah, and New Mexico were closely arranged (figure 4), while the rest of the American West states showed a dissimilar composition to any other state, being isolated in ordination space. Alaska and Idaho were the farthest states from each other, and therefore, the most dissimilar. The experiential and emotional human-carnivore connections fitted well to the ordination (figures 4(a) and (b)). Although some states were dissimilar in terms of human-carnivore relations and ecosystem services mentioned, they shared similar composition of mentions of experiential human-carnivore connections (e.g. Alaska, Oregon, and Montana; figure 4(a)). In contrast, states located away from each other also had a dissimilar composition of mentions of emotional connections (e.g. Alaska, Colorado and Oregon; figure 4(b)).

Social and ecological factors influencing the research on human-carnivore coexistence, tolerance, ecosystem services and conflicts

Regarding the number of mentions of coexistence in the reviewed research, we found that the Ursidae family yielded the strongest negative effect but exhibited the highest uncertainty as compared to any other variable (figure 5(a)). Federal and private land showed positive and negative effects, respectively, and low uncertainty (i.e. posterior probability that the credible intervals overlapped zero were 0.069).

Regarding the number of articles that mentioned or evaluated tolerance in human-carnivore relations research in the American West, we found that several variables can be considered important based on a 95% credible interval (figure 5(b)). Research focused on Canidae, mentions of local and manager/academia social actors, non-lethal management actions, community development programs, regulating services and human–human conflicts had a positive influence on the likelihood to research tolerance toward carnivores, while research focused on Felidae or Ursidae,



The range of distances reached between states of the American West. States were arranged so that the distances between them were as close to the observed differences between the human-carnivore coexistence mentioned, tolerance mentioned or evaluated, number of ecosystem services and number of conflicts reported by articles. A shorter distance between states means greater similarity between them, whereas a longer distance corresponds to a greater dissimilarity. N = 120 (i.e. total number of articles carried out across states; note that one article can be assigned to different states). For this ordination, the number of experiential (a) and emotional (b) humannature connections was fitted as contour lines using penalized splines (Oksanen *et al* 2019). The number of connections and the NMDS axis 1 and 2 coordinates were used as a response variable and predictors, respectively. R^2 represents the goodness of fit and the *P*-value indicates the significance of the nonlinear term.

mentions of non-local social actors and conflicts related to human safety negatively influence the number of articles that mentioned or evaluated tolerance. The number of articles researching the Canidae family, mentions of local actors, manager/academia and community development programs per state positively influenced the number of ecosystem services mentioned. However, the number of articles focused on Ursidae family showed negative effects on the number of ecosystem services mentioned (figure 5(c)). Percentages of federal and private land were the most important variables for the number of human-carnivore conflicts mentioned, with positive and negative effects, respectively. The posterior probability that the credible intervals overlapped zero was 0.050. Finally, mentions of manager/academia actors as well as lethal management actions positively influenced the number of conflicts mentioned (figure 5(d)).




Figure 5. Estimated effects (\pm 95% credible intervals) of the social-ecological factors (i.e. predictor variables) that influence (a) human-carnivore coexistence mentioned, (b) tolerance mentioned or evaluated, (c) number of ecosystem services mentioned and (d) number of human-carnivore conflicts mentioned in articles per state of the American West. The effects represent the change in the log of the response variable when the predictor increases by one standard deviation.

Discussion

What are the current research trends of humancarnivore relations in the American West?

This research shows a growing interest of the scientific community on the human-carnivore relations in the American West (figure 1(a)). This iconic region encompasses several social (i.e. population growth) and ecological (i.e. aridity, topography) characteristics that reveal the complex relationships between humans and carnivores (Jones *et al* 2019). To date, human-carnivore research has mostly focused on mountains and temperate forests in Colorado and Montana, regions where co-occurrence between humans and large-bodied carnivores (e.g. bears and wolves, figure 1(d)) exist due to extensive farmland activities and recreational use. This finding is consistent with a global assessment of the literature on human-carnivore relations research by Lozano *et al* (2019)

that shows how scientific interest is biased towards large and charismatic carnivore species. This result is also consistent with previous research that shows the influence of charismatic vertebrate species on the conservation research agenda (e.g. Clark and May 2002, Martín-López *et al* 2009).

Our results indicated a possible bias with regard to the diversity of actors and the diversity of management actions mentioned to mitigate conflicts. On one hand, we found that farmers (subsistence and commercial) and property owners are less mentioned than environmental managers (figure 2(a)). Despite many studies that suggest the importance to engage and integrate all social actors involved in conflicts as a way to achieve coexistence and tolerance toward carnivores (Treves *et al* 2006, Marchani *et al* 2019), our results showed that the engagement of actors in human-carnivore research is limited. On the other hand, non-lethal management (i.e. the use of deterrents and zoning livestock, figure 2(b)) is the most reported strategy to mitigate conflicts. However, the effectiveness of different management practices to reduce human-carnivore conflict has been poorly addressed in the studies. The emphasis on non-lethal management actions aligns with previous research that showed the relevance of these measures to mitigate conflicts with carnivores (Eklund *et al* 2017, Moreira-Arce *et al* 2018). Our findings also support previous research that identified educational programs and non-lethal measures as the most reported management actions (figure 2(b)). In addition, several studies showed that educational programs and non-lethal measures are successful strategies for fostering coexistence (Nyhus *et al* 2003, Fernández-Gil *et al* 2016, Lozano *et al* 2019).

Finally, we found that research on human-carnivore relations in the American West is biased towards conflicts, mainly damages to food (figure 3(c)). We also found that when research focuses on ecosystem services, it mainly addresses cultural ecosystem services (figure 3(a)). These findings are consistent with the global assessment of human-carnivore relations research conducted by Lozano *et al* (2019).

What factors are related to mentioning coexistence and tolerance toward carnivores in the research of the American West?

Research on coexistence, tolerance, ecosystem services and conflicts was determined by state-specific social and ecological factors. Bayesian modeling identified percentage of federal/private land, carnivore family, social actors, and management action, as those social and ecological factors explaining how coexistence, tolerance, conflicts and services are addressed in literature (figure 5).

Percentage of federal and private lands exerted a positive and negative effect on the probability of mentioning coexistence in carnivore research, respectively. A large portion of federal land in the US is concentrated in the American West, which has been the subject of debate and controversy about different policies and practices regarding the multiple uses of public lands (e.g. land sparing versus land sharing) (Crespin and Simonetti 2019). On the one hand, a positive effect of federal lands on articles mentioning coexistence is consistent with conservation efforts in states dominated by federal lands, where traditional strategies for wildlife conservation have segregated human activities from remnants of wilderness to avoid further human intervention. On the other hand, the negative effect of private lands suggests the importance of promoting coexistence across states with multi-use landscapes, where space limitation for carnivores demands balancing nature preservation and livelihood protection (Crespin and Simonetti 2019).

Articles mentioning or evaluating tolerance were found to be positively related with the Canidae family (i.e. wolves and coyotes), local actors, managers/ academia, non-lethal management actions, community development and human-human conflicts (figure 5(a)). This result can be explained in two ways. First, a large body of literature documents a general positive attitude toward wolves and wolf recovery by the general public (e.g. Browne-Nuñez et al 2015, Killion et al 2019). This tendency has also been shown towards coyotes (George et al 2016). Second, tolerance towards carnivores depends on the type of actor and the management actions suggested. Former research has emphasized the urgent need of conservation planning models for wildlife that integrate the involvement of multiple social actors in decision-making (Kansky et al 2014). Finally, species such as bears and cougars are frequently related to attacks on humans (Penteriani et al 2016, Smith and Herrero 2018, Bombieri et al 2019). This is consistent with factors here identified and shown to have a negative effect on tolerance (i.e. Ursidae and Felidae families, non-local actors and damage to human safety), and can be interpreted as knowledge gaps to advance in the study of tolerance towards carnivores.

A roadmap for advancing coexistence with carnivores in the American West

Based on this study, we propose that future research on human-carnivore relations in the American West should advance knowledge in four main domains: (1) beneficial contributions of carnivores to people (ecosystem services), (2) social-ecological approaches to determine key factors underpinning beneficial and detrimental contributions, including causes of conflicts, (3) consideration of multiple social actors affected by or involved in the management of carnivores, and (4) cross-state collaborative management.

First, this research shows that conflicts with carnivores are much more frequently mentioned in literature than the variety of ecosystem services they provide to people (figure 3). Indeed, ecosystem services provided by carnivores were reported only in 38% of articles. Neglecting ecosystem services provided by carnivores, both in scientific research and outreach activities, can undermine attempts to foster human tolerance for carnivores, which is a critical component of coexistence (Peterson et al 2010, Pooley et al 2017, Ceausu et al 2018, Lozano et al 2019). Therefore, we call for a shift in mindset that recognizes the dual role of carnivores as providers of both ecosystem services and disservices to humans. To promote this shift, we encourage the scientific community to further explore the ecosystem services provided by carnivores to society. For example, a recent review indicated that predators and scavengers can directly benefit humans by reducing disease prevalence, reducing the abundance of species that can injure people (e.g. vehicledeer collisions), increasing agricultural output, and removing organic waste (O'Bryan et al 2018). This goal might also be reached by revisiting pioneering

M Expósito-Granados et al

research conducted in Wester USA on the role of carnivores and particularly top-predator in ecosystem functioning (i.e. trophic cascades, Beschta and Ripple 2009) that translate in key regulating ecosystem services in rewilding landscapes (Kuijper et al 2016). New analytical frameworks have been developed to evaluate tradeoffs in ecosystem services and disservices from carnivores to different recipients (Ceausu et al 2018), although parameterizing these frameworks with empirical data across sites and species are much needed. This idea of the dual role of biodiversity as provider of beneficial and detrimental contributions to people has been acknowledged by the Intergovernmental Platform on Biodiversity and Ecosystem Services under the paradigm of 'nature's contributions to people' (Díaz et al 2018).

Second, we found that current research interest on beneficial and detrimental contributions provided by carnivores in the American West differs according to social and ecological characteristics. Although our results do not indicate a causative relationship between carnivores' contributions and predictor variables, they should be interpreted as different research interest in the study of human-carnivore relations. Therefore, while current research interest on beneficial contributions (or ecosystem services) showed a positive effect on the probability of mentioning species of Canidae and multiple social actors, we found that current research on detrimental contributions (or conflicts) was positively associated with the probability of mentioning species of Ursidae family (figure 5(c)). Thus, future research on human-bear relations needs to further investigate the causes of the lack of research on the beneficial contributions of bears to humans, and needs to consider a wider range of social actors. In addition, we suggest that research on human-carnivore relations develop and analyze a standard, comprehensive set of social and ecological factors, wherever possible, to allow for more direct comparisons across sites and over time. Several studies provide an excellent foundation for enumerating and refining those factors (Lischka et al 2018). Our findings on how different social and ecological characteristics led to differential research interest on ecosystem services and conflicts supports previous calls to apply social-ecological approaches in order to uncover the multiple beneficial and detrimental contributions provided by carnivores to people (Ceausu et al 2018, Lozano *et al* 2019, Jones *et al* 2019).

Third, we found that highly relevant actors, such as farmers and decision-makers are less represented in research than others, such as managers, the general public and rural residents (figure 2(a)). Human-carnivore research that overlooks the diversity of social actors involved in human-carnivore management can perpetuate and escalate the conflict with carnivores and create new social conflicts (Hartel *et al* 2019). To effectively promote coexistence, future research therefore should address the social causes underpinning conflicts, including human-human conflicts, the latter especially can be important but have often overlooked (Dickman 2010, Young et al 2010, Draheim et al 2015). In addition, previous research also shows that the lack of communication among actors involved in specific management strategies leads to ineffective management (Lute and Gore 2014, Browne-Nuñez et al 2015). To create long-term trustful communication between those social actors relevant for the management of human-carnivore relations, future research should promote participatory and transdisciplinary approaches in which multiple actors engage in the design and implementation of a coexistence strategy in the American West (Pooley et al 2017, Hovardas 2018, Lozano et al 2019). Specific methods include collaborative learning, mental models, discursive approaches, and structure decision making (Chan et al 2012, Ban et al 2013). In addition, Hartel et al (2019) proposes to go beyond the simple inclusion of multiple actors and research the deeper levels of values and norms that underpin the actors' actions and behavior. By incorporating norms and values, but also through sustained collaboration with local actors, transdisciplinary approaches can therefore contribute to the long-term viability of humancarnivore coexistence (Hartel et al 2019).

Finally, our results also indicate how land management across states of the American West seems to influence how often conflicts with carnivores are discussed (i.e. more conflicts mentioned on federal lands) (figure 5(d)). Federal lands prevail in the American West, however, agencies that manage these lands, like the Bureau for Land Management and the US Forest Service, have different missions and management approaches, making it very difficult to develop a coordinated strategy for reducing human-carnivore conflicts across federal lands under different jurisdictions and in different states. These obstacles require future research on how to build platforms for collaboration among key actors and institutions across states in order to foster coexistence in the American West. Although the formation of cross-state platforms might lead to new conflicts (Redpath et al 2017), such platforms could also represent new institutions by which multiple actors can embrace the challenge of coexisting with carnivores and engage to find shared solutions (see also Smith et al 2016, Hartel et al 2019). In this context, our proposal goes beyond including different sectors and actors since it aims to promote cross-state and transboundary collaborative management in the American West.

Conclusions

Our study has demonstrated that current research on human-carnivore relations in the American West has several knowledge gaps. These knowledge gaps include: (1) knowledge about the ecosystem services provided by carnivores, particularly regulating and provisioning, (2) knowledge on the social roots that underpin intolerance and coexistence, (3) information on how relevant actors, such as farmers and decisionmakers, relate with carnivores, and (4) effectiveness of different management practices to reduce humancarnivore conflicts and foster coexistence. Based on these findings, we call for a research agenda that applies social-ecological and transdisciplinary approaches to understand and manage human-carnivore relations in the American West. This agenda, in turn, should focus on four main themes: (1) the dual role of carnivores as providers of both beneficial and detrimental contributions to people, (2) social-ecological factors that underpin the provision of beneficial and detrimental contributions, (3) inclusion of diverse actors affected by carnivores or involved in their management, and (4) cross-state collaborative management. This agenda should be revised under an adaptive management framework in a context of rewilding and global change.

Acknowledgments

MEG was supported by a research contract from Sistema Nacional de Garantía Juvenil co-funded by the Social European Fund and the Junta de Andalucía, Spain (PID_UAL_2018/001). AJC, NHC, and JMRM were funded by the NSF Idaho EPSCoR Program under award number IIA-1301792 and OIA-1757324. MM and AFM were supported by Ramón y Cajal research contracts from the MINECO (RYC-2015-19231 and RYC-2016-21114, respectively), ZMR by a postdoctoral contract from the Generalitat Valenciana (APOSTD/2019/016) and ACA by a postdoctoral contract of Programa Viçent Mut of Govern Balear, Spain (PD/039/2017).

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID iDs

Neil H Carter ^(b) https://orcid.org/0000-0002-4399-6384

Juan M Requena-Mullor (1) https://orcid.org/0000-0002-5120-7947

Zebensui Morales-Reyes D https://orcid.org/0000-0002-4529-8651

References

Agarwala M, Kumar S, Treves A and Naughton-Treves L 2010 Paying for wolves in Solapur, India and Wisconsin, USA: comparing compensation rules and practice to understand the goals and politics of wolf conservation *Biol. Conserv.* **143** 2945–55

- Arbieu U, Grünewald C, Martín-López B, Schleuning M and Böhning-Gaese K 2017 Large mammal diversity matters for wildlife tourism in Southern African Protected Areas: insights for management *Ecosyst. Serv.* **31** 481–90
- Ban N C *et al* 2013 A social-ecological approach to conservation planning: embedding social considerations *Front. Ecol. Environ.* **11** 194–202
- Bangs E and Shivik J 2001 Managing Wolf conflict with livestock in the northwestern United States *Carnivore Damage Prevention News* **3** 2–5
- Besag J, York J and Mollié A 1991 Bayesian image restoration, with two applications in spatial statistics *Ann. Inst. Stat. Math.* **43** 1–20
- Beschta R L and Ripple W J 2009 Large predators and trophic cascades in terrestrial ecosystems of the western United States *Biol. Conservation* **142** 2401–14
- Blangiardo M and Cameletti M 2015 Spatial and Spatio-Temporal Bayesian Models with R–INLA(New York: Wiley)
- Bombieri G *et al* 2019 Brown bear attacks on humans: a worldwide perspective *Sci. Rep.* **9** 8573
- Borrini-Feyerabend G and Hill R 2015 Governance of the conservation of nature *Protected Area Governance and Management* (Camberra: Anu Press) pp 169–206
- Braczkowski A R, O'Bryan C J, Stringer M J, Watson J E M, Possingham H P and Beyer H L 2018 Leopards provide public health benefits in Mumbai, India *Front. Ecol. Environ.* 16 176–82
- Bradley E H, Robinson H S, Bangs E E, Kunkel K, Jimenez M D, Gude J A and Grimm T 2015 Effects of wolf removal on livestock depredation recurrence and wolf recovery in Montana, Idaho, and Wyoming J. Wildl. Manage. **79** 1337–46
- Browne-Nuñez C, Treves A, Macfarland D, Voyles Z and Turng C 2015 Tolerance of wolves in Wisconsin: a mixed-methods examination of policy effects on attitudes and behavioral inclinations *Biol. Conserv.* **189** 59–71
- Bruskotter J T 2013 The predator pendulum revisited: social conflict over wolves and their management in the western United States *Wildl. Soc. Bull.* **37** 674–9
- Bruskotter J T and Wilson R S 2014 Determining where the wild things will be: using psychological theory to find tolerance for large carnivores *Conserv. Lett.* **7** 158–65
- Bruskotter J T *et al* 2017 Modernization, risk, and conservation of the world's largest carnivores *BioScience* **67** 646–55
- Carter N H and Linnell J D C 2016 Co-adaptation is key to coexisting with large carnivores? *Trends Ecol. Evol.* **31** 575–8

Carter N H, Viña A, Hull V, McConnell W J, Axinn W, Ghimire D and Liu J 2014 Coupled human and natural systems approach to wildlife research and conservation *Ecol. Soc.* **19**43

- Ceausu S, Graves R, Killion A, Svenning J and Carter N H 2018 Governing trade-offs in ecosystem services and disservices to achieve human-wildlife coexistence *Conserv. Biol.* **33** 543–53
- Chan K M *et al* 2012 Where are cultural and social in ecosystem services? A framework for constructive engagement *BioScience* **62** 744–56
- Clark J A and May R M 2002 Taxonomic bias in conservation research *Science* 297 191–2
- Clarke K R 1993 Non-parametric multivariate analyses of changes in community structure *Aust. J. Ecol.* **18** 117–43
- Crespin S J and Simonetti J A 2019 Reconciling farming and wild nature: Integrating human–wildlife coexistence into the land-sharing and land-sparing framework *Ambio* 48 131–8
- Darimont C T, Paquet P C, Treves A, Artelle K A and Chapron G 2018 Political populations of large carnivores *Conserv. Biol.* 32 747–9
- Díaz S et al 2018 Assessing nature's contributions to people Science 359 270–2
- Dickman A J 2010 Complexities of conflict: the importance of considering social factors for effectively resolving humanwildlife conflict *Anim. Conserv.* **13** 458–66
- Draheim M M, Madden F, McCarthy J B and Parsons E C M 2015 Human-Wildlife Conflict: Complexity in the Marine Environment (Oxford: Oxford University Press)

Dressel S, Ericsson G and Sandström C 2018 Mapping socialecological systems to understand the challenges underlying wildlife management *Environ. Sci. Pol.* 84 105–12

- Eklund A, López-Bao J V, Tourani M, Chapron G and Frank J 2017 Limited evidence on the effectiveness of interventions to reduce livestock predation by large carnivores *Sci. Rep.* 7 2097
- Folke C *et al* 2011 Reconnecting to the biosphere *AMBIO J. Hum. Environ.* **40** 719–38 Frank B, Glikman J A and Marchini S 2019 *Human-Wildlife*
- Interactions: Turning Conflict into Coexistence (Cambridge: Cambridge University Press)
- Fernández-Gil A, Naves J, Ordiz A, Quevedo M, Revilla E and Delibes M 2016 Conflict misleads large carnivore management and conservation: brown bears and wolves in Spain *PLoS One* **11** e0151541
- George K A, Slagle K M, Wilson R S, Moeller S J and Bruskotter J T 2016 Changes in attitudes toward animals in the United States from 1978 to 2014 *Biol. Conserv.* **201** 237–42
- Gore M L, Knuth B A, Curtis P D and Shanahan J 2006 Stakeholder perceptions of risk associated with human-black bear conflicts in New York's adirondack park campgrounds: implications for theory and practice *Wildl. Soc. Bull.* **34** 36–43
- Harris N C and Dunn R R 2010 Using host associations to predict spatial patterns in the species richness of the parasites of North American carnivores *Ecol. Lett.* **12** 1411–8
- Hartel T, Scheele B, Vanak A, Rozylowicz L, Linnell J and Ritchie E 2019 Mainstreaming human and large carnivore coexistence through institutional collaboration *Conserv. Biol.* **33** 1256–65
- Hovardas T 2018 Addressing human dimensions in large carnivore conservation and management *Large Carnivore Conservation and Management (Human Dimensions)* ed T Hovardas (London: Routledge) pp 3–18
- Inskip C and Zimmermann A 2009 Human-felid conflict: a review of patterns and priorities worldwide *Oryx* **43** 18–34
- Ives C D, Abson D J, Wehrden H V, Dorninger C, Klaniecki K and Fischer J 2018 Reconnecting with nature for sustainability Sustain. Sci. 13 1389–97
- Ives C D *et al* 2017 Human-nature connection: a multidisciplinary review *Curr. Opin. Environ. Sustain.* 26–27 106–13
- Jones K *et al* 2019 What is the American West and why does it matter?: a social-ecological system perspective on research and management *Environ. Res. Lett.* **14** 115008
- Kansky R, Kidd M and Knight A T 2014 Meta-analysis of attitudes toward damage-causing mammalian wildlife *Conserv. Biol.* 28 924–38
- Kellert S R, Black M, Rush C R and Bath A J 1996 Human culture and large carnivore conservation in North America *Conserv. Biol.* **10** 977–90
- Killion A K, Melvin T, Lindquist E and Carter N H 2019 Tracking a half century of media reporting on gray wolves *Conserv. Biol.* **33** 645–54
- Kruskal J B 1964 Multidimensional scaling by optimizing goodness of fit to a nonmetric hypotheses *Psychometrika* **29** 1–27
- Kuijper D P et al 2016 Paws without claws? Ecological effects of large carnivores in anthropogenic landscapes Proc. R. Soc. B 283
- Linnell J, Swenson J E and Andersen R 2001 Predators and people: conservation of large carnivores is possible at high human densities if management policy is favourable *Anim. Conserv.* **4** 345–9
- Lischka S A, Teel T L, Johnson H E, Reed S E, Breck S, Carlos A D and Crooks K R 2018 A conceptual model for the integration of social and ecological information to understand humanwildlife interactions *Biol. Conserv.* **225** 80–7
- Lozano J et al 2019 Human-carnivore relations: a systematic review Biol. Conserv. 237 480–92
- Lute M and Gore M 2014 Stewardship as a path to cooperation? Exploring the role of identity in intergroup conflict among Michigan wolf stakeholders *Hum. Dimens. Wildl.* **19** 267–79 Lute M L, Navarrete C D, Nelson M P and Gore M L 2016 Moral
- dimensions of human-wildlife conflict *Conserv. Biol.* **30** 1200–11 MA (Millennium Ecosystem Assessment) 2005 *Ecosystems and*
- Human Well-being: Synthesis (Washington, DC: Island Press)

- Marchani S, Ferraz K M P M B, Zimmermann A, Guimarães-Luiz T, Morato R, Correa P L P and Macdonald D W 2019 Planning for coexistence in a complex human-dominated world *Human– Wildlife Interactions: Turning Conflict into Coexistence* ed B Frank *et al* (Cambridge: Cambridge University Press) pp 414–38
- Martín-López B, Montes C, Ramírez L and Benayas J 2009 What drives policy decision-making related to species conservation? *Biol. Conserv.* **142** 1370–80
- Miller J B 2015 Mapping attack hotspots to mitigate humancarnivore conflict: approaches and applications of spatial predation risk modeling *Biodivers. Conserv.* 24 1–25
- Moleón M, Sánchez-Zapata J A, Margalida A, Carrete M, Owen-Smith N and Donázar J A 2014 Humans and scavengers: the evolution of interactions and ecosystem services *Bioscience* **64** 394–403
- Morales-Reyes Z, Martín-López B, Moleón M, Mateo-Tomás P, Botella F, Margalida A, Donázar J A, Blanco G, Pérez I and Sánchez-Zapata J A 2018 Farmer perceptions of the ecosystem services provided by scavengers: what, who, and to whom *Conserv. Lett.* 11 e12392
- Moreira-Arce D, Ugarte C S, Zorondo-Rodríguez F and Simonetti J A 2018 Management tools to reduce carnivorelivestock conflicts: current gap and future challenges *Rangeland Ecol. Manage.* 71 389–94
- Morzillo A T, Mertig A G, Garner N and Liu J 2007 Resident attitudes toward black bears and population recovery in East Texas *Hum. Dimens. Wildl.* **12** 417–28
- Morzillo A T, Mertig A G, Hollister J W, Garner N and Liu J 2010 Socioeconomic factors affecting local support for black bear recovery strategies *Environ. Manage.* **45** 1299–311
- Nyhus P, Fischer H, Madden F and Osofsky S 2003 Taking the bite out of wildlife damage: the challenges of wildlife compensation schemes *Conserv. Pract.* 4 37–43
- O'Bryan C J, Braczkowski A R, Beyer H L, Carter N H, Watson J E M and McDonald-Madden E 2018 The contribution of predators and scavengers to human wellbeing *Nat. Ecol. Evol.* **2** 229–36
- Oksanen J 2019 Vegan: Community Ecology Package. R package version 2.5-5 (https://CRAN.R-project.org/package=vegan)
- Penteriani V *et al* 2016 Human behaviour can trigger large carnivore attacks in developed countries *Sci. Rep.* **6** 20552
- Peterson M N, Birckhead J L, Leong K, Peterson M J and Peterson T R 2010 Rearticulating the myth of human-wildlife conflict *Conserv. Lett.* **3**74–82
- Pooley S P *et al* 2017 An interdisciplinary review of current and future approaches to improving human-predator relations *Conserv. Biol.* **31** 513–23
- Pullin A S and Stewart G B 2006 Guidelines for Systematic Review in Conservation and Environmental Management *Conserv. Biol.* **20** 1647–56
- R Core Team 2019 R: A language and environment for statistical computing (Vienna: R Foundation for Statistical Computing) (https://R-project.org/)
- Redpath S *et al* 2017 Don't forget to look down—collaborative approaches to predator conservation *Biol. Rev.* **92** 2157–63
- Ripple W J et al 2014 Status and ecological effects of the world's largest carnivores Science 343 1241484
- Rue H, Sara Martino S and Chopin N 2009 Approximate bayesian inference for latent gaussian models using integrated nested Laplace approximations (with discussion) J. R. Stat. Soc. B 71 319–92
- Sarchet B A 2005 Survey and analysis of Wyoming livestock producers Masters Thesis University of Wyoming, Laramie, WY
- Slagle K M, Bruskotter J T, Singh A S and Schmidt R H 2017 Attitudes toward predator control in the United States: 1995 and 2014 J. Mammal. **98** 7–16
- Smith D W, White P, Stahler D, Wydeven A P and Hallac D 2016 Managing wolves in the Yellowstone area: balancing goals across jurisdictional boundaries *Wildl. Soc. Bull.* **40** 436–45
- Smith J B, Nielsen C and Hellgren E C 2014 Illinois resident attitudes toward recolonizing large carnivores J. Wildl. Manage. 78 930–43

- Smith T S and Herrero S 2018 Human-bear conflict in Alaska: 1880–2015 Wildl. Soc. Bull. 42 254–63
- Treves A 2009 Hunting for large carnivore conservation J. Appl. Ecol. 46 1350–6
- Treves A and Karanth K U 2003 Human-carnivore conflict and perspectives on carnivore management worldwide *Conserv. Biol.* 17 1491–9
- Treves A and Naughton-Treves L 2005 Evaluating lethal control in the management of human-wildlife conflict *People and Wildlife: Conflict or Coexistence*? ed R Woodroffe *et al* (Cambridge: Cambridge University Press) pp 86–106
- Treves A, Naughton-Treves L, Harper E, Mladenoff D, Rose R, Sickley T and Wydeven A 2004 Predicting human-carnivore conflict: a spatial model derived from 25 years of data on wolf predation on livestock *Conserv. Biol.* **18** 114–25
- Treves A, Wallace R B, Naughton-Treves L and Morales A 2006 Comanaging human–wildlife conflicts: a review *Hum. Dimens. Wildl.* 11 383–96

- van Eeden L M *et al* 2018 Carnivore conservation needs evidencebased livestock protection *PLoS Biol.* **16** e2005577
- Vincent C H, Hanson L A and Argueta C N 2017 Federal land ownership: overview and data *CRS Report* R42346 (Washington, DC: Congressional Research Service) 1–25
- Willemen L, Cottam A J, Drakou E G and Burgess N D 2015 Using social media to measure the contribution of red list species to the nature-based tourism potential of African protected areas *PLoS One* **10** e0129785
- Young J C, Marzano M, White R M, McCracken D L, Redpath S M, Carss D N, Quine C P and Watt A D 2010 The emergence of biodiversity conflicts from biodiversity impacts: characteristics and management strategies *Biodivers. Conserv.* **19** 3973–90
- Young J K, Ma Z, Laudati A and Berger J 2015 Human-carnivore interactions: lessons learned from communities in the American West *Hum. Dimens. Wildl.* **20** 349–66
- Ziegltrum G J and Nolte D L 2000 Black bear forest damage in Washington state, USA: economic, ecological, social aspects *Ursus* **12** 169–72