

Formulation of *Capra Hircus* Feed to Utilize *Artocarpus Heterophyllus* Leaves and Palm Acid Oil (PAO)

I Farah Amalina¹, J Muhammad Haziq¹, A R Abdul Syukor¹, A H Mohd Rashid²,
K A Nadzirah Izzati¹

¹ Faculty of Civil Engineering & Earth Resources, Universiti Malaysia Pahang, 26300 Gambang, Pahang, Malaysia.

² Faculty of Industrial Management, Universiti Malaysia Pahang, 26300 Gambang, Pahang, Malaysia.

*Corresponding Author: syukor@ump.edu.my, abdsyukor@gmail.com

Abstract. Palm Acid Oil (PAO) is the by-product from physical refining product of Palm Oil Mill Effluent (POME). PAO is the product from the last portion of converted Crude Palm Oil (CPO) in the end of that consist free fatty acid (FFA) >50%. The chemical properties in the PAO can make it be a part of animal feed as it is low in free fatty acid (FFA) and cheaper than CPO [1]. This study is focus on reducing the negative impact to the environment by utilizing the waste of POME, jackfruit leaves and soybean waste. Three formulations were tested on goats to determine the most suitable formulation. The formulations have different amount of each material. The nutrient in each formulation was calculated based on feeding guides by Department of Veterinary entitled nutrient composition of Malaysian feed materials and guides to feeding of cattle and goats [3]. Three goats were involved in this study. Two goats were fed with the formulations. Meanwhile, the other one was fed with its regular feeds which is Napier grass Taiwan and commercial pellet that act as a control. The result obtained will be compared with the control goat. The feed intake of each goat was recorded. As results, the suitable formulation of the composition of waste material had been choose as they can increase the weight and performance of Asian domestic goats. The uses of waste proved which is development of waste to be animal feed to meet the green technology and Malaysia Industrial revolution in livestock industries [2].

1. Introduction

The palm oil industry has developed as one top industry in Malaysia since the beginning of 1970. In facts, Malaysia is one of the world's largest producer of palm oil with more than 5 million ha of land under oil palm. Currently, Malaysia accounts for 39% of world palm oil production and 44% of world palm oil exportation. It is shown that Malaysia has played an important role in fulfilling the growing global need for oils and fats sustainably. The production of palm oil results in the generations of large quantities of polluted wastewater referred to as POME. POME is the waste in the form of liquid that discharges from palm oil extraction in the palm oil mill. The residue left from the purification of CPO contains various dirt and suspended solids. Besides, it finds that the residue is highly polluting materials due to increases of biological oxygen demand (BOD), low PH and colloidal nature. So, it

Note N03 Importance of Dry Matter Intake Estimating Dry Matter Intake 1 . Measure How Much of Each Feed Indicators of Adequate / Inadequate Daily Dry Matter Intake Useful Rules of Thumb.”

- [12] Rahman and Mohammad Mijanur 2014. “*Feed Intake and Growth Performance of Goats Supplemented with Soy Waste.*” *Pesquisa Agropecuaria Brasileira* **49(7)**: 554–58.
- [13] Rahman and M M 2014. “*Feed Intake and Growth Performance of Goats Offered Napier Grass (Pennisetum Purpureum) Supplemented with Concentrate Pellet and Soya Waste.*” *Sains Malaysiana* **43(7)**: 967–71.
- [14] Van and Do Thi Thanh 2006. *Some Animal and Feed Factors Affecting Feed Intake, Behaviour and Performance of Small Ruminants.* *J Dairy Sci.* 74:3583–3597. 10.3168/jds. S0022-0302(91)78551-2
- [15] Escala, Emely J and Lolito C Bestil 2014. “*Supplementation of Jackfruit Concentrate for Early Weaning of Kids.*” **1(2)**: 221-24
- [16] Dong, Nguyen Thi Kim, K. Elwinger, J E Lindberg, and R B Ogle 2005. “*Effect of Replacing Soybean Meal with Soya Waste and Fish Meal with Ensiled Shrimp Waste on the Performance of Growing Crossbred Ducks.*” *Asian-Australasian Journal of Animal Sciences* **18(6)**: 825-34
- [17] Makkar, H. P. S.; Tran, G.; Heuzé, V.; Giger-Reverdin, S.; Lessire, M.; Lebas, F.; Ankers, P., 2016. Seaweeds for livestock diets: A review. *Anim. Feed Sci. Technol.*, 212: 1-17
- [18] Ainie Kuntom, Wai-Lin Slew and Yew-Ai Tan 1994. *Chemistry and Technology Division, Palm Oil Research Institute of Malaysia, Ministry of Primary Industry, Kuala Lumpur, Malaysia*
- [19] Teresa Banaszkiwicz 2011. “*Nutritional Value of Soybean Meal.*” In book: *Soybean and Nutrition*
- [20] Chan, D K S, and Cheung S F 1998. “*Tannins and human health.*” Department of Microbiology and Molecular Cell Sciences, University of Memphis, TN 38152, USA.
- [21] Shuhong Li, Dan Zhu, Kejuan Li, Yingnan Yang, Zhongfang Lei, and Zhenya Zhang 2013. “*Soybean Curd Residue: Composition, Utilization, and Related Limiting Factors.*” Graduate School of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8572, Japan
- [22] Laura B Harthan and Debbie J R Cherney 2017. “*Okara as a protein supplement affects feed intake and milk composition of ewes and growth performance of lambs.*” Published in *Animal nutrition* 2017.

Acknowledgement

This study was supported by Universiti Malaysia Pahang (UMP), and Ministry of Higher Education (MOHE), Malaysia. The authors are grateful to Faculty of Civil Engineering and Earth Resources (FKASA), UMP for the fully support for this research project.