

## Università degli Studi di Padova

## Padua Research Archive - Institutional Repository

Use of WWI photos for quantitative reconstructions of glaciers along the Italian-Austrian front

Original Citation:
Availability: This version is available at: 11577/3322740 since: 2020-01-23T10:37:35Z
Publisher:
Published version: DOI:
Terms of use: Open Access
This article is made available under terms and conditions applicable to Open Access Guidelines, as described at http://www.unipd.it/download/file/fid/55401 (Italian only)

## EGU 2020 Vienna | Austria | 3–8 May 2020

GI1.3 - The use of historical images in geosciences Co-organized by EOS9/CL2/CR2/GM2/HS13/NH6

## Use of WWI photos for quantitative reconstructions of glaciers along the Italian-Austrian front

Carturan L.<sup>1,2</sup>, Bondesan A.<sup>3,6</sup>, Carton A.<sup>1</sup>, Cazorzi F.<sup>4</sup>, Cucchiaro S.<sup>4</sup>, De Marco J.<sup>4</sup>, Piermattei L.<sup>5</sup>

<sup>1</sup> Dipartimento di Geoscienze, Università degli Studi di Padova, Italy
 <sup>2</sup> Dipartimento Territorio e Sistemi Agro-Forestali, Università degli Studi di Padova, Italy
 <sup>3</sup> Dipartimento di Scienze Storiche, Geografiche e dell'Antichità, Università degli Studi di Padova, Italy
 <sup>4</sup> Dipartimento di Scienze AgroAlimentari, Ambientali e Animali, Università degli Studi di Udine, Italy
 <sup>5</sup> Physical Geography, Catholic University of Eichstaett-Ingolstadt, Eichstaett, Germany
 <sup>6</sup> Department of Military Geography, University of Stellenbosch, South Africa (Research Fellow)

The knowledge of past fluctuations of glaciers is the key for understanding their dynamics and their climaterelated evolution. Glacier mass balance and length changes are the two metrics normally used for reconstructing past fluctuations series of glaciers. However, length change measurements series are often discontinuous and require validation, whereas mass balance measurements are available for only few glaciers worldwide and only for the latest decades.

In the context of glacier reconstructions, other sources of information such as historical-archival, glacioarchaeological and geomorphological data are of critical importance, because they enable the completion and validation of direct measurement series and their extension into the past, providing spatial and temporal constrains

A unique source of unexploited historical information dating back to the First World War (WWI, 1915-1918) exist for many glaciers in the Eastern Italian Alps. This information mainly consists of old photos, which however are spread over a multitude of sources and in many cases are not yet digitized.

We propose a workflow that enable extracting quantitative information from terrestrial photographs taken during the WWI period, aimed at the reconstruction of glacier area, volume and firn lines by means of the monoplotting technique. This method relies on the availability of high-resolution digital elevation models, which became recently available over wide areas thanks to LiDAR and aerial photogrammetry. This work presents the methods applied, and the results obtained, on several case studies in the Adamello-Presanella, Ortles-Cevedale, Dolomites, and Julian Alps.