

Classification of some nilpotent class of Leibniz superalgebras

L.M. Camacho, J.R. Gómez, R.M. Navarro, B.A. Omirov

Abstract

The aim of this work is to present the description of Leibniz superalgebras up to isomorphism with characteristic sequence $(n|m-1, 1)$ and nilindex $n+m$.

2000 MSC: 17A32, 17B30

Keywords: Lie superalgebras, Leibniz superalgebras, nilindex, characteristic sequence.

1 Introduction

During many years the theory of Lie superalgebras has been actively studied by many mathematicians and physicists. Many works have been devoted to them but few have dealt with nilpotent Lie superalgebras. Recent works [5]–[9] have studied the problems of description of some classes of nilpotent Lie superalgebras. It is well known that Lie superalgebras are a generalization of Lie algebras [10]. In the same way, the notion of Leibniz algebras can be generalized by means of Leibniz superalgebras. The elementary properties of Leibniz superalgebras were obtained in [1]. The description of case of maximal nilindex for nilpotent Leibniz superalgebras (nilpotent Leibniz superalgebras distinguished by the feature of being single-generated) is not difficult and was done in [1]. However, the next stage is very problematic, it consists of Leibniz superalgebras with dimensions even and odd parts equal to n and m , respectively, and of nilindex $n+m$. It should be noted that such Lie superalgebras were classified in [8]. Due to the great difficulty in solving the problem, some restrictions on the characteristic sequence are added. The experience of using the characteristic sequence in Lie and Leibniz algebras (even in Lie superalgebras) leads us to choose a restriction on this invariant. Since graded non-commutative identity in non Lie Leibniz superalgebras does not hold, we usually have to solve many technical tasks when describing Leibniz superalgebras, [3].

In a similar way to Leibniz algebras and Lie superalgebras cases, it is possible to define the notions of null-filiform and filiform Leibniz superalgebras [2], [5] as superalgebras with characteristic sequences $(n|m)$ and $(n-1, 1|m)$ respectively. We should take into account that the superalgebras in [7] show that all null-filiform superalgebras have nilindex $n+m$ (except for a unique superalgebra of maximal nilindex) and there exist filiform superalgebras which also have nilindex $n+m$ for some n, m . In the present paper we investigate Leibniz superalgebras with the characteristic sequence $C(L) = (n|m-1, 1)$ and with nilindex equal to $n+m$.

In this paper all spaces and superalgebras are considered over the complex number field.

2 Preliminaries

We recall the definition of Leibniz superalgebras.

Definition 2.1. A \mathbb{Z}_2 -graded vector space $L = L_0 \oplus L_1$ is called a Leibniz superalgebra if it is equipped with a product $[-, -]$ which satisfies the following conditions:

$$[L_\alpha, L_\beta] \subseteq L_{\alpha+\beta(\text{mod}2)} \quad \text{for all } \alpha, \beta \in \mathbb{Z}_2,$$

$$[x, [y, z]] = [[x, y], z] - (-1)^{\alpha\beta} [[x, z], y] \quad \text{graded Leibniz identity}$$

for all $x \in L$, $y \in L_\alpha$, $z \in L_\beta$, $\alpha, \beta \in \mathbb{Z}_2$.

Note that if in L the identity $[x, y] = -(-1)^{\alpha\beta} [y, x]$ (where $x \in L_\alpha$, $y \in L_\beta$) holds, then the graded Leibniz and graded Jacobi identities coincide. Thus, Leibniz superalgebras are a generalization of Lie superalgebras.

Let us anote an example of non Lie Leibniz superalgebras, which generalize the construction of non Lie Leibniz algebras [11].

Let $A = A_0 \oplus A_1$ be an associative superalgebra over a field F and $D : A \rightarrow A$ be a F -linear map satisfying the condition:

$$D(a(Db)) = DaDb = D((Da)b)$$

for all $a, b \in A$. If in the vector space A we define the new product:

$$\langle a, b \rangle_D := a(Db) - (-1)^{\alpha\beta} D(b)a$$

for $a \in A_\alpha$, $b \in A_\beta$, A becomes a Leibniz superalgebra.

Let us introduce some notations

$$\mathfrak{R}(L) = \{R_x \mid x \in L\},$$

$$Leib^{n,m} = \{L = L_0 \oplus L_1 \mid \dim L_0 = n, \dim L_1 = m\}.$$

It is not difficult to see that the set $\mathfrak{R}(L)$ will be a Lie superalgebra with the following multiplication:

$$\langle R_a, R_b \rangle := R_a R_b - (-1)^{\alpha\beta} R_b R_a$$

for all $R_a \in \mathfrak{R}(L)_\alpha$, $R_b \in \mathfrak{R}(L)_\beta$.

Let $V = V_0 \oplus V_1$, $W = W_0 \oplus W_1$ be two \mathbb{Z}_2 -graded spaces. We say that a linear map $f : V \rightarrow W$ has degree α (denoted as $\deg(f) = \alpha$), if $f(V_\beta) \subseteq W_{\alpha+\beta}$ for all $\beta \in \mathbb{Z}_2$.

Definition 2.2. Let L and L' be Leibniz superalgebras. A linear map $f : L \rightarrow L'$ is called a homomorphism of Leibniz superalgebras if

1. f preserves the grading, i.e. $f(L_0) \subseteq L'_0$ and $f(L_1) \subseteq L'_1$ ($\deg(f) = 0$);
2. $f([x, y]) = [f(x), f(y)]$ for all $x, y \in L$.

Moreover, if f is one-to-one then it is called an isomorphism of Leibniz superalgebras L and L' .

For a given Leibniz superalgebra L we define a descending central sequence as follows:

$$L^1 = L, \quad L^{k+1} = [L^k, L^1], \quad k \geq 1.$$

Definition 2.3. A Leibniz superalgebra L is called nilpotent, if there exists $s \in \mathbb{N}$ such that $L^s = 0$. The minimal number s with this property is called index of nilpotency (nilindex) of the superalgebra L .

Definition 2.4. The set $R(L) = \{z \in L \mid [L, z] = 0\}$ is called the right annihilator of a superalgebra L .

Using the Leibniz graded identity it is not difficult to see that $R(L)$ is an ideal of the superalgebra L . Moreover, elements of the form $[a, b] + (-1)^{\alpha\beta}[b, a]$ ($a \in L_\alpha$, $b \in L_\beta$) belong to $R(L)$.

The description of Leibniz superalgebras of maximal nilindex is represented in the following theorem.

Theorem 2.1. [1]. Let L be an n -dimensional Leibniz superalgebra with maximal index of nilpotency. Then L is isomorphic to one of the following two non isomorphic superalgebras:

$$\begin{cases} [e_i, e_1] = e_{i+1}, & 1 \leq i \leq n-1 \\ [e_i, e_1] = e_{i+1}, & 1 \leq i \leq n-1 \\ [e_i, e_2] = 2e_{i+2}, & 1 \leq i \leq n-2 \end{cases}$$

where the omitted products are zero.

It should be noted that for the second superalgebra when $n + m$ is even, we have $m = n$ and if $n + m$ is odd then $m = n + 1$. Moreover, it is clear that the Leibniz superalgebra has the maximal nilindex if and only if it is one-generated.

We define the characteristic sequence as in [5].

Let $L = L_0 \oplus L_1$ be a nilpotent Leibniz superalgebra. For an arbitrary element $x \in L_0$, the operator of right multiplication R_x is a nilpotent endomorphism of the space L_i , where $i \in \{0, 1\}$. Let us denote by $C_i(x)$ ($i \in \{0, 1\}$) the descending sequence of the dimensions of Jordan blocks of the operator R_x . Consider the lexicographical order on the set $C_i(L_0)$.

Definition 2.5. A sequence

$$C(L) = \left(\max_{x \in L_0 \setminus [L_0, L_0]} C_0(x) \mid \max_{\tilde{x} \in L_0 \setminus [L_0, L_0]} C_1(\tilde{x}) \right)$$

is said to be the characteristic sequence of the Leibniz superalgebra L .

Similarly to [6] (corollary 3.0.1) it can be proved that the characteristic sequence is invariant under isomorphisms.

3 Description of Leibniz superalgebras with characteristic sequence $(n \mid m - 1, 1)$ and nilindex $n + m$

The following theorem gives us the location of the generators of the Leibniz superalgebra from $Leib^{n,m}$ with characteristic sequence $(n \mid m - 1, 1)$ and nilindex $n + m$.

Theorem 3.1. *Let $L = L_0 \oplus L_1$ be a Leibniz superalgebra from the variety $Leib^{n,m}$ with characteristic sequence $(n \mid m - 1, 1)$ and nilindex $n + m$. Then L is two-generated and they belong to L_1 .*

Proof. Since the nilindex of L is $n + m$, then superalgebra L is two-generated. From the definition of characteristic sequence we can conclude that there exists a basis $\{y_1, y_2, \dots, y_m\}$ of L_1 such that the operator $R_{x_1|L_1}$ in this basis has one of the following forms:

$$\begin{pmatrix} J_{n-1} & 0 \\ 0 & J_1 \end{pmatrix}, \quad \begin{pmatrix} J_1 & 0 \\ 0 & J_{n-1} \end{pmatrix}.$$

By a change of the basis elements $\{y_1, y_2, \dots, y_m\}$, we can assume that the operator $R_{x_1|L_1}$ has the first form.

Since $C(L_0) = (n)$, then from [2] (example 1) we have that L_0 is a zero-filiform Leibniz algebra. Without loss of generality we may suppose that

$$\begin{cases} [x_i, x_1] = x_{i+1}, & 1 \leq i \leq n - 1, \\ [y_j, x_1] = y_{j+1}, & 1 \leq j \leq m - 2, \\ [y_m, x_1] = [y_m, x_1] = 0, \end{cases}$$

where $\{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_m\}$ is the basis of the superalgebra L .

From these multiplications we deduce that $\{x_2, x_3, \dots, x_n\}$ lie in the right annihilator of the superalgebra L and generators can be selected as a linear combination of elements of the set $\{x_1, y_1, y_m\}$.

It is clear that two generators can not lie in L_0 . Let us suppose the opposite assertion to the assertion of the theorem, i.e. one generator element lies in L_0 and the second lies in L_1 . Then we can choose as generators the elements $x_1, Ay_1 + By_m$.

Case 1. Let us suppose $A = 0$. Then x_1 and y_m can be chosen as generators. Hence

$$L^2 = \{x_2, x_3, \dots, x_n, y_1, y_2, \dots, y_{m-1}\}.$$

Introduce the notations

$$[y_1, y_m] = \sum_{k=2}^n \beta_k x_k, \quad [x_1, y_m] = \sum_{s=1}^{m-1} \alpha_s y_s.$$

Consider the products

$$[x_i, [y_m, x_1]] = [[x_i, y_m], x_1] - [[x_i, x_1], y_m], \quad 1 \leq i \leq n.$$

On the other hand, $[x_i, [y_m, x_1]] = 0$.

Therefore

$$\begin{aligned} [x_i, y_m] &= \sum_{k=1}^{m-i} \alpha_k y_{k-1+i}, & 1 \leq i \leq \min\{n, m-1\}, \\ [x_i, y_m] &= 0, & \min\{n, m-1\} < i \leq n. \end{aligned}$$

Thus, we obtain $y_1 \notin \langle [x_i, y_m] \rangle$, $2 \leq i \leq n$ and since $y_1 \in L^2$, then $\alpha_1 \neq 0$. Consider the products

$$[y_j, [y_m, x_1]] = [[y_j, y_m], x_1] - [[y_j, x_1], y_m], \quad 1 \leq j \leq m-2.$$

On the other hand we have $[y_j, [y_m, x_1]] = 0$, $1 \leq j \leq m-2$.

Therefore

$$\begin{aligned} [y_j, y_m] &= \sum_{s=2}^{n+1-j} \beta_s x_{s-1+j}, & 1 \leq j \leq \min\{m-1, n-1\}, \\ [y_j, y_m] &= 0, & \min\{n-1, m-1\} < j \leq m-1. \end{aligned}$$

Consider the equalities

$$[x_1, [y_m, y_m]] = 2[[x_1, y_m], y_m] = 2\left[\sum_{s=1}^{m-1} \alpha_s y_s, y_m\right] = \sum_{s=1}^{m-1} \alpha_s \sum_{t=2}^{n+1-s} \beta_t x_{t-1+s}.$$

Since $[y_m, y_m] \in \langle x_2, x_3, \dots, x_n \rangle$ then $[y_m, y_m] \in R(L)$ and hence $[x_1, [y_m, y_m]] = 0$. Therefore,

$$\sum_{s=1}^{m-1} \alpha_s \sum_{t=2}^{n+1-s} \beta_t x_{t-1+s} = 0.$$

Comparing the coefficient at the basic elements x_i , $2 \leq i \leq n$, we obtain $\beta_i = 0$, $2 \leq i \leq n$. And we have $L^3 = \{x_3, x_4, \dots, x_n, y_2, y_3, \dots, y_{m-1}\}$, but it follows that the index of nilpotency of the superalgebra L is smaller than $n+m$. Thus, we have contradiction with assumption $A=0$.

Case 2. $A \neq 0$. Then we can take as generators x_1 and y_1 . Hence, $L^2 = \{x_2, x_3, \dots, x_n, y_2, y_3, \dots, y_m\}$.

Let us define the notations

$$[x_i, y_1] = \sum_{k=2}^m \alpha_{i,k} y_k, \quad 1 \leq i \leq n, \quad [y_j, y_1] = \sum_{s=2}^n \beta_{j,s} x_s, \quad 1 \leq j \leq m. \quad (3.1)$$

Let us suppose $x_2 \notin L^3$. Then $L^3 = \{x_3, x_4, \dots, x_n, y_2, y_3, \dots, y_m\}$ and there exist some i_0, j_0 ($2 \leq i_0, j_0$) such that $\alpha_{i_0,2} \alpha_{j_0,m} - \alpha_{j_0,2} \alpha_{i_0,m} \neq 0$ and $\alpha_{i_0,2} \alpha_{j_0,m} \neq 0$.

Since $[x_{i_0}, y_1] = \sum_{k=2}^m \alpha_{i_0,k} y_k \in R(L)$ and $\alpha_{i_0,2} \neq 0$ then multiplying from the right side by x_1 enough times, we obtain $\{y_3, y_4, \dots, y_{m-1}\} \subseteq R(L)$.

Therefore

$$[x_{i_0,2}, y_1] - \sum_{k=3}^{m-1} \alpha_{i_0,k} y_k = \alpha_{i_0,2} y_2 + \alpha_{i_0,m} y_m \in R(L),$$

$$[x_{j_0,2}, y_1] - \sum_{s=3}^{m-1} \alpha_{j_0,s} y_s = \alpha_{j_0,2} y_2 + \alpha_{j_0,m} y_m \in R(L).$$

Since $\alpha_{i_0,2} \alpha_{j_0,m} - \alpha_{j_0,2} \alpha_{i_0,m} \neq 0$, then we have $y_2, y_m \in R(L)$, i.e. $\{y_2, y_3, \dots, y_m\} \subseteq R(L)$ and hence $[x_1, y_1] \in R(L)$.

From equality

$$[[x_{i-1}, x_1], y_1] = [x_{i-1}, [x_1, y_1]] + [[x_{i-1}, y_1], x_1],$$

we obtain

$$\begin{aligned} [x_i, y_1] &= \sum_{k=2}^{m-i} \alpha_{1,k} y_{k+i-1}, & 2 \leq i \leq \min\{n, m-2\}, \\ [x_i, y_1] &= 0, & \min\{n, m-2\} < i \leq n, \end{aligned}$$

from which we have a contradiction to the assumption $\alpha_{i_0,2} \neq 0$.

Therefore $x_2 \in L^3$ and x_2 is a linear combination of products $[y_i, y_1]$, $1 \leq i \leq m$, hence $[x_2, y_1] = \sum_{i=1}^m \gamma_i [[y_i, y_1], y_1]$. Using the Leibniz graded identity and $[y_1, y_1] \in R(L)$ one can easily see that $[x_2, y_1] = 0$.

Thus, we have $L^3 = \{x_2, x_3, \dots, x_n, Cy_2 + Dy_m, y_3, \dots, y_{m-1}\}$.

Since $[x_2, y_1] = 0$, then $Cy_2 + Dy_m \in L^3$ should be expressed by linear combinations of products $[x_i, y_1]$, $3 \leq i \leq n$. Therefore, $Cy_2 + Dy_m \in L^5$.

Moreover, if $C \neq 0$, then $Cy_2 + Dy_m \in R(L)$. Then multiplying from the right side by x_1 enough times, we obtain $\{y_3, y_4, \dots, y_{m-1}\} \subseteq R(L)$.

Since $[x_2, y_1] = 0$ and $Cy_2 + Dy_m \in L^3$ then there exist t_0 , $t_0 \geq 3$, such that $[x_{t_0}, y_1] = c_2(Ay_2 + By_m) + c_3y_3 + \dots + c_{m-1}y_{m-1}$, $c_2 \neq 0$.

Applying the Leibniz graded identity we obtain

$$[x_1, \underbrace{[\dots [[y_1, x_1], x_1], \dots, x_1]}_{t_0-1\text{-times}}] = (-1)^{t_0-1} c_2 (Cy_2 + Dy_m) + \{y_3, y_4, \dots, y_{m-1}\}. \quad (3.2)$$

As L is nilpotent, we have existence $s \in \mathbb{N}$, such that $y_{t_0} \in L^s \setminus L^{s+1}$.

If $t_0 < m$, then $[x_1, y_{t_0}] = (-1)^{t_0-1} c_2 (Cy_2 + Dy_m) + \{y_3, y_4, \dots, y_{m-1}\} \in L^{s+1}$ and multiplying this equality $(t_0 - 2)$ times from the right side by x_1 , we obtain that $(-1)^{i_0-1} c_2 Cy_{t_0} \in L^{s+t_0-1}$. The inequality $s + t_0 - 1 > s$ contradicts the condition $y_{t_0} \in L^s \setminus L^{s+1}$ and hence we obtain $C = 0$.

If $t_0 \geq m$, then $[x_1, y_{t_0}] = 0$. From (3.2) we again obtain $C = 0$.

Thus, $L^3 = \{x_2, x_3, \dots, x_n, y_3, \dots, y_{m-1}, y_m\}$.

Consider the following subcases.

Case 2.1 Let $\alpha_{1,2} \neq 0$. Suppose that $x_2 \in L^l \setminus L^{l+1}$ for some l ($3 \leq l \leq m$). Then

$$L^l = \{x_2, x_3, \dots, x_n, y_l, \dots, y_{m-1}, y_m\}.$$

$$L^{l+1} = \{x_3, x_4, \dots, x_n, y_l, \dots, y_{m-1}, y_m\}.$$

Since $x^2 \in L^l \setminus L^{l+1}$, we obtain $\beta_{l-1,2} \neq 0$, i.e.

$$[y_{l-1}, y_1] = \sum_{k=2}^n \beta_{l-1,k} x_k.$$

The equality $[x_2, y_1] = 0$, deduce $y_l \in L^{l+2}$ and $L^{l+2} = \{x_4, x_5, \dots, x_n, y_l, \dots, y_{m-1}, y_m\}$.

Therefore $[y_l, y_1] = \sum_{k=4}^n \beta_{l,k} x_k$, i.e. $\beta_{l,2} = \beta_{l,3} = 0$

In notation (3.1) by induction on j for any value of i one can prove the following equality:

$$[y_i, y_j] = \sum_{k=0}^{\min\{i+j-1, m-1\}-i} (-1)^k C_{j-1}^k \sum_{t=2}^{n-j+k+1} \beta_{i+k,t} x_{t+j-k-1}, \quad (3.3)$$

where $1 \leq j \leq m-1$, $1 \leq i \leq m-1$.

From (3.3) we have $[y_2, y_l] = \beta_{l-1,2} x_3 + x_4, x_5, \dots, x_n$

Consider the equalities

$$\begin{aligned} [x_1, [y_1, y_l]] &= [[x_1, y_1], y_l] + [[x_1, y_l], y_1] = \\ &= \sum_{k=2}^m \alpha_{1,k} [y_k, y_l] + \sum_{s=l+1}^m \gamma_{1,s} [y_k, y_1] = \alpha_{1,2} \beta_{l-1,2} x_3 + \{x_4, \dots, x_n\}. \end{aligned}$$

On the other hand

$$[x_1, [y_1, y_l]] = 0.$$

So $\alpha_{1,2} \beta_{l-1,2} x_3 = 0$, i.e. we have a contradiction with supposition $\alpha_{1,2} \neq 0$.

Case 2.1. If $\alpha_{1,2} = 0$. Then $[y_1, x_1] + [x_1, y_1] = y_2 + \alpha_{1,3} y_3 + \alpha_{1,4} y_4 + \dots + \alpha_{1,m} y_m \in R(L)$ and then multiplying from the right side by x_1 enough times, we obtain $y_2 + \alpha_{1,m} y_m, y_3, \dots, y_{m-1} \in R(L)$.

Since $y_m \in L^3$, there exists $i_0 \geq 2$ such that

$$[x_{i_0}, y_1] = \alpha_{i_0,3} y_3 + \alpha_{i_0,4} y_4 + \dots + \alpha_{i_0,m} y_m, \quad \alpha_{i_0,m} \neq 0.$$

From $[x_{i_0}, y_1] \in R(L)$, we have $\alpha_{i_0,3} y_3 + \alpha_{i_0,4} y_4 + \dots + \alpha_{i_0,m} y_m \in R(L)$. Therefore $y_2, y_m \in R(L)$.

Consider the equalities

$$\begin{aligned} [y_j, [y_1, x_1]] &= [[y_j, y_1], x_1] - [[y_j, x_1], y_1] = \left[\sum_{k=2}^{n+1-j} \beta_{j,k} x_{k-1+j}, x_1 \right] - [y_{j+1}, y_1] = \\ &= \sum_{k=2}^{n-j} \beta_{j,k} x_{k+j} - [y_{j+1}, y_1]. \end{aligned}$$

On the other hand we have $[y_j, [y_1, x_1]] = [y_j, y_2] = 0$.

Therefore

$$[y_{j+1}, y_1] = \sum_{k=2}^{n-j} \beta_{1,2} x_{k+j}, \quad 1 \leq j \leq \min\{n-2, m-2\},$$

$$[y_{j+1}, y_1] = 0 \quad \min\{n-2, m-2\} \leq j \leq m-2.$$

From these we conclude that x_2 cannot be expressed via linear combination of $\{[y_j, y_1]\}$, $2 \leq j \leq m$. Therefore $x_2 \notin L^3$, it is a contradiction with assumption $x_2 \in L^3$.

Thus, the supposition of the case when one generator lies in L_0 and the second lies in L_1 leads to a contradiction, therefore both generators belong to L_1 and the proof of the theorem is completed. \square

The following lemma is an important technical result in our description.

Lemma 3.1. *Let $L = L_0 \oplus L_1$ be a Leibniz superalgebra from the variety $\text{Leib}^{n,m}$ with characteristic sequence $(n \mid m-1, 1)$ and nilindex $n+m$. Then there exists $y \in L_1$ such that the superalgebra $\langle y \rangle$ is isomorphic to the superalgebra from Theorem 2.1 and $m \in \{n+1, n+2\}$.*

Proof. From Theorem 3.1 we have that both generators are in L_1 . Let $\{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_m\}$ be the basis of L such that y_1, y_m are the generators and

$$[x_i, x_1] = x_{i+1}, \quad 1 \leq i \leq n-1, \quad [y_j, x_1] = y_{j+1}, \quad 1 \leq j \leq m-2,$$

$$\begin{aligned} [y_1, y_1] &= \sum_{i=1}^n a_i x_i, & [y_m, y_1] &= \sum_{i=1}^n b_i x_i, \\ [y_1, y_m] &= \sum_{i=1}^n c_i x_i, & [y_m, y_m] &= \sum_{i=1}^n d_i x_i. \end{aligned}$$

Since $[y_j, x_1] = y_{j+1}$, $1 \leq j \leq m-2$, then x_1 cannot be generated via multiplications of the elements $\{y_2, y_3, \dots, y_{m-1}\}$. Therefore $x_1 \in \langle [y_1, y_1], [y_m, y_1], [y_1, y_m], [y_m, y_m] \rangle$ and hence $(a_1, b_1, c_1, d_1) \neq (0, 0, 0, 0)$.

Let us suppose that $a_1 \neq 0$. Then making the change

$$x'_i = \sum_{k=1}^{n+1-i} a_k x_{k+i-1}, \quad 1 \leq i \leq n, \quad y'_j = a_1^{i-1} y_j, \quad 1 \leq j \leq m-1$$

we obtain

$$[x_i, x_1] = x_{i+1}, \quad 1 \leq i \leq n-1, \quad [y_j, x_1] = y_{j+1}, \quad 1 \leq j \leq m-2, \quad [y_1, y_1] = x_1.$$

It is clear that the subsuperalgebra $\langle y_1 \rangle = \{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_{m-1}\}$ is one-generated and therefore it has maximal index of nilpotency. Hence from Theorem 2.1 we have that either $m = n+1$ or $m = n+2$.

Let us suppose now that $a_1 = 0$.

Consider the product

$$[y_m, [y_m, x_1]] = [[y_m, y_m], x_1] - [[y_m, x_1], y_m].$$

Then $[y_m, y_m] = d_n x_n$ and $d_i = 0$, $1 \leq i \leq n-1$. It leads to $(b_1, c_1) \neq (0, 0)$.

From $[y_m, y_1] - [y_1, y_m] = \sum_{i=1}^n (b_i - c_i) x_i \in R(L)$ and $x_j \in R(L)$, $2 \leq j \leq n$ we obtain that $(b_1 - c_1) x_1 \in R(L)$, but $x_1 \notin R(L)$ and hence $b_1 = c_1 \neq 0$.

By the following change of basis:

$$x'_1 = \sum_{k=1}^n b_k x_k, \quad x'_{i+1} = [x'_i, x'_1], \quad 1 \leq i \leq n-1, \quad y'_j = y_j, \quad 1 \leq j \leq m$$

we can assume $[y_m, y_1] = x_1$.

From the products

$$\begin{aligned} [y_m, [y_i, x_1]] &= [[y_m, y_i], x_1] - [[y_m, x_1], y_i], \\ [y_m, [y_1, y_1]] &= 2[[y_m, y_1], y_1] = 2[x_1, y_1] \end{aligned}$$

we obtain $[y_m, y_{i+1}] = x_{i+1}$, $1 \leq i \leq m-2$, $[x_1, y_1] = 0$.

Since $[y_1, x_1] + [x_1, y_1] \in R(L)$ then $y_2 \in R(L)$, but this contradicts $[y_m, y_2] = x_2$ and therefore the case $a_1 = 0$ is not possible either. \square

The existence of an adapted basis with conditions of Lemma 3.1 in case of $m = n+1$ is described in the following lemma.

Lemma 3.2. *Let L be a Leibniz superalgebra from the variety $\text{Leib}^{n,m}$ with characteristic sequence $(n \mid m-1, 1)$ and nilindex $n+m$. Then, in case $m = n+1$, there exists a basis $\{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_{n+1}\}$ of L in which the products have the following form:*

$$\begin{aligned} [x_i, x_1] &= x_{i+1}, \quad 1 \leq i \leq n-1, & [y_j, x_1] &= y_{j+1}, \quad 1 \leq j \leq n-1, \\ [x_i, y_1] &= \frac{1}{2}y_{i+1}, \quad 1 \leq i \leq n-1, & [y_j, y_1] &= x_j, \quad 1 \leq j \leq n, \\ [y_{n+1}, y_{n+1}] &= \gamma x_n, & [x_i, y_{n+1}] &= \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^{n+1-i} \beta_k y_{k-1+i}, \quad 1 \leq i \leq \lfloor \frac{n-1}{2} \rfloor, \\ [y_1, y_{n+1}] &= -2 \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta_k x_{k-1} + \beta x_n, & [y_j, y_{n+1}] &= -2 \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^{n+2-j} \beta_k x_{k-2+j}, \quad 2 \leq j \leq \lfloor \frac{n+1}{2} \rfloor. \end{aligned}$$

Proof. From Lemma 3.1 and Theorem 2.1 we have that there exists a basis $\{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_{n+1}\}$ in which products have the following form:

$$\begin{aligned} [x_i, x_1] &= x_{i+1}, \quad 1 \leq i \leq n-1, & [y_j, x_1] &= y_{j+1}, \quad 1 \leq j \leq n-1, \\ [x_i, y_1] &= \frac{1}{2}y_{i+1}, \quad 1 \leq i \leq n-1, & [y_j, y_1] &= x_j, \quad 1 \leq j \leq n, \\ [y_1, y_{n+1}] &= \sum_{k=1}^n \alpha_k x_k, & [x_1, y_{n+1}] &= \sum_{s=2}^n \beta_s y_s. \end{aligned}$$

From equalities:

$$\begin{aligned} [y_{n+1}, [y_1, x_1]] &= [[y_{n+1}, y_1], x_1] - [[y_{n+1}, x_1], y_1], \\ [y_{n+1}, [x_1, y_1]] &= [[y_{n+1}, x_1], y_1] - [[y_{n+1}, y_1], x_1], \\ [y_{n+1}, y_{i+1}] &= [y_{n+1}, [y_i, x_1]] = [[y_{n+1}, y_i], x_1] - [[y_{n+1}, x_1], y_i], \quad 1 \leq i \leq n-1, \end{aligned}$$

we have $[y_{n+1}, y_2] = [[y_{n+1}, y_1], x_1]$ and $[y_{n+1}, y_2] = -2[[y_{n+1}, y_1], x_1]$, it follows that

$$[y_{n+1}, y_1] = b x_n \text{ and } [y_{n+1}, y_i] = 0 \text{ for } 2 \leq i \leq n.$$

Consider the equality:

$$[y_{n+1}, [y_{n+1}, x_1]] = [[y_{n+1}, y_{n+1}], x_1] + [[y_{n+1}, x_1], y_{n+1}],$$

then $[y_{n+1}, y_{n+1}] = \gamma x_n$.

Considering the graded identities:

$$[y_j, [y_{n+1}, x_1]] = [[y_j, y_{n+1}], x_1] - [[y_j, x_1], y_{n+1}],$$

$$[x_i, [y_{n+1}, x_1]] = [[x_i, y_{n+1}], x_1] - [[x_i, x_1], y_{n+1}],$$

we have

$$[y_j, y_{n+1}] = \sum_{k=1}^{n+1-j} \alpha_k x_{k-1+j}, \quad 2 \leq j \leq n,$$

$$[x_i, y_{n+1}] = \sum_{s=2}^{n+1-i} \beta_s y_{s-1+i}, \quad 2 \leq i \leq n-1.$$

From the chain of equalities

$$0 = [y_1, [y_{n+1}, y_1]] = [[y_1, y_{n+1}], y_1] + [[y_1, y_1], y_{n+1}] =$$

$$= \left[\sum_{k=1}^n \alpha_k x_k, y_1 \right] + [x_1, y_{n+1}] = \frac{1}{2} \sum_{k=1}^{n-1} \alpha_k y_{k+1} + \sum_{s=2}^n \beta_s y_s,$$

we obtain $\alpha_i = -2\beta_{i+1}$, $1 \leq i \leq n-1$.

Substituting these relations in the chain of equalities

$$0 = [y_1, [y_{n+1}, y_{n+1}]] = 2[[y_1, y_{n+1}], y_{n+1}],$$

we get $\beta_i = 0$, $2 \leq i \leq \lfloor \frac{n+2}{2} \rfloor$. Thus, we obtain the multiplications of the superalgebra as in the assertion of the lemma. \square

For convenience, we will denote the superalgebra from the family of Lemma 3.2 as $L\left(\gamma, \beta_{\lfloor \frac{n+4}{2} \rfloor}, \beta_{\lfloor \frac{n+4}{2} \rfloor + 1}, \dots, \beta_n, \beta\right)$.

Using the properties of adapted basis we obtain necessary and sufficient conditions when two arbitrary superalgebras from the family of Lemma 3.2 are isomorphic.

Theorem 3.2. *Two superalgebras $L\left(\gamma, \beta_{\lfloor \frac{n+4}{2} \rfloor}, \beta_{\lfloor \frac{n+4}{2} \rfloor + 1}, \dots, \beta_n, \beta\right)$ and*

$L'\left(\gamma', \beta'_{\lfloor \frac{n+4}{2} \rfloor}, \beta'_{\lfloor \frac{n+4}{2} \rfloor + 1}, \dots, \beta'_n, \beta'\right)$ are isomorphic if and only if there exist $a_1, a_{n+1}, b_{n+1} \in$

\mathbb{C} such that the following conditions hold:

for odd n :

$$\begin{cases} b_{n+1}^2 \gamma = \gamma' a_1^{2n}, \\ b_{n+1} \beta_j = a_1^{2j-3} \beta'_j, \quad \lfloor \frac{n+4}{2} \rfloor \leq j \leq n, \\ a_{n+1} b_{n+1} \gamma + a_1 b_{n+1} \beta = a_1^{2n} \beta' + 4\beta'_{\lfloor \frac{n+4}{2} \rfloor} a_1^{2\lfloor \frac{n+1}{2} \rfloor - 1} a_{n+1} \beta_{\lfloor \frac{n+4}{2} \rfloor}, \end{cases}$$

for even n :

$$\begin{cases} b_{n+1}^2 \gamma = \gamma' a_1^{2n}, \\ b_{n+1} \beta_j = a_1^{2j-3} \beta'_j, \quad \lfloor \frac{n+4}{2} \rfloor \leq j \leq n, \\ a_{n+1} b_{n+1} \gamma + a_1 b_{n+1} \beta = a_1^{2n} \beta'. \end{cases}$$

Proof. Let us make a general change of generator elements in the form:

$$y'_1 = \sum_{i=1}^{n+1} a_i y_i, \quad y'_{n+1} = \sum_{j=1}^{n+1} b_j y_j,$$

where the rank $\begin{pmatrix} a_1 & a_2 & \cdots & a_{n+1} \\ b_1 & b_2 & \cdots & b_{n+1} \end{pmatrix} = 2$.

We express the new basis $\{x'_1, x'_2, \dots, x'_n, y'_1, y'_2, \dots, y'_{n+1}\}$ of the superalgebra $L' \left(\gamma', \beta'_{\lfloor \frac{n+4}{2} \rfloor}, \beta'_{\lfloor \frac{n+4}{2} \rfloor + 1}, \dots, \beta'_n, \beta' \right)$ with respect to the old basis $\{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_{n+1}\}$.

Then for element x'_1 we have

$$\begin{aligned} x'_1 = [y'_1, y'_1] &= a_1 \sum_{i=1}^n a_i x_i - 2a_1 a_{n+1} \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta_k x_{k-1} + \\ &+ a_1 a_{n+1} \beta x_n - 2a_{n+1} \sum_{i=2}^{\lfloor \frac{n+1}{2} \rfloor} a_i \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^{n+2-i} \beta_k x_{k+i-2} + a_{n+1}^2 \gamma x_n. \end{aligned}$$

The expression of x'_{t+1} ($1 \leq t \leq \lfloor \frac{n-1}{2} \rfloor$) will be as follows:

$$x'_{t+1} = [x'_t, x'_1] = a_1^{2t+1} \sum_{i=1}^{n-t} a_i x_{i+t} - 2a_1^{2t} a_{n+1} \sum_{i=1}^{\lfloor \frac{n+1}{2} \rfloor - t} a_i \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^{n+2-t-i} \beta_k x_{k+t+i-2}.$$

And for x'_{t+1} ($\lfloor \frac{n+1}{2} \rfloor \leq t \leq n-1$) we have

$$x'_{t+1} = [x'_t, x'_1] = a_1^{2t+1} \sum_{i=1}^{n-t} a_i x_{t+i}.$$

For basis elements y'_i of the space L'_1 basis we have:

$$y'_t = [y'_{t-1}, x'_1] = a_1^{2(t-1)} \sum_{j=1}^{n+1-t} a_j y_{t+j-1}, \quad 2 \leq t \leq n.$$

Consider the equalities:

$$\begin{aligned} [y'_{n+1}, y'_{n+1}] &= b_1 \sum_{i=1}^n b_i x_i - 2b_1 b_{n+1} \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta_k x_{k-1} + b_1 b_{n+1} \beta x_n - \\ &- 2b_{n+1} \sum_{i=2}^{\lfloor \frac{n+1}{2} \rfloor} b_i \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^{n+2-i} \beta_k x_{k+i-2} + b_{n+1}^2 \gamma x_n = \gamma' x'_n = \gamma' a_1^{2n} x_n. \end{aligned}$$

Comparing the coefficients we obtain the following restrictions:

$$b_1 = b_2 = \dots = b_{\lfloor \frac{n-1}{2} \rfloor} = 0, \quad (3.4)$$

$$-2b_{n+1}b_{\lfloor \frac{n+1}{2} \rfloor}\beta_{\lfloor \frac{n+4}{2} \rfloor} + b_{n+1}^2\gamma = \gamma'a_1^{2n}. \quad (3.5)$$

Using the chain of the equalities:

$$[y'_{n+1}, y'_1] = a_1 \sum_{i=\lfloor \frac{n+1}{2} \rfloor}^n b_i x_i + a_{n+1}b_{n+1}\gamma x_n - 2a_{n+1}b_{\lfloor \frac{n+1}{2} \rfloor}\beta_{\lfloor \frac{n+4}{2} \rfloor}x_n = 0,$$

the equalities (3.4) - (3.5) and comparing the coefficients to the basis elements we obtain:

$$\begin{cases} b_1 = b_2 = b_{\lfloor \frac{n-1}{2} \rfloor} = b_{\lfloor \frac{n+1}{2} \rfloor} = \dots = b_{n-1} = 0, \\ b_{n+1}^2\gamma = \gamma'a_1^{2n}, \\ a_1b_n + a_{n+1}b_{n+1}\gamma = 0. \end{cases} \quad (3.6)$$

Therefore, $y'_{n+1} = b_n y_n + b_{n+1} y_{n+1}$.

Consider the products:

$$\begin{aligned} [x'_1, y'_{n+1}] &= \left[a_1 \sum_{i=1}^n a_i x_i - 2a_1 a_{n+1} \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta_k x_{k-1} + a_1 a_{n+1} \beta x_n - \right. \\ &\quad \left. - 2a_{n+1} \sum_{i=2}^{\lfloor \frac{n+1}{2} \rfloor} a_i \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^{n+2-i} \beta_k x_{k+i-2} + a_{n+1}^2 \gamma x_n, b_n y_n + b_{n+1} y_{n+1} \right] = \\ &= a_1 b_{n+1} \sum_{i=1}^{\lfloor \frac{n-1}{2} \rfloor} a_i \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^{n+1-i} \beta_k y_{k+i-1}, \\ [x'_1, y'_{n+1}] &= \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta'_k y'_k = \beta'_{\lfloor \frac{n+4}{2} \rfloor} a_1^{2\lfloor \frac{n+4}{2} \rfloor - 2} \left(a_1 y_{\lfloor \frac{n+4}{2} \rfloor} + a_2 y_{\lfloor \frac{n+4}{2} \rfloor + 1} + \dots + a_{n+1 - \lfloor \frac{n+4}{2} \rfloor} y_n \right) + \\ &\quad + \beta'_{\lfloor \frac{n+4}{2} \rfloor + 1} a_1^{2\lfloor \frac{n+4}{2} \rfloor} \left(a_1 y_{\lfloor \frac{n+4}{2} \rfloor + 1} + a_2 y_{\lfloor \frac{n+4}{2} \rfloor + 2} + \dots + a_{n - \lfloor \frac{n+4}{2} \rfloor} y_n \right) + \dots + \\ &\quad + \beta'_n a_1^{2n-1} y_n = \sum_{\lfloor \frac{n+4}{2} \rfloor}^n \beta'_k a_1^{2(k-1)} \sum_{j=1}^{n+1-k} a_j y_{k+j-1}. \end{aligned}$$

From which, comparing the coefficients we have the following restrictions:

$$b_{n+1}\beta_j = a_1^{2j-3}\beta'_j, \quad \left\lfloor \frac{n+4}{2} \right\rfloor \leq j \leq n.$$

Consider the following product on the one hand:

$$\begin{aligned} [y'_1, y'_{n+1}] &= [a_1 y_1 + a_2 y_2 + \dots + a_{n+1} y_{n+1}, b_n y_n + b_{n+1} y_{n+1}] = \\ &= -2a_1 b_{n+1} \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta_k x_{k-1} + a_1 b_{n+1} \beta x_n - 2b_{n+1} \sum_{i=2}^{\lfloor \frac{n+1}{2} \rfloor} a_i \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^{n+2-i} \beta_k x_{k+i-2} + a_{n+1} b_{n+1} \gamma x_n \end{aligned}$$

and on the other hand, let us consider the followings product in the case of an odd n .

$$\begin{aligned} [y'_1, y'_{n+1}] &= -2 \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta'_k x'_{k-1} + \beta' x'_n = -2\beta'_{\lfloor \frac{n+4}{2} \rfloor} \left(a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 1)} x_{\lfloor \frac{n+4}{2} \rfloor - 1} + \right. \\ &+ a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 2) + 1} a_2 x_{\lfloor \frac{n+4}{2} \rfloor} + \dots + a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 2) + 1} a_{n - \lfloor \frac{n+4}{2} \rfloor + 2} x_n - 2a_1^{2\lfloor \frac{n-1}{2} \rfloor + 1} a_{n+1} \beta_{\lfloor \frac{n+4}{2} \rfloor} x_n \left. \right) - \\ &- 2\beta'_{\lfloor \frac{n+4}{2} \rfloor + 1} \left(a_1^{2\lfloor \frac{n+4}{2} \rfloor} x_{\lfloor \frac{n+4}{2} \rfloor} + a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 1) + 1} a_2 x_{\lfloor \frac{n+4}{2} \rfloor + 1} + \dots + a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 1) + 1} a_{n - \lfloor \frac{n+4}{2} \rfloor + 1} x_n \right) - \\ &- \dots - 2\beta'_n (a_1^{2n-2} x_{n-1} + a_1^{2n-3} a_2 x_n) + \beta' a_1^{2n} x_n = -2 \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta'_k a_1^{2k-3} \sum_{i=1}^{n-k+2} a_i x_{k+i-2} + \\ &+ 4\beta'_{\lfloor \frac{n+4}{2} \rfloor} a_1^{2\lfloor \frac{n-1}{2} \rfloor + 1} a_{n+1} \beta_{\lfloor \frac{n+4}{2} \rfloor} x_n + \beta' a_1^{2n} x_n. \end{aligned}$$

In the case of an even n , for the product $[y'_1, y'_{n+1}]$ we have:

$$\begin{aligned} [y'_1, y'_{n+1}] &= -2 \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta'_k x'_{k-1} + \beta' x'_n = -2\beta'_{\lfloor \frac{n+4}{2} \rfloor} \left(a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 1)} x_{\lfloor \frac{n+4}{2} \rfloor - 1} + \right. \\ &+ a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 2) + 1} a_2 x_{\lfloor \frac{n+4}{2} \rfloor} + \dots + a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 2) + 1} a_{n - \lfloor \frac{n+4}{2} \rfloor + 1} x_n \left. \right) - 2\beta'_{\lfloor \frac{n+4}{2} \rfloor + 1} \left(a_1^{2\lfloor \frac{n+4}{2} \rfloor} x_{\lfloor \frac{n+4}{2} \rfloor} + \right. \\ &+ a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 1) + 1} a_2 x_{\lfloor \frac{n+4}{2} \rfloor + 1} + \dots + a_1^{2(\lfloor \frac{n+4}{2} \rfloor - 1) + 1} a_{n - \lfloor \frac{n+4}{2} \rfloor + 1} x_n \left. \right) - \dots - 2\beta'_n (a_1^{2n-2} x_{n-1} + \\ &+ a_1^{2n-3} x_n) + \beta' a_1^{2n} x_n = -2 \sum_{k=\lfloor \frac{n+4}{2} \rfloor}^n \beta'_k a_1^{2k-3} \sum_{i=1}^{n-k+2} a_i x_{k+i-2} + \beta' a_1^{2n} x_n. \end{aligned}$$

Comparing the coefficients we obtain the following restrictions:

when n is odd

$$\begin{cases} b_{n+1} \beta_j = a_1^{2j-3} \beta'_j, & \lfloor \frac{n+4}{2} \rfloor \leq j \leq n, \\ a_{n+1} b_{n+1} \gamma + a_1 b_{n+1} \beta = a_1^{2n} \beta' + 4\beta'_{\lfloor \frac{n+4}{2} \rfloor} a_1^{2\lfloor \frac{n+1}{2} \rfloor - 1} a_{n+1} \beta_{\lfloor \frac{n+4}{2} \rfloor} \end{cases} \quad (3.7)$$

when n is even

$$\begin{cases} b_{n+1}\beta_j = a_1^{2j-3}\beta'_j, & \lfloor \frac{n+4}{2} \rfloor \leq j \leq n, \\ a_{n+1}b_{n+1}\gamma + a_1b_{n+1}\beta = a_1^{2n}\beta'. \end{cases} \quad (3.8)$$

It is not difficult to check that considering other multiplications we have either restrictions (3.7)-(3.8) or identity.

Note that from (3.6) we have $b_n = \frac{-a_{n+1}b_{n+1}\gamma}{a_1}$.

Thus, combining the restrictions (3.6), (3.7) and (3.8) it follows the proof of the theorem. \square

Introduce the operators which are similar like k -dimensional vectors:

$$V_{j,k}^0(\alpha_1, \alpha_2, \dots, \alpha_k) = (0, \dots, 0, 1, \delta \sqrt{\delta^{j+1}} S_{m,j}^{j+1} \alpha_{j+1}, \delta \sqrt{\delta^{j+2}} S_{m,j}^{j+2} \alpha_{j+2}, \dots, \delta \sqrt{\delta^k} S_{m,j}^k \alpha_k);$$

$$V_{j,k}^1(\alpha_1, \alpha_2, \dots, \alpha_k) = (0, \dots, 0, 1, S_{m,j}^{j+1} \alpha_{j+1}, S_{m,j}^{j+2} \alpha_{j+2}, \dots, S_{m,j}^k \alpha_k);$$

$$V_{j,k}^2(\alpha_1, \alpha_2, \dots, \alpha_k) = (0, \dots, 0, 1, S_{m,2j+1}^{2(j+1)+1} \alpha_{j+1}, S_{m,2j+1}^{2(j+2)+1} \alpha_{j+2}, \dots, S_{m,2j+1}^{2k+1} \alpha_k);$$

$$V_{k+1,k}^0(\alpha_1, \alpha_2, \dots, \alpha_k) = V_{k+1,k}^1(\alpha_1, \alpha_2, \dots, \alpha_k) = V_{k+1,k}^2(\alpha_1, \alpha_2, \dots, \alpha_k) = (0, 0, \dots, 0);$$

$$\begin{aligned} W_{s,k}(0, \dots, 0, 1, S_{m,j}^{j+1} \alpha_{j+1}, S_{m,j}^{j+2} \alpha_{j+2}, \dots, S_{m,j}^k \alpha_k) = \\ = (0, \dots, 0, 1, 0, \dots, 0, 1, S_{m,j}^{j+1} \alpha_{s+j+1}, S_{m,j}^{j+2} \alpha_{s+j+2}, \dots, S_{m,j}^{k-s} \alpha_k), \end{aligned}$$

$$W_{k+1-j,k}(0, \dots, 0, 1, 0, \dots, 0) = (0, \dots, 0, 1, 0, \dots, 0)$$

where $k \in \mathbb{C}$, $\delta = \pm 1$, $1 \leq j \leq k$, $1 \leq s \leq k - j$, $S_{m,t} = \cos \frac{2\pi m}{t} + i \sin \frac{2\pi m}{t}$ ($m = 0, 1, \dots, t - 1$).

Theorem 3.2 allows us to classify the Leibniz superalgebras from the variety $Leib^{n,m}$ with characteristic sequence $(n \mid m - 1, 1)$, nilindex $n + m$ and $m = n + 1$.

Theorem 3.3. *Let L be a Leibniz superalgebra of variety $Leib^{n,m}$ with characteristic sequence $(n \mid m - 1, 1)$, nilindex $n + m$ and $m = n + 1$. Then L is isomorphic to one of the following pairwise non isomorphic superalgebras:*

if n is odd (i.e. $n = 2q - 1$):

$$L(1, \delta \beta_{q+1}, V_{j,q-2}^0(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n), 0), \quad \beta_{q+1} \neq \pm \frac{1}{2}, \quad 1 \leq j \leq q - 1,$$

$$L(1, \beta_{q+1}, V_{j,q-1}^0(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n, \beta)), \quad \beta_{q+1} = \pm \frac{1}{2}, \quad 1 \leq j \leq q,$$

$$L(0, 1, V_{j,q-2}^0(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n), 0), \quad 1 \leq j \leq q - 1,$$

$$L(0, 0, W_{s,q-1}(V_{j,q-1}^1(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n, \beta))), \quad 1 \leq j \leq q - 1, \quad 1 \leq s \leq q - j,$$

$$L(0, 0, \dots, 0);$$

if n is even (i.e. $n = 2q$):

$$\begin{aligned} &L(1, V_{j,q-1}^2(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n), 0), \quad 1 \leq j \leq q, \\ &L(0, W_{s,q}(V_{j,q}^1(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n, \beta))), \quad 1 \leq j \leq q, \quad 1 \leq s \leq q+1-j, \\ &L(0, 0, \dots, 0). \end{aligned}$$

Proof. Consider n is odd, i.e. $n = 2q - 1$, where $q \in \mathbb{N}$. From Theorem 3.2 we have the following restrictions:

$$\begin{cases} b_{n+1}^2 \gamma = \gamma' a_1^{2n}, \\ b_{n+1} \beta_j = a_1^{2j-3} \beta'_j, \quad q+1 \leq j \leq n, \\ a_{n+1} b_{n+1} \gamma + a_1 b_{n+1} \beta = a_1^{2n} \beta' + 4\beta'_{q+1} a_1^n a_{n+1} \beta_{q+1}, \end{cases}$$

for which we consider all possible cases.

Case 1. Let $\gamma \neq 0$. Then taking $b_{n+1} = \pm \frac{a_1^n}{\sqrt{\gamma}}$, we obtain $\gamma' = 1$. Substituting the value of b_{n+1} in other restrictions we obtain equalities:

$$\begin{aligned} \beta'_{q+1+j} &= \pm \frac{\beta_{q+1+j}}{a_1^{2j} \sqrt{\gamma}}, \quad 0 \leq j \leq q-2, \\ \beta' &= \pm \frac{a_{n+1}(\gamma - 4\beta_{q+1}^2) + a_1 \beta}{a_1^n \sqrt{\gamma}}. \end{aligned}$$

Case 1.1. If $\gamma - 4\beta_{q+1}^2 \neq 0$, then putting $a_{n+1} = -\frac{a_1 \beta}{\gamma - 4\beta_{q+1}^2}$, we have $\beta' = 0$ and

$$\beta'_{q+1+j} = \pm \frac{\beta_{q+1+j}}{\sqrt{\gamma}} a_1^{-2j} \text{ for } 0 \leq j \leq q-2.$$

If $\beta_{q+1+j} = 0$ for any $j \in \{1, \dots, q-2\}$, then $\beta'_{q+1+j} = 0$ and we obtain the superalgebras:

$$L(1, \delta \beta_{q+1}, 0, \dots, 0), \quad \delta = \pm 1.$$

If $\beta_{q+2} = \beta_{q+3} = \dots = \beta_{q+t} = 0$ and $\beta_{q+t+1} \neq 0$ for some $t \in \{1, 2, \dots, q-2\}$. Then taking $a_1^{-2t} = \pm \frac{\sqrt{\gamma}}{\beta_{q+t+1}}$ (i.e. $a_1^{-2} = \sqrt{\pm 1}^t \sqrt{\left| \frac{\sqrt{\gamma}}{\beta_{q+t+1}} \right|} (\cos \frac{\varphi}{t} + i \sin \frac{\varphi}{t}) (\cos \frac{2\pi m}{t} + i \sin \frac{2\pi m}{t})$, where $\varphi = \arg \left(\frac{\sqrt{\gamma}}{\beta_{q+t+1}} \right)$, $m = 0, 1, \dots, t-1$), we obtain:

$$\beta'_{q+t+1} = 1 \text{ and } \beta'_{q+t+j} = \pm \sqrt{\pm 1}^j S_{m,t}^j \beta_{q+t+j}, \quad m = 0, 1, \dots, t-1.$$

So, in this case we have the following superalgebras:

$$L(1, \delta \beta_{q+1}, V_{j,q-2}^0(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n), 0), \quad \beta_{q+1} \neq \pm \frac{1}{2}, \quad 1 \leq j \leq q-1, \quad \delta = \pm 1.$$

Case 1.2. If $\gamma - 4\beta_{q+1}^2 = 0$, then $\beta'_{q+1} = \pm \frac{1}{2}$ and we have

$$\beta'_{q+1+j} = \pm \frac{\beta_{q+1+j}}{\sqrt{\gamma}} a_1^{-2j}, \quad 1 \leq j \leq q-2, \quad (3.9)$$

$$\beta' = \pm \frac{\beta}{\sqrt{\gamma}} a_1^{-n+1}. \quad (3.10)$$

If we assume that $j = q - 1$ in restriction (3.9), then we obtain $\beta'_{2q} = \pm \frac{\beta_{2q}}{\sqrt{\gamma}} a_1^{-2(q-1)}$. Since $n = 2q - 1$, then $-2(q - 1) = -n + 1$, i.e. we formally have restriction (3.10) and therefore restriction (3.10) can be considered as a particular case of restriction (3.9) when $j = q - 1$.

Furthermore, as in case 1.1, we obtain the following superalgebras:

$$L(1, \beta_{q+1}, V_{j,q-1}^0(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n, \beta)), \quad \beta_{q+1} = \pm \frac{1}{2}, \quad 1 \leq j \leq q.$$

Case 2. $\gamma = 0$. Then $\gamma' = 0$ and

$$b_{n+1}\beta_{q+1+j} = a_1^{2q-1+2j}\beta'_{q+1+j}, \quad 0 \leq j \leq q - 2,$$

$$a_1 b_{n+1} \beta = a_1^{2n} \beta' + 4\beta'_{q+1} a_1^n a_{n+1} \beta_{q+1}.$$

Case 2.1. $\beta_{q+1} \neq 0$. Then taking $b_{n+1} = \frac{a_1^n}{\beta_{q+1}}$ and $a_{n+1} = \frac{b_{n+1}\beta}{4a_1^{n-1}\beta_{q+1}}$, we have

$$\beta'_{q+1} = 0, \beta' = 0 \text{ and } \beta'_{q+1+j} = \frac{\beta_{q+1+j}}{\beta_{q+1}} a_1^{-2j}, \quad 1 \leq j \leq q - 2.$$

Furthermore, as in case 1.1, we obtain the superalgebras:

$$L(0, 1, V_{j,q-2}^1(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n), 0), \quad 1 \leq j \leq q - 1.$$

Case 2.2. $\beta_{q+1} = 0$. Then $\beta'_{q+1} = 0$ and

$$b_{n+1}\beta_{q+1+j} = a_1^{2q-1+2j}\beta'_{q+1+j}, \quad 1 \leq j \leq q - 2, \quad (3.11)$$

$$b_{n+1}\beta = a_1^{2n-1}\beta'. \quad (3.12)$$

If we assume that $j = q - 1$, in restriction (3.11), then we obtain $b_{n+1}\beta_{2q} = a_1^{2n-1}\beta'_{2q}$. Since $n = 2q - 1$, then $2q - 1 + 2(q - 1) = 4q - 3 = 2n - 1$, and then we formally obtain restriction (3.12) and, therefore restriction (3.12) can be considered as a particular case of (3.11) when $j = q - 1$.

Case 2.2.1 $\beta_{q+2} = \beta_{q+3} = \dots = \beta_{q+t} = 0$ and $\beta_{q+t+1} \neq 0$ for some $t \in \{1, 2, \dots, q - 1\}$. Then if we choose $b_{n+1} = \frac{a_1^{2q-1+2t}}{\beta_{q+1+t}}$, we obtain $\beta'_{q+1+t} = 1$ and

$$\beta'_{q+1+j} = \frac{\beta_{q+1+j}}{\beta_{q+1+t}} a_1^{-2(j-t)}, \quad t + 1 \leq j \leq q - 1.$$

Thus, in this case we have the superalgebras:

$$L(0, 0, W_{s,q-1}(V_{j,q-1}^1(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n, \beta))), \quad 1 \leq j \leq q - 1, \quad 1 \leq s \leq q - j.$$

Case 2.2.2 $\beta_{q+j+1} = 0$ for any j ($1 \leq j \leq q - 1$). Then we obtain superalgebra:

$$L(0, 0, \dots, 0).$$

Consider the case of even n , i.e. $n = 2q$ for some $q \in \mathbb{N}$.

Then from Theorem 3.2 we have the following restrictions:

$$\begin{cases} b_{n+1}^2 \gamma = \gamma' a_1^{2n}, \\ b_{n+1} \beta_{q+2+j} = a_1^{2q+2j+1} \beta'_{q+2+j}, \quad 0 \leq j \leq q-2, \\ a_{n+1} b_{n+1} \gamma + a_1 b_{n+1} \beta = a_1^{2n} \beta'. \end{cases}$$

Case 1. $\gamma \neq 0$. Then taking $b_{n+1} = \pm \frac{a_1^n}{\sqrt{\gamma}}$ and $a_{n+1} = -\frac{a_1 \beta}{\gamma}$ we obtain $\gamma' = 1$;

$$\beta'_{q+2+j} = \pm \frac{\beta_{q+2+j}}{a_1^{2j+1} \sqrt{\gamma}} \quad (0 \leq j \leq q-2) \text{ and } \beta' = 0.$$

If $\beta_{q+2+j} = 0$ for any j ($0 \leq j \leq q-2$), then we have superalgebra:

$$L(1, 0, \dots, 0).$$

If $\beta_{q+2} = \beta_{q+3} = \dots = \beta_{q+t+1} = 0$ and $\beta_{q+t+2} \neq 0$ for some t ($1 \leq t \leq q-2$), then putting $a_1^{-(2t+1)} = \pm \frac{\sqrt{\gamma}}{\beta_{q+2+t}}$ (i.e. $a_1^{-1} = \pm \sqrt[2t+1]{\left| \frac{\sqrt{\gamma}}{\beta_{q+2+t}} \right|} (\cos \frac{\varphi}{2t+1} + i \sin \frac{\varphi}{2t+1}) (\cos \frac{2\pi m}{2t+1} + i \sin \frac{2\pi m}{2t+1})$, where $\varphi = \arg \left(\frac{\sqrt{\gamma}}{\beta_{q+2+t}} \right)$, $m = 0, 1, \dots, 2t$), and substituting the value of a_1^{-1} in other restrictions we obtain

$$\begin{aligned} \beta'_{q+2+j} &= \pm \frac{\beta_{q+2+j}}{\sqrt{\gamma}} \left(\pm \sqrt[2t+1]{\left| \frac{\sqrt{\gamma}}{\beta_{q+2+j}} \right|} \left(\cos \frac{\varphi}{2t+1} + i \sin \frac{\varphi}{2t+1} \right) S_{m, 2j+1} \right)^{2j+1} = \\ &= \beta_{q+2+j} S_{m, 2j+1}^{2j+1}, \quad t+1 \leq j \leq q-2. \end{aligned}$$

Thus, in this case we have the following superalgebras:

$$L(1, V_{j, q-1}^2(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n), 0), \quad 1 \leq j \leq q.$$

Case 2. $\gamma = 0$. Then $\gamma' = 0$ and $\beta'_{j+2+q} = \frac{b_{n+1} \beta_{q+2+j}}{a_1^{2q+1+2j}}$, $0 \leq j \leq q-2$, $\beta' = \frac{b_{n+1} \beta}{a_1^{2n-1}}$.

Note that in this case β' also can be considered as a particular case of β'_{j+2+q} for $j = q-1$.

If $\beta_{q+2+j} = 0$ for any j ($0 \leq j \leq q-1$), then we have the superalgebra:

$$L(0, 0, 0, \dots, 0).$$

If $\beta_{q+2} = \beta_{q+3} = \dots = \beta_{q+t+1} = 0$ and $\beta_{q+t+2} \neq 0$ for some t ($1 \leq t \leq q-1$). Then putting $b_{n+1} = \frac{a_1^{2q+2t+1}}{\beta_{q+2+t}}$, we obtain:

$$\beta'_{q+2+t} = 1, \quad \beta'_{q+2+j} = \frac{\beta_{q+2+j}}{\beta_{q+2+t}} a_1^{-2(j-t)} \quad (t+1 \leq j \leq q-1).$$

As in case 2.2.1 for odd n , we obtain superalgebras:

$$L(0, W_{s, q}(V_{j, q}^1(\beta_{q+2}, \beta_{q+3}, \dots, \beta_n, \beta))), \quad 1 \leq j \leq q, \quad 1 \leq s \leq q+1-j.$$

□

The existence of an adapted basis under the conditions of Lemma 3.1 for $m = n + 2$ is represented in the following lemma.

Lemma 3.3. *Let L be a Leibniz superalgebra of variety $\text{Leib}^{n,m}$ with characteristic sequence $(n \mid m - 1, 1)$, nilindex $n + m$ and $m = n + 2$. Then, there exists a basis $\{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_{n+2}\}$ of L in which the multiplication has the following form:*

$$\begin{aligned} [x_i, x_1] &= x_{i+1}, \quad 1 \leq i \leq n-1, & [y_j, x_1] &= y_{j+1}, \quad 1 \leq j \leq n, \\ [x_i, y_1] &= \frac{1}{2}y_{i+1}, \quad 1 \leq i \leq n, & [y_j, y_1] &= x_j, \quad 1 \leq j \leq n, \\ [x_i, y_{n+2}] &= \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+2-i} \beta_k y_{k-1+i}, \quad 1 \leq i \leq \lfloor \frac{n}{2} \rfloor, & [y_j, y_{n+2}] &= -2 \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+2-j} \beta_k x_{k-2+j}, \quad 1 \leq j \leq \lfloor \frac{n}{2} \rfloor. \end{aligned}$$

Proof. The proof of this lemma is analogous to the proof of Lemma 3.2. \square

Let us denote the superalgebra from the family of Lemma 3.3 by $L(\beta_{\lfloor \frac{n+5}{2} \rfloor}, \beta_{\lfloor \frac{n+5}{2} \rfloor + 1}, \dots, \beta_{n+1})$.

The condition of isomorphism of two superalgebras is represented in the following theorem.

Theorem 3.4. *Two superalgebras $L(\beta_{\lfloor \frac{n+5}{2} \rfloor}, \beta_{\lfloor \frac{n+5}{2} \rfloor + 1}, \dots, \beta_{n+1})$ and $L'(\beta'_{\lfloor \frac{n+5}{2} \rfloor}, \beta'_{\lfloor \frac{n+5}{2} \rfloor + 1}, \dots, \beta'_{n+1})$ are isomorphic if and only if there exist $a_1, b_{n+2} \in \mathbb{C}$ such that the following conditions hold:*

$$b_{n+2}\beta_j = a_1^{2j-3}\beta'_j, \quad \lfloor \frac{n+5}{2} \rfloor \leq j \leq n+1.$$

Proof. By a change of basis the generators of the new basis are expressed by

$$y'_1 = \sum_{i=1}^{n+2} a_i y_i, \quad y'_{n+1} = \sum_{j=1}^{n+2} b_j y_j,$$

where the rank $\begin{pmatrix} a_1 & a_2 & \dots & a_{n+2} \\ b_1 & b_2 & \dots & b_{n+2} \end{pmatrix} = 2$, this allows us to express the elements of the new basis $\{x'_1, x'_2, \dots, x'_n, y'_1, y'_2, \dots, y'_{n+2}\}$ of the superalgebra $L'(\beta'_{\lfloor \frac{n+5}{2} \rfloor}, \beta'_{\lfloor \frac{n+5}{2} \rfloor + 1}, \dots, \beta'_{n+1})$ with respect to the elements of old basis $\{x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_{n+2}\}$ as:

$$\begin{aligned} x'_1 &= [y'_1, y'_1] = a_1 \sum_{k=1}^n a_k x_k - 2a_{n+2} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} a_i \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+2-i} \beta_k x_{k-2+i}; \\ x'_{t+1} &= [x'_t, x'_1] = a_1^{2t+1} \sum_{k=1}^{n-t} a_k x_{t+k} - 2a_1^{2t} a_{n+2} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor - t} a_i \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+2-t-i} \beta_k x_{k+t-2+i}, \quad 1 \leq t \leq \lfloor \frac{n-2}{2} \rfloor; \end{aligned}$$

$$x'_{t+1} = [x'_t, x'_1] = a_1^{2t+1} \sum_{k=1}^{n-t} a_k x_{t+k}, \quad \left\lfloor \frac{n}{2} \right\rfloor \leq t \leq n-1;$$

$$y'_t = [y'_{t-1}, x'_1] = a_1^{2(t-1)} \sum_{i=1}^{n+2-t} a_i y_{t-1+i}, \quad 2 \leq t \leq n+1.$$

If we consider the products

$$[x'_i, x'_1] = \frac{1}{2} y'_{i+1}, \quad 1 \leq i \leq n, \quad [y'_t, y'_1] = x'_t, \quad 1 \leq t \leq n$$

we find no restrictions.

Consider the chain of the equalities:

$$[y'_{n+2}, y'_{n+2}] = b_1 \sum_{k=1}^n b_k x_k - 2b_{n+2} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} b_i \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+2-i} \beta_k x_{k-2+i} = 0.$$

Comparing the coefficients of the basis elements in the last equality we obtain the following restrictions:

$$b_i = 0, \quad 1 \leq i \leq \left\lfloor \frac{n}{2} \right\rfloor. \quad (3.13)$$

From the following equalities we obtain:

$$[y'_{n+2}, y'_1] = \left[\sum_{i=\lfloor \frac{n}{2} \rfloor + 1}^{n+2} b_i y_i, \sum_{j=1}^{n+2} a_j y_j \right] = a_1 \sum_{k=\lfloor \frac{n}{2} \rfloor + 1}^n b_k x_k = 0$$

restrictions and summing them with (3.13) we obtain $b_i = 0, 1 \leq i \leq n$.

Therefore we have $y'_{n+2} = b_{n+1} y_{n+1} + b_{n+2} y_{n+2}$.

Consider the multiplications defining the parameters:

$$[x'_1, y'_{n+2}] = \left[a_1 \sum_{k=1}^n a_k x_k - 2a_{n+2} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} a_i \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+2-i} \beta_k x_{k-2+i}, b_{n+1} y_{n+1} + b_{n+2} y_{n+2} \right] =$$

$$= a_1 b_{n+2} \sum_{j=1}^{\lfloor \frac{n-2}{2} \rfloor + 1} a_j \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+2-j} \beta_k y_{k+j-1}; \quad (3.14)$$

$$[x'_1, y'_{n+2}] = \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+1} \beta'_k y'_k = \beta'_{\lfloor \frac{n+5}{2} \rfloor} a_1^{2\lfloor \frac{n+5}{2} \rfloor - 2} \left(a_1 y_{\lfloor \frac{n+5}{2} \rfloor} + a_2 y_{\lfloor \frac{n+5}{2} \rfloor + 1} + \dots + \right.$$

$$\left. + a_{n+2 - \lfloor \frac{n+5}{2} \rfloor} y_{n+1} \right) + b'_{\lfloor \frac{n+5}{2} \rfloor + 1} a_1^{2\lfloor \frac{n+5}{2} \rfloor} \left(a_1 y_{\lfloor \frac{n+5}{2} \rfloor + 1} + a_2 y_{\lfloor \frac{n+5}{2} \rfloor + 2} + \dots + a_{n+1 - \lfloor \frac{n+5}{2} \rfloor} y_{n+1} \right) +$$

$$+ \dots + \beta'_{n+1} a_1^{2n+1} y_{n+1} = \sum_{k=\lfloor \frac{n+5}{2} \rfloor}^{n+1} \beta'_k a_1^{2(k-1)} \sum_{j=1}^{n+2-k} a_j y_{k+j-1}. \quad (3.15)$$

From (3.14) and (3.15) we obtain restrictions:

$$b_{n+2}\beta_j = a_1^{2j-1}\beta'_j, \quad \left\lfloor \frac{n+5}{2} \right\rfloor \leq j \leq n+1. \quad (3.16)$$

If we consider other multiplications, then we obtain restrictions (3.16) or identities. \square

The description up to isomorphism of a family from Lemma 3.3 is represented in the following theorem.

Theorem 3.5. *Let L be a Leibniz superalgebra of variety $\text{Leib}^{n,m}$ with characteristic sequence $(n \mid m-1, 1)$, nilindex $n+m$ and $m = n+2$. Then L is isomorphic to one of the following pairwise non isomorphic superalgebras:*

$$L \left(W_{s, n+2 - \lfloor \frac{n+5}{2} \rfloor} \left(V_{j, n+2 - \lfloor \frac{n+5}{2} \rfloor}^1 \left(\beta_{\lfloor \frac{n+5}{2} \rfloor}, \beta_{\lfloor \frac{n+5}{2} \rfloor + 1}, \dots, \beta_{n+1} \right) \right) \right),$$

where $1 \leq j \leq n+2 - \left\lfloor \frac{n+5}{2} \right\rfloor$, $1 \leq s \leq n+3 - \left\lfloor \frac{n+5}{2} \right\rfloor - j$,

$$L(0, 0, \dots, 0).$$

Proof. From Theorem 3.4 we have the following restrictions:

$$b_{n+2}\beta'_{\lfloor \frac{n+5}{2} \rfloor + j} = a_1^{2\lfloor \frac{n+5}{2} \rfloor + 2j - 3} \beta'_{\lfloor \frac{n+5}{2} \rfloor + j}, \quad 0 \leq j \leq n+1 - \left\lfloor \frac{n+5}{2} \right\rfloor.$$

As $\left\lfloor \frac{n+5}{2} \right\rfloor \approx q+2$, for $n = 2q$ or $n = 2q-1$, then we obtain:

$$b_{n+2}\beta_{q+j+2} = a_1^{2q+2j+1} \beta'_{q+j+2}, \quad 0 \leq j \leq n+1 - q - 2.$$

The proof of this theorem is complete by using the same arguments as in the proof of Theorem 3.3 for even case. \square

Thus, Theorems 3.3 and 3.5 complete the classifications (up to isomorphism) of Leibniz superalgebras with characteristic sequence $C(L) = (n \mid m-1, 1)$ and nilindex $n+m$.

Acknowledgments. The last named author would like to acknowledge the hospitality of the University of Sevilla (Spain). He was supported by the grants INTAS - 04-83-3035 and NATO-Reintegration ref. CBP.EAP.RIG.983169.

References

- [1] Albeverio S., Ayupov. Sh.A., Omirov B.A. On nilpotent and simple Leibniz algebras. Comm. in Algebra, 2005, vol. 33, No. 1, p. 159-172
- [2] Ayupov Sh.A., Omirov B.A., On some classes of nilpotent Leibniz algebras. Sib. Math. J., 2001, vol. 42, No. 1, p. 18-29.

- [3] Ayupov Sh.A., Omirov B.A., Khudoyberdiyev A. Kh., The classification of filiform Leibniz superalgebras of nilindex $n+m$. *Acta Mathematica Sinica*, to appear 2009.
- [4] Bordeman M., Gómez J.R., Khakimdjano Yu., Navarro R.M., Some deformations of nilpotent Lie superalgebras. *J. Geometry and Physics*, (2007), 57(5), 1391-1403.
- [5] Gilg M. On the classification of filiform Lie superalgebras. *Proceeding of the I Colloquium on Lie Theory and Applications (Vigo)*, 2000, p. 83-87.
- [6] Gilg M. *Super-algèbres*: PhD thesis. University of Haute Alsace, 2000. 126 p.
- [7] Gómez J.R., Navarro R.M., Omirov B.A. On nilpotent Leibniz superalgebras. Preprint MA1-02-XI06 Universidad de Sevilla, 2006, pp. 24 (arXiv.math.RA/0611723).
- [8] Gómez J.R., Khakimdjano Yu., Navarro R.M. Some problems concerning to nilpotent Lie superalgebras. *J. Geom. Phys.* (2004), vol. 51, No. 4, p. 473-486.
- [9] Gómez J.R., Khakimdjano Yu., Navarro R.M. Infinitesimal deformations of the Lie superalgebras. *J. Geom. Phys.* (2008), p. 849-859.
- [10] Kac V.G. Lie superalgebras, *Advances in Math.* (1977), vol. 26(1), p. 8-96.
- [11] Loday J.-L. Une version non commutative des algèbres de Lie: les algèbres de Leibniz. *Enseign. Math.*, 1993, vol. 39, p. 269-293.

LUISA M. CAMACHO, JOSÉ R. GÓMEZ. Dpto. Matemática Aplicada I. Universidad de Sevilla. Avda. Reina Mercedes, s/n. 41012 Sevilla. (Spain), e-mail: *lcamacho@us.es*, *jrgomez@us.es*

ROSA M. NAVARRO. Dpto. de Matemáticas, Universidad de Extremadura, Cáceres (Spain), e-mail: *rnavarro@unex.es*

BAKHROM A. OMIROV. Institute of Mathematics and Information Technologists, Uzbekistan Academy of Science, F. Hodjaev str. 29, 100125, Tashkent (Uzbekistan), e-mail: *omirovb@mail.ru*