

University Libraries University of Nevada, Las Vegas

Physics & Astronomy Faculty Publications

Physics and Astronomy

1-3-2020

Relation Between Gravitational Mass and Baryonic Mass for Non-**Rotating and Rapidly Rotating Neutron Stars**

He Gao Beijing Normal University, gaohe@bnu.edu.cn

Shun-Ke Ai University of Nevada, Las Vegas

Zhou-Jian Cao Beijing Normal University

Bing Zhang University of Nevada, Las Vegas, bing.zhang@unlv.edu

Zhen-Yu Zhu Xiamen University Follow this and additional works at: https://digitalscholarship.unlv.edu/physastr_fac_articles

🔮 Part of the Cosmology, Relativity, and Gravity Commons, and the Stars, Interstellar Medium and the See next page for additional authors

Repository Citation

Gao, H., Ai, S., Cao, Z., Zhang, B., Zhu, Z., Li, A., Zhang, N., Bauswein, A. (2020). Relation Between Gravitational Mass and Baryonic Mass for Non-Rotating and Rapidly Rotating Neutron Stars. Frontiers in Physics, 15(2), 1-10. Frontiers Media.

http://dx.doi.org/10.1007/s11467-019-0945-9

This Article is protected by copyright and/or related rights. It has been brought to you by Digital Scholarship@UNLV with permission from the rights-holder(s). You are free to use this Article in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s) directly, unless additional rights are indicated by a Creative Commons license in the record and/ or on the work itself.

This Article has been accepted for inclusion in Physics & Astronomy Faculty Publications by an authorized administrator of Digital Scholarship@UNLV. For more information, please contact digitalscholarship@unlv.edu.

Authors

He Gao, Shun-Ke Ai, Zhou-Jian Cao, Bing Zhang, Zhen-Yu Zhu, Ang Li, Nai-Bo Zhang, and Andreas Bauswein



Relation between gravitational mass and baryonic mass for non-rotating and rapidly rotating neutron stars

He Gao^{1,†}, Shun-Ke Ai^{2,1}, Zhou-Jian Cao¹, Bing Zhang², Zhen-Yu Zhu^{3,4}, Ang Li³, Nai-Bo Zhang⁵, Andreas Bauswein⁶

¹Department of Astronomy, Beijing Normal University, Beijing 100875, China

²Department of Physics and Astronomy, University of Nevada Las Vegas, NV 89154, USA

³Department of Astronomy, Xiamen University, Xiamen 361005, China

⁴Institute for Theoretical Physics, Max-von-Laue-Strasse 1, 60438 Frankfurt, Germany

⁵Shandong Provincial Key Laboratory of Optical Astronomy and Solar-Terrestrial Environment, School of Space Science and

Physics, Institute of Space Sciences, Shandong University, Weihai 264209, China

⁶GSI Helmholtzzentrum für Schwerionenforschung, Planckstraße 1, 64291 Darmstadt, Germany

Corresponding author. E-mail: [†]gaohe@bnu.edu.cn

Received November 4, 2019; accepted November 26, 2019

With a selected sample of neutron star (NS) equations of state (EOSs) that are consistent with the current observations and have a range of maximum masses, we investigate the relations between NS gravitational mass M_q and baryonic mass M_b , and the relations between the maximum NS mass supported through uniform rotation $(M_{\rm max})$ and that of nonrotating NSs $(M_{\rm TOV})$. We find that for an EOS-independent quadratic, universal transformation formula $(M_b = M_q + A \times M_q^2)$, the best-fit A value is 0.080 for non-rotating NSs, 0.064 for maximally rotating NSs, and 0.073 when NSs with arbitrary rotation are considered. The residual error of the transformation is $\sim 0.1 M_{\odot}$ for non-spin or maximum-spin, but is as large as $\sim 0.2 M_{\odot}$ for all spins. For different EOSs, we find that the parameter A for non-rotating NSs is proportional to $R_{1.4}^{-1}$ (where $R_{1.4}$ is NS radius for $1.4M_{\odot}$ in units of km). For a particular EOS, if one adopts the best-fit parameters for different spin periods, the residual error of the transformation is smaller, which is of the order of $0.01 M_{\odot}$ for the quadratic form and less than $0.01 M_{\odot}$ for the cubic form $(M_b = M_g + A_1 \times M_g^2 + A_2 \times M_g^3)$. We also find a very tight and general correlation between the normalized mass gain due to spin $\Delta m \equiv (M_{\rm max} - M_{\rm TOV})/M_{\rm TOV}$ and the spin period normalized to the Keplerian period \mathcal{P} , i.e., $\log_{10} \Delta m = (-2.74 \pm 0.05) \log_{10} \mathcal{P} + \log_{10} (0.20 \pm 0.01)$, which is independent of EOS models. These empirical relations are helpful to study NS-NS mergers with a long-lived NS merger product using multi-messenger data. The application of our results to GW170817 is discussed.

Keywords gravitational waves

1 Introduction

The structure of neutron stars (NSs) depends on the poorly understood physical properties of matter under extreme conditions, especially the equation-of-state (EOS) of matter at the nuclear density [1, 2]. One way to diagnose the NS EOS is to constrain the NS maximum mass $M_{\rm TOV}$ of nonrotating stellar models. For instance, observing an NS with a sufficiently large mass could set an interesting lower limit on the NS maximum mass, thus ruling out a set of soft EOS models. Up to now, the largest well-measured NS mass is ~ $2.01 M_{\odot}$ for PSR J0348+0432 [3].

Binary neutron star (BNS) mergers have been suggested to be good targets for constraining the NS maximum mass in two different approaches: i) For one particular BNS merger event, if the gravitational mass of the BNS merger remnant, its rotational properties, and its life time until black-hole formation can be inferred from gravitational wave (GW) and electromagnetic (EM) observations, one can directly infer the NS maximum mass [4–8]; ii) Supposing that the internal X-ray plateau followed by a very rapid decay as observed in some SGRBs signals the formation of a long-lived, supramassive NS after the merger [9–11], statistical analyses on historical SGRB X-ray afterglow data can be used to estimate the fraction for different BNS merger products, thus placing constraint on the NS maximum mass [11–15].

For both approaches, it is important to estimate the

^{*}arXiv: 1905.03784.

remnant mass for the merger product. In NS problems, two masses, i.e., the baryonic mass M_b and the gravitational mass M_g , are usually discussed. The former (M_b) is theoretically relevant, since it is directly connected to the mass of the iron core of the progenitor massive star. In the problems of NS-NS mergers, it is the baryonic mass that is conserved. The latter (M_a) , on the other hand, is directly measured from observations, and is smaller than M_b due to the subtraction of the binding energy. Studying the general relationship between M_q and M_b becomes ever more important, as the accuracy of NS mass measurements becomes sufficiently precise to i) set interesting lower limits on the NS maximum mass [1, 16]; ii) to study the NS initial mass function [17–19]; iii) to study the neutrino emission from core collapse supernovae [20–23], and so on.

In order to calculate the remnant mass of BNS mergers, the conversion between the gravitational mass and baryonic mass for both non-rotating and rapidly rotating NSs is needed. Specifically speaking, it is generally assumed that NS-NS mergers conserve the baryonic mass, with a small fraction, $M_{\rm ej} \lesssim 10^{-1} M_{\odot}$, ejected during the merger [e.g., 24–28]. On the other hand, GW observations can only provide the total gravitational mass of the system with the infinite binary separation, $M_{a,tot}$. In order to calculate the mass of the merger remnant, one needs to first convert $M_{g,tot}$ to total baryonic mass $M_{b,tot}$, then convert $M_{b,tot} - M_{ej}$ back to the remnant gravitational mass $M_{a,\text{rem}}$. According to the observed galactic NS binary population, before the merger, the relatively low spin period is expected for both NSs in the binary [29, for details]. Therefore, the relation between M_a and M_b for nonrotating NSs should be enough in converting $M_{q,tot}$ into $M_{b,tot}$. However, since the newborn central remnant must be rapidly spinning, if the remnant is a uniformly rotating NS, the conversion from M_b to M_g for a rapidly rotating NS becomes essential¹⁾. Finally, the maximum gravitational mass is significantly enhanced by rapid rotation. For some constraints on $M_{\rm TOV}$, the relation between the maximum NS mass supported through uniform rotation (marked as $M_{\rm max}$) and $M_{\rm TOV}$ is also required.

In this work, we consider rigidly rotating stellar equilibrium models (treated by numerical relativity methods) and aim to find relations between M_g and M_b , and between $M_{\rm max}$ and $M_{\rm TOV}$, for a selected sample of EOSs, which are consistent with current observations (i.e., $M_{\rm TOV}$ is larger than $\sim 2.01 M_{\odot}$) and have a range of maximum masses. We apply our results to GW170817 to estimate its merger remnant mass. Here we only consider rigidly rotating non-magnetized NSs at zero temperature. In the literature, the relationships between M_g and M_b or be-

tween binding energy $M_b - M_g$ and M_g have been investigated by many authors, most of which have focused on nonrotating non-magnetized NSs at zero temperature [e.g., 17, 20, 31, 32], with some studying rotating NSs [e.g., 33]. The comparisons between our results and previous results are discussed in detail. NS structural quantities in the differential rotation phase are not touched in this work, which has been discussed in previous works [e.g., 33–37].

2 NS structure equations and NS EOS

We consider the equilibrium equations for a stationary, axially symmetric, rigid rotating NS, within a fully general relativistic framework. The spacetime metric can be written in the form

$$ds^{2} = -e^{2\nu}dt^{2} + r^{2}\sin^{2}\theta B^{2}e^{-2\nu}(d\phi - \omega dt)^{2} +e^{2\alpha}(dr^{2} + r^{2}d\theta^{2}),$$
(1)

where the potentials ν , B, ω , and α only depend on r and θ , and have the following asymptotic decay [38]:

$$\nu = -\frac{M}{r} + \frac{B_0 M}{3r^3} + \nu_2 P_2(\cos \theta) + \mathcal{O}\left(\frac{1}{r^4}\right),$$

$$B = 1 + \frac{B_0}{r^2} + \mathcal{O}\left(\frac{1}{r^4}\right),$$

$$\omega = \frac{2I\Omega}{r^3} + \mathcal{O}\left(\frac{1}{r^4}\right),$$
(2)

where M, I and Ω are the NS mass, moment of inertia, and angular frequency, respectively. B_0 and ν_2 are real constants.

We describe the interior of the NS as a perfect fluid, whose energy-momentum tensor reads

$$T^{\mu\nu} = (\rho + P)u^{\mu}u^{\nu} + Pg^{\mu\nu}, \qquad (3)$$

where ρ and P are the energy density and the pressure, and u^{μ} is the 4-velocity. Given a particular NS EOS, we use the public code RNS [39] to solve the field equations for the rotating NS.

Our selection of realistic (tabulated) EOSs (as listed in Table 1) are SLy [40], WFF1 [41], WFF2 [41], AP3 [42], AP4 [42], BSK21 [43], DD2 [44], MPA1 [45], MS1 [46], MS1b [46], with $M_{\rm TOV}$ ranging from $2.05 M_{\odot}$ to $2.78 M_{\odot}$.

3 Relation between M_g and M_b

Given a particular EOS, the baryonic mass (M_b) and gravitational mass (M_g) for a rigid rotating NS are determined by the values of central energy density (ρ_c) and spin period (P). In order to figure out the relationship between M_b and M_g , for each EOS, we calculate a series of M_b and M_g for different spin periods. We first present the results for the non-spin case and the maximally-spinning

¹⁾It is worth noticing that besides knowing the relations between M_g and M_b , one still faces some problems such as extracting the ejecta mass information from EM observations or estimating the initial spin period of the rigidly rotating NS, especially when significant angular momentum loss (e.g., due to the strong viscous spin down [30]) is considered.

	$M_{ m TOV}$ (M_{\odot})	$R_{1.4}$ (km)	A^*	A_1^*	A_2^*	A^+	A_1^+	A_2^+	P_k (ms)	$\stackrel{\alpha}{\left(10^{-10}\mathrm{s}^{-\beta}\right)}$	β	a	b
SLy	2.05	11.69	0.083	0.041	0.023	0.066	0.042	0.011	0.55	2.311	-2.734	0.086	0.209
wff1	2.14	10.40	0.102	0.046	0.029	0.076	0.050	0.012	0.47	3.111	-2.729	0.106	0.279
wff2	2.20	11.1	0.092	0.043	0.026	0.070	0.046	0.011	0.50	2.170	-2.694	0.096	0.270
ap4	2.22	11.36	0.090	0.045	0.023	0.069	0.046	0.010	0.51	2.095	-2.721	0.094	0.251
Bsk21	2.28	12.55	0.079	0.039	0.020	0.061	0.038	0.010	0.60	1.958	-2.799	0.083	0.257
ap3	2.39	12.01	0.087	0.046	0.019	0.066	0.047	0.007	0.55	1.883	-2.780	0.091	0.288
dd2	2.42	13.12	0.077	0.046	0.014	0.059	0.044	0.006	0.65	2.269	-2.811	0.079	0.259
mpa1	2.48	12.41	0.082	0.046	0.017	0.063	0.046	0.007	0.59	2.584	-2.693	0.088	0.284
ms1	2.77	14.70	0.069	0.042	0.010	0.053	0.038	0.005	0.72	5.879	-2.716	0.072	0.263
ms1b	2.78	14.46	0.070	0.043	0.011	0.054	0.040	0.005	0.71	4.088	-2.770	0.074	0.283

 Table 1
 Characteristic parameters for various EOSs.

*Best fit vales for non-spinning NS cases.

⁺Best fit values for Keplerian spinning NS cases.

(Keplerian) case. We then sample the spin period P as $\{1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 3.0, 4.0\}$ P_k , where P_k is the Keplerian period corresponding to $M_{g,max}$. We consider both relations describing all the EOSs and the relations for individual EOSs.

3.1 Non-rotating NSs

For a non-rotating NS, transformation of the baryonic NS mass to gravitational mass is commonly approximated using the quadratic formula

$$M_b = M_q + A \times M_a^2,\tag{4}$$

where A is a constant, which is quoted as 0.075 by Timmes et al. [17]. Throughout the paper, M_b and M_g are in units of M_{\odot} . Here we treat A as a free parameter in order to find out its best-fit value for each adopted EOS (results are collected in Table 1). If each EOS adopting its own best fit A value, the residual error of the transformation between M_g and M_b is in order of a few $10^{-2}M_{\odot}$, see Fig. 1 for details. Putting (M_g, M_b) results for all adopted EOS together, the overall best fit A value is 0.080. In this case, the residual error of the transformation from M_g to M_b is as large as $0.12M_{\odot}$ and the residual error from M_b to M_g is as large as $0.09M_{\odot}$. By comparison, when A is adopted as 0.075, the residual error of the transformation from M_g to M_b is as large as $0.14M_{\odot}$ and the residual error from M_b to M_g is as large as $0.11M_{\odot}$.

In order to further reduce the residual error of the transformation, we test a new cubic formula

$$M_b = M_g + A_1 \times M_g^2 + A_2 \times M_g^3.$$
 (5)

We determine the best fit values of A_1 and A_2 for each adopted EOS (collected in Table 1). If each EOS adopts its own best-fit values, the residual error of the transformation between M_q and M_b is in the order of a few $10^{-3}M_{\odot}$, which is almost one order of magnitude better than the quadratic cases. See Fig. 1 for details. Putting all EOSs together, the overall best fit A_1 and A_2 values are 0.0729 and 0.0032. In this case, the residual error of the transformation from M_g to M_b is as large as $0.12M_{\odot}$ and the residual error from M_b to M_g is as large as $0.09M_{\odot}$, which are in the same order as the quadratic case.

In the literature, some universal relations between the binding energy $M_b - M_g$ and the neutron star's compactness have been proposed [31, 32], which are also applicable for transformation between M_b and M_g . Here we find that for the quadratic transformation, one has $A \times R_{1.4} \approx 1$ for all adopted EOS, where A is the best fit value and $R_{1.4}$ is the NS radius (in unit of km) for $1.4M_{\odot}$. We thus propose a new universal relation

$$M_b = M_g + R_{1.4}^{-1} \times M_g^2, \tag{6}$$

with 1.8% relative error. We note that in our proposed relation, only $R_{1.4}$ is invoked in the relation, instead of using the compactness number proposed in previous works [e.g., 31, 32]. The accuracy is similar to the previous ones in terms of the residual error of the transformation, but our proposed relation is more practical since it does not involve the calculations of the radii corresponding to different masses. In case that the binding energy $M_b - M_g$ is measured independently, such a universal relation may be helpful to constrain $R_{1.4}$, which helps to distinguish NS EOSs [31].

3.2 Maximally rotating NSs

It is generally believed that the NS produced from NS-NS mergers may have an initial spin period close to the Keplerian period, which is supposed to be the minimum spin period of a rigidly rotating NS with a certain M_b . Also, M_g in this case is known as the maximum gravitational



Fig. 1 The relation between the baryonic mass (M_b) and gravitational mass (M_g) for a non-rotating NS and the residual error for different transformation formulae. Different colors denote different EOSs. The solid and dashed lines represent the best fitting results with the quadratic and cubic formulae, respectively, with the best-fit parameters adopted for each EOS.

mass for that M_b . We still apply the quadratic and cubic formulae to fit the relation of M_b vs. M_q . For the quadratic formula, it is worth noting that if A = 0.075 is adopted, the residual error of the transformation from M_q to M_b could be up to $0.22M_{\odot}$, and that of the transformation from M_b to M_g could be up to $0.15M_{\odot}$, which is imprecise to estimate the remnant mass of NS-NS mergers. Here we find that in the maximal-rotation case, the overall best fitting value is A = 0.064 with the residual errors $0.11 M_{\odot}$ and $0.08 M_{\odot}$, respectively, for transformation from M_q to M_b and vice versa. The newly proposed coefficient could significantly reduce the residual error of the transformation. For the cubic formula, the overall best fitting value are $A_1 = 0.0561$ and $A_2 = 0.0033$. But it cannot reduce the residual error effectively compared with the quadratic formula.

When we adopt the best-fit parameters for each EoS, the residual error of transformation between M_g and M_b is in the order of a few $10^{-2}M_{\odot}$ for the quadratic formulae and $10^{-3}M_{\odot}$ for the cubic formulae. See details in Fig. 2.

Similar to the non-spin case, we find $A \times R_{1.4} \approx 0.78$ for the quadratic formula. We thus have a universal formula in the maximally rotation case as

$$M_b = M_q + 0.78R_{1.4}^{-1} \times M_q^2 \tag{7}$$

with a 1.3% relative error.

3.3 General NSs with arbitrary rotation

For individual EOSs, we also fit their M_b and M_q relations with both the quadratic and cubic formulae for the situations with different rotation periods. We find following conclusions which could be applied to all adopted EOSs: i) The relation between M_b and M_g is different for different spin periods. ii) If one adopts the best-fit parameters for each spin period, the cubic formula is almost one order of magnitude better than the quadratic formula in terms of the residual error of the transformation. For instance, the residual error of the transformation from M_q to M_b is up to $0.06M_{\odot}$ for the quadratic case and is up to $0.01M_{\odot}$ for the cubic case. iii) Putting together the (M_q, M_b) results for different spin periods, with the overall best-fit values, the cubic formula is no better than the quadratic formula. For instance, the residual error of the transformation from M_q to M_b is up to $0.068 M_{\odot}$ for the quadratic case and is up to $0.062 M_{\odot}$ for the cubic case. For a comparison, when A is fixed to 0.075, the residual error of the transformation from M_q to M_b is as large as $0.11 M_{\odot}$. The best-fit values for the quadratic and cubic formulae are collected in Table 2.

Considering (M_g, M_b) results for different EOSs and different spin periods together, the overall best-fit value is A = 0.073 for the quadratic formula and $A_1 = 0.078$ and $A_2 = -0.0018$ for the cubic formula. In this case, the



Fig. 2 The relation between the baryonic mass (M_b) and gravitational mass (M_g) for a maximally rotating NS and the residual error for different transformation formulae.

Table	2	Best	fitting	$\operatorname{results}$	for	$\operatorname{different}$	\mathbf{EoSs}	with	different	spin	periods.
-------	----------	------	---------	--------------------------	-----	----------------------------	-----------------	------	-----------	-----------------------	----------

	SLy			wff1			wff2		ap4			Bsk21			
	A	A_1	A_2	A	A_1	A_2	A	A_1	A_2	A	A_1	A_2	A	A_1	A_2
$P=1.3P_k$	0.072	-0.318	0.177	0.085	-0.172	0.112	0.079	-0.327	0.171	0.077	-0.214	0.123	0.068	-0.179	0.101
$P=1.4P_k$	0.073	-0.112	0.087	0.084	-0.058	0.066	0.079	-0.111	0.083	0.077	-0.075	0.067	0.068	-0.058	0.054
$P = 1.5P_k$	0.072	-0.046	0.058	0.085	-0.020	0.050	0.080	-0.049	0.058	0.078	-0.029	0.049	0.067	-0.017	0.038
$P = 1.6P_k$	0.073	-0.016	0.045	0.086	0.002	0.042	0.079	-0.017	0.045	0.077	-0.004	0.039	0.068	0.001	0.032
$P=1.7P_k$	0.073	0.000	0.038	0.086	0.015	0.037	0.079	0.000	0.039	0.078	0.009	0.034	0.068	0.012	0.028
$P=1.8P_k$	0.073	0.011	0.034	0.086	0.024	0.033	0.079	0.012	0.034	0.078	0.018	0.031	0.069	0.019	0.025
$P=1.9P_k$	0.073	0.018	0.031	0.088	0.028	0.032	0.080	0.019	0.031	0.079	0.023	0.029	0.069	0.024	0.023
$P=2.0P_k$	0.074	0.022	0.029	0.088	0.033	0.030	0.081	0.024	0.030	0.079	0.028	0.026	0.069	0.028	0.022
$P=3.0P_k$	0.076	0.038	0.023	0.091	0.047	0.026	0.083	0.041	0.024	0.082	0.043	0.022	0.072	0.037	0.019
$P = 4.0P_k$	0.077	0.041	0.022	0.093	0.049	0.026	0.085	0.044	0.024	0.083	0.046	0.022	0.073	0.039	0.019
		ap3			dd2			mpa1			ms1			ms1b	
	A	ap3 A_1	A_2	A	$\frac{\mathrm{dd}2}{A_1}$	A_2	A	$\frac{mpa1}{A_1}$	A_2	A	$ms1$ A_1	A_2	A	ms1b A_1	A_2
$P = 1.3P_k$	A 0.072	ap3 A_1 -0.292	A_2 0.140	A 0.064	dd2 A_1 -0.100	A_2 0.064	A 0.069	mpa1 A_1 -0.112	A_2 0.070	A 0.058	ms1 A_1 -0.095	A_2 0.052	A 0.059	ms1b A_1 -0.104	A_2 0.055
$\boxed{\begin{array}{c} P=1.3P_k\\ \hline P=1.4P_k \end{array}}$	A 0.072 0.073	ap3 A_1 -0.292 -0.095	A_2 0.140 0.068	A 0.064 0.064	$ dd2 A_1 -0.100 -0.028 $	A_2 0.064 0.038	A 0.069 0.070	mpa1 A_1 -0.112 -0.039	A_2 0.070 0.043	A 0.058 0.058	ms1 A_1 -0.095 -0.030	A_2 0.052 0.031	A 0.059 0.059	ms1b A_1 -0.104 -0.031	A ₂ 0.055 0.032
$P = 1.3P_k$ $P = 1.4P_k$ $P = 1.5P_k$	A 0.072 0.073 0.073	$ \begin{array}{c} \text{ap3} \\ A_1 \\ -0.292 \\ -0.095 \\ -0.033 \end{array} $	A_2 0.140 0.068 0.045	A 0.064 0.065	$ dd2 A_1 -0.100 -0.028 -0.002 $	A2 0.064 0.038 0.029	A 0.069 0.070 0.070	$\begin{array}{c} {\rm mpa1} \\ A_1 \\ -0.112 \\ -0.039 \\ -0.005 \end{array}$	$ \begin{array}{c} A_2 \\ 0.070 \\ 0.043 \\ 0.032 \end{array} $	A 0.058 0.058 0.058	ms1 A_1 -0.095 -0.030 -0.006	$ \begin{array}{c} A_2 \\ 0.052 \\ 0.031 \\ 0.024 \end{array} $	A 0.059 0.059 0.059	ms1b A_1 -0.104 -0.031 -0.004	$ \begin{array}{c} A_2 \\ 0.055 \\ 0.032 \\ 0.023 \\ \end{array} $
$\begin{array}{c} \hline P = 1.3 P_k \\ \hline P = 1.4 P_k \\ \hline P = 1.5 P_k \\ \hline P = 1.6 P_k \end{array}$	A 0.072 0.073 0.073 0.073	$\begin{array}{c} {\rm ap3} \\ A_1 \\ -0.292 \\ -0.095 \\ -0.033 \\ -0.006 \end{array}$	$\begin{array}{c} A_2 \\ 0.140 \\ 0.068 \\ 0.045 \\ 0.035 \end{array}$	A 0.064 0.065 0.065	$\begin{array}{c} {\rm dd2} \\ A_1 \\ -0.100 \\ -0.028 \\ -0.002 \\ 0.013 \end{array}$	$ \begin{array}{c} A_2 \\ 0.064 \\ 0.038 \\ 0.029 \\ 0.023 \end{array} $	A 0.069 0.070 0.070 0.070	$\begin{array}{c} {\rm mpa1} \\ A_1 \\ -0.112 \\ -0.039 \\ -0.005 \\ 0.010 \end{array}$	$ \begin{array}{c} A_2 \\ 0.070 \\ 0.043 \\ 0.032 \\ 0.027 \\ \end{array} $	A 0.058 0.058 0.058 0.058	ms1 A_1 -0.095 -0.030 -0.006 0.007	$ \begin{array}{c} A_2 \\ 0.052 \\ 0.031 \\ 0.024 \\ 0.020 \\ \end{array} $	A 0.059 0.059 0.059 0.059	ms1b A ₁ -0.104 -0.031 -0.004 0.009	$ \begin{array}{c} A_2 \\ 0.055 \\ 0.032 \\ 0.023 \\ 0.019 \\ \end{array} $
$P = 1.3P_k$ $P = 1.4P_k$ $P = 1.5P_k$ $P = 1.6P_k$ $P = 1.7P_k$	A 0.072 0.073 0.073 0.073 0.074	$\begin{array}{c} {\rm ap3} \\ A_1 \\ -0.292 \\ -0.095 \\ -0.033 \\ -0.006 \\ 0.008 \end{array}$	$\begin{array}{c} A_2 \\ 0.140 \\ 0.068 \\ 0.045 \\ 0.035 \\ 0.030 \end{array}$	A 0.064 0.065 0.065 0.066	$\begin{array}{c} \rm dd2 \\ A_1 \\ -0.100 \\ -0.028 \\ -0.002 \\ 0.013 \\ 0.022 \end{array}$	$\begin{array}{c} A_2 \\ 0.064 \\ 0.038 \\ 0.029 \\ 0.023 \\ 0.020 \end{array}$	A 0.069 0.070 0.070 0.070 0.071	$\begin{array}{c} mpa1 \\ A_1 \\ -0.112 \\ -0.039 \\ -0.005 \\ 0.010 \\ 0.019 \end{array}$	$\begin{array}{c} A_2 \\ 0.070 \\ 0.043 \\ 0.032 \\ 0.027 \\ 0.023 \end{array}$	A 0.058 0.058 0.058 0.058 0.059	$\begin{array}{c} {\rm ms1} \\ \\ A_1 \\ \\ -0.095 \\ \\ -0.030 \\ \\ -0.006 \\ \\ 0.007 \\ \\ 0.015 \end{array}$	$\begin{array}{c} A_2 \\ 0.052 \\ 0.031 \\ 0.024 \\ 0.020 \\ 0.017 \end{array}$	A 0.059 0.059 0.059 0.059 0.059	$\begin{array}{c} {\rm ms1b} \\ A_1 \\ -0.104 \\ -0.031 \\ -0.004 \\ 0.009 \\ 0.017 \end{array}$	$\begin{array}{c} A_2 \\ 0.055 \\ 0.032 \\ 0.023 \\ 0.019 \\ 0.017 \end{array}$
$\begin{array}{c} \hline P = 1.3P_k \\ \hline P = 1.4P_k \\ \hline P = 1.5P_k \\ \hline P = 1.6P_k \\ \hline P = 1.7P_k \\ \hline P = 1.8P_k \end{array}$	A 0.072 0.073 0.073 0.073 0.073 0.074	$\begin{array}{c} {\rm ap3} \\ A_1 \\ -0.292 \\ -0.095 \\ -0.033 \\ -0.006 \\ 0.008 \\ 0.017 \end{array}$	$\begin{array}{c} A_2 \\ 0.140 \\ 0.068 \\ 0.045 \\ 0.035 \\ 0.030 \\ 0.027 \end{array}$	A 0.064 0.065 0.065 0.065 0.066	$\begin{array}{c} \rm dd2 \\ A_1 \\ -0.100 \\ -0.028 \\ 0.013 \\ 0.022 \\ 0.027 \end{array}$	$\begin{array}{c} A_2 \\ 0.064 \\ 0.038 \\ 0.029 \\ 0.023 \\ 0.020 \\ 0.019 \end{array}$	A 0.069 0.070 0.070 0.070 0.070 0.071	$\begin{array}{c} mpa1 \\ A_1 \\ -0.112 \\ -0.039 \\ -0.005 \\ 0.010 \\ 0.019 \\ 0.026 \end{array}$	$\begin{array}{c} A_2 \\ 0.070 \\ 0.043 \\ 0.032 \\ 0.027 \\ 0.023 \\ 0.021 \end{array}$	A 0.058 0.058 0.058 0.058 0.059 0.059	$\begin{array}{c} {\rm ms1} \\ A_1 \\ -0.095 \\ -0.030 \\ 0.007 \\ 0.015 \\ 0.020 \end{array}$	$\begin{array}{c} A_2 \\ 0.052 \\ 0.031 \\ 0.024 \\ 0.020 \\ 0.017 \\ 0.016 \end{array}$	A 0.059 0.059 0.059 0.059 0.059 0.059 0.059 0.060	$\begin{array}{c} {\rm ms1b} \\ A_1 \\ -0.104 \\ -0.031 \\ 0.009 \\ 0.017 \\ 0.023 \end{array}$	$\begin{array}{c} A_2 \\ 0.055 \\ 0.032 \\ 0.023 \\ 0.019 \\ 0.017 \\ 0.015 \end{array}$
$\begin{array}{c} P = 1.3 P_k \\ P = 1.4 P_k \\ \hline P = 1.5 P_k \\ P = 1.6 P_k \\ \hline P = 1.7 P_k \\ \hline P = 1.8 P_k \\ \hline P = 1.9 P_k \end{array}$	A 0.072 0.073 0.073 0.073 0.074 0.074	$\begin{array}{c} {\rm ap3} \\ A_1 \\ -0.292 \\ -0.095 \\ -0.033 \\ -0.006 \\ 0.008 \\ 0.017 \\ 0.024 \end{array}$	$\begin{array}{c} A_2 \\ 0.140 \\ 0.068 \\ 0.045 \\ 0.035 \\ 0.030 \\ 0.027 \\ 0.024 \end{array}$	A 0.064 0.065 0.065 0.065 0.066 0.066	$\begin{array}{c} \rm dd2 \\ A_1 \\ -0.100 \\ -0.028 \\ 0.013 \\ 0.022 \\ 0.027 \\ 0.031 \end{array}$	$\begin{array}{c} A_2 \\ 0.064 \\ 0.038 \\ 0.029 \\ 0.023 \\ 0.020 \\ 0.019 \\ 0.017 \end{array}$	A 0.069 0.070 0.070 0.070 0.071 0.071 0.072	$\begin{array}{c} mpa1 \\ A_1 \\ -0.112 \\ -0.039 \\ 0.010 \\ 0.010 \\ 0.019 \\ 0.026 \\ 0.031 \end{array}$	$\begin{array}{c} A_2 \\ 0.070 \\ 0.043 \\ 0.032 \\ 0.027 \\ 0.023 \\ 0.021 \\ 0.020 \end{array}$	A 0.058 0.058 0.058 0.058 0.059 0.059 0.060	$\begin{array}{c} {\rm ms1} \\ A_1 \\ -0.095 \\ -0.030 \\ 0.007 \\ 0.007 \\ 0.015 \\ 0.020 \\ 0.024 \end{array}$	$\begin{array}{c} A_2 \\ 0.052 \\ 0.031 \\ 0.024 \\ 0.020 \\ 0.017 \\ 0.016 \\ 0.015 \end{array}$	$\begin{array}{c} A \\ 0.059 \\ 0.059 \\ 0.059 \\ 0.059 \\ 0.060 \\ 0.060 \\ 0.060 \\ 0.061 \end{array}$	$\begin{array}{c} {\rm ms1b} \\ A_1 \\ -0.104 \\ -0.031 \\ 0.009 \\ 0.017 \\ 0.023 \\ 0.027 \end{array}$	$\begin{array}{c} A_2 \\ 0.055 \\ 0.032 \\ 0.023 \\ 0.019 \\ 0.017 \\ 0.015 \\ 0.014 \end{array}$
$\begin{tabular}{ c c c c c } \hline P &= 1.3 P_k \\ \hline P &= 1.4 P_k \\ \hline P &= 1.5 P_k \\ \hline P &= 1.6 P_k \\ \hline P &= 1.7 P_k \\ \hline P &= 1.8 P_k \\ \hline P &= 1.9 P_k \\ \hline P &= 2.0 P_k \\ \hline \end{tabular}$	A 0.072 0.073 0.073 0.073 0.074 0.074 0.074	$\begin{array}{c} {\rm ap3} \\ A_1 \\ -0.292 \\ -0.095 \\ -0.033 \\ -0.006 \\ 0.008 \\ 0.017 \\ 0.024 \\ 0.029 \end{array}$	$\begin{array}{c} A_2 \\ 0.140 \\ 0.068 \\ 0.045 \\ 0.035 \\ 0.030 \\ 0.027 \\ 0.024 \\ 0.023 \end{array}$	A 0.064 0.065 0.065 0.066 0.066 0.066 0.066	$\begin{array}{c} \rm dd2 \\ A_1 \\ -0.100 \\ -0.028 \\ 0.013 \\ 0.022 \\ 0.027 \\ 0.031 \\ 0.034 \end{array}$	$\begin{array}{c} A_2 \\ 0.064 \\ 0.038 \\ 0.029 \\ 0.023 \\ 0.020 \\ 0.019 \\ 0.017 \\ 0.017 \end{array}$	A 0.069 0.070 0.070 0.070 0.071 0.071 0.072 0.073	$\begin{array}{c} mpa1 \\ A_1 \\ -0.112 \\ -0.039 \\ 0.010 \\ 0.010 \\ 0.026 \\ 0.031 \\ 0.033 \end{array}$	$\begin{array}{c} A_2 \\ 0.070 \\ 0.043 \\ 0.032 \\ 0.027 \\ 0.023 \\ 0.021 \\ 0.020 \\ 0.019 \end{array}$	A 0.058 0.058 0.058 0.058 0.059 0.059 0.060	$\begin{array}{c} {\rm ms1} \\ A_1 \\ -0.095 \\ -0.030 \\ 0.007 \\ 0.007 \\ 0.015 \\ 0.020 \\ 0.024 \\ 0.027 \end{array}$	$\begin{array}{c} A_2 \\ 0.052 \\ 0.031 \\ 0.024 \\ 0.020 \\ 0.017 \\ 0.016 \\ 0.015 \\ 0.015 \end{array}$	$\begin{array}{c} A \\ 0.059 \\ 0.059 \\ 0.059 \\ 0.059 \\ 0.060 \\ 0.060 \\ 0.060 \\ 0.061 \\ 0.061 \end{array}$	$\begin{array}{c} {\rm ms1b} \\ A_1 \\ -0.104 \\ -0.031 \\ 0.009 \\ 0.017 \\ 0.023 \\ 0.027 \\ 0.030 \end{array}$	$\begin{array}{c} A_2 \\ 0.055 \\ 0.032 \\ 0.023 \\ 0.019 \\ 0.017 \\ 0.015 \\ 0.014 \\ 0.013 \end{array}$
$\begin{array}{c} P = 1.3 P_k \\ \hline P = 1.4 P_k \\ P = 1.5 P_k \\ \hline P = 1.6 P_k \\ P = 1.7 P_k \\ \hline P = 1.8 P_k \\ \hline P = 1.9 P_k \\ \hline P = 2.0 P_k \\ \hline P = 3.0 P_k \end{array}$	A 0.072 0.073 0.073 0.073 0.074 0.074 0.075	$\begin{array}{c} {\rm ap3} \\ A_1 \\ -0.292 \\ -0.095 \\ -0.033 \\ -0.006 \\ 0.008 \\ 0.017 \\ 0.024 \\ 0.029 \\ 0.043 \end{array}$	$\begin{array}{c} A_2 \\ 0.140 \\ 0.068 \\ 0.045 \\ 0.035 \\ 0.030 \\ 0.027 \\ 0.024 \\ 0.023 \\ 0.018 \end{array}$	A 0.064 0.065 0.065 0.066 0.066 0.066 0.067	$\begin{array}{c} \mathrm{dd2} \\ A_1 \\ -0.100 \\ -0.028 \\ 0.013 \\ 0.022 \\ 0.027 \\ 0.031 \\ 0.034 \\ 0.043 \end{array}$	$\begin{array}{c} A_2 \\ 0.064 \\ 0.038 \\ 0.029 \\ 0.023 \\ 0.020 \\ 0.019 \\ 0.017 \\ 0.017 \\ 0.014 \end{array}$	A 0.069 0.070 0.070 0.071 0.071 0.072 0.073	$\begin{array}{c} mpa1 \\ A_1 \\ -0.112 \\ -0.039 \\ -0.005 \\ 0.010 \\ 0.019 \\ 0.026 \\ 0.031 \\ 0.033 \\ 0.044 \end{array}$	$\begin{array}{c} A_2 \\ 0.070 \\ 0.043 \\ 0.032 \\ 0.027 \\ 0.023 \\ 0.021 \\ 0.020 \\ 0.019 \\ 0.016 \end{array}$	A 0.058 0.058 0.058 0.058 0.059 0.060 0.060	$\begin{array}{c} {\rm ms1} \\ A_1 \\ -0.095 \\ -0.030 \\ 0.007 \\ 0.007 \\ 0.015 \\ 0.020 \\ 0.024 \\ 0.027 \\ 0.037 \end{array}$	$\begin{array}{c} A_2 \\ 0.052 \\ 0.031 \\ 0.024 \\ 0.020 \\ 0.017 \\ 0.016 \\ 0.015 \\ 0.015 \\ 0.012 \end{array}$	$\begin{array}{c} A \\ 0.059 \\ 0.059 \\ 0.059 \\ 0.059 \\ 0.060 \\ 0.060 \\ 0.061 \\ 0.061 \\ 0.064 \end{array}$	$\begin{array}{c} {\rm ms1b} \\ A_1 \\ -0.104 \\ -0.031 \\ 0.009 \\ 0.017 \\ 0.023 \\ 0.027 \\ 0.030 \\ 0.039 \end{array}$	$\begin{array}{c} A_2 \\ 0.055 \\ 0.032 \\ 0.023 \\ 0.019 \\ 0.017 \\ 0.015 \\ 0.014 \\ 0.013 \\ 0.011 \end{array}$

residual error of the transformation from M_g to M_b is up to $0.20M_{\odot}$ for the quadratic case and is up to $0.19M_{\odot}$ for the cubic case. It is interesting to note that the overall best fit A value is very close to 0.075 as proposed in Ref. [17].

For a given EOS, the quadratic transformation parameter A is a rough function of the spin period (see Fig. 3),

$$A \approx a \mathrm{e}^{-\frac{b}{p}},\tag{8}$$

where $\mathcal{P} \equiv P/P_k$ is the characterized NS spin period normalized to the Keplerian period. The best-fit values for aand b are collected in Table 1. Interestingly, we find that $a \times R_{1.4} \approx 1$ and $b \approx 1/4$ for all the adopted EOSs. We thus have a new universal relation for a rotating NS,

$$M_b = M_g + R_{1.4}^{-1} e^{-\frac{1}{4\mathcal{P}}} \times M_g^2, \tag{9}$$

with a 3.3% relative error. This equation is consistent with Eq. (7) and Eq. (6) when $\mathcal{P} = 1$ and $\mathcal{P} \to \infty$ are adopted. Two EOS related parameters $R_{1.4}$ and P_k are invoked in this relation, which may be constrained if M_b and the binding energy $M_b - M_g$ for a NS could be independently measured.

4 M_{max} and M_{TOV} relation

It has been proposed to parameterize M_{max} as a function of the spin period of the central star [12],

$$M_{\rm max} = M_{\rm TOV} (1 + \alpha P^{\beta}), \tag{10}$$

where P is the spin period of the NS in units of second. Using the RNS code, we calculate the numerical values for α and β for our adopted EOSs. The results are presented in Table 1.



Fig. 3 Relation between the quadratic transformation coefficient A and the spin period of the NS. The dashed lines represent the best fitting results with formula $A \approx a e^{-\frac{b}{P}}$, with the fitting uncertainty showing in the lower panels.



Fig. 4 (a) The correlation between α and β for our selected EOSs. The solid and dashed lines present the regression line and its 2- σ region. The correlation coefficient between α and β is 0.75. (b) The correlation between the normalized mass gain due to spin $\Delta m \equiv (M_{\text{max}} - M_{\text{TOV}})/M_{\text{TOV}}$ and the normalized spin period $\mathcal{P} \equiv P/P_k$. The thin colored solid lines represent the results when specific α and β values are adopted for each EOS. The thick black solid and dashed lines represent the universal relation proposed by Eq. (12) and its 2- σ region.

We find that α and β are not independent of each other (see Fig. 4 for details). If we define

$$\Delta m \equiv (M_{\rm max} - M_{\rm TOV})/M_{\rm TOV} = \alpha P^{\beta}, \qquad (11)$$

as the normalized mass gain due to spin, as shown in Fig. 4, a very tight and general correlation between Δm and \mathcal{P} could be found as

$$\log_{10}\Delta m = (-2.74 \pm 0.05)\log_{10}\mathcal{P} + \log_{10}(0.20 \pm 0.01),$$
(12)

which is essentially independent of the EOS models. It is worth noticing that when $\mathcal{P} = 1$, i.e., when NS spin period equals to the Keplerian period, one has $\Delta m \simeq 0.2$, or $M_{\text{max}} \simeq 1.2M_{\text{TOV}}$, which is well consistent with the previous numerical simulation results [47–49].

5 Conclusion and discussion

In this work, we solved the field equations for the rotating NS with a selected sample of EOSs. For each EOS, we calculated a series of M_b and M_g for different spin periods. Our results could be summarized as follows:

- For non-rotating NSs, if one intends to apply an EOSindependent universal quadratic or cubic transformation formula to all the EOSs, one has the best-fit formulae $M_b = M_g + 0.080 M_g^2$ and $M_b = M_g + 0.0729 M_g^2 + 0.0032 M_g^3$. The residual error of the transformation from M_g to M_b is as large as $0.12 M_{\odot}$, and the residual error from M_b to M_g is as large as $0.09 M_{\odot}$. There is no advantage for the higher order formula. However, for individual EOSs, if one adopts its own best-fit values for the coefficients, the cubic formula is much better than the quadratic formula, with the residual error of the transformation less than $0.01 M_{\odot}$.
- For maximally rotating NSs, if one intends to apply an EOS-independent universal quadratic or cubic transformation formula to all the EOSs, one has the best-fit formulae $M_b = M_g + 0.064M_g^2$ and $M_b = M_g + 0.0595M_g^2 + 0.0017M_g^3$. The residual error of the transformation from M_g to M_b is as large as $0.10M_{\odot}$, and the residual error from M_b to M_g is as large as $0.07M_{\odot}$. There is no advantage for the higher order formula. Also, for individual EOSs, if one adopts its own best-fit values for the coefficients, the cubic formula is much better than the quadratic formula, with the residual error of the transformation less than $0.01M_{\odot}$.
- For general NSs with arbitrary rotation, the relation between M_b and M_q is different for different spin periods. If one intends to apply an EOS-independent universal quadratic or cubic transformation formula to all the EOSs for all spin periods, one has M_b = $M_g + 0.073 M_q^2$ and $M_b = M_g + 0.078 M_q^2 + 0.002 M_q^3$. The residual error of the transformation is up to $0.22M_{\odot}$. There is still no advantage for the higher order formula. Given an EOS, if one intends to apply an EOS-independent universal quadratic or cubic transformation formula to all spin periods, the residual error of the transformation is up to $0.068 M_{\odot}$. Again no advantage for the higher order formula. However, if one adopts the best-fit parameters for each spin period, the cubic transformation residual error is less than $0.01 M_{\odot}$, almost one order of magnitude better than the quadratic formula.
- For quadratic transformation, we find three EOS insensitive relations: $M_b = M_g + R_{1.4}^{-1} \times M_g^2$ for non-rotating NSs, $M_b = M_g + 0.78R_{1.4}^{-1} \times M_g^2$ for maximally rotating NSs, and more generally, $M_b =$

 $M_g + R_{1.4}^{-1} e^{-\frac{1}{4P}} \times M_g^2$ for rotating NSs with spin period $P = \mathcal{P} \times P_k$. In principle, the first two relations for non-rotating NSs (which may be also applied to slowly spinning NSs) and maximally rotating NSs could be used to give constraints on the NS radius $R_{1.4}$, and hence, the NS equation of state, once M_b and M_g for a NS could be independently measured [31]. It becomes less straightforward for the relation including spin, since two EOS related parameters $R_{1.4}$ and P_k are invoked and the NS spin period may be difficult to determine.

• With our calculations, we also find a very tight and general correlation between the normalized mass gain due to spin $\Delta m \equiv (M_{\rm max} - M_{\rm TOV})/M_{\rm TOV}$ and normalized spin period \mathcal{P} as $\log_{10}\Delta m = (-2.74 \pm 0.05)\log_{10}\mathcal{P} + \log_{10}(0.20 \pm 0.01)$, which is independent of EOS models. Note that this universal relation is valid only for rigidly rotating NSs.

In Table 3, we summarize all the transformation formulae between M_b and M_q proposed in previous works and in this work. For a fair comparison, we calculate the the maximum and average residual error of each transformation formula with the same $M_b - M_a$ sample (third column of Table 3). In general, the novel findings of this work compared with previous ones include: i) We study in detail how M_b and M_a are correlated for rapidly rotating NSs and derived the residual errors of the proposed transformation formulae; ii) The accuracy of our newly proposed universal relation between M_b and M_q is similar to the previous ones, but only $R_{1.4}$ is invoked in our relation instead of using the compactness number proposed in previous works [e.g., 31, 32]. As a result, our proposed relation is more practical since it does not involve the calculations of the NS radii corresponding to different NS masses; iii) For non-rotating NSs, if one adopts its own best-fit coefficients for individual EOSs (provided in Table 2), our proposed cubic transformation formula is much better than the quadratic formula, with the residual error of the transformation less than $0.01 M_{\odot}$.

It has been proposed that binary neutron star merger events could give tight constraints on the NS maximum mass, as long as we could calculate the mass of the merger remnant and justify whether the remnant is a BH or a long-lasting NS. For instance, it has been claimed that the multi-messenger observations of GW170817 could provide an upper bound on $M_{\rm TOV}$ [6–8], e.g., $M_{\rm TOV} \lesssim 2.16 M_{\odot}$ in Ref. [6]. However, this result sensitively depends on the assumption that the merger remnant of GW170817 is a not-too-long-lived hypermassive NS. This is still subject to debate, see supporting arguments by Margalit et al. [6-8] and counter opinion by Yu *et al.* [50-53]. This is due to the lack of GW detection in the post-merger phase and the insufficient capability to distinguish the merger product using electromagnetic (EM) observations only. If the low-significance flare-like feature at 155 days in the X-ray

	Non-rotating NS		Rotating NS	N EoSs	
	$M_b - M_g$ relation	Residual error ^A	$M_b - M_g$ relation	${\rm Residual}\;{\rm error}^{\rm A}$	
Lattimer & Yahil (1989) [20]	$M_b = M_g + 0.084 \times M_g^2$	4.1% (1.7%)	_	_	7
Timmes <i>et al.</i> (1996) [17]	$M_b = M_g + 0.075 \times M_g^2$	5.8% (1.5%) -		_	0^{B}
Lattimer & Prakash (2001) [31]	$ \begin{array}{l} (M_b-M_g)/M_g=0.6\beta/(1-0.5\beta), \\ \text{where } \beta=GM_g/Rc^2 \end{array} $	2.8% (0.92%)	_	-	14
Coughlin <i>et al.</i> (2017) [32]	$M_b/M_g = 1 + 0.89 \times (M_g/R)^{1.2}$	$2.6\% \ (0.56\%)$	_	_	19
	_	_	Maximally rotating NSs	-	_
This work	$M_b = M_g + R_{1.4}^{-1} \times M_g^2$	1.8% (1.1%)	$M_b = M_g + 0.078 R_{1.4}^{-1} \times M_g^2$	1.3%~(0.74%)	10
	$M_b = M_g + 0.080 \times M_g^2$	5.0%~(1.5%)	$M_b = M_g + 0.064 \times M_g^2$	3.3%~(1.2%)	10
	$M_b = M_g + A_1 \times M_g^2 + A_2 \times M_g^3$	0.45%~(0.16%)	$M_b = M_g + 0.056 \times M_g^2 + 0.003 \times M_g^3$	3.5%~(1.0%)	10
	_	-	General NSs with arbitrary rotation	-	-
	_	_	$M_b = M_g + R_{1.4}^{-1} \mathrm{e}^{-\frac{1}{4\mathcal{P}}} \times M_g^2$	3.3%~(1.0%)	10
	_	_	$M_b = M_g + 0.073 \times M_g^2$	6.0%~(1.6%)	10
	_	_	$M_b = M_g + 0.078 \times M_g^2 + 0.002 \times M_g^3$	4.0% (0.95%)	10

Table	3	$M_b - M_g$	relations	proposed	$_{\mathrm{in}}$	the	literature.
-------	---	-------------	-----------	----------	------------------	-----	-------------

^AThe maximum residual error of the transformation between M_b and M_g , with the average residual error showing in the brackets.

^BTimmes et al. [17] claims that the M_b-M_g relations used in their paper is based on private communications with J. M. Lattimer.

afterglow of GW170817 as claimed in Ref. [53] is true, or if the argument by Li *et al.* [51] that an additional energy injection from the merger product is required to interpret the blue component of AT2017gfo (GW170817 optical counterpart) is valid, the central remnant of GW170817 has to be a long-lived NS. The constraints on $M_{\rm TOV}$ would be reversed to a lower bound. Note that before GW170817 was detected, it was proposed that the statistical observational properties of Swift SGRBs favors NS EOS with $M_{\rm TOV}$ greater than 2.3 M_{\odot} [14, 15], if the cosmological NS mass distribution follows that observed in the BNS systems in our Galaxy.

In the case that the merger remnant of GW170817 was a long-lasting, rigidly rotating NS, the remnant gravitational mass could be estimated as follows: the total gravitational mass of the binary system is determined as $m_{g,1} + m_{g,2} = 2.74 M_{\odot}$ based on the inspiral phase GW signal [29]. The mass ratio of the binary is bound to 0.7–1 under the low dimensionless NS spin prior. We consider both the size and direction of the uncertainty. Without any prior for the NS EOS, the total baryonic mass could be estimated as²¹ $3.04 M_{\odot}$ by using the EOS-independent universal quadratic formula for non-rotating NSs (i.e., $M_b =$ $M_g + 0.08 \times M_g^2$). The uncertainty for the transformation is $[-0.05 M_{\odot}, +0.10 M_{\odot}]$. With the EM counterpart observations, the ejecta mass for GW170817 is estimated as

 $\sim 0.06 M_{\odot}$, with uncertainty in the order of $10^{-2} M_{\odot}$ [54, and reference therein]. In this case, the baryonic mass for the merger remnant could be estimated as $2.98M_{\odot}$, with conservative uncertainty ~ $[-0.06M_{\odot}, +0.11M_{\odot}]$. If the remnant is a rapidly rotating NS, although it is difficult to determine its initial spin period, we can transform its baryonic mass to the gravitational mass as $2.52 M_{\odot}$, by using the EOS-independent universal quadratic formula for arbitrary rotating NSs (i.e., $M_b = M_g + 0.073 \times M_q^2$), with the transformation uncertainty as $[-0.12M_{\odot}, +0.14M_{\odot}]$. Putting together all the uncertainties in the transformation and the ejecta mass, the gravitational remnant mass should be estimated as $M_{g,\text{rem}} = 2.52^{+0.25}_{-0.18}$. If we assume the initial spin period of the rigid rotating NS produced by NS-NS merger is always the Kepler period corresponding to its baryonic mass, the EOS-independent universal quadratic formula for maximally rotating NSs (i.e., $M_b = M_g + 0.064 \times M_q^2$) should be adopted. The gravitational remnant mass is thus estimated as $2.56 M_{\odot}$, with transformation uncertainty as $[-0.08M_{\odot}, 0.08M_{\odot}]$. Putting together all the uncertainties, the remnant mass should be estimated as $M_{g,\text{rem}} = 2.56^{+0.19}_{-0.14}$. Recently, Radice et al. [30] proposed that given an EOS, the initial spin period of the merger remnant could be estimated based on the value of its baryonic mass. In this case, if we have some strong prior for the NS EOS, the transformation uncertainty between M_b and M_g could be reduced to the order of $0.01 M_{\odot}$, if one applies the cubic transformation formula with the best fit values for a specific EOS with a specific initial spin period. The overall uncertainty

²⁾The total baryonic mass is highly insensitive to the binary mass ratio. For instance, the total baryonic mass for GW170817 is $3.04\pm 0.10 M_{\odot}$ for $m_{g,1}/m_{g,2} = 1$ and $3.05\pm 0.10 M_{\odot}$ for $m_{g,1}/m_{g,2} = 0.7$.

for this case is mainly contributed by the ejecta mass uncertainty, which might be further reduced in the future when a larger sample of EM counterparts for binary NS mergers are observed.

Acknowledgements This work was supported by the National Natural Science Foundation of China under Grant Nos. 11722324, 11603003, 11633001, 11690024, and 11873040, the Strategic Priority Research Program of the Chinese Academy of Sciences, Grant No. XDB23040100 and the Fundamental Research Funds for the Central Universities. A. B. acknowledges support by the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Programme under grant agreement No. 759253 and by the Sonderforschungsbereich SFB 881 "The Milky WaySystem" (Subproject A10) of the German Research Foundation (DFG).

References

- 1. J. M. Lattimer, The nuclear equation of state and neutron star masses, Annu. Rev. Nucl. Part. Sci. 62(1), 485 (2012)
- A. K. Harding, The neutron star zoo, *Front. Phys.* 8(6), 679 (2013)
- J. Antoniadis, P. C. C. Freire, N. Wex, T. M. Tauris, R. S. Lynch, M. H. van Kerkwijk, M. Kramer, C. Bassa, V. S. Dhillon, T. Driebe, J. W. T. Hessels, V. M. Kaspi, V. I. Kondratiev, N. Langer, T. R. Marsh, M. A. McLaughlin, T. T. Pennucci, S. M. Ransom, I. H. Stairs, J. van Leeuwen, J. P. W. Verbiest, and D. G. Whelan, A massive pulsar in a compact relativistic binary, *Science* 340(6131), 1233232 (2013)
- S. Lawrence, J. G. Tervala, P. F. Bedaque, and M. C. Miller, An upper bound on neutron star masses from models of short gamma-ray bursts, *Astrophys. J.* 808(2), 186 (2015)
- C. L. Fryer, K. Belczynski, E. Ramirez-Ruiz, S. Rosswog, G. Shen, and A. W. Steiner, The fate of the compact remnant in neutron star mergers, *Astrophys. J.* 812(1), 24 (2015)
- B. Margalit and B. D. Metzger, Constraining the maximum mass of neutron stars from multi-messenger observations of GW170817, Astrophys. J. 850(2), L19 (2017)
- L. Rezzolla, E. R. Most, and L. R. Weih, Using gravitational-wave observations and quasi-universal relations to constrain the maximum mass of neutron stars, *Astrophys. J.* 852(2), L25 (2018)
- M. Ruiz, S. L. Shapiro, and A. Tsokaros, GW170817, general relativistic magnetohydrodynamic simulations, and the neutron star maximum mass, *Phys. Rev. D* 97(2), 021501 (2018)
- A. Rowlinson, P. T. O'Brien, N. R. Tanvir, B. Zhang, P. A. Evans, N. Lyons, A. J. Levan, R. Willingale, K. L. Page, O. Onal, D. N. Burrows, A. P. Beardmore, T. N. Ukwatta, E. Berger, J. Hjorth, A. S. Fruchter, R. L. Tunnicliffe, D. B. Fox, and A. Cucchiara, The unusual X-ray emission of the short Swift GRB 090515: Evidence

for the formation of a magnetar? Mon. Not. R. Astron. Soc. 409(2), 531 (2010)

- A. Rowlinson, P. T. O'Brien, B. D. Metzger, N. R. Tanvir, and A. J. Levan, Signatures of magnetar central engines in short GRB light curves, *Mon. Not. R. Astron. Soc.* 430(2), 1061 (2013)
- H. J. Lü, B. Zhang, W. H. Lei, Y. Li, and P. Lasky, The millisecond magnetar central engine in short GRBs, *Astrophys. J.* 805(2), 89 (2015)
- P. D. Lasky, B. Haskell, V. Ravi, E. J. Howell, and D. M. Coward, Nuclear equation of state from observations of short gamma-ray burst remnants, *Phys. Rev. D* 89(4), 047302 (2014)
- V. Ravi and P. D. Lasky, The birth of black holes: neutron star collapse times, gamma-ray bursts and fast radio bursts, Mon. Not. R. Astron. Soc. 441(3), 2433 (2014)
- H. Gao, B. Zhang, and H. J. Lü, Constraints on binary neutron star merger product from short GRB observations, *Phys. Rev. D* 93(4), 044065 (2016)
- A. Li, B. Zhang, N. B. Zhang, H. Gao, B. Qi, and T. Liu, Internal X-ray plateau in short GRBs: Signature of supramassive fast-rotating quark stars? *Phys. Rev. D* 94(8), 083010 (2016)
- J. M. Lattimer and M. Prakash, Neutron star observations: Prognosis for equation of state constraints, *Phys. Rep.* 442(1–6), 109 (2007)
- F. X. Timmes, S. E. Woosley, and T. A. Weaver, The neutron star and black hole initial mass function, *Astrophys. J.* 457, 834 (1996)
- C. L. Fryer, K. Belczynski, G. Wiktorowicz, M. Dominik, V. Kalogera, and D. E. Holz, Compact remnant mass function: Dependence on the explosion mechanism and metallicity, *Astrophys. J.* 749(1), 91 (2012)
- O. Pejcha and T. A. Thompson, The landscape of the neutrino mechanism of core-collapse supernovae: Neutron star and black hole mass functions, explosion energies, and nickel yields, *Astrophys. J.* 801(2), 90 (2015)
- 20. J. M. Lattimer and A. Yahil, Analysis of the neutrino events from supernova 1987A, Astrophys. J. 340, 426 (1989)
- A. Burrows, D. Klein, and R. Gandhi, The future of supernova neutrino detection, *Phys. Rev. D* 45(10), 3361 (1992)
- J. Gava, J. Kneller, C. Volpe, and G. C. McLaughlin, Dynamical collective calculation of supernova neutrino signals, *Phys. Rev. Lett.* 103(7), 071101 (2009)
- 23. G. Camelio, A. Lovato, L. Gualtieri, O. Benhar, J. A. Pons, and V. Ferrari, Evolution of a proto-neutron star with a nuclear many-body equation of state: Neutrino luminosity and gravitational wave frequencies, *Phys. Rev. D* 96(4), 043015 (2017)
- 24. S. Rosswog, T. Piran, and E. Nakar, The multimessenger picture of compact object encounters: Binary mergers versus dynamical collisions, *Mon. Not. R. Astron. Soc.* 430(4), 2585 (2013)



- 25. A. Bauswein, S. Goriely, and H. T. Janka, Systematics of dynamical mass ejection, nucleosynthesis, and radioactively powered electromagnetic signals from neutron-star mergers, *Astrophys. J.* 773(1), 78 (2013)
- 26. K. Hotokezaka, K. Kiuchi, K. Kyutoku, H. Okawa, Y. Sekiguchi, M. Shibata, and K. Taniguchi, Mass ejection from the merger of binary neutron stars, *Phys. Rev. D* 87(2), 024001 (2013)
- 27. R. Fernández, D. Kasen, B. D. Metzger, and E. Quataert, Outflows from accretion discs formed in neutron star mergers: effect of black hole spin, *Mon. Not. R. Astron. Soc.* 446(1), 750 (2015)
- C. Y. Song, T. Liu, and A. Li, Outflows from black hole hyperaccretion systems: Short and long-short gamma-ray bursts and "quasi-supernovae", *Mon. Not. R. Astron. Soc.* 477(2), 2173 (2018)
- B. P. Abbott, et al. (LIGO Scientific Collaboration and Virgo Collaboration), GW170817: Observation of gravitational waves from a binary neutron star inspiral, *Phys. Rev. Lett.* 119, 161101 (2017)
- D. Radice, A. Perego, S. Bernuzzi, and B. Zhang, Longlived remnants from binary neutron star mergers, *Mon. Not. R. Astron. Soc.* 481(3), 3670 (2018)
- 31. J. M. Lattimer and M. Prakash, Neutron star structure and the equation of state, *Astrophys. J.* 550(1), 426 (2001)
- M. Coughlin, T. Dietrich, K. Kawaguchi, S. Smartt, C. Stubbs, and M. Ujevic, Toward rapid transient identification and characterization of kilonovae, *Astrophys. J.* 849(1), 12 (2017)
- 33. G. Bozzola, N. Stergioulas, and A. Bauswein, Universal relations for differentially rotating relativistic stars at the threshold to collapse, *Mon. Not. R. Astron. Soc.* 474(3), 3557 (2018)
- 34. A. M. Studzińska, M. Kucaba, D. Gondek-Rosińska, L. Villain, and M. Ansorg, Effect of the equation of state on the maximum mass of differentially rotating neutron stars, *Mon. Not. R. Astron. Soc.* 463(3), 2667 (2016)
- D. Gondek-Rosińska, I. Kowalska, L. Villain, M. Ansorg, and M. Kucaba, A New View on the Maximum Mass of Differentially Rotating Neutron Stars, *Astrophys. J.* 837(1), 58 (2017)
- 36. A. Bauswein and N. Stergioulas, Semi-analytic derivation of the threshold mass for prompt collapse in binary neutron-star mergers, *Mon. Not. R. Astron. Soc.* 471(4), 4956 (2017)
- 37. L. R. Weih, E. R. Most, and L. Rezzolla, On the stability and maximum mass of differentially rotating relativistic stars, *Mon. Not. R. Astron. Soc.* 473(1), L126 (2018)
- 38. E. M. Butterworth and J. R. Ipser, On the structure and stability of rapidly rotating fluid bodies in general relativity (I): The numerical method for computing structure and its application to uniformly rotating homogeneous bodies, *Astrophys. J.* 204, 200 (1976)
- N. Stergioulas and J. L. Friedman, Comparing models of rapidly rotating relativistic stars constructed by two numerical methods, *Astrophys. J.* 444, 306 (1995)

- 40. F. Douchin and P. Haensel, A unified equation of state of dense matter and neutron star structure, Astronomy and Astrophysics 380, 151 (2001)
- R. B. Wiringa, V. Fiks, and A. Fabrocini, Equation of state for dense nucleon matter, *Phys. Rev. C* 38(2), 1010 (1988)
- A. Akmal and V. R. Pandharipande, Spin-isospin structure and pion condensation in nucleon matter, *Phys. Rev.* C 56(4), 2261 (1997)
- S. Goriely, N. Chamel, and J. M. Pearson, Further explorations of Skyrme–Hartree–Fock–Bogoliubov mass formulas (XII): Stiffness and stability of neutron-star matter, *Phys. Rev. C* 82(3), 035804 (2010)
- 44. S. Typel, G. Röpke, T. Klähn, D. Blaschke, and H. H. Wolter, Composition and thermodynamics of nuclear matter with light clusters, *Phys. Rev. C* 81(1), 015803 (2010)
- H. Müther, M. Prakash, and T. L. Ainsworth, The nuclear symmetry energy in relativistic Brueckner-Hartree-Fock calculations, *Phys. Lett. B* 199(4), 469 (1987)
- H. Müller and B. D. Serot, Relativistic mean-field theory and the high-density nuclear equation of state, *Nucl. Phys. A.* 606(3–4), 508 (1996)
- G. B. Cook, S. L. Shapiro, and S. A. Teukolsky, Rapidly rotating neutron stars in general relativity: Realistic equations of state, *Astrophys. J.* 424, 823 (1994)
- 48. J. P. Lasota, P. Haensel, and M. A. Abramowicz, Fast rotation of neutron stars, *Astrophys. J.* 456, 300 (1996)
- C. Breu and L. Rezzolla, Maximum mass, moment of inertia and compactness of relativistic stars, Mon. Not. R. Astron. Soc. 459(1), 646 (2016)
- Y. W. Yu, L. D. Liu, and Z. G. Dai, A long-lived remnant neutron star after GW170817 inferred from its associated kilonova, Astrophys. J. 861(2), 114 (2018)
- 51. S. Z. Li, L. D. Liu, Y. W. Yu, and B. Zhang, What powered the optical transient AT2017gfo associated with GW170817? Astrophys. J. 861(2), L12 (2018)
- 52. S. Ai, H. Gao, Z. G. Dai, X. F. Wu, A. Li, B. Zhang, and M.Z. Li, The allowed parameter space of a long-lived neutron star as the merger remnant of GW170817, Astrophys. J. 860(1), 57 (2018)
- 53. L. Piro, E. Troja, B. Zhang, G. Ryan, H. van Eerten, R. Ricci, M. H. Wieringa, A. Tiengo, N. R. Butler, S. B. Cenko, O. D. Fox, H. G. Khandrika, G. Novara, A. Rossi, and T. Sakamoto, A long-lived neutron star merger remnant in GW170817: Constraints and clues from Xray observations, *Mon. Not. R. Astron. Soc.* 483(2), 1912 (2019)
- 54. B. D. Metzger, Welcome to the multi-messenger era! Lessons from a neutron star merger and the landscape ahead, arXiv: 1710.05931 (2017)