

1 **IMMUNOHISTOCHEMICAL EXAMINATION OF IMMUNE CELLS IN ADIPOSE**
2 **TISSUE OF RAINBOW TROUT (*ONCORHYNCHUS MYKISS*) FOLLOWING**
3 **INTRAPERITONEAL VACCINATION**

4

5 Kimberly A. Veenstra^a, Ayham Alnabulsi^b, Lincoln Tubbs^c, Juliette Ben Arous^d,
6 Christopher J. Secombes^a *

7

8 ^a Scottish Fish Immunology Research Centre, Institute of Biological and
9 Environmental Sciences, University of Aberdeen, Tillydrone Avenue, Aberdeen,
10 AB24 2TZ, UK

11

12 ^b Vertebrate Antibodies Ltd., Zoology Building, Tillydrone Avenue, Aberdeen, AB24
13 2TZ, UK

14

15 ^c Elanco Animal Health, Food Animal Vaccines R&D, 2500 Innovation Way,
16 Greenfield, IN 46140, USA

17

18

19 ^d Seppic, 50 Boulevard National, La Garenne-Colombes, Paris 92250, France

20

21

22

23 * Corresponding author at: Institute of Biological and Environmental Sciences,
24 University of Aberdeen, Tillydrone Avenue, Aberdeen AB24 2TZ, UK. Tel: +44
25 (0)1224 272872; fax: +44 (0)1224 274429.

26

27 [✉] Current address: Laboratory of Fish Immunology, Institute of Infectology, Friedrich-
28 Loeffler-Institut, Südufer 10, Greifswald-Insel Riems 17493, Germany

29

30 Email addresses: kimberly.veenstra@fli.de (K. Veenstra),
31 ayham.alnabulsi.09@aberdeen.ac.uk (A. Alnabulsi),
32 juliette.benarous@airliquide.com (J. Ben Arous), tubbs_lincoln@elanco.com (L.
33 Tubbs), c.secombes@abdn.ac.uk (C. Secombes).

34

35

36

37

38

39

40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72

Abstract

Mammalian perivisceral adipose has been shown to play an important role in the regulation of the peritoneal immune responses. Recently it has been demonstrated that peritoneal antigens are collected by leukocytes within the visceral adipose mass, and a broad range of immunomodulatory genes are differentially expressed in adipose tissue after intraperitoneal vaccination in rainbow trout. To assess the immune cell component in adipose, immunohistochemical analysis was used to examine B-cell, T-cell and antigen presenting cell (APC) numbers and distribution in rainbow trout adipose tissue 24 and 72 h post vaccination in comparison to control fish. The results of this study support previous work on mammals with omental milky spots in naïve fish found to contain APCs and T-cells which then increased in size, number and complexity following vaccination. It suggests that following peritoneal stimulation the visceral adipose mass in fish likely plays an important role in vaccine antigen uptake and presentation by APCs, as well as subsequent T-cell activation and differentiation.

Keywords:

- rainbow trout
- adipose tissue
- immunohistochemistry
- cell markers
- vaccination
- milky spots
- APC
- T-cell
- B-cell

73

74

75

76 **1. Introduction**

77

78 Oil-adjuvanted vaccines used in aquaculture are injected directly into the peritoneal
79 cavity, which in mammals and fish contains a wide range of immune cells (van Vugt
80 et al., 1996; Rapoport et al., 1999; Williams et al., 2010; Mathis et al., 2013). While the
81 resident cell population can vary between teleost species (Meseguer et al., 1993;
82 Tumbol et al., 2009), the composition in rainbow trout (*Oncorhynchus mykiss*) is
83 dominated by myeloid and lymphocyte cells (Afonso et al., 1997, 1998; Korytář et al.,
84 2013). The injection of vaccines (or other inflammatory agents) into the peritoneal
85 cavity of fish generates a rapid change in composition as well as an increase in the
86 number of cells present (Afonso et al., 1998, 2000; do Vale et al., 2002; Mutoloki et
87 al., 2006; Korytář et al., 2013; Noia et al., 2014; Brietzke et al., 2015), although foreign-
88 body inflammatory reactions can be maintained in the cavity for several months post-
89 vaccination in salmonids (Midtlyng, 1996a, 1996b; Poppe & Breck, 1997; Mutoloki et
90 al., 2004, 2010; Koppang et al., 2005; Evensen et al., 2005; Noia et al., 2014;
91 Villumsen et al., 2015).

92 Mammalian perivisceral adipose (also referred to as the omentum) has been
93 shown to influence and be influenced by adjacent and embedded lymphocytes, and
94 plays an important role in the regulation of peritoneal immune responses (Rangel-
95 Moreno et al., 2009). The visceral adipose mass is also capable of capturing bacteria
96 and other antigenic particulates from the peritoneal cavity (Cui et al., 2002; Ha et al.,
97 2006; Rangel-Moreno et al., 2009), and promoting immunity against them (Rangel-
98 Moreno et al., 2009). Immune cells and numerous pro-inflammatory, anti-inflammatory
99 and immune-modulating proteins and peptides (including cytokines) have been
100 identified in mammalian adipocytes (Rangel-Moreno et al., 2009; Schäffler &
101 Schölmerich, 2010; Chandra et al., 2011). Omental milky spots (MS) contain antigen
102 presenting cells (APCs), T- and B-cells and are thought to play a key role in the
103 transition of leukocytes from blood through the omentum to the peritoneal cavity
104 and back (Carlow et al., 2009).

105 Pignatelli et al. (2014) demonstrated that peritoneal antigens are collected by
106 leukocytes in rainbow trout visceral adipose. These leukocytes transcribe marker
107 genes for different leukocyte subpopulations, and are likely responsible for the
108 secretion of a range of immune cytokines (Pignatelli et al., 2014). The establishment
109 of a mature adipocyte phenotype has been shown to be associated with high activity
110 of immune genes in Atlantic salmon (*Salmo salar*) (Todorčević et al., 2010), and
111 teleost adipocytes have been shown to constitutively express pro-inflammatory
112 cytokines and genes relating to the interferon response (Todorčević et al., 2010;
113 Pignatelli et al., 2014). Alongside evidence demonstrating that rainbow trout visceral
114 adipose is capable of responding to viruses (Pignatelli et al., 2014), bacteria, and pro-
115 inflammatory cytokines (Veenstra et al., 2018), it can be concluded that teleost
116 adipose is an immunologically active tissue. Furthermore, the work of Veenstra et al.
117 (2017) established that a broad range of immunomodulatory genes are differentially
118 expressed in adipose tissue after intraperitoneal (ip) injection of oil-adjuvanted
119 bacterial vaccines and revealed a relationship between adipose tissue immune
120 function and the development of vaccine-induced adhesions.

121 Since it has been suggested that cellular mechanisms occurring immediately
122 post-vaccination within adipose tissue may contribute to the development of adhesions
123 and potentially be involved in the adaptive immune response (Veenstra et al., 2017),
124 in the present study we assessed immune cell distribution in rainbow trout visceral
125 adipose tissue following injection of an oil-adjuvanted vaccine into the peritoneal
126 cavity, using immunohistochemistry. The results of this work showed that MS in naïve
127 fish contain APCs and T-cells and that following an ip administration of oil-adjuvanted
128 vaccines MS increase in number, size and complexity and are associated with vaccine
129 remnants. Overall the results of this work suggest that the visceral adipose mass in
130 fish likely plays an important role in the uptake and presentation of vaccine antigens
131 and subsequent T-cell activation and differentiation following peritoneal stimulation.

132

133 **2. Methodology**

134

135 A total of 12 juvenile rainbow trout weighing approximately 60g (College Mill Trout
136 Farm, Perthshire, U.K.) were maintained in 400L tanks at the University of Aberdeen

137 aquarium facility supplied with recirculating freshwater at 14°C. Fish were fed *ad*
138 *libitum* daily with commercial pellets (EWOS) and were acclimated for at least two
139 weeks prior to vaccination. All trials were carried out in compliance with the Animals
140 (Scientific Procedures) Act 1986 by a UK Home Office license holder and approved
141 by the ethics committee at the University of Aberdeen. Fish were anaesthetised by
142 immersion with 2-phenoxyethanol (Fluka) and each fish injected intraperitoneally (ip)
143 with either 0.1 mL of phosphate buffered saline (PBS) or a water-in-oil adjuvanted
144 vaccine posterior to the pelvic girdle. The aqueous phase of the vaccine was a
145 formalin-killed whole-cell *A. salmonicida* bacterin (pre-inactivation titre of 1.55×10^9
146 cfu/mL) suspended in BHI Media and provided by Elanco Animal Health Ltd. (Victoria,
147 P.E.I., Canada) while the oil phase was comprised of Montanide™ ISA 761 VG
148 (Seppic, France). The water-in-oil emulsions was prepared at a 70:30 oil:water ratio
149 48h prior to vaccination using a high shear mixer (IKA Ultra Turrax Tube Drive) and
150 was tested for stability prior to use.

151 Visceral adipose located around the internal organs was harvested from freshly
152 killed trout (n=3 per treatment group per time point) at 24 and 72 h post injection (hpi).
153 These timings were chosen based on the previous study of Veenstra et al. (2017),
154 where the transcript response of immune genes was studied in adipose tissue at 3, 14
155 and 28 days post-vaccination. In that study gene modulation was already maximal at
156 day 3 in the majority of cases, and so here that timing was included together with an
157 earlier time point to assess whether changes were occurring before this. The tissue
158 was stored in Bouin's Solution (Sigma) for 18 h, washed 3x in PBS, then left in PBS
159 for 3-5 h. Samples were then stored in 70% ethanol (Sigma) before being embedded
160 in paraffin and sectioned at 5µm onto silane-coated glass slides (Microscopy and
161 Histology Core Facility, University of Aberdeen). Immunohistochemistry for each
162 antibody (Table 1) was performed using reagents from the REAL Dako Envision
163 detection kit (Dako UK Ltd) using a Dako autostainer (Dako) as described previously
164 (Alnabulsi et al., 2017) at the Department of Pathology, NHS Grampian Biorepository
165 (Aberdeen, UK). The antibodies used included a B (IgM) and T (CD3) cell marker, and
166 two markers of antigen presenting cells (APCs), MHC-II and CLEC4T1. In the case of
167 the APC markers CLEC4T1 is related to DC-SIGN (see discussion). Primary antibody
168 dilutions used for immunohistochemistry are described in Table 1. The sections were

169 evaluated by light microscopy using a Zeiss Axioscop 40 (Microscopy and Histology
170 Core Facility, University of Aberdeen).

171

172 **Table 1:** Antibodies used for immunohistochemical analysis.

Antibody	Type	Dilution	Dilutant	Reference/Source
IgM (4C10)	Monoclonal	1:5	Dako antibody dilutant	Thuvander et al., 1987
CD3- $\gamma\delta$	Monoclonal	1:15	Dako antibody dilutant	Vertebrate Antibodies Ltd
MHC-II β	Rabbit polyclonal	1:200	PBST*	Vertebrate Antibodies Ltd
CLEC4-T1	Rabbit polyclonal	1:500	Dako antibody dilutant	Johansson et al., 2016

173 * PBST = Phosphate Buffered Saline with Tween 20 (Sigma)

174

175 3. Results

176

177 The number of CLEC4T1, MHC-II, CD3 and IgM positive cells was found to vary
178 between treatment groups and time points. These results indicate that changes in
179 expression and distribution of APCs, T- and B-cells occur in rainbow trout adipose
180 tissue following vaccination.

181 CLEC4T1 staining was observed in areas analogous to the centre of clostridial
182 MS located on the periphery of the omentum tissue and in cells encircling apoptotic
183 adipocytes- crown like structures (CLS) in the naïve fish (Fig. 1A & 1C). At 24 and 72
184 hpi, large quantities of CLEC4T1 positive cells were observed infiltrating the adipose
185 tissue, primarily associated with areas of vaccine-induced cellular damage (Fig. 1B)
186 as well as strongly presenting within newly developed clostridial MS within the adipose
187 tissue (Fig. 1D).

188 MHC-II positive cells in naïve fish was observed in macrophage-like cells
189 located within adipocyte junctions and in some cell clusters (Fig. 2A & 2C), but were
190 not associated with CLS (Fig. 2C). The largest amount of anti-MHC-II staining was
191 observed at 24 hpi in the vaccinated group and was associated with granulomatous
192 cell clusters and areas of vaccine-infiltration (Fig. 2B). By 72 hpi the quantity of MHC-

193 Il positive cells decreased in the vaccinated group, but staining of small clusters of
194 mononuclear cells within the adipose tissue and associated with MS were still
195 apparent (Fig. 2D).

196 CD3 was detectable in the adipose tissue of naïve fish, in the cytoplasm of
197 single mononuclear cells found in cell clusters within adipose (Fig. 3A), and in some
198 structures analogous to clostridial MS found on the periphery of adipose tissue (Fig.
199 3C). In vaccinated fish at 24 hpi the staining appeared much stronger, and was present
200 in an increased number of peripheral MS, as well as newly developed clostridial MS
201 structures throughout the tissue (Fig. 3B). By 72 hpi staining was still clearly present
202 within MS of vaccinated fish, although weaker than seen in the 24 hpi fish. In
203 vaccinated fish, CD3 positive- MS were associated with CLS (Fig. 3D). The increase
204 in CD3 positive stained structures in cells located within milky spots can be observed
205 in greater detail in Figure 4.

206 IgM positive cells were not found in the control fish (Fig. 5A & 5C). However,
207 following vaccination cells staining positive for IgM could be observed within adipocyte
208 junctions at 24 hpi (Fig. 5B). Staining was still present but weaker at 72 hpi in individual
209 cells, occasionally associated with MS (Fig. 5D). Staining was also present within
210 blood vessels in the vaccinated groups, presumed to be soluble IgM in the blood (Fig.
211 5D).

212

213 **4. Discussion**

214

215 As teleost adipose has been found capable of sequestering antigens from the
216 peritoneal cavity (Pignatelli et al., 2014), and immune-related genes were
217 transcriptionally upregulated as early as 72 h after ip vaccination (Veenstra et al.,
218 2017), in this study we aimed to further characterize the relationship between vaccine-
219 induced stimulation of the peritoneal cavity and adipose immune cell response up to
220 72 h post injection via immunohistochemical analysis. The results of this study were
221 found to be broadly similar to what has been described previously in regards to
222 mammalian adipose clostridial milky spots (MS). MS associated with peritoneal
223 adipose tissue (omentum) have been described in a number of species (Mixer, 1941)
224 including fish (Pignatelli et al., 2014). They have been shown to contain macrophages,

225 APCs, T- and B-cells (Carlow et al., 2009) and to have important biological functions
226 within the peritoneal cavity (Beelen, 1991; Shimotsuma et al., 1993; Takemori et al.,
227 1995, Lenzi et al., 1996; van Vugt et al., 1996) and omentum (Carlow et al., 2009;
228 Rangel-Moreno et al., 2009), acting as a gateway through which circulating cells,
229 antigens, particulates and pathogens are collected from the peritoneal cavity to
230 promote a variety of immune responses (Beelen et al., 1980a, 1980b; Cranshaw &
231 Leak, 1990). Following stimulation in mammals, the increases in the number and size
232 of MS occur alongside an influx of leukocytes within MS (van Vugt et al., 1996), as
233 appeared to be happening in the current study. It is worth noting that viral stimulation
234 did not alter the size or number of MS in rainbow trout adipose (Pignatelli et al., 2014).

235 Dendritic cells (DCs) and macrophages are regarded as the key APCs of the
236 immune system and play an important role in the transition of innate immunity to
237 adaptive immunity. In mammals MS are considered to be the site of origin of peritoneal
238 macrophage precursors (Lee & Lee, 2014). An influx of macrophages into the
239 peritoneal cavity has been described in salmonids following stimulation (Afonso et al.,
240 1998; Jørgensen et al., 2008). C-type lectin (CLEC) domain family 4-T1 is a rainbow
241 trout transmembrane protein thought to be closely related to the well-characterised
242 CLEC4 family protein CD209 / DC-SIGN (Johansson et al., 2016). It, along with MHC
243 class-II proteins are found on DCs/ macrophages and help present extracellular
244 antigens to CD4 positive cells and to promote the rapid activation of T- and B-cells
245 (Carlow et al., 2009). The lack of MHC-II positive staining cells within MS supports
246 previous observations in trout (Pignatelli et al., 2014), however the presence of
247 CLEC4T1 positive cells within these structures demonstrates that APCs (potentially
248 DC or macrophage precursors) are present within MS in naïve fish, in accordance with
249 previous work on mice (Bertola et al., 2012). Additionally, crown-like structures (CLS),
250 described as clusters of macrophages surrounding dead adipocytes in obese
251 mammalian adipose (Murano et al., 2008; Noia et al., 2014), were observed to be
252 strongly CLEC4T1 positive in naïve and vaccinated rainbow trout. As the results in the
253 present study showed that there was little to no overlap in staining patterns of
254 CLEC4T1 and MHC-II, it indicates that within trout adipose tissue these markers are
255 expressed on distinct cell populations at these time points. The key function of
256 immature DCs is capturing and processing antigens which trigger full maturation, and
257 in time leads to the assembly of antigen-MHC-II complexes which are capable of

258 stimulating T-cells (Banchereau & Steinman, 1998; Geijtenbeek et al., 2000; Engering
259 et al., 2002). As it has been demonstrated that bacteria can stimulate DC maturation
260 (Sallusto & Lanzavecchia 1995; Winzler et al., 1997), it is likely that in teleosts APCs
261 preferentially begin production/maturation of CLEC4T1 to facilitate ingestion and
262 presentation of foreign substances with MHC-II complexes playing a larger role at a
263 later time point than studied here.

264 Lymphocytes are the second major cellular component of normal mammalian
265 MS (Shimotsuma et al., 1991, 1993; Krist et al., 1995). More recent studies (Rangel-
266 Moreno et al., 2009; Carlow et al., 2009) show that the omentum can support the
267 activation of CD4 and CD8 positive lymphocytes and mount T cell-dependent B-cell
268 responses to peritoneal antigens. CD3 is part of the T-cell receptor complex on the
269 cell surface which aids activation of naïve T cells (Guy & Vignali, 2009) and is in
270 rainbow trout considered a good pan-T-cell marker (Leal et al., 2016). The present
271 study reveals the presence of CD3 positive cells in clostridial MS on the periphery of
272 adipose tissue in naïve fish, in distinct areas separate to CLEC4T1 positive cells within
273 MS. An increase in staining intensity was observed at 24 hpi (which reduced by 72
274 hpi) in MS, which supports work in mammals showing that the omentum effectively
275 operates as a site for early antigen presentation, with a rapid turnover of lymphocytes
276 (Carlow et al., 2009). As CD3 positive MS were also found to be associated with
277 CLEC4T1 positive CLS, it strongly advocates that following vaccination APCs play a
278 large role in antigen uptake, presentation and subsequent T-cell activation in trout
279 adipose tissue MS.

280 Immunoglobulin (Ig) M is the most ancient and prevalent Ig in fish. It can be
281 expressed on the surface of B-cells or secreted as an antibody. In this study no
282 evidence of IgM positive staining in milky spots was observed, in agreement with work
283 on rainbow trout by Pignatelli et al. (2014) but in contrast to mammalian studies
284 (Rangel-Moreno et al., 2009). Pignatelli et al. (2014) identified IgM positive cells in the
285 interstitial space between adipocytes within visceral adipose and Ballesteros et al.
286 (2013) found that IgM transcript level could be increased in adipose in response to
287 oral vaccination. The present study found evidence of IgM positive cells in interstitial
288 spaces in naïve fish which increased in number following vaccination.

289

290 In conclusion, the immunohistochemical results of this paper show that naïve
291 teleost MS contain APCs (CLEC4T1 positive cells) and T-cells (CD3 positive cells).
292 Following the administration of an ip oil-adjuvanted vaccine, MS in rainbow trout
293 adipose increased in number, size and complexity and may play a significant role in
294 T-cell activation and differentiation via APCs.

295

296 **Acknowledgements**

297 KAV was supported by ELANCO and SEPPIC to undertake a PhD program at the
298 University of Aberdeen. KAV would like to thank Dr. Abdo Alnabulsi and Dr. Jason
299 Holland for their assistance with immunohistochemistry.

300

301 *Declarations of interest:* none

302

303 **References**

304 Afonso, A., Ellis, A. E. & Silva, M. T. (1997). The leucocyte population of the
305 unstimulated peritoneal cavity of rainbow trout (*Oncorhynchus mykiss*).
306 *Fish & Shellfish Immunology*, 7: 335-348.

307

308 Afonso, A., Lousada, S., Silva, J., Ellis, A. E. & Silva, M. T. (1998). Neutrophil and
309 macrophage responses to inflammation in the peritoneal cavity of rainbow trout
310 *Oncorhynchus mykiss*. A light and electron microscopic cytochemical study.
311 *Diseases of Aquatic Organisms*, 34: 27-37.

312

313 Afonso, A., Macedo, P. M., Ellis, A. E. & Silva, M. T. (2000). Glycogen granules in
314 resting and inflammatory rainbow trout phagocytes—an ultrastructural study.
315 *Diseases of Aquatic Organisms*, 42: 101-110.

316

317 Alnabulsi, A., Swan, R., Cash, B., Alnabulsi, A. & Murray, G. I. (2017). The
318 differential expression of omega-3 and omega-6 fatty acid metabolising enzymes in
319 colorectal cancer and its prognostic significance. *Br J Cancer.*, 116(12): 1612-1620.
320 doi: 10.1038/bjc.2017.135.

321

322 Ballesteros, N. A., Castro, R., Abos, B., Saint-Jean, S. S. R., Pérez-Prieto, S. I. &
323 Tafalla, C. (2013). The pyloric caeca area is a major site for IgM+ and IgT+ B-cell
324 recruitment in response to oral vaccination in rainbow trout. *PLOS ONE*, 8(6):
325 e66118. <https://doi.org/10.1371/journal.pone.0066118>

326

327 Banchereau, J. & Steinman, R. M. (1998). Dendritic cells and the control of
328 immunity. *Nature*, 392: 245–252. doi:10.1038/32588

329

330 Beelen, R. H. J., Fluitsma, D. M. & Hoefsmit, E. C. M. (1980a). The cellular
331 composition of omentum milky spots and the ultrastructure of milky spot
332 macrophages and reticulum cells. *J. Reticuloendothel. Soc.*, 28: 585-599.
333

334 Beelen, R. H. J., Fluitsma, D. M. & Hoefsmit, E. C. M. (1980b). Peroxidatic activity of
335 mononuclear phagocytes developing in omentum milky spots. *J. Reticuloendothel.*
336 *Soc.*, 28: 601-609.
337

338 Beelen, R. H. J. (1991). The greater omentum: physiology and immunological
339 concepts. *Neth J Surg*, 43: 145-149.
340

341 Bertola, A., Ciucci, T., Rousseau, D., Bourlier, V., Duffaut, C., Bonnafous, S., Blin-
342 Wakkach, C., Anty, R., Iannelli, A., Gugenheim, J., Tran, A., Bouloumié, A., Gual, P.
343 & Wakkach, A. (2012). Identification of adipose tissue dendritic cells correlated with
344 obesity-associated insulin-resistance and inducing Th17 responses in mice and
345 patients. *Diabetes*, 61(9): 2238-47. doi: 10.2337/db11-1274.
346

347 Brietzke, A., Korytář, T., Jaros, J., Köllner, B., Goldammer, T., Seyfert, H. M. &
348 Rebl, A. (2015). *Aeromonas salmonicida* infection only moderately regulates
349 expression of factors contributing to toll-like receptor signaling but massively
350 activates the cellular and humoral branches of innate immunity in rainbow trout
351 (*Oncorhynchus mykiss*). *Journal of Immunology Research*, 2015(901015): 1-16.
352 doi:10.1155/2015/901015
353

354 Carlow, D. A., Gold, M. R. & Ziltener, H. J. (2009). Lymphocytes in the peritoneum
355 home to the omentum and are activated by resident dendritic cells. *J Immunol.*,
356 183(2): 1155-65. doi:10.4049/jimmunol.0900409
357

358 Castro, R., Abos, B., Gonzalez, L., Granja, A. G. & Tafalla, C. (2017). Expansion and
359 differentiation of IgM+ B-cells in the rainbow trout peritoneal cavity in response to
360 different antigens. *Developmental and Comparative Immunology*, 70: 119-127.
361

362 Chandra, A., Srivastava, R. K., Kashyap, M. P., Kumar, R., Srivastava, R. N. & Pant,
363 A. B. (2011). The anti-inflammatory and antibacterial basis of human omental
364 defense: Selective expression of cytokines and antimicrobial peptides. *PLOS ONE*,
365 6(5): e20446. <https://doi.org/10.1371/journal.pone.0020446>
366

367 Cranshaw, M. L. & Leak, L. V. (1990). Milky spots of the omentum: a source of
368 peritoneal cells in the normal and stimulated animal. *Arch Histol Cytol*, 53(Suppl):
369 165-177.
370

371 Cui, L., Johkura, K., Liang, Y., Teng, R., Ogiwara, N., ... Sasaki, K. (2002).
372 Biodefense function of omental milky spots through cell adhesion molecules and
373 leukocyte proliferation. *Cell Tissue Res.*, 310: 321-330.
374

375 do Vale, A., Afonso, A. & Silva, M. T. (2002). The professional phagocytes of sea
376 bass (*Dicentrarchus labrax* L.): cytochemical characterisation of neutrophils and
377 macrophages in the normal and inflamed peritoneal cavity. *Fish & Shellfish*
378 *Immunology*, 13: 183-198.
379

380 Engering, A., Geijtenbeek, T. B. H., van Vliet, S. J., ...-van Kooyk, Y. (2002). The
381 dendritic cell-specific adhesion receptor DC-SIGN internalizes antigen for
382 presentation to T cells. *J. Immunol.*, 168(5): 2118-2126.
383 doi.org/10.4049/jimmunol.168.5.2118
384

385 Evensen, Ø., Brudeseth, B.E. & Mutoloki, S. (2005). The vaccine formulation and its
386 role in inflammatory processes in fish--effects and adverse effects. *Developments in*
387 *Biologicals*, 121: 117-25.
388

389 Geijtenbeek, T. B., Torensma, R., van Vliet, S. J.-Figdor, C. G. (2000).
390 Identification of DC-SIGN, a novel dendritic cell-specific ICAM-3 receptor that
391 supports primary immune responses. *Cell*, 100: 575-585.
392

393 Guy, C. S. & Vignali, D. A. (2009). Organization of proximal signal initiation at the
394 TCR:CD3 complex. *Immunol. Rev.*, 232(1): 7-21. doi: 10.1111/j.1600-
395 065X.2009.00843.x.
396

397 Ha, S., Tsuji, M., Suzuki, K., Meek, B., Yasuda, N., Kaisho, T. & Fagarasan, S.
398 (2006). Regulation of B1 cell migration by signals through Toll-like receptors. *Journal*
399 *of Experimental Medicine*, 203(11): 2541-2550.
400 <https://doi.org/10.1084/jem.20061041>
401

402 Johansson, P., Wang, T., Collet, B., Corripio-Miyar, Y., Monte, M. M., Secombes,
403 C. J. & Zou, J. (2016). Identification and expression modulation of a C-type lectin
404 domain family 4 homologue that is highly expressed in monocytes/macrophages in
405 rainbow trout (*Oncorhynchus mykiss*). *Dev Comp Immunol.*, 54(1): 55-65. doi:
406 10.1016/j.dci.2015.08.005.
407

408 Jørgensen, S. M., Afanasyev, S. & Krasnov, A. (2008). Gene expression analyses in
409 Atlantic salmon challenged with infectious salmon anemia virus reveal differences
410 between individuals with early, intermediate and late mortality. *BMC Genomics*, 9:
411 179. doi:10.1186/1471-2164-9-179
412

413 Krist, L. F., Eestermans, I. L., Steenbergen, J. J., Hoefsmit, E. C., Cuesta, M. A.,
414 Meyer, S. & Beelen, R. H. (1995). Cellular composition of milky spots in the human
415 greater omentum: an immunochemical and ultrastructural study. *Anat Rec*, 241: 163-
416 174.
417

418 Koppang, E. O., Haugarvoll, E., Hordvik, I., Aune, L. & Poppe, T. T. (2005). Vaccine-
419 associated granulomatous inflammation and melanin accumulation in Atlantic
420 salmon, *Salmo salar* L., white muscle. *Journal of Fish Diseases*, 28(1): 13-22.
421 <https://doi.org/10.1111/j.1365-2761.2004.00583.x>
422

423 Korytář, T., Jaros, J., Verleih, M., Rebl, A., Kotterba, G., Kühn, C., Goldammer, T. &
424 Köllner, B. (2013). Novel insights into the peritoneal inflammation of rainbow trout
425 (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology*, 35(4): 1192-1199.
426 <https://doi.org/10.1016/j.fsi.2013.07.032>
427

428 Leal, E., Granja, A. G., Zarza, C. & Tafalla, C. (2016). Distribution of T cells in
429 rainbow trout (*Oncorhynchus mykiss*) skin and responsiveness to viral infection.
430 *PLoS ONE* 11(1): e0147477. <https://doi.org/10.1371/journal.pone.0147477>
431

432 Lee, B-C. & Lee, J. (2014). Cellular and molecular players in adipose tissue
433 inflammation in the development of obesity-induced insulin resistance. *J Biochimica*
434 *et Biophysica Acta, Molecular Basis of Disease*, 1842(3): 446-462.
435 <https://doi.org/10.1016/j.bbadis.2013.05.017>
436

437 Lenzi, H. L., Oliveira, D. N., Pelajo-Machado, M., Borojevic, R. & Lenzi, J. A. (1996).
438 Coelom-associated lymphomyeloid tissue (milky spots): site of lymphoid and
439 myelomonocytic cell generation. *Braz J Med Biol Res*, 29: 19-24.
440

441 Mathis, D. (2013). Immunological goings-on in visceral adipose tissue. *Cell*
442 *Metabolism*, 17(6): 851–859. doi:10.1016/j.cmet.2013.05.008
443

444 Meseguer, J., Esteban, M. A., Munoz, J., Lopezruiz, A. (1993). Ultrastructure of the
445 peritoneal-exudate cells of seawater teleosts, seabream (*Sparus aurata*) and sea
446 bass (*Dicentrarchus labrax*). *Cell and Tissue Research*, 273: 301-307.
447

448 Midtlyng, P.J., Reitan, L.J. & Speilberg, L. (1996a). Experimental studies on the
449 efficacy and side-effects of intraperitoneal vaccination of Atlantic salmon (*Salmo*
450 *salar* L.) against furunculosis. *Fish & Shellfish Immunology*, 6(5): 335–350.
451 doi:10.1006/fsim.1996.0034
452

453 Midtlyng, P.J., Reitan, L.J., Lillehaug, A. & Ramstad, A. (1996b). Protection, immune
454 responses and side effects in Atlantic salmon (*Salmo salar* L.) vaccinated against
455 furunculosis by different procedures. *Fish & Shellfish Immunology*, 6(8): 599–613.
456 doi:10.1006/fsim.1996.0055
457

458 Mixer, R. L. (1941). On macrophagal foci (“milky spots”) in the pleura of different
459 mammals, including man. *Am J Anat*, 69: 159-186.
460

461 Murano, I., Barbatelli, G., Parisani, V., Latini, C., Muzzonigro, G., Castellucci, M. &
462 Cinti, S. (2008). Dead adipocytes, detected as crown-like structures, are prevalent in
463 visceral fat depots of genetically obese mice. (2008). *The Journal of Lipid Research*,
464 49: 1562-1568. doi: 10.1194/jlr.M800019-JLR200
465

466 Mutoloki, S., Alexandersen, S. & Evensen, Ø. (2004). Sequential study of antigen
467 persistence and concomitant inflammatory reactions relative to side-effects and
468 growth of Atlantic salmon (*Salmo salar* L.) following intraperitoneal injection with oil-
469 adjuvanted vaccines. *Fish & Shellfish Immunology*, 16(5): 633–644.
470 <https://doi.org/10.1016/j.fsi.2003.10.002>
471

472 Mutoloki, S., Reite, O.B., Brudeseth, B., Tverdal, A. & Evensen, Ø. (2006). A
473 comparative immunopathological study of injection site reactions in salmonids
474 following intraperitoneal injection with oil-adjuvanted vaccines. *Vaccine*, 24(5): 578–
475 588. doi:10.1016/j.vaccine.2005.08.070
476

477 Mutoloki, S., Cooper, G. A., Marjara, I. S., Koop, B. F. & Evensen, Ø. (2010). High
478 gene expression of inflammatory markers and IL-17A correlates with severity of
479 injection site reactions of Atlantic salmon vaccinated with oil-adjuvanted vaccines.
480 *BMC Genomics*, 11(1): 336. <https://doi.org/10.1186/1471-2164-11-336>
481

482 Noia, M., Domínguez, B., Leiro, J., Blanco-Méndez, J., Luzardo-Álvarez, A. &
483 Lamas, J. (2014). Inflammatory responses and side effects generated by several
484 adjuvant-containing vaccines in turbot. *Fish & Shellfish Immunology*, 38(1): 244–254.
485 doi:10.1016/j.fsi.2014.03.020
486

487 Pignatelli, J., Castro, R., González Granja, A., Abós, B., González, L., Jensen, L.B. &
488 Tafalla, C. (2014). Immunological characterization of the teleost adipose tissue and
489 its modulation in response to viral infection and fat-content in the diet. *PLoS ONE*,
490 9(10): e110920. doi:10.1371/journal.pone.0110920
491

492 Poppe, T. T. & Breck, O. (1997). Pathology of Atlantic salmon *Salmo salar*
493 intraperitoneally immunised with oil adjuvanted vaccine. A case report
494 *Diseases of Aquatic Organisms*, 29: 219-226.
495

496 Rangel-Moreno, J., Moyron-Quiroz, J. E., Carragher, D. M., Kusser, K., Hartson, L.,
497 et al. (2009). Omental milky spots develop in the absence of lymphoid tissue-inducer
498 cells and support B and T cell responses to peritoneal antigens. *Immunity* 30: 731–
499 743.
500

501 Rapoport, J., Hausmann, M. J. & Chaimovitz, C. (1999). The peritoneal immune
502 system and continuous ambulatory peritoneal dialysis. *Nephron*, 81: 373-380.
503

504 Sallusto, F. & Lanzavecchia, A. (1995). Dendritic cells use macropinocytosis and
505 the mannose receptor to concentrate antigen to the MHC class II compartment.
506 Downregulation by cytokines and bacterial products. *J. Exp. Med.*, 182: 389–400.
507

508 Schäffler, A. & Schölmerich, J. (2010). Innate immunity and adipose tissue biology.
509 *Trends in Immunology*, 31(6): 228–235. <https://doi.org/10.1016/j.it.2010.03.001>
510

511 Shimotsuma, M., Takahashi, T., Kawata, M. & Dux, K. (1991). Cellular subsets of the
512 milky spots in the human great omentum. *Cell Tissue Res*, 264: 599-601.
513

514 Shimotsuma, M., Shields, J. W., Simpson-Morgan, M. W., Sakuyama, A., Shirasu,
515 M., Hagiwara, A. & Takahashi, T. (1993). Morpho- physiological function and role of
516 omental milky spots as omentum-associated lymphoid tissue (OALT) in the
517 peritoneal cavity. *Lymphology*, 26: 90-101.
518

519 Takemori, N., Hirai, K., Onodera, R., Saito, N. & Namiki, M. (1995). Light and
520 electron microscope study of splenoportal milky spots in New Zealand black mice:
521 comparison between splenoportal milky spots and aberrant spleens. *J Anat*, 186:
522 287-299.
523

524 Thuvander, A., Hongslo, T., Jansson, E. & Sundquist, B. (1987). Duration of
525 protective immunity and antibody titres measured by ELISA after vaccination of
526 rainbow trout, *Salmo gairdneri* Richardson, against vibriosis. *J Fish Dis*, 10: 479-486.

527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568

Tumbol, R. A., Baiano, J. C. F. & Barnes, A. C. (2009). Differing cell population structure reflects differing activity of percoll-separated pronephros and peritoneal leucocytes from barramundi (*Lates calcarifer*). *Aquaculture*, 292: 180-188.

Todorčević, M., Škugor, S., Krasnov, A. & Ruyter, B. (2010). Gene expression profiles in Atlantic salmon adipose-derived stromo-vascular fraction during differentiation into adipocytes. *BMC Genomics* 11(1): 39.

van Vugt, E., Van Rijthoven, E. A., Kamperdijk, E. W. & Beelen, R. H. (1996). Omental milky spots in the local immune response in the peritoneal cavity of rats. *Anat Rec*, 244: 235-245.

Veenstra, K. A., Wang, T., Alnabulsi, A., Douglas, A., Russell, K. S., Tubbs, L., Ben Arous, J., Secombes, C. J. (2017). Analysis of adipose tissue immune gene expression after vaccination of rainbow trout with adjuvanted bacterins reveals an association with side effects. *Molecular Immunology*, 88: 89-98.
<https://doi.org/10.1016/j.molimm.2017.05.026>

Veenstra, K. A., Wangkahart, E., Wang, T., Tubbs, L., Ben Arous, J. & Secombes, C. J. (2018). Rainbow trout (*Oncorhynchus mykiss*) adipose tissue undergoes major changes in immune gene expression following bacterial infection or stimulation with pro-inflammatory molecules. *Developmental and Comparative Immunology*, 81: 83-94. DOI: [10.1016/j.dci.2017.11.001](https://doi.org/10.1016/j.dci.2017.11.001)

Villumsen, K., Koppang, E. O. & Raida, M. K. (2015). Adverse and long-term protective effects following oil-adjuvanted vaccination against *Aeromonas salmonicida* in rainbow trout. *Fish & Shellfish Immunology*, 42(1): 193-203.
<https://doi.org/10.1016/j.fsi.2014.09.024>

Walker, F. C. & Rogers, A. W. (1961). The greater omentum as a site of antibody synthesis. *Br J Exp Pathol.*, 42: 222-31.

Williams, J. C., Wagner, N. J., Earp, H. S., Vilen, B. J., Matsushima, G. K. (2010). Increased hematopoietic cells in the mertk^{-/-} mouse peritoneal cavity: a result of augmented migration. *J. Immunol.*, 184: 6637-6648.

Winzler, C., Rovere, P., Rescigno, M., Ricciardi-Castagnoli, P. (1997). Maturation stages of mouse dendritic cells in growth factor-dependent long-term cultures. *J. Exp. Med.*, 185: 317-328.

569 Figure Legends

570

571 **Figure 1:** Arrow heads (red) point to representative positive staining of CLEC4T1 in
572 rainbow trout adipose tissue. A: 24 hpi unvaccinated; B: 24 hpi vaccinated; C: 72 hpi
573 unvaccinated; D: 72 hpi vaccinated (star = vaccine remnant).

574

575 **Figure 2:** Arrow heads (red) point to representative positive staining of MHC-II in
576 rainbow trout adipose tissue. A: 24 hpi unvaccinated; B: 24 hpi vaccinated; C: 72 hpi
577 unvaccinated; D: 72 hpi vaccinated (star = vaccine remnant).

578

579 **Figure 3:** Arrow heads (red) point to representative positive staining of CD3- $\gamma\delta$ in
580 rainbow trout adipose tissue. A: 24 hpi unvaccinated; B: 24 hpi vaccinated; C: 72 hpi
581 unvaccinated; D: 72 hpi vaccinated.

582




583 **Figure 4:** CD3- $\gamma\delta$ positive stained cells in a rainbow trout adipose tissue milky spot
584 at 24 h post-vaccination.





585





586 **Figure 5:** Arrow heads (red) point to representative positive staining of IgM in
587 rainbow trout adipose tissue. Arrow head (black) show staining in blood vessels. A:
588 24 hpi unvaccinated; B: 24 hpi vaccinated; C: 72 hpi unvaccinated; D: 72 hpi
589 vaccinated.


590

591









20 μm

