

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

1-6-2020

Adaptive capacity in ecosystems

David G. Angeler

Hannah Fried-Petersen

Craig R. Allen

Ahjond Garmestani

Dirac L. Twidwell Jr

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Botany Commons](#), [Horticulture Commons](#), [Other Plant Sciences Commons](#), and the [Plant Biology Commons](#)

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

David G. Angeler, Hannah Fried-Petersen, Craig R. Allen, Ahjond Garmestani, Dirac L. Twidwell Jr, H.E. Birgé, W. Chuang, V.M. Donovan, T. Eason, C.P. Roberts, S.M. Sundstrom, and C.L. Wonkka



EPA Public Access

Author manuscript

Adv Ecol Res. Author manuscript; available in PMC 2020 January 06.

About author manuscripts

Submit a manuscript

Published in final edited form as:

Adv Ecol Res. 2019 ; 60: 1–24. doi:10.1016/bs.aecr.2019.02.001.

Adaptive capacity in ecosystems

David G. Angeler¹, Hannah Fried-Petersen¹, Craig R. Allen², Ahjond Garmestani³, Dirac Twidwell⁴, H.E. Birgé⁵, W. Chuang³, V.M. Donovan⁴, T. Eason³, C.P. Roberts⁴, S.M. Sundstrom⁵, C.L. Wonkka⁴

¹Swedish University of Agricultural Sciences, Department of Aquatic Sciences and Assessment, Box 7050, SE-750 07 Uppsala, Sweden

²U.S. Geological Survey, Nebraska Cooperative Fish and Wildlife Research Unit, University of Nebraska -- Lincoln, School of Natural Resources, Lincoln, Nebraska, USA

³National Risk Management Research Laboratory, U.S. Environmental Protection Agency, Cincinnati, OH 45268, USA

⁴University of Nebraska, Department of Agronomy & Horticulture, Keim Hall, Lincoln, Nebraska, USA 66583-0915

⁵University of Nebraska -- Lincoln, School of Natural Resources, Lincoln, Nebraska, USA

Summary

Understanding the adaptive capacity of ecosystems to cope with change is crucial to management. However, unclear and often confusing definitions of adaptive capacity make application of this concept difficult. In this paper, we revisit definitions of adaptive capacity and operationalize the concept. We define adaptive capacity as the latent potential of an ecosystem to alter resilience in response to change. We present testable hypotheses to evaluate complementary attributes of adaptive capacity that may help further clarify the components and relevance of the concept. Adaptive sampling, inference and modeling can reduce key uncertainties incrementally over time and increase learning about adaptive capacity. Such improvements are needed because uncertainty about global change and its effect on the capacity of ecosystems to adapt to social and ecological change is high.

Keywords

adaptive capacity; resilience; management; ecological memory; ecological traits; global change

Introduction

Future global environmental sustainability requires research that integrates human-nature interactions with sustainable practices to foster ecosystem regimes that are desirable (Kates et al. 2011). Ecosystems are subject to stresses (increasing intensification of agriculture, increasingly over-appropriated water supplies, and climate change), and these stresses are

dangerously approaching the planetary boundaries of sustainable use of natural resources (Rockström et al. 2008). The ability of ecosystems to adapt to these changes is limited. Eventually, ecosystems may undergo *regime shifts* (for definition of terms in italics see Box1) to alternate species assemblages and ecosystem functioning at local, regional, and even global scales (Hughes et al. 2013). The outcomes of regime shifts are highly uncertain, potentially having substantial negative effects on human health, security and welfare (Horner-Dixon 1991; McMichael et al 2008). Therefore, it is important to determine the capacity of ecosystems to adapt to swiftly-changing social-ecological baselines towards a future without historical analogue, and how management and conservation can contribute to this adaptation.

The concept of *adaptive capacity* has been rapidly assimilated in the social sciences and transdisciplinary social-ecological research (Gunderson 2000; Folke et al. 2003), with multiple attempts made to formalize its meaning. Adaptive capacity is related to resilience (Holling 1973) and panarchy (Gunderson and Holing 2002), which has taken center stage in the effort to understand ecosystem dynamics during change. The concept of adaptive capacity has, in parallel with the transdisciplinary development of resilience theory, helped to diversify the meanings and definitions of systems undergoing change (Gallopín 2006). Adaptive capacity has been mainly used qualitatively in climate change, vulnerability and a risk/disaster management context in the social sciences and varies between different contexts and systems (Adger et al. 2007). Similarly, in the ecological sciences, *adaptation*, *adaptedness*, *adaptability* and *adaptive capacity*, terms with different meanings, have often been used interchangeably (Gallopín 2006, Smit and Wandel 2006). Consequently, operationalizing the concept of adaptive capacity, and by extension resilience theory for application and management, has been difficult, because of a loss of clarity and loose, incorrect and often normative use of these disparate concepts (Brand and Jax 2007; Angeler and Allen 2016). Misuse of terms can have significant negative impacts, because resilience and adaptive capacity are being used to help guide responses to natural disasters. Further, assessments of ecosystems that drive international research priorities depend on a comprehensive understanding of these concepts (Smit and Wandel 2006).

Because the concept of adaptive capacity is muddled with multiple meanings, its current use often makes it indistinguishable from resilience. In this paper, our goal is to clearly define the concept of adaptive capacity in ecosystems with the aim of differentiating it from similar concepts, particularly *ecological resilience*. Since approaches for operationalization and quantification of these concepts are needed, we describe components of adaptive capacity in ecosystems and discuss how they might mitigate and direct ecological response to ongoing environmental change. Further, we identify a research agenda to test hypotheses related to adaptive capacity and the ability of ecosystems to cope with environmental change.

Definitions and formalization

Much of the terminology and definitions used in the ecological adaptive capacity context has a Darwinian adaptation focus on species and populations. This is reflected in the currently most comprehensive definition of adaptive capacity (Beever et al. 2015; Nicotra et al. 2015). These authors define adaptive capacity for species and populations as a combination of

evolutionary potential, dispersal ability, life-history traits and phenotypic plasticity, which are influenced by genetic, epigenetic, and behavioral and acclimation processes.

This definition is well aligned with the broad use of the term adaptation in ecology, which is defined as an organism's ability to cope with environmental changes in order to survive and reproduce (Smit and Wandel 2006). The term adaptation itself is often used interchangeably with the term *adaptability*, which, as defined in biology, means the ability to become adjusted and to live and reproduce under a certain range of environmental conditions (Conrad 1983). Another term, *adaptedness*, has a more specific meaning than adaptation or adaptability. Dobzhansky (1968) defined adaptedness as the adaptive traits (structure, function and behavior of an organism) that are crucial for an organism to thrive in an environment. Adaptedness embraces species- or population-specific adaptation to a certain range of environmental conditions. Adaptedness is therefore context dependent and not a generic property as adaptability or adaptation would suggest. That is, high adaptedness does not necessarily mean high adaptability because a species may be highly adapted to a special and constant environment but have little capacity to adapt to other environments or to changes in its environment (Gallopín 2006). For example, a cold-stenothermic mayfly may thrive in an arctic stream, but it does not have the necessary adaptation to live in tropical lakes or to keep pace with warming of the arctic stream environment. Adaptedness can be tested through reciprocal transplant experiments to assess phenotypic fitness to local ecological niches.

Despite the dominant focus of adaptive capacity on lower levels of biological hierarchical organization in the literature, the term is increasingly used as an ecosystem property, recognizing that the ability of ecosystems to cope with disturbances is limited and that regime shifts can occur. Gunderson (2000) defined adaptive capacity as a system property, where adaptive capacity modifies ecological resilience (or "*basin of attraction*"). This definition is very similar to the earlier definition of ecological resilience: "Resilience is a measure of the persistence of systems and of their ability to absorb change and disturbance and still maintain the same relationships between populations or state variables" (Holling 1973).

Underlying ecological resilience is the capacity of ecosystems to undergo regime shifts, meaning that ecosystems can exist in more than one regime (Holling 1973). Gunderson and Holling (2002) defined ecological resilience as "the magnitude of disturbance that can be absorbed before the system changes its structure by changing the variables and processes that control behavior". Similarly, in a recent overview of resilience definitions, Angeler and Allen (2016) refer to ecological resilience as "a measure of the amount of change needed to change an ecosystem from one set of processes and structures to a different set of processes and structures".

Ecological resilience encompasses broader systems dynamics by considering both adaptation within, and shifts between, alternative basins of attraction (i.e., *alternative regimes*). The distinction between single *vs* multiple regimes helps distinguish adaptive capacity from ecological resilience. Adaptive capacity focuses on dynamics within a specific regime, and therefore adaptive capacity is a subset of ecological resilience, which is

explicitly concerned with dynamics both within and between regimes. Thus, similar to the view of Gunderson (2000), ecosystem adaptive capacity can be formalized and defined as follows (Figure 1, Box 1): *Adaptive capacity is the latent potential of an ecosystem to alter resilience in response to change*. In particular, adaptive capacity is the capability of an ecological system or other complex system to alter its basin of attraction in response to change such that the current regime is maintained.

Considering adaptive capacity as a subset of ecological resilience has applied relevance. Research is increasingly focused on the assessment of early warning signals of impending regime shifts (e.g., Dakos et al. 2015; Spanbauer et al. 2014, 2016), with the goal of employing management intervention if appreciable signals are detected (Batt et al. 2016). Although such studies implicitly consider an exhaustion of adaptive capacity, underlying mechanisms are not fully accounted for. However, scrutinizing adaptive capacity may provide such a mechanistic understanding. We discuss components of adaptive capacity and forward hypotheses to test these.

Components of adaptive capacity

Adaptive capacity as a latent potential of ecosystems is comprised of components that are dynamically interlinked (Table 1):

Ecological memory

The composition and distribution of organisms, their interactions in space and time and their life-history experience with environmental fluctuations contribute to *ecological memory* (Nyström and Folke 2004). Ecological memory has been defined as “the capability of the past states or experiences of a community to influence the present or future ecological responses of the community” (Zhong-Yu and Hai 2011). Specifically, ecological memory comprises all structural and functional features of ecological communities, which have been shaped by the interaction of past disturbances (natural and anthropogenic), spatial aspects (dispersal, habitat connectivity), biological interactions (competition, predation), evolutionary (speciation, extinctions, anagenesis, random mutations) and phylogenetic processes. This memory of ecological communities allows for a “learning process” (Carpenter et al. 2001). From this learning at the community level patterns and processes emanate, which enable ecosystems to prepare for and respond to future disturbances. This highlights that aspects of ecological memory compartmentalize by scales of space and time. Adaptive capacity explicitly accounts for pattern-process relationships of ecological memory that operate within and across the hierarchy of biological organization (i.e., they contribute to *cross-scale resilience*) (Table 1).

Cross-scale interactions

Ecosystems are hierarchically organized and have distinct patterns of structure, function, and processes that are compartmentalized by spatiotemporal scales. Considering cross-scale interactions is important because the impact of disturbance in ecosystems can be scale-specific (Pickett & White, 1985; Nash et al., 2014). That is, if disturbances affect components of ecological memory at one scale, other components at other scales might

buffer the disturbances in the entire ecosystem to maintain functioning and resilience (Peterson *et al.*, 1998; Allen et al., 2005; Allen & Holling 2008). This buffering ability is critical to the understanding of the latent potential of adaptive capacity because it can be expected that adaptive capacity to absorb disturbances and maintain ecosystem regimes increases with the buffering ability conferred through cross-scale interactions. This buffering ability can be further explored through assessments of functional ecosystem characteristics.

Ecological functioning

Ecosystem reactions to disturbances, including buffering, rely on functional responses to perturbations, which in turn depends on the diversity of traits (e.g., reproductive phenology, seed bank potential, colonization and dispersal abilities [*functional diversity*]) that provide a range of response patterns to disturbances (i.e., *response diversity*) (Elmqvist et al., 2003). A recent study on coral-reefs (Nash et al. 2016) and a meta-analysis of forest resilience (Cole et al. 2014) support the importance of response diversity and cross-scale resilience after disturbances. Additionally, response-effect trait patterns and ecological network structure influence response diversity and ecosystem service provisioning (Mori et al. 2013, Oliver et al. 2015, Schleuning et al. 2015). In addition to diversity, redundancies of functional traits (*functional redundancy*) are important to stabilize processes (e.g., primary production, decomposition) and feedbacks, and therefore contribute to the resilience of an ecosystem (Folke et al. 2004). Assessing the distribution, diversity and redundancy of functional traits within and across spatiotemporal scales can therefore be used as a measurable surrogate for adaptive capacity, and may provide an indicator of the erosion of adaptive capacity as a result of environmental change (Laliberté et al. 2010). Important in such assessments is the consideration of rare species.

Rare species

Mouillot et al. (2013) found that rare species in alpine meadows, coral reefs, and tropical forests supported functional trait combinations that were not represented by abundant species. This suggests that if rare species go extinct with ongoing environmental change, negative effects on ecosystem processes may ensue with a subsequent loss of adaptive capacity. Such effects may occur even if biodiversity associated with abundant species is high (Mouillot et al. 2013).

The importance of rare species is also evident in their ability to replace dominant species after perturbation and maintain ecological functions in the system, which in turn contributes to ecological resilience (Walker et al. 1999). For instance, rare shrub species with larger root crowns than dominant species were able to compensate for the loss of dominant shrub species to mechanical disturbance by re-sprouting prolifically, thus maintaining a shrub-dominated system despite disturbance (Wonkka et al. 2016). This example shows that rare species may contribute an important but, to some extent, unpredictable degree of adaptive capacity to ecosystem change.

Assessing adaptive capacity

The integration of scales, and functional and structural features between abundant and rare species offers a means to assess adaptive capacity. Resilience assessments have used discontinuity approaches to objectively identify the scaling structure present in ecosystems (Angeler *et al.*, 2016). The discontinuity analyses have so far shown promising results in assessing resilience of aquatic and terrestrial ecosystems (Angeler *et al.* 2016) and also other complex systems (economic, anthropological, social-ecological; Garmestani *et al.* 2005; Garmestani *et al.* 2009; Sundstrom *et al.* 2014). Discontinuity analysis may therefore also be useful in assessing the adaptive capacity of ecosystem regimes. The implementation of this approach will be examined from an adaptive capacity assessment point of view.

Because information about ecosystems is frequently limited adaptive capacity can be assessed following a recently proposed hypothesis-testing framework for quantifying ecological resilience (Baho *et al.* 2017). This evaluation comprises initial assessments of specific facets of adaptive capacity and then tests and recalibrates hypotheses iteratively to increase knowledge and provide learning opportunities about its general adaptive capacity.

Surrogates of adaptive capacity can be evaluated using simple measures of ecological *stability* (*resistance, persistence, variability, and engineering resilience*) (Donohue *et al.* 2013), biodiversity (Magurran 2004), and resilience (Angeler *et al.* 2016). The stability aspects can be evaluated for structural and functional variables (e.g., diversity, abundance, evenness, community composition, functional redundancies and diversity and process rates) within and across scales.

The initial step for quantifying adaptive capacity builds on Carpenter *et al.* (2001) to test for the “adaptive capacity of what to what”. However, testing for specific aspects of adaptive capacity may not be representative of the general adaptive capacity of an ecosystem. This is because there is limited surrogacy of metrics when assessing ecological responses to stressors (Johnson and Hering 2009). In addition, focusing on specified adaptive capacity can be problematic because managing adaptive capacity of particular parts of an ecosystem, especially in terms of managing for predictable outcomes of disturbances or provision of ecosystem services, may cause the system to lose adaptive capacity or resilience in other ways (Carpenter *et al.*, 2015). Specified assessments of adaptive capacity shall therefore be regarded as an initial step towards assessing the broader systemic or general adaptive capacity of an ecosystem.

It follows that assessing and managing for general adaptive capacity will require the simultaneous assessment of a range of variables to cover generic system properties and create possibilities for integral, resilience-based ecosystem management, which is difficult for most ecosystems.

Hypothesis testing to clarify adaptive capacity concepts

We suggest that this problem can be overcome by implementing adaptive monitoring and management. For this purpose, posing hypotheses that test premises of adaptive capacity are helpful (Table 2). We propose hypotheses that are not mutually exclusive and are well

aligned with our adaptive capacity definition, allowing for the evaluation of its attributes in a logical, iterative sequence. These hypotheses can be tested using quantifiable stability and resilience measures (Angeler *et al.*, 2016) based on multiple lines of evidence (e.g., taxa across distinct trophic levels; Burthe *et al.*, 2016). Most hypotheses can be framed specifically from a management perspective to facilitate the quantification of adaptive capacity without sacrificing the complexity inherent in management-related assessments. Also, most of our proposed hypotheses are supported by empirical observations (examples in Table 2), suggesting implementation of our quantification framework with ecological realism.

Hypothesis testing is carried out sequentially, which first objectively identifies key species in ecosystems that might serve as sentinels of system change (Angeler *et al.* 2016a). Next, sampling can be adapted to select appropriate spatial and/or temporal scales for monitoring to account for the cross-scale structure present in the system (Thompson and Seber 1996). This can contribute to pattern identification following population responses of sentinel species to disturbances. Incorporation of genetic, evolutionary, molecular and physiological variables and the measurement of process rates in monitoring can increase inference about ecosystem change, providing information for recalibration of management hypotheses. Monitoring can be refined by subsequently recalibrating hypotheses in an adaptive process that first focuses on reducing Type II errors (identifying false negatives) prior to reduce uncertainty sufficiently such that subsequent analyses can focus on Type I error (identifying false positives) reduction (adaptive inference, Holling and Allen 2002). Type II errors can be reduced by assessing adaptive capacity attributes (e.g., cross-scale and within scale structure and associated functional diversity redundancy) when ecological information of the ecosystem is limited. This can be done with the analysis of temporal snapshots, which are often the only resource available to managers. Subsequently, monitoring can be designed, implemented and sequentially modified to successively reduce Type I errors; that is, by improving knowledge of a broader range of adaptive capacity characteristics that need to be sampled over time (e.g., how fast is recovery after a disturbance). Such recalibrations can target functional assessments of sentinel species to change and, in further iterations, be extended to other taxa. This type of hypothesis testing builds on adaptive management (Allen *et al.*, 2011), sampling (Thompson and Seber 1996), modeling (Uden *et al.*, 2015) and inference (Holling & Allen, 2002). It allows revealing, refining, understanding and ultimately managing general ecosystem adaptive capacity, while increasing learning and reducing uncertainty. In this process, experiments can be designed that sequentially recalibrate strategies based on the outcomes of previous experiments and from which decisions about further data generation and monitoring can be made (Figure 2).

Managing adaptive capacity

Our suggested hypotheses are very general at this stage, but they can provide an initial step to inform management. First, managing for adaptive capacity may help maintain ecosystems in a regime desirable for humans (Allen *et al.* 2011). In this case maintaining adaptive capacity is fundamental for managing ecosystems away from critical thresholds (Batt *et al.* 2016). Crucial to managing for adaptive capacity is the consideration of cross-scale interactions across hierarchical levels and temporal scales. Upholding our premises while

testing hypotheses iteratively can be useful for designing management interventions to foster the adaptive capacity of a specific desired regime. A combination of ecological and technological approaches might be necessary to this end (Rist et al. 2014). Some of these approaches, e.g. assisted translocations or introductions of species invasions to compensate for lost crucial functions in an ecosystem (Chaffin et al. 2016b), are debated and thus potentially limited in their application to management. Second, when an ecosystem is in an undesired regime, management can reduce adaptive capacity, induce a shift towards a more desired regime and foster the adaptive capacity of this new regime (Chaffin et al. 2016a) (Figure 2). There is, for instance, a rich body of literature on lake biomanipulation, which exemplifies transformation of degraded lakes into more desirable systems (Hansson et al. 1998).

The biomanipulation example is useful because while it offers possible management options, it also highlights potential limitations when managing for adaptive capacity. Lake biomanipulation has adopted a series of management interventions, based on ecological (food web manipulations) and technological interventions (water column aeration, sediment dredging or lining, nutrient precipitation). However, lessons from biomanipulation have shown that these solutions can incur short- to long-term costs that may not be tenable for most systems.

It is clear that a series of ecological, resource and ethical issues may currently complicate the translation of a solid body of theory on adaptive capacity to its management on the ground. Current environmental policy further limits the implementation of resilience to management (Green et al. 2015).

Conclusion

This paper suggests a way forward to enhance our ability to explicitly define and reduce uncertainties and promote more holistic and effective modeling, management and monitoring of adaptive capacity. In addition to testing premises and hypotheses of adaptive capacity, defining highly-related concepts will aid in continuing toward operationalizing adaptive capacity. For instance, although we have defined how adaptive capacity relates to the width of the basin of attraction in the iconic ball-and-cup heuristic, methods for defining the actual basin of attraction are few and lack rigorous testing (Gunderson et al. 2000). Estimating the basin of attraction would allow estimates of adaptive capacity to move beyond point estimates comparable only between subsequent measures and toward a direct estimate of adaptive capacity relative to the system's potential adaptive capacity (Carpenter et al. 2001). Defining the basin of attraction would also allow the buffer that the current level of adaptive capacity provides against system transformations to be estimated, and it would allow transformative elements (e.g. invasive species) to be distinguished from elements that contribute to adaptive capacity (e.g. rare species; Elmqvist et al. 2003, Folke et al. 2010). However, new methods for detecting spatial regimes and discontinuous resource aggregations show promise for delineating basins of attraction in space and over time (Angeler et al. 2016, Allen et al. 2016, Sundstrom et al. 2017).

Uncertainty will not be eliminated completely or immediately, but it can be reduced incrementally while an ecosystem is monitored, modeled and managed over time. Explicit learning during this process can overcome common management problems, such as delayed action under uncertainty (Conroy et al., 2011), prioritization of limited financial resources (Stewart-Koster, Olden & Johnson, 2015), and the limited coordination in governance of natural resources (Cumming et al., 2013). Such improvements are needed because of uncertainty about global change impacts on an ecosystem's ability to absorb disturbances. An improved understanding of adaptive capacity can ultimately help to facilitate ecosystem management within current ecological, economic and ethical constraints. Our approach to assess adaptive capacity provides insight into the challenges to account for ecological complexity in ecosystem management. It particularly highlights enormous resource needs to the practical implementation and pinpoints persistent problems for closing gaps between science, policy, and management (Garmestani and Benson 2013).

Acknowledgements

This study was funded through the August T. Larsson Foundation of the Faculty of Natural Resources of the Swedish University of Agricultural Sciences and grants from the Swedish Research Councils FORMAS (2014-1193) and VR (2014-5828), and The United States Department of Defence, SERDP (RC-2510). The Nebraska Cooperative Fish and Wildlife Research Unit is jointly supported by a cooperative agreement between the U.S. Geological Survey, the Nebraska Game and Parks Commission, the University of Nebraska-Lincoln, the United States Fish and Wildlife Service and the Wildlife Management Institute. The views expressed in this paper are those of the authors and do not necessarily represent the views or policies of the U.S. Environmental Protection Agency. This research was performed while WC held NRC Research Associateship awards at the U.S. EPA.

References

- Adger WN, Agrawala S, Mirza MMQ, Conde C, O'Brien K, Pulhin J, Pulwarty R, Smit B, Takahashi K (2007) Assessment of adaptation practices, options, constraints and capacity. *Climate Change 2007: Impacts, Adaptation and Vulnerability* (eds Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, Hanson CE), pp. 1–22. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK.
- Allen CR, Gunderson L, Johnson AR (2005) The use of discontinuities and functional groups to assess relative resilience in complex systems. *Ecosystems* 8, 958–966.
- Allen CR & Holling CS (2008) *Discontinuities in ecosystems and other complex systems*. Columbia University Press, New York.
- Allen CR, Angeler DG, Cumming GS, Folke C, Twidwell D, & Uden DR (2016). Quantifying spatial resilience. *Journal of Applied Ecology*, 53(3), 625–635.
- Allen CR, Fontaine JJ, Pope KL, Garmestani AS (2011) Adaptive management for a turbulent future. *J Environ Manag* 92:1339–1345
- Angeler DG and Allen CR 2016 Quantifying resilience. *Journal of Applied Ecology* 53(3): 617–624.
- Angeler DG, Allen CR, Barichievy C, Eason T, Garmestani AS, Graham NAJ, Granholm D, Gunderson L, Knutson M, Nash KL, Nelson RJ, Nyström M, Spanbauer TE, Stow CA & Sundstrom SM (2016a) Management applications of discontinuity theory. *Journal of Applied Ecology* 53(3): 688–698.
- Angeler DG, Allen CR, Garmestani AS, Gunderson LH & Linkov I (2016b) Panarchy use in environmental science for risk and resilience planning. *Environment, Systems and Decisions* 10.1007/s10669-016-9605-6
- Baho DL, Drakare S, Johnson RK, Allen CR, Angeler DG (2014) Similar resilience attributes in lakes with different management practices. *PLoS ONE* 9(3): e91881. doi:10.1371/journal.pone.0091881 [PubMed: 24618720]

- Baho DL, Allen CR, Garmestani A, Fried-Petersen HB, Renes SE, Gunderson LH & Angeler DG A quantitative framework for assessing ecological resilience. *Ecology and Society* 22(3):17 10.5751/ES-09427-220317
- Beever EA, O'Leary J, Mengelt C, West JM, Julius S, Green N, Magness D, Petes L, Stein B, Nicotra AB, Hellmann JJ, Robertson AL, Staudinger MD, Rosenberg AA, Babij E, Brennan J, Schuurman GW and Hofmann GE (2015) Improving conservation outcomes with a new paradigm for understanding species' fundamental and realized adaptive capacity. *Conservation Letters*. doi: 10.1111/conl.12190
- Bellingham PJ, Tanner EVJ & Healy JR (1995). Damage and responsiveness of Jamaican montane tree species after disturbance by a hurricane. *Ecology* 76, 2562–2580.
- Biggs R, Carpenter SR, & Brock WA (2009). Turning back from the brink: detecting an impending regime shift in time to avert it. *Proceedings of the National Academy of Sciences USA* 106, 826–831.
- Boucher DH, Vandermeer JH, Mallona MA, Zamora N & Perfecto I (1994). Resistance and resilience in a directly regenerating rainforest: Nicaraguan trees of the Vochysiaceae after Hurricane Joan. *Forest Ecology and Management* 68, 127–136.
- Brand FS & Jax L (2007) Focusing the meaning(s) of resilience: resilience as a descriptive concept and a boundary object. *Ecology and Society*, 12, 23
- Burthe SJ, Henrys PA, Mackay EB, Spears BM, Campbell R, Carvalho L, Dudley B, Gunn IDM, Johns DG, Maberly SC, May L, Newell MA, Wanless S, Winfield IJ, Thackeray SJ and Daunt F (2016), Do early warning indicators consistently predict nonlinear change in long-term ecological data?. *J Appl Ecol*, 53: 666–676.
- Carpenter S, Walker B, Anderies JM & Abel N (2001) From metaphor to measurement: resilience of what to what? *Ecosystems*, 4, 765–781.
- Carpenter SR, and Brock WA 2008 Adaptive capacity and traps. *Ecology and Society* 13(2): 40 [online] URL: <http://www.ecologyandsociety.org/vol13/iss2/art40/>
- Carpenter SR, Brock WA, Folke C, van Nes EH and Scheffer M 2015 Allowing variance may enlarge the safe operating space for exploited ecosystems. *PNAS* 112(46): 14384–14389 [PubMed: 26438857]
- Chaffin BC, Garmestani AS, Gunderson L, Harm Benson M, Angeler DG, Arnold CA, Cosens B, Craig RK, Ruhl JB & Allen CR (2016a) Transformative environmental governance. *Annual Review of Resources and Environment* 41: (Volume publication date November 2016)
- Chaffin BC, Garmestani AS, Angeler DG, Herrmann DL, Hopton ME, Kolasa J, Nyström M, Sendzimir J, Stow CA & Allen CR (2016) Adaptive governance, biological invasions and ecological resilience. *Journal of Environmental Management* doi:10.1016/j.jenvman.2016.04.040
- Cole LE, Bhagwat SA, & Willis KJ (2014). Recovery and resilience of tropical forests after disturbance. *Nature communications*, 5.
- Conrad M (1983) *Adaptability, the significance of variability from molecule to ecosystem*. Plenum Press, New York.
- Conroy MJ, Runge MC, Nichols JD, Stodola KW, Cooper RJ (2011) Conservation in the face of climate change: the role of alternative models, monitoring, and adaptation in confronting and reducing uncertainty. *Biol Conserv* 144:1204–1213
- Crowl TA, Crist TO, Parmenter RR, Belovsky G, Lugo AE (2008) The spread of invasive species and infectious disease as drivers of ecosystem change. *Front Ecol Environ* 6:238–246
- Cumming GS, Olsson P, Chapin FS III, Holling CS (2013) Resilience, experimentation, and scale mismatches in social-ecological landscapes. *Landsc Ecol* 28:1139–1150
- Dakos V, Scheffer M, van Nes EH, Brovkin V, Petoukhov V, & Held H (2008). Slowing down as an early warning signal for abrupt climate change. *Proceedings of the National Academy of Sciences*, 105(38), 14308–14312.
- Donohue I, Petchey OL, Montoya JM, Jackson AL, McNally L, Viana M, Healy K, Lurgi M, O'Connor NE & Emmerson MC (2013) On the dimensionality of ecological stability. *Ecology Letters*, 16, 421–429. [PubMed: 23419041]

- Eason T, Garmestani AS, Stow CA, Rojo C, Alvarez-Cobelas M, Cabezas H (2016), Managing for resilience: an information theory-based approach to assessing ecosystems. *Journal of Applied Ecology*, 53: 656–665.
- Elmqvist T, Folke C, Nyström M, Peterson G, Bengtsson J, Walker B, and Norberg J 2003 Response diversity, ecosystem change and resilience. *Frontiers in Ecology and the Environment* 1: 488–494.
- Folke C, Colding J, & Berkes F (2003). Synthesis: building resilience and adaptive capacity in social-ecological systems. *Navigating social-ecological systems: Building resilience for complexity and change*, 9(1), 352–387.
- Folke C, Carpenter S, Walker B, Scheffer M, Elmqvist T, Gunderson L, & Holling CS (2004). Regime shifts, resilience, and biodiversity in ecosystem management. *Annual Review of Ecology, Evolution, and Systematics*, 557–581.
- Folke C, Carpenter SR, Walker B, Scheffer M, Chapin T & Rockström J (2010) Resilience thinking: integrating resilience, adaptability and transformability. *Ecology and Society* 15(4): 20.
- Gallopín GC (2006). Linkages between vulnerability, resilience, and adaptive capacity. *Global environmental change*, 16(3), 293–303.
- Garmestani AS, Allen CR, and Gunderson L 2009 Panarchy: discontinuities reveal similarities in the dynamic system structure of ecological and social systems. *Ecology and Society* 14(1): 15 [online] URL:<http://www.ecologyandsociety.org/vol14/iss1/art15/>
- Garmestani AS, and H Benson M 2013 A framework for resilience-based governance of social-ecological systems. *Ecology and Society* 18(1):9 10.5751/ES-05180-180109
- Green OO, Garmestani AS, Allen CR, Ruhl JB, Arnold CA, Gunderson LH, Graham N, Cosens B, Angeler DG, Chaffin BC & Holling CS (2015) Barriers and bridges to the integration of social-ecological resilience and law. *Frontiers in Ecology and the Environment* 13: 332–337.
- Gunderson LH 2000 Ecological resilience--in theory and application. *Annual Review of Ecological Systems* 31:425–439.
- Gunderson LH, and Holling CS, editors. 2002 *Panarchy: understanding transformations in human and natural systems*. Island Press, Washington, D.C., USA.
- Hansson L-A, Annadotter H, Bergman E, Hamrin SF, Jeppesen E, Kairesalo T, Luokkanen E, Nilsson P-A, Söndergaard M & Strand J (1998) Biomanipulation as an application of food chain theory: constraints, synthesis, and recommendations for temperate lakes. *Ecosystems*, 1, 558–574.
- Holling CS, 1973 Resilience and stability of ecological systems. *Annual Review of Ecology and Systematics*, 4, 1–23.
- Holling CS, Allen CR (2002) Adaptive inference for distinguishing credible from incredible patterns in nature. *Ecosystems* 5:319–328.
- Hooper DU, Adair EC, Cardinale BJ, Byrnes JE, Hungate BA, Matulich KL, et al. (2012). A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature*, 486(7401), 105–108. [PubMed: 22678289]
- Horner-Dixon TF (1991) On the threshold: Environmental changes as causes of acute conflict. *International Security* 16:76–116
- Hughes TP, Carpenter S, Rockström J, Scheffer M, and Walker B 2013 Multiscale regime shifts and planetary boundaries. *Trends in Ecology and Evolution* 28:389–295. [PubMed: 23769417]
- Johnson RK and Hering D (2009) Response of taxonomic groups in streams to gradients in resource and habitat characteristics. *Journal of Applied Ecology*, 46: 175–186.
- Kates RW, Clark WC, Corell R, Hall JM, Jaeger CC, Lowe I, McCarthy JJ, Schellnhuber HJ, et al. (2011) *Sustainability Science*. *Science* 292:641–642
- Kühnel S & Blüthgen N (2015). High diversity stabilizes the thermal resilience of pollinator communities in intensively managed grasslands. *Nature Communications* 6, art7989.
- Labarte E, Wells JA, DeClerck F, Metcalfe DJ, Catterall CP, Queiroz C, et al. (2010). Land-use intensification reduces functional redundancy and response diversity in plant communities. *Ecology Letters*, 13(1), 76–86. [PubMed: 19917052]
- Magurran AE (2004). *Measuring biological diversity*. Oxford: Blackwell.
- McMichael AJ, Friel S, Nyong A, Corvalan C (2008) Global environmental change and health: impacts, inequalities, and the health sector. *Brit Med J* 336:191–194 [PubMed: 18219041]

- Mori AS, Furukawa T, and Sasaki T 2013 Response diversity determines the resilience of ecosystems to environmental change. *Biological Reviews* 88:349–364. [PubMed: 23217173]
- Mouillot D, Bellwood DR, Barlato C et al. (2013). Rare species support vulnerable functions in high-diversity ecosystems. *PLoS Biology* 11(5), e1001569 Doi:10.1371/journal.pbio.1001569 [PubMed: 23723735]
- Mumby PJ, Chollett I, Bozec YM & Wolff NH (2014). Ecological resilience, robustness and vulnerability: how do these concepts benefit ecosystem management?. *Current Opinion in Environmental Sustainability* 7, 22–27
- Nash KL, Allen CR, Angeler DG, Barichiev C, Eason T, Garmestani AS, Graham NAJ, Granholm D, Knutson M, Nelson RJ, Nyström M, Stow CA and Sundstrom SM (2014), Discontinuities, cross-scale patterns, and the organization of ecosystems. *Ecology*, 95: 654–667. [PubMed: 24804450]
- Nash KL, Graham NAJ, Jennings S, Wilson SK, Bellwood DR (2016). Herbivore cross-scale redundancy supports response diversity and promotes coral reef resilience. *Journal of Applied Ecology* 53, 646–655.
- Nicotra AB, Beever EA, Robertson AL, Hofmann GE and O’Leary J (2015), Assessing the components of adaptive capacity to improve conservation and management efforts under global change. *Conservation Biology*, 29: 1268–1278. [PubMed: 25926277]
- Nyström M (2006). Redundancy and response diversity of functional groups: implications for the resilience of coral reefs. *AMBIO* 35, 30–35. [PubMed: 16615697]
- Nyström M, & Folke C (2001). Spatial resilience of coral reefs. *Ecosystems*, 4(5), 406–417.
- Oliver TH, Heard MS, Isaac NJB, Roy DB, Procter D, Eigenbrod F, Freckleton R, Hector A, Orme CDL, Petchey OL, et al. 2015 Biodiversity and the resilience of ecosystem services. *Trends in Ecology and Evolution* 30:673–684 [PubMed: 26437633]
- Peterson GD (2002). Contagious disturbance, ecological memory, and the emergence of landscape pattern. *Ecosystems*, 5(4), 329–338
- Pickett STA & White PS (1985). *The ecology of natural disturbance and patch dynamics*. Academic Press, New York White
- Resilience Alliance. (2015) Resilience Alliance - Adaptive Capacity. <http://www.resalliance.org/adaptive-capacity>
- Rist L, Felton A, Nyström M, Troell M, Sponseller RA, Bengtsson J, Österblom H, Lindborg R, Tidåker P, Angeler DG, Milestad R & Moen J (2014) Applying resilience thinking to production systems. *Ecosphere* 5(6): art73.
- Rockström J, Steffen W, Noone K, Persson Å, Chapin FS III, et al E 2009 Planetary boundaries: exploring the safe operating space for humanity. *Ecology and Society* 14(2):32.
- Rodrigues ASL & Brooks TM (2007) Shortcuts for biodiversity conservation planning: the effectiveness of surrogates. *Annual Review of Ecology, Evolution, and Systematics*, 38, 713–737.
- Scheffer M & Carpenter SR (2003) Catastrophic regime shifts in ecosystems: linking theory to observation. *Trends in Ecology and Evolution*, 18, 648–656.
- Schleuning M, Fründ J, and García D 2015 Predicting ecosystem functions from biodiversity and mutualistic networks: an extension of trait-based concepts to plant- animal interactions. *Ecography* 38:380–392
- Smit B, & Wandel J (2006). Adaptation, adaptive capacity and vulnerability. *Global environmental change*, 16(3), 282–292.
- Spanbauer TL, Allen CR, Angeler DG, Eason T, Fritz SC, Garmestani AS et al. (2014). Prolonged instability prior to a regime shift. *PLoS ONE*, 9(10), e108936. [PubMed: 25280010]
- Spanbauer TL, Allen CR, Angeler DG, Eason T, Fritz SC, Garmestani AS et al. (2016). Body size distributions signal a regime shift in a lake ecosystem. *Proceedings of the Royal Society B* 283, 20160249 [PubMed: 27335415]
- Spears BM, Ives SC, Angeler DG, Allen CR, Birk S, Carvalho L, Cavers S, Daunt F, Morton RD, Pocock MJO, Rhodes G, Thackeray SJ (2015): Effective management of ecological resilience – are we there yet? *Journal of Applied Ecology* 52: 1311–1315.
- Stewart-Koster B, Olden JD, Johnson PTJ (2015) Integrating landscape connectivity and habitat suitability to guide offensive and defensive invasive species management. *J Appl Ecol* 52:366–378

- Sundstrom SM, Angeler DG, Garmestani AS, García JH, & Allen CR (2014). Transdisciplinary application of cross-scale resilience. *Sustainability*,6(10), 6925–6948.
- Sundstrom SM, Eason T, Nelson RJ, Angeler DG, Allen CR, Barichiev C, Garmestani AS, Graham NAJ, Granholm D, Gunderson L, Knutson M, Nash KL, Nyström M, Spanbauer T & Stow CA (2017) Detecting spatial regimes in ecosystems. *Ecology Letters* 20(1): 19–32 [PubMed: 28000431]
- Thompson SK, Seber GAF (1996) Adaptive sampling. Wiley, New York
- Thompson JN, Reichman OJ, Morin PJ, Polis GA, Power ME, Sterner RW et al. (2001). Frontiers of Ecology. *BioScience*,51(1), 15–24.
- Uden DR, Allen CR, Angeler DG, Corral L & Fricke KA (2015) Adaptive invasive species distribution models: A framework for modeling incipient invasions. *Biological Invasions* 17(19): 2831–2850.
- Walker B, Holling CS, Carpenter SR & Kinzig A (2004) Resilience, adaptability and transformability in social–ecological systems. *Ecology and Society*, 9(2), 5
- Walker B, Kinzig A, & Langridge J (1999). Original articles: plant attribute diversity, resilience, and ecosystem function: the nature and significance of dominant and minor species. *Ecosystems*, 2(2), 95–113.
- Wonkka CL, Twidwell D, West JB, & Rogers WE (2016). Shrubland resilience varies across soil types: implications for operationalizing resilience in ecological restoration. *Ecological Applications*, 26, 128–145. [PubMed: 27039515]
- Zhong-Yu S, & Hai R (2011). Ecological memory and its potential applications in ecology: A review. *Yingyong Shengtai Xuebao*, 22(3).

Box 1.**Definitions****Adaptive capacity**

Latent property of an ecological system (or other complex system) to respond to disturbances in a manner that maintains it within its current basin of attraction by altering the depth and/or breadth of that basin (Figure 1).

Contrasted with**Adaptation**

Alterations in the structure or function of an organism due to natural selection by which the organism becomes better fitted to survive and reproduce in its environment

Adaptability

ability to become adapted to live and reproduce under a particular range of environmental conditions

Adaptedness

adaptive traits (structural, functional, and behavioural), that are necessary for an organism to thrive in a particular environment

Ecological Resilience

ecological resilience is the capacity of a system to absorb disturbance to avoid a regime shift (multiple equilibrium focus), and a measure of the amount of disturbance a system can withstand before collapsing.

Engineering Resilience

return time to equilibrium after disturbance

Alternative State/Regime

a potential alternate configuration in terms of the structural and functional composition, processes, and feedbacks of a system

Basin of Attraction (stability domain)

configuration in terms of the abundance, composition, and processes of a system in which the system tends to remain

Cross-scale Resilience

resilience in ecological systems is enhanced when functional traits are diverse within scales and reinforced across scales

Ecological Memory

The collective representation of functional and structural attributes in an ecosystem that has been shaped by the systems disturbance history

Stability

Stability is a system characteristic whereby system variables remain unchanged following disturbance. Adaptive capacity can increase stability, but system components can fluctuate (therefore being unstable) while still remaining within the range of values that signify a particular state, and therefore, a system can be somewhat unstable while still possessing high adaptive capacity

Persistence

duration of species existence before it becomes extinct (either locally or globally)

Resistance

the external force or pressure needed to displace a system by a certain amount.

Variability

inverse of ecological stability; fluctuation in ecosystem parameters over time.

Functional Diversity

Diversity of reproductive phenology, seed bank potential, colonization and dispersal abilities, and other traits. This can enhance adaptive capacity by increasing functional redundancy and response diversity (see below).

Functional Redundancy

Existence of more than one species or process delivering the same ecological function. This contributes to adaptive capacity in ecosystems by providing buffering for loss of function due to extinction.

Response Diversity

Response diversity is variability in combinations of traits that provides a range of response patterns to disturbances and therefore increases the overall adaptive capacity in the system.

Regime Shift

persistent change in structure, function, and feedbacks of an ecosystem




Figure 1: Schematics illustrating high (A) and low (B) adaptive capacity. Adaptive capacity as a latent potential is shown by the dotted lines and the lengths of arrows that surround the basins of attraction that represent ecological resilience (left drawings). Drawings on the right show how high and low adaptive capacity can translate in rebound or a regime shift after a disturbance.




Figure 2: Reiterative testing, recalibrating, and refining of explicit hypotheses of adaptive capacity within an adaptive management, inference and modeling framework. The approach first recognizes patterns (reducing risk of type II error) and then refines knowledge about patterns reiteratively (reducing risk of type I error) to meet adaptive or transformative management objectives and reduce uncertainty.

Table 1:

Factors that contribute to ecological memory and that mediate adaptive capacity across different scales of biological organization. Note: the table is not exhaustive and meant only to highlight the complexity of factors influencing adaptive capacity.

Hierarchy of organization and selected traits
<ul style="list-style-type: none"> • Sub-individual <ul style="list-style-type: none"> ○ Matching physiological conditions to fluctuating inputs or internal demands (allostasis) (Carpenter and Brock, 2008). ○ Genetic, epigenetic and molecular processes (e.g., mutation). • Individual <ul style="list-style-type: none"> ○ Phenotypic plasticity. ○ Learning and dispersal ability. ○ Behavior ○ Adaptive evolution related to genetic diversity and evolutionary rates. ○ Links between life-history traits, phenotypic plasticity, and evolutionary potential. • Population <ul style="list-style-type: none"> ○ Heritable life history characteristics: generation time, reproductive capacity, migration, habitat selection, genome size, survival characteristics (resting stages; hibernation, estivation), generalist vs. specialist species. ○ Population structure. ○ Metapopulation dynamics. • Community <ul style="list-style-type: none"> ○ Taxonomic diversity. ○ Functional diversity (redundancy, response diversity). ○ Strength of species interactions. ○ Metacommunity dynamics (colonization and dispersal abilities). ○ Founder effects. ○ Priority effects. ○ Dormancy (resting eggs and propagule banks) and bet-hedging strategies. • Ecosystem <ul style="list-style-type: none"> ○ Interaction of and connection between abiotic and biotic elements in feedback loops (balancing and reinforcing or negative and positive). ○ Changing shapes of basin of attraction/stability landscape (topography, soils, landforms) • Biome <ul style="list-style-type: none"> ○ Biogeographical distributions of native and invasive species. ○ Phylogenetic dynamics. ○ Evolutionary, disturbance and climate histories.

Table 2:

Premises of resilience components and simple, management-relevant, mutually non-exclusive hypotheses. These serve as a starting point for reiterative hypothesis testing and can be refined, modified and adapted to specific ecosystems during monitoring.

<i>Ecological regime</i>	<i>Adaptive capacity</i>	
	Premise 1 (stable)	Premise 2 (erodes)
<i>Hypotheses</i>		
1. Adaptive capacity surrogates	High persistence, resistance and recovery; low variability.	Slowed down recovery, decreasing resistance and persistence, and higher variability.
2. Ecological memory	High structural and functional redundancy and diversity	Decreasing redundancy and diversity.
3. Key/deterministic/stochastic species	Contribute to ecological memory.	Increasing extinctions
4. Response diversity	High	Reduced
5. Functional compensation	High	Decreasing
6. Functional trait distributions	High diversity and redundancy within and across scales	Decreasing diversity and redundancy within and across scales.
<i>Support</i>	Allen et al. (2005); Angeler <i>et al.</i> (2016); Bellingham et al. (1995); Baho et al. (2014); Boucher et al. (1994); Kühnel & Blüthgen (2015); Moullot et al. (2013); Nash et al. (2016); Walker et al. (1999)	Carpenter & Brock (2006); Dakos et al. (2008); Hooper et al. (2012); Mumby et al. (2014); Nystrom (2006); Nash et al. (2016); Spanbauer et al. (2016)
<i>Management implication</i>	Maintain regime	1) Adaptive management and governance to stave off regime shifts. 2) Scenario planning and transformative governance if adaptive capacity gets exhausted.