



Eddleston, M., Karunaratne, A., & Gunnell, D. (2019). Response to Bayer regarding pesticide suicides. *Clinical Toxicology*.
<https://doi.org/10.1080/15563650.2019.1708092>

Peer reviewed version

Link to published version (if available):
[10.1080/15563650.2019.1708092](https://doi.org/10.1080/15563650.2019.1708092)


[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Taylor and Francis at <https://www.tandfonline.com/doi/full/10.1080/15563650.2019.1708092>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



Response to Bayer regarding pesticide suicides

Journal:	<i>Clinical Toxicology</i>
Manuscript ID	Draft
Manuscript Type:	Letter
Keywords:	Pesticides, Suicides, Regulation
Abstract:	

SCHOLARONE™
Manuscripts

1
2
3 **Dear Sir**
4

5 We thank Dr Dunn and colleagues from Bayer Crop Science for responding to our recent article
6 estimating the total number of pesticide suicides that have occurred since the Green Revolution.
7

8 We did not question the Green Revolution per se, only its introduction of highly hazardous pesticides
9 into communities completely unable to use or store them safely. Much less hazardous pesticides could
10 have been used, either initially or by the late 1960s when the immense harm that they were causing
11 had already become apparent [1, 2]. It was their easy availability at times of (completely predictable)
12 stress that resulted in deaths.
13
14

15 It is great to learn that the pesticide industry is now concerned about suicide. This was not the case for
16 a long time when the major emphasis was on 'proper' use, blaming all problems including suicide on
17 'misuse' [3, 4]. As we stated, pesticide suicides should be considered a Schilling category 4
18 occupational condition. It is not that these compounds are 'misused', they just could never have been
19 used safely by these small scale farmers, as so eloquently highlighted more than 50 years ago by
20 Vethanayagam [1].
21
22

23 If the industry was really concerned about suicide, it would immediately stop producing all WHO Hazard
24 Class I pesticides, in particular phorate which can still be bought by non-professionals in concentrated
25 formulations in India. This pesticide has the same rat oral LD50 as the nerve agent sarin (both 0.5
26 mg/kg). Does the industry think it reasonable that people are given bottles of 'sarin-equivalent' toxicity
27 pesticides and urged to use them 'safely'? Would this be accepted in the USA or Europe?
28
29

30 And if suicide was a priority, then the industry would actively take paraquat (case fatality >40% [5]) out
31 of small-scale use and work with communities to find other, ideally non-chemical alternatives.
32

33 In the continued absence of action by industry to accept its role in pesticide suicides and to stop
34 producing such highly hazardous pesticides, it is left to governments to decide how best to protect their
35 population with legislation that removes problematic pesticides from use [6]. Government must also
36 prevent the entry of highly hazardous pesticides across borders. However, many do not have sufficient
37 resources. Coordinated action across government, industry and international agreements is required to
38 prevent the easy availability of such pesticides.
39
40

41 We did not state that there were no farmer suicides before 1960, only that there were very few pesticide
42 suicides. Farmer suicides cannot be conflated with pesticide suicides (the subject of our article). They
43 are but a small (although highly visible) part of the problem. In India, for example, where farmer
44 suicides are a major political issue, there are about 15,000 suicides of land-owning (therefore nearly
45 always male) farmers each year [7], around half by pesticide ingestion. But there are an estimated
46 230,000 suicides in India each year [8], with around 70,000 (30% of all suicides) by pesticide ingestion
47 (9.3 times the number of 'farmer pesticide suicides') [8, 9]. Unfortunately, these deaths - of family
48 members, of children, and non-farming households who can access pesticides - receive little attention.
49
50

51 One key point to remember is that suicide impulses are transient and may last only minutes or hours.
52 The absence of a lethal method (such as a gun or highly hazardous pesticide in the house) during that
53 period massively increases the chance that the person will use a less lethal method (such as medicine
54 ingestion), thereby surviving, or not harm themselves at all. This saves lives, as clearly shown by the
55 data from Sri Lanka and other Asian countries that have carried out pesticide regulation to prevent
56 suicides [10].
57
58
59
60

References

1. Vethanayagam AV. 'Folidol' (parathion) poisoning. *BMJ*. 1962;986-987.
2. Indian Council of Agricultural Research. Report of the Special Committee On Harmful Effects Of Pesticides. New Delhi, ICAR. 1967.
3. Brown R, Clapp M, Dyson J, et al. Paraquat in perspective. *Outlooks on Pest Management*. 2004;Dec 2004:259-267.
4. Mishara BL. Report on the International Workshop on Secure Access to Pesticides in conjunction with the Annual Congress of the International Association for Suicide Prevention, Durban, South Africa, Monday September 12, 2005 (sponsored by Syngenta Crop Protection). <http://www.iasp.info/pdf/papers/Mishara.pdf> (accessed Apr 20, 2009)2005.
5. Dawson AH, Eddleston M, Senarathna L, et al. Acute human lethal toxicity of agricultural pesticides: a prospective cohort study. *PLoS Med*. 2010;7:e1000357.
6. Gunnell D, Knipe D, Chang SS, et al. Prevention of suicide with regulations aimed at restricting access to highly hazardous pesticides: a systematic review of the international evidence. *Lancet Glob Health*. 2017;5:e1026-e1037.
7. Merriott D. Factors associated with the farmer suicide crisis in India. *J Epidemiol Glob Health*. 2016;6:217-227.
8. India State-Level Disease Burden Initiative Suicide Collaborators. Gender differentials and state variations in suicide deaths in India: the Global Burden of Disease Study 1990-2016. *Lancet Public Health*. 2018;3:e478-e489.
9. Mew EJ, Padmanathan P, Konradsen F, et al. The global burden of fatal self-poisoning with pesticides 2006-15: Systematic review. *J Affect Disord*. 2017;219:93-104.
10. World Health Organization, Food and Agriculture Organization of the United Nations. Preventing suicide: a resource for pesticide registrars and regulators. ed. Geneva: WHO; 2019.