

SUPPLEMENTARY MATERIAL

related to :

On giant shoulders: How a seamount affects the microbial community composition of seawater and sponges

Kathrin Busch¹, Ulrike Hanz², Furu Mienis², Benjamin Müller³, Andre Franke⁴,
Emyr Martyn Roberts⁵, Hans Tore Rapp⁵, Ute Hentschel^{1,6}

¹GEOMAR Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20, 24105 Kiel, Germany

²NIOZ Royal Netherlands Institute for Sea Research and Utrecht University, 1790 AB Den Burg, Texel, The Netherlands


³University of Amsterdam, Science Park 904, P.O. Box 94248, Amsterdam, The Netherlands

⁴Institute of Clinical Molecular Biology (IKMB), Rosalind-Franklin-Straße 12, 24105 Kiel, Germany

⁵University of Bergen, Department of Biological Sciences and K.G. Jebsen Centre for Deep-sea Research, P.O. Box 7803, 5020 Bergen, Norway

⁶Christian-Albrechts University of Kiel, Düsternbrooker Weg 20, 24105 Kiel, Germany

Correspondence to: Ute Hentschel (uhentschel@geomar.de)



Supplementary Figure S1 Richness of seawater microbial communities for the mid-water and near-bed water samples, across the determined clusters. Faith Phylogenetic Diversity (Faith's PD) is plotted as alpha-diversity measure. Mid-water clusters are coloured in orange (MW1) and yellow (MW2), while near-bed water clusters are marked in red (BW1), dark blue (BW2), light blue (BW3), and purple (BW4).

Supplementary Table S1 Overview of pairwise-comparisons (PERMANOVAs) across pelagic microbial micro-habitats based on weighted UniFrac matrices of microbial communities for both, seawater and the three sponge species. All possible pairwise combinations were tested. *L. complicata* and *S. rosea* were not present in all near-bed water clusters. Four biological replicates per near-bed water cluster were tested for every sponge species.

Group 1	Group 2	pseudo-F	p-value	
MW1	MW2	22.49	0.001	*
BW1	BW2	27.7	0.004	*
BW1	BW3	11.79	0.002	*
BW1	BW4	15.07	0.003	*
BW2	BW3	8.52	0.001	*
BW2	BW4	15.56	0.001	*
BW3	BW4	10.98	0.001	*
<i>G. hentscheli</i> BW1	<i>G. hentscheli</i> BW2	5.77	0.032	*
<i>G. hentscheli</i> BW1	<i>G. hentscheli</i> BW3	8.07	0.028	*
<i>G. hentscheli</i> BW1	<i>G. hentscheli</i> BW4	16.71	0.030	*
<i>G. hentscheli</i> BW2	<i>G. hentscheli</i> BW3	4.57	0.039	*
<i>G. hentscheli</i> BW2	<i>G. hentscheli</i> BW4	14.41	0.035	*
<i>G. hentscheli</i> BW3	<i>G. hentscheli</i> BW4	5.57	0.028	*
<i>L. complicata</i> BW1	<i>L. complicata</i> BW2	35.93	0.034	*
<i>S. rosea</i> BW1	<i>S. rosea</i> BW2	1.88	0.035	*
<i>S. rosea</i> BW1	<i>S. rosea</i> BW3	1.71	0.174	
<i>S. rosea</i> BW2	<i>S. rosea</i> BW3	1.18	0.36	
all MW	all BW	46.72	0.001	*

Supplementary Table S2 Averages and standard errors of biogeochemical parameters for each mid-water and near-bed water cluster.

	SPM [mg L ⁻¹]	DIC [μmol L ⁻¹]	SiO₄⁻ [μmol L ⁻¹]	PO₄³⁻ [μmol L ⁻¹]	NH₄⁺ [μmol L ⁻¹]	NO₃⁻ [μmol L ⁻¹]	NO₂⁻ [μmol L ⁻¹]	O₂ [mL L ⁻¹]	Depth [m]
MW1	NA	2246.00 ± 6.99	5.72 ± 0.10	0.88 ± 0.02	NA	12.84 ± 0.08	0.02 ± 0.02	6.90 ± 0.04	400 ± 0
MW2	2.22 ± 1.39	2292.14 ± 53.64	5.63 ± 0.08	0.84 ± 0.01	0.11 ± 0.03	13.08 ± 0.11	0.02 ± 0.01	6.75 ± 0.03	400 ± 0
significance mid-water								*	
BW1	0.49 ± NA	NA	6.00	0.90	0.10	13.00	0.00	6.99 ± NA	575
BW2	NA	2265.67 ± 7.69	8.40 ± 0.31	0.97 ± 0.03	NA	14.20 ± 0.06	0.00 ± 0.00	6.71 ± 0.02	919 ± 106
BW3	1.87 ± 1.08	2260.40 ± 12.35	7.87 ± 0.42	0.95 ± 0.02	0.17 ± 0.03	14.03 ± 0.15	0.03 ± 0.02	6.84 ± 0.07	922 ± 142
BW4	0.95 ± 0.33	2248.00	10.65 ± 1.42	0.95 ± 0.03	0.10 ± 0.00	14.78 ± 0.52	0.05 ± 0.03	6.48 ± 0.10	1836 ± 376
significance near-bed water			*			*		*	*