

## Mammal Community Structure Analysis

Kris Kovarovic<sup>1</sup>, Kari Lintulaakso<sup>2</sup>, Denise F. Su<sup>3</sup>

<sup>1</sup>*Department of Anthropology, Durham University, South Road, Durham DH1 3LE, United Kingdom ([kris.kovarovic@durham.ac.uk](mailto:kris.kovarovic@durham.ac.uk))*

<sup>2</sup>*Department of Geosciences and Geography, University of Helsinki, Gustaf Hällströmin katu 2a, 00014 University of Helsinki, Finland ([kari.lintulaakso@gmail.com](mailto:kari.lintulaakso@gmail.com))*

<sup>3</sup>*Department of Paleobotany and Paleoecology, Cleveland Museum of Natural History, 1 Wade Oval Drive, Cleveland, OH, United States of America ([dsu@cmnh.org](mailto:dsu@cmnh.org))*

### Abstract

Fundamentally rooted in Odum's (1953) niche concept, mammal community studies are based on the understanding that each resident species reveals information about its environment through its adaptations to specific resources and landscape features. Ecologists view the community's profile of strategies for exploiting particular spatial and dietary niches; a quantitative summary of these strategies when compared across locales from a variety of habitat types demonstrates striking similarities in the communities that live in similar habitats regardless of their location. Recognising that communities can be compared across space, paleoecologists implemented community studies across time in an effort to reconstruct past environments. This synecological approach to paleoenvironmental reconstruction may be thought of as holistic, since it is not restricted to a single mammal family. However, thorough explorations of how fossil and extant communities differ have revealed significant dissimilarities brought about by the taphonomic history of paleontological assemblages. Techniques have been developed for addressing differences between the modern comparative community sample and the paleontological sample to which it is compared, but recent research conducted by both neo- and paleoecologists has suggested that there are unappreciated differences between modern habitats, as well.

### Key Words

community analysis, ecological diversity, ecomorphology, niche, taxon-free, synecology

### Running Head

Mammal Community Analysis

## INTRODUCTION

Mammalian remains are often abundant at paleontological sites around the world and represent a critical source of information about the Earth's biological past. Individual species and entire assemblages can reveal the unique characteristics of the mammalian community in a given time and place, contributing to our understanding of how life forms evolved in different settings over geological time.

Many classic post-Darwin paleontological studies focused largely on morphological descriptions of mammalian remains for the sake of understanding species and, by extension, their reconstructed evolutionary relationships. Influential mid-20<sup>th</sup> century developments in modern ecological theory, however, spurred something of a revolution in mammalian paleosciences. A step-change in thinking arose with the extension of Odum's (1953) classic niche concept into the work of paleontologists interested generally in the interaction between the environment and mammalian evolution and the reconstruction of ancient environments. Paleontologists came to appreciate more fully that mammals could provide information about their habitats and studies utilizing their remains in this manner added to a growing corpus of paleoecological literature.

There are numerous ways that mammals form an integral part of paleoecological research, from isotopic work (see Higgins, this volume) to ecomorphology (see Barr, this volume), but the focus here is on the use of entire communities, referred to variously as community structure or ecological diversity analysis among other monikers. We review the theoretical underpinning for this synecological approach and its scientific development, consider the pros and cons of this in comparison to other available methods, and look ahead to further refinements that will allow us to elucidate ever more nuanced information about ancient environments.

## **ECOLOGICAL THEORY: THE UNDERPINNING OF COMMUNITY STRUCTURE ANALYSIS**

### **The Niche Concept**

Eugene P. Odum is often credited with being one of, if not *the*, founder of the modern science of ecology. He was certainly not the only person in pre- or post-Darwin times to discuss connections between mammals and their environments or to overtly outline the nature of relationships between the two (such as predator-prey interactions). However, Odum was a prime mover in an influential mid-20<sup>th</sup> century shift in ecological thinking. The general principles he championed are still persuasive and, despite some having been shown to be flawed in some respects, his work in the 1950s can be described almost as “contagious” in many scientific circles, particularly in debates surrounding the utility of evolutionary versus ecosystem ecology. One area where his intellectual input contributed greatly to the development of paleoecology is in the ecological niche concept, which is described in the 2<sup>nd</sup> edition (1959) of his popular 1953 text *Fundamentals of Ecology* as “the position or status of an organism within its community and ecosystem resulting from the organism's structural adaptations, physiological responses, and specific behavior (inherited and/or learned)” (Odum 1959, pg. 27). This he contrasted to a habitat, the physical space occupied by a species, employing an analogy that has endured for decades: the habitat is the species’ “address”, but the niche is its “profession”.

The very language used here conveys something that is dynamic rather than static. It suggests that a niche is defined by relationships between individual species and between species and their physical environments and, furthermore, that these relationships can change over time. Indeed, by emphasising that *structural adaptations* are a part of the very definition of a niche, an evolutionary time scale is invoked. Bodily structures are adaptations and, as

such, they can tell us something about the niche(s) to which an organism was adapted (Harrison 1962). This isn't a circular concept, so much as an empirical approach which enables us to work "backwards" as paleontologists often do by comparing what can be observed in the present with remains from the past; in this case, the hard tissue adaptive structures of the skeleton and dentition.

Although ecologically meaningful mammalian structural adaptations (also known as "ecomorphologies", a concatenation of the term "ecological functional morphologies") may be explicitly studied to infer past environmental conditions (see Barr, this volume), community analysis uses information derived from these adaptations, as well as other proxies, to classify individual species into categories that broadly describe their preferred niches. The concept of a niche as expressed by Odum is also particularly pertinent here, as it makes clear that individual species have a place – their "position or status" – within a community, as well as within the ecosystem as a whole. The implicit idea is that the entire community can yield a greater amount of information about its environment than individual taxa, which occupy a limited range of niches, and therefore cannot provide a composite picture. Simply put, the whole is greater than the sum of its parts.

*Mammalian adaptations to the niche:* The exploitable niches that are available in a given habitat are conditioned by a combination of both biotic and abiotic factors, such as the different kinds of vegetation (trees, grass, herbs, shrubs, etc.) and the particular qualities of the soils in which they grow (drainage, nutrients, texture, inorganic composition, etc.) (Lomolino et al., 2006). These characteristics dictate what types of mammals can live in a given place and how they behave – a locale with nutrient-rich, moist but well-drained soil supporting heavy vegetation cover will have a very different mammalian community than a locale with dry, sandy soils and sparse vegetation cover. For example, tropical and subtropical rain forests or similar biomes with high annual average temperature and

precipitation, where fruit resources are available year round, support more pure frugivorous mammals than areas with seasonal precipitation, which have higher numbers of herbivorous taxa (Pineda-Munoz et al., 2016).

A habitat can be divided into two distinct niches that are relevant for paleocology: a spatial niche, which relates to the physical space that a species moves around in during its life cycle, and the trophic niche, which relates to the diet a species relies on to meet its nutritional and energy requirements. All mammals need a place to live and food to eat, thus they are adapted to (or “fill”) a particular spatial and trophic niche.

Soft tissue structures are every bit as adaptive as hard tissue structures, but they are of limited use to paleontologists given their resistance to fossilization. However, the preservation of skeletal and dental remains, although impacted by numerous taphonomic factors, yields a wealth of material from the past that can be related back to spatial and trophic niche exploitation. Mammals negotiate and utilize space with different locomotor repertoires evidenced by the morphology of their postcranial skeleton, particularly elements of the fore and hindlimb, which vary according to the substrate to which the species is adapted (e.g., Kappelman 1988).

Teeth, in comparison to postcranial elements, provide evidence regarding the trophic niche to which a species is adapted. Animals have specific dietary requirements and, thus, dental morphologies that are indicative of a carnivorous, insectivorous, frugivorous or herbivorous diet. Combination niche exploitation strategies are particularly common in herbivores, which may be further defined as browsers (preferring dicots) or grazers (preferring grass) and, quite frequently, a mix of the two which often relates to the seasonal availability of food sources. It is also common for herbivores on the browsing end of the spectrum to consume fruits and for insectivores to consume small mammals and/or fruit. True omnivores are rare and the morphology of their dentition tends to point to a specific

combination of dietary strategies rather than an equal reliance on all food resources (Pimm & Lawton, 1978; Yodzis, 1984).

*Community structure:* The “structure” in a community structure analysis is essentially a summary of niche exploitation profiles. The primary interest is how the mammalian species fill trophic and spatial niches with their dietary and locomotor adaptations, although in many cases, body size is also considered as ecologically relevant and given the same consideration as the two primary niches profiles (Andrews et al. 1979). Summaries can be helpfully visualised in histograms, which demonstrate quite effectively a key concept of this analytical approach to environmental reconstruction: habitats with a similar array of available niches have similar community structure profiles regardless of their geographical location. Figure 1, originally published by Andrews et al. (1979) expresses this in a series of histograms organized according to five broad habitat types. Profiles for the relative proportion of species in taxonomic orders, average body mass categories and locomotor and trophic niche categories are presented. By comparing these profiles across the habitat categories, it is easy to see where differences between them occur in their mammalian faunas. For example, compare the lowland and montane forest locomotor profiles with those from non-forest habitats. Species in the large ground mammal category (LGM), which are entirely terrestrial, are found in much greater numbers in non-forests (approximately 50% of the fauna), whereas forests have a higher proportion of species classified as small ground mammals (SGM), which practice some form of arboreality in addition to exploiting the terrestrial niche. These data are from 23 modern localities in Africa, but the patterns can also be viewed on a broader geographic scale in datasets that extend to multiple regions and continents (Andrews 1996).

INSERT FIGURE 1 HERE; WIDTH = 2 COLUMNS

See, for instance, the similarities in the profiles of the Central American tropical forest and Chinese paratropical forest in Figure 2 (data from Andrews & Humphrey, 1999).

Although separated by some 14,000 km and an 80 million year evolutionary history, these profiles are visibly similar with approximately equal numbers of species in each category. The differences appear relatively minor; Chinese paratropical locales have a small number of aerial species whereas they are lacking in the sampled Central American communities, which themselves have a slightly higher proportion of terrestrial taxa. But, overall their general patterns are the same. This structural similarity in their faunal communities are indicative of the fact that all forests present a similar array of exploitable niches for resident species to fill.

INSERT FIGURE 2 HERE; WIDTH = 2 COLUMNS

### **Significance for Understanding Paleocology**

Paleoenvironmental community studies are fundamentally rooted in the present, based on observations of mammalian behavior in habitats where they are known to live. Demonstrating the existence of similarities in patterns of community structure in broad habitat categories was largely based on extant communities (e.g., Karr 1971, Fleming 1973), but an obvious next step for paleoecologists was to extend this principle to the fossil record. While observations of past mammalian dietary and locomotor habits cannot be made, their postcranial and dental remains provide evidence for their adaptive niches, from which community structure profiles can be constructed.

Andrews and colleagues (1979) developed the community structure technique for interpreting palaeoecology based on early studies of species diversity in fossil and modern avian and mammalian communities (Kurtén 1952, Harrison 1962, Simpson 1964; Karr 1971). They noted that these early studies (Simpson 1964; Karr 1971) showed changes in the diversity of animal and plant species following a north-south gradient on different continents such that as latitudes decreased, the number of species increased. Additionally, the structure of the communities, i.e., distribution of body sizes and dietary and locomotor adaptations, also changed with latitude (Fleming 1973). They showed that community structures vary

significantly with habitat type within a single latitudinal zone, but that when both latitude and habitat type are similar, the structure of the communities under study are constant, regardless of differences in continent or species number and composition. They proposed that the habitat of a faunal community can be determined based on its community structure irrespective of species composition and that this predictive quality has direct application for paleoecology. By establishing patterns of community structure for a series of modern faunal communities and comparing them to those calculated for fossil communities, it is possible to make direct comparisons and inferences regarding the habitats from which the fossil assemblages may have come. This methodology was quickly integrated by many researchers into studies of the paleoecology of fossil hominoid-bearing faunal communities (e.g. Andrews and Van Couvering 1979; Van Couvering 1980; Nesbit Evans et al. 1981) and is regularly incorporated into paleoecological studies of mammalian fossil sites today (e.g., Reed 1997; Kovarovic et al. 2002; Flynn et al. 2003; Su and Harrison 2007; Croft et al. 2008; Su et al., 2009; Meloro and Kovarovic 2013).

## **PALEOENVIRONMENTAL RECONSTRUCTION**

Bearing in mind the theoretical considerations of community structure analysis, there are also practical aspects to consider when implementing this method as a form of paleoenvironmental reconstruction. Decisions must be made regarding the selection of extant localities for the comparative datasets and how the mammalian community niche exploitation profiles are compiled and categorized. The issue of categorization is particularly critical as neither mammalian niche exploitation behavior nor habitats are discrete entities, existing instead on a spectrum of related variables. But, the method requires that species are classified according to their preferred niches and localities to their broad habitat or biome,



which reduces the representation of their natural biological diversity. A balance must be struck between using a classification scheme that is nuanced enough to provide useful environmental interpretations and one that is also applicable to fossil species for which behavior is reconstructed from proxies rather than observed directly as in the modern world.

### **Extant Comparative Sample Selection**

Assembling a dataset of extant localities to which the target paleontological locality is compared requires a careful and often lengthy process of identifying natural localities (i.e., not significantly altered by human habitation or activity) associated with complete mammalian species lists (Andrews et al. 1979). Many faunal lists for modern localities were compiled for specific research questions or conservation projects and it is often the case that they record only the small or large fauna or species in particular families. Only localities for which faunal lists are considered complete are viable for community structure analyses, although establishing completeness itself may be difficult. Fortunately, with the advent of electronic databases and data sharing requirements, there are a number of resources that can be useful in this regard; for example, the database compiled by the National Center for Ecological Analysis and Synthesis (NCEAS) workshop on Mammalian Communities, which provides a great deal of information compiled by experts on the fauna present at a number of localities around the world (see Badgley et al. 2001, van Dam et al. 2001, Damuth et al. 2002).

In addition to identifying suitable localities, they should encompass the entire spectrum of ecological possibilities. It is possible that certain major biomes are not included, as dictated by the specific research question or location of the paleontological site that is being subjected to the environmental reconstruction. There may be reasonable grounds for excluding temperate and Arctic tundra communities, for example, if the paleocommunity is

derived from an equatorial location where other indicators suggest that it was not physiognomically similar to tundra (i.e. desert in the tropical realm).

Gross vegetation physiognomy is, in fact, one of the best defining characteristics for any habitat classification system. There are many systems available in the literature, some of which span the entire range of global biomes, but they are often based on specific vegetation species associations (White 1983; Eiten 1992; Lawesson 1994). But, just as a community structure analysis does not account for the taxonomic identity of the mammalian species, it also cannot account for the species of vegetation in the habitats. Structural variables such as the number of canopies, crown height, overall tree, bush or shrub cover and the presence of grassy or herbaceous ground cover are more relevant. Of course, such systems can be rather coarse, lumping several habitat types into broad groups that obscure ecological nuances that exist between them. The system used by Andrews et al. (1979) in Figure 1 considers only four categories including one for all woodlands and bushlands which, while being rather reductive, also presents a series of rules that make it reasonably easy to place each locality into a habitat group. The more nuanced the habitat classification system, the more information can be extracted from a paleocommunity, but the harder it becomes to categorize the habitat groups in the first place. They exist on a continuous spectrum of vegetation cover so the difficulty is in knowing where to draw the line between them to create discrete, analyzable categories. For this reason, paleoecologists have drawn on current research and descriptions of extant vegetation communities and habitats (particularly White 1983 and Olson et al. 1983, 1985) to devise clearly defined classification systems that can be applied to the paleontological record (Andrews et al. 1979, Reed 1997, Andrews & Humphrey 1999). Table 1 (originally from Kovarovic et al. 2002) summarizes the habitat categories of 22 modern localities used in a community structure analysis of the Upper Ndolanya Beds of Laetoli, a Pliocene hominin-bearing site. The habitat type column represents the broad

classification system employed, while a more detailed description of each locality provides the basis for the categorization and aids in subsequent interpretations.

INSERT TABLE 1 HERE

### **Fossil Assemblage Suitability**

Not all paleontological localities can be subjected to a community structure analysis. The assemblage needs to have preserved a reasonably complete portion of the mammalian fauna that existed. Although it can be expected that most sites will suffer some amount of taphonomic loss dependent on a variety of ecological and depositional conditions, an assemblage that is noticeably lacking in overall species numbers, or richness, is not a good candidate for this methodology (Andrews 1996). It is difficult to judge how complete a community is in the fossil record, but some basic protocols can be applied for determining if an assemblage is suitable.

One point to consider is the number of unique taxa required purely from the standpoint of multivariate statistical robusticity. From this perspective, even an assemblage with a low number of species, as few as five or six (Louys 2007; Louys et al. 2009), can be used. However, communities with such low species richness are obviously depauperate when compared to the richness observed in natural modern communities. Theoretically speaking, habitats that are structurally and topographically complex provide a greater number of niches to exploit and therefore have greater richness, such as forests and complex woodlands (Reed 2008). In the tropical realm it is uncommon to find communities with fewer than 40 taxa, approached only by communities inhabiting quite open environments (e.g. Serengeti Plains, Andrews & Nesbit Evans 1979, Andrews et al. 1979). While this might be a useful threshold to consider given that the number is reflected by extant faunas, it is an unreasonable expectation of many fossil communities, which tend to average lower than 40 taxa (see, for example, Andrews & Nesbit Evans 1979) for reasons pertaining to taphonomy, the intensity

of scientific sampling (it is known that bigger samples tend to have more species, Su 2016) and our ability to correctly identify all individual fossil species from fragmentary remains. In an extensive exploration of the species richness required to effectively distinguish between modern habitats, Louys et al (2009) found that this differed by broad habitat categories as defined by vegetation structure: closed, mixed and open. They provide empirical support for what makes intuitive sense, richness is generally correlated with broad habitat type, so it is expected that a greater number of species might be required to distinguish between those that are taxonomically richer. A minimum of 32 species was identified as the threshold for distinguishing between communities adapted for closed and mixed habitats, whereas only 12 were required to distinguish between open and mixed and 8 for open and closed.

### **Ecovaryable categories**

Community structure analysis considers three variables that reflect the niche exploitation of the species, also termed *ecovaryables*. They are body size, locomotor adaptation, and feeding preference, and their expression provides information on how the species of a mammalian community fills the spatial and trophic niches of the ecosystem.

*Body Size (Mass)*: Body size is one of the most fundamental biological characteristics of mammals. It has been shown to correlate significantly with other aspects of biology and behavior and is an important factor in space utilization for an organism (see Hopkins, this volume, for detailed discussion and other references). Various studies have shown a correlation between body size distributions of mammalian communities and their environments (e.g. Legendre 1986, 1989; Flynn et al. 2003) although it appears to be the least discriminating of the three ecovaryables (Andrews 1996; Reed 1997; Lintulaakso & Kovarovic 2016). Body size classification in extant mammals is relatively straightforward, done by calculating the mean of the species' weight range. Reconstructing body size in fossil

mammals is more complicated and is based on the consistent isometric and allometric relationships between various skeletal dimensions and body mass (e.g., Gould 1975, Alexander 1977, Fortelius 1990, Gingerich et al. 1982, Scott 1990, Legendre 1986, references in Damuth and McFadden 1990). For the most part, because community structure analysis relies on the mammalian species list and species identification is dependent on cranio-dental material, the methodology uses dental dimensions as body size proxies. It is an added advantage that teeth are the most identifiable and preserved element in the mammalian fossil record and its various dimensions are known to correlate with body size, among them molar surface ratio and toothrow length (Gould 1975, Gingerich et al. 1982, Fortelius 1990, Hopkins 2008, this volume). It is worth noting that although there are many factors that can influence the robustness of body size estimation (see Hopkins this volume for detailed discussion), precision is not as crucial in community structure analysis because species are categorized in a range of body weights that compensates for any inexactness in the predication. Body size categories vary from study to study and can have as few as three categories (<1kg, 1-45kg, >45kg, Andrews et al. 1979) to as many as eight categories (<100g, 100-1000g, 1-10kg, 10-45kg, 45-90kg, 90-180kg, 180-360kg, >360kg, Kovarovic et al. 2002). The division of body size categories is arbitrary to a certain extent, but it is important to consider the body size distribution of the target paleocommunity so that body size categories are not overly or under-divided.

*Locomotor adaptation (substrate categories):* Locomotor adaptations are generally considered in terms of the substrate or spatial niche for which the organisms are adapted (Harrison 1962; Eisenberg 1981). Assignment of a locomotor class should be based on the morphological adaptations of the limb bones that allow the species to move within its niche. However, there are instances, particularly within a fossil assemblage where it is often not possible to associate isolated postcranial elements to taxonomically-identified cranio-dental

materials, when it is necessary to categorize a species based on its known locomotor adaptation or substrate from ecomorphological studies on specimens of the same species from other sites or from taxonomic analogy as a last resort.

Locomotor categories are relatively broad and emphasize ecological niche structure, the animals in a community are categorized to correspond to the physical layers in the environment (Andrews et al. 1979). The categories proposed by Andrews et al. (1979) and commonly used in community structure analyses are: (1) aerial, the space above the vegetation to which aerial mammals, such as bats, are restricted; (2) arboreal, mammals that are found solely in the upper canopy or small branch zone of trees; (3) scansorial, mammals that are found in the middle and lower canopy and often climb up and down trees; (4) terrestrial-scansorial, terrestrial mammals that are not restricted to the ground, but often utilize the lower branches of trees; (5) terrestrial, mammals that are restricted to the ground; (6) fossorial, mammals that burrow in the ground; (7) aquatic, mammals that are found in water. Combination niche categories are also possible, e.g. terrestrial-arboreal or terrestrial-aquatic, as some species equally exploit the tree canopy and the ground, or underground dens and the land surface, albeit often for different sorts of activities like eating, sleeping and rearing young. There are also specialist niche exploitation categories, such as cursorial, which indicates a species that is well-adapted for running long distances over relatively open ground although they are generally considered as part of the terrestrial category. Later studies have further combined the arboreal and scansorial categories and re-named terrestrial-scansorial to terrestrial-arboreal and included all mammals that use both substrates (e.g., Reed 1997, Su and Harrison 2007), while others further refined the categories to accommodate the suite of specializations that allow the animal to exploit more than one ecological zone (e.g. Kovarovic et al. 2002). The less refined the categories, the easier it will be to assign fossil taxa to the appropriate category; however, it may also result in a decreased ability to distinguish between

faunal communities. Most studies also do not include the aerial category in their analysis due to the paucity of aerial mammals in the fossil record (Andrews et al. 1979, Reed 1997, Kovarovic et al. 2002, Su and Harrison 2007, among others).

*Dietary adaptations (trophic categories):* Trophic categories are assigned on the basis of the primary dietary adaptation of the animal, usually inferred from tooth morphology (specialization of molar cusps, degree of hypsodonty, canine morphology and reduction, see Evans, this volume), wear patterns (mesowear and microwear, see Green and Croft, this volume), and/or carbon stable isotope composition (see Higgins, this volume). The use of multiple proxies to infer the diet of fossil mammals can sometimes result in contradicting reconstructions where the lines of evidence do not align, for example carbon stable isotope data suggest that the species is eating both browse and graze even though its teeth are hypsodont (e.g., alcelaphin species from Laetoli, see Kingston and Harrison 2007). This is indicative of the difference between fundamental niche and realized niche (Hutchinson 1957), and, for the most part, categorization is based on what the species is actually eating at that particular site rather than what the species may have been adapted for over the course of its evolution. Furthermore, in some cases so little is preserved of the species or is understood about its ecological behavior that we default to the behavior of its relatives (extinct or extant) or “average” what we know about behavior if the taxon is identified only to a broad group such as genus or family.

The trophic or dietary categories commonly used include: (1) carnivore, animals that consume meat as a majority of its diet irrespective of taxonomic classification and can be further subdivided into animals that consume only meat, animals that consume meat and bone, animals that consume meat and invertebrates; (2) insectivore, animals that eat mostly insects irrespective of taxonomic classification; (3) grazers, animals that mostly eat grass; (4) fresh grass grazers, animals that eat specific types of grass from edaphic (i.e. where drainage

is impeded and there is often seasonal flooding) grasslands, (5) browsers, animals that mostly eat products of dicot bushes and trees; (6) mixed feeders, animals that eat both dicot and monocot leaves; (7) frugivore, animals that eat mostly fruits and can be further refined to animals that eat fruit and leaf and animals that eat fruit and insects (but often is not because it is difficult to distinguish these two categories based on ecomorphology); (8) omnivores, animals whose diet is wide-ranging and consists of food from more than two trophic categories; (9) root and tuber eater, animals that specialize in roots and tubers. The use of categories vary across studies, the difference is primarily in the degree of refinement in category definition and the use of combination categories (e.g., carnivore vs. carnivore-meat, carnivore-bone, carnivore-invertebrate, see Reed 1997 vs. Kovarovic 2002). As is the case with choosing locomotor or substrate categories, it will be easier to assign fossil taxa to broader categories, particularly if the ecology of the fossil species is not well known; however, this may also result in greater overlap in the community structure of faunal communities and make it more difficult to distinguish between them.

### **Multivariate statistics for a multivariate problem**

Since its inception, analyses of community structure have included univariate, bivariate, and multivariate comparisons. Profiles of community structure in histograms (for examples see Andrews et al. 1979a, Nesbit Evans et al. 1981, Kovarovic et al. 2002) and bivariate plots of ecovariables (for examples see Reed 1997, 2008, Kovarovic et al. 2002) are useful interpretative tools and can demonstrate ecological relationships, for example, the proportions of frugivorous mammals is inversely proportional to that of grazing mammals, with forest habitats having more frugivorous species and open habitats having more grazing species (see Figure 3). Univariate statistical tests are also appropriate when testing the null hypothesis that mean proportions of ecovariables are the same between habitat categories. For example, see Reed (2008) for her use of single factor ANOVA to test the null hypothesis



followed by the “Tukey honestly significant difference for uneven sample sizes” post-hoc test for differences between pairs of habitats for each ecovisible.

INSERT FIGURE 3 HERE; WIDTH = 1 COLUMN

However, community structure is fundamentally a multivariate problem and requires robust multivariate statistical techniques. It is important to note that multivariate analysis does not actually provide environmental data, rather, it organizes the data by ordination or classification in such way as to allow for visualization of the data and environmental interpretations (Andrews 1996). Different multivariate analyses have been used in studies of community structure, they are summarized as follows. (1) *Hierarchical clustering analysis*, a method of cluster analysis where a hierarchy of clusters is built using a measure of distance between pairs of observations and typically shown in a dendrogram (for examples of its application see Andrews 1996, Kovarovic et al. 2002, Su 2011). (2) *Principal components analysis* is an ordination technique where the data is transformed into a smaller number of uncorrelated variables called principal components. The first principal component contains as much of the variability of the data as possible and each remaining component contains as much of the remaining variability as possible (Zuur et al. 2007). The results are usually presented as a bivariate plot of its first and second principal components, which should contain most of the variance of the data (for examples of its application see Kovarovic et al. 2002, Flynn et al. 2003, Su and Harrison 2007). (3) *Principal coordinates analysis* is another ordination technique that is similar to PCA but uses a distance matrix instead of data transformation (Zuur et al. 2007); it is sometimes used in preference to principal components analysis because correlations within the ecological data matrix are generally weak (Andrews 1996, also see this reference for an example of its application). This analysis can be followed by a canonical analysis (discriminant analysis when the interest is in group membership and correlation analysis when the interest is in the relationships with environmental variables)

that allows for constrained ordination on the basis of any distance or dissimilarity measure (Anderson and Willis, 2003, see Lintulaakso & Kovarovic 2016 for an example of its application). (4) *Non-metric multidimensional scaling* is a non-metric ordination method designed to circumvent the assumption of linearity in principal coordinates analysis. Instead, it uses rank order information in a dissimilarity matrix to carry out iterative multidimensional scaling that produces differences in distribution and clustering that is analogous to the components in principal components analysis (Andrews 1996, also see this reference for an example of its application). (5) *Correspondence analysis* is a geometric technique used to visualize the associations between the levels of a two-way contingency table; rows and columns of the table are displayed as points in a low-dimensional space that is consistent with their associations in the table (Greenacre 1984). For examples that illustrate the application of this technique see Greenacre and Vrba (1984), Flynn et al. (2003), Reed (2008), Kovarovic et al. (2013), Su and Haile-Selassie (in review). (6) *Discriminant function analysis* is a statistical technique used to predict a categorical dependent variable (i.e., group membership) by one or more continuous or binary independent predictor variables. See Croft et al. (2008) and Su and Haile-Selassie (in review) for examples of the application of this technique and Kovarovic et al. (2011) for a review of some important considerations of this method.

The choice of which multivariate analysis to use is dependent on the data as the analyses often have different assumptions (see referred citations above for each method for computational details) and the questions being asked by the researcher. They are not mutually exclusive and are often used in conjunction with each other. With analyses that deal with proportional data, as is the case with community structure data, transformation of the raw data is often necessary. Sokal and Rohlf (1995) proposed using the arcsine square root transformation for these types of data and this has been widely adopted since then by ecologists and paleoecologists alike (Warton and Hui 2011). There are other types of

transformation that can be used (e.g. logit transformation; Warton and Hui, 2011); the choice of transformation and, importantly, its necessity, is determined by the research question and the assumptions made about the data (Clarke & Warwick, 1994; Kindt and Coe, 2005).

## **STRENGTHS OF COMMUNITY ANALYSIS**

The usefulness of the community structure approach to reconstructing past environments is multi-faceted. It makes use of what is often ample and easily accessible data (fossil remains), it is grounded in a particular theoretical context, the main principles of which (i.e. adaptation, niche exploitation) can be clearly observed in the modern natural world so comparative data is plentiful and reliable, and it is based on a complete mammalian community as defined by the ecological relationships that are demonstrated to exist between species and their environments. There are, of course, some challenges to utilizing the approach that we detail later in this chapter, but here we outline some of the more nuanced advantages to selecting this method over others, using examples to illustrate.

### **An All-Inclusive Approach**

Community structure analysis is holistic, in that it can include both micro- and macromammals. This is desirable because they relate to different aspects of their environment, including scale, and therefore provide complementary information on niche exploitation. Smaller taxa tend to more regularly exploit fruits and seeds and they are able to more efficiently exploit the arboreal and underground substrates in comparison to larger, generally more terrestrial species. Small species are also prey animals for larger carnivores reliant on meat consumption, while smaller species with carnivorous diets tend to eat a combination of small animal protein and insects, or insects in combination with other

resources. These differences suggest that eliminating smaller body mass categories from an analysis would also eliminate important ecological information provided by the many frugivorous, insectivorous, fossorial and arboreal taxa present. These niches, in essence, would be under-represented and discrimination of habitat types would be based on a limited array of niches, thus blurring the boundaries between them.

However, many paleontological sites present a taphonomic bias against the preservation of smaller-bodied species which can make their inclusion analytically challenging (see **Analytical Challenges** below, e.g. Kovarovic et al. 2002). Additionally, there are some taxonomic units traditionally left out of this type of analysis for other practical and ecological reasons. Volant mammals (i.e. Chiroptera) are excluded because of their rarity in the fossil record (Andrews et al. 1979, Reed 1998, Kovarovic et al. 2002, Mendoza et al. 2004, Louys et al. 2011). Despite the group displaying diverse ecological niche adaptations within it (bats vary considerably in body mass and diet across a wide spectrum) which would contribute to the discrimination of modern habitats, they are of limited usefulness in the fossil record in this regard. Mammals restricted entirely to aquatic environments are also often left out of the analysis, such as Cetacea and Sirenia, because it is the terrestrial habitat that is generally of interest at paleontological sites (although marine mammal community analyses can be conducted in their own right, see for example: Schick et al., 2011). The inclusion of an aquatic or semi-aquatic category is often to allow for categorisation of mammals that are partly adapted for an aquatic lifestyle, but still exploit the terrestrial environment in some fashion, such as hippopotami, otters, beavers and related species. This allows some element of water availability to be implicitly considered, without direct reference to the geological evidence.

### **“Taxon-Free”: Ecological Versus Species Diversity**

When taken as a sum of available niches and the proportion of species that exploit them as is the case in a community structure analysis, a paleohabitat can be viewed without specific reference to the identity of the resident species so, in a sense, this method may be thought of as “taxon-free”. This is perhaps one of the most important aspects of the method, since paleospecies can be difficult to classify taxonomically (see discussion in **Analytical Challenges** below). Taxonomic considerations in a paleocommunity under study are thus restricted to the determination of how many species are present (admittedly, not always in and of itself an easy task), not their exact identities or phylogenetic relationships, which is a somewhat separate area of study and one that is under frequent review for many families. What matters for a community structure analysis is what the individual species are doing as evidenced by their adaptations to particular niches. Diversity in terms of the ecological relationships and niche exploitation strategies employed by a community are far more relevant than species diversity alone.

Being unencumbered by taxonomy removes the reliance on “indicator species” in paleoecological reconstructions. This approach, which is implicitly imbued with the principle of taxonomic uniformitarianism, was a favoured analytical tool in earlier paleoecological work (e.g., Coe 1980; Vrba 1980; Greenacre & Vrba 1984). Much rigorous research went into determining which species are ecologically consistent in the modern world with the idea that they were likely to have behaved the similarly in the past; one oft-used example of this approach is the Antilopine-Alcelaphine Criterion developed by Vrba and colleagues (Vrba 1980, Greenacre & Vrba 1984) and applied to a number of sites in the East and South African paleoanthropological record (Kappelman 1984, Potts 1988, Shipman & Harris 1988). The AAC method was based on observations of extant bovid tribes and their tendency towards the consistent exploitation of specific trophic and spatial niches. Some tribes, such as the antilopines and alcelaphines, are known to prefer the resources of open, arid habitats where

they comprise 60% or more of the bovid fauna in these environments. When compared to the proportion of species in other tribal pairs such as Bovini and Reduncini, representing wetter, closed habitats, and Tragelaphini plus Aepycerotini (the impala, *Aepyceros melampus*), indicating dry and closed habitats, this criterion was understood to be a powerful predictor of paleoenvironment, particularly gross vegetation physiognomy. However, despite strong evidence that ecological similarities in closely related bovid taxa may be the result of evolutionary constraint and canalization of traits, not all tribes have had such stable ecological strategies over the same time scales (Sponheimer & Lee-Thorp 2003; Faith et al. 2014, Behrensmeyer 2015; Cerling et al 2015), or are known to be comprised of species that are evolutionarily sensitive to ecological change, such as the alcelaphine hartebeest, *Alcelaphus buselaphus* (e.g. Flagstad et al. 2001) or the impala and its extinct relatives in the genus *Aepyceros* (Faith et al. 2014). For reasons such as these, indicator species are approached with some caution. The AAC method, for example, has not been abandoned, but is used more conservatively and almost always in conjunction with other proxies, such as non-mammalian indicators, stable isotopes or depositional conditions (e.g. Schrenk et al 1995; Sponheimer & Lee-Thorp 2003; Campisano & Feibel 2007; Bedaso et al 2013). Community structure analyses are also often discussed in light of the taxonomic composition of the target community with particular reference to specific indicator taxa, but this is used to inform or support the main interpretations derived from the community analysis itself (e.g. Su 2016).

## **Time and Space**

Mammalian communities, being sensitive to ecological conditions, are especially useful when reconstructing habitats across a broad stretch of time or space (e.g. Fortelius et al. 2002). This is borne out by research in extant communities, particularly in small mammal communities, which experience turnover and re-assembly rapidly enough that they can be

observed over historical time periods or after a known environmental event that shifted habitat distributions, as well as on spatial scales where observations of presence/absence and ecological behavior are easily made (e.g Williams et al. 2002, Thibault et al. 2004, Thibault & Brown 2008). Macromammal evidence is less well established, but also suggests that communities change quickly in response to shifts in climate and that these changes are recorded in assemblages of skeletal and dental remains on modern landscapes (e.g. Western & Behrensmeyer 2009).

In paleontological locations where the fauna is known to have changed taxonomically over time, a full community analysis can bring greater insight to the ecological conditions that supported these different communities. Good examples of this approach are from the paleoanthropological sites of Laetoli and Olduvai in the southern Rift Valley in northern Tanzania where a significant amount of paleoecological work has been conducted in an effort to understand the environmental context of three hominin genera throughout the Pliocene and Pleistocene: *Australopithecus*, *Paranthropus* and *Homo*.

At Laetoli, *Australopithecus afarensis* is found in the 3.6-3.8 Ma Upper Laetoli Beds and *Paranthropus aethiopicus* in the 2.66 Ma Upper Ndolanya Beds (Leakey & Harris 1987, Harrison 2011). Community structure analyses indicated that the older beds were more wooded (Andrews 1989, Reed 1997) than early interpretations based on indicator species or single-family studies that emphasised the presence of terrestrial and grazing taxa (Leakey & Harris 1987). The composite mammalian community instead suggests that a range of habitats and resources were present, including grass cover to supported the grazing guild, but the proportion of frugivores, for example, indicates that fruiting vegetation cover was present, as well (Su & Harrison 2007, 2015). Both the overall locomotor and trophic niche profiles are most similar to modern Serengeti woodland habitats (Andrews 1989). In the same location approximately one million years later, the community structure looks somewhat different,

notably in the preponderance of terrestrial and grazing species at the expense of browsers and taxa relying on the arboreal substrate (Kovarovic et al. 2002). Even when the greater taphonomic loss of small-bodied species in this stratum is accounted for, the large mammal community most closely resembles a semi-arid bushland habitat (Kovarovic et al. 2002). This apparent trend towards increased aridity is also seen elsewhere in the region. Thirty-six km north of Laetoli, where the extensive deposits of Olduvai Gorge record the evolutionary history of hominins in great detail, the 2.0-1.8 Ma Bed I sediments (Deino 2012) indicate a change in climatic conditions towards greater aridity and the mammal community suggests a shift in the vegetation towards more open habitats at the end of the sequence (Fernandez-Jalvo et al. 1998). However, later in time, the mammalian community structure suggests that even in light of continued increasing aridity (Cerling & Hay 1986), the woody vegetation cover and the mammals it supported had stabilized. Although studies had long acknowledged differences in the Bed II fauna above and below the 1.74 Ma Lemuta Member (Leakey 1971, Hay 1976, 1990, Gentry & Gentry 1978), community structure analyses comparing Lower Bed II, where *Paranthropus* and *Homo habilis* remains are known, to Upper Bed II deposits in which *Paranthropus* persists and *Homo erectus* emerges, demonstrated that the faunal differences are most likely explained by taphonomy as both of the Bed II mammal communities were woodland-adapted (Kovarovic et al. 2013).

Work at Olduvai reminds us that mammal communities vary not only over time, but over space. A long programme of research at the site, particularly in Bed I and lowermost Bed II, has been focused on assessing landscape use of resident hominins (Blumeschne & Peters 1998, Blumeschne et al. 2003, 2012a,b), which necessarily considers both habitat and geology. Recursions of the paleo-lake Olduvai and periods where springs and river-fed wetlands were the main water sources shaped the landscape and modified habitat availability intermittently (Ashley et al. 2009, Stanistreet 2012). These hydrological phases would have



supported different vegetation complexes across the landscape. A full-scale spatial community structure analysis has yet to be undertaken, but faunal composition does differ spatially (Cushing 2002) and it can be hypothesized that areas of higher relief, where drainage was good and woody growth was possible, supported different communities from those located in areas of poorer drainage and marshy or otherwise more open habitats.

One difficulty in assessing community structure on a landscape scale is that in addition to the fact that mammals are not static users of their habitats and move around considerably, taphonomic conditions may not allow full communities to have been preserved at each place of interest, forcing researchers to “average” the fauna across a large area (see discussion in **Analytical Challenges** below). However, some sites such as Aramis, Ethiopia where *Ardipithecus ramidus* remains are known at 4.4 Ma, demonstrate that it may be possible to provide a more refined interpretation of community structure variation across a landscape (White et al. 2009). Here the mammals identified along a 7 km transect situated on a southeasterly gradient, vary in their functional adaptations to both the trophic and spatial niche. The entire mammalian community’s niche exploitation profile was not considered for both niches, but the selected macromammals studied are suggestive of full community differences inferring open habitats in the southeast and woodlands, with which *Ardipithecus* is associated, elsewhere.

## **ANALYTICAL CHALLENGES**

### **Locality Size and Habitat Homogeneity**

One issue to consider when conducting community structure analysis concerns the size and scale of the areas from which modern faunal communities derive. Very often, they are entire national parks or even entire regions of a country, which are likely to encompass

multiple habitat types. This results in faunal lists with species that are not actually found in the same area or habitat in nature, but artificially appear on the same faunal list due to the large scale of the faunal survey in question (e.g. Wilson 1975; Happold 1987). This makes it difficult to directly compare fossil and modern faunal communities because, often, fossil sites are much more constrained in size than the area from which modern comparative faunal communities are derived (Su and Harrison 2007; Louys et al. 2009) and also decreases the resolution of the interpretations possible. Thus, it is important that modern localities are chosen carefully to minimize latitudinal variation and to represent uniform topographical zones and vegetation types (Andrews et al. 1979a) although, in theory, the size of the area should not be a significant consideration as long as the habitat is homogeneous across the area covered. The most straightforward way to deal with this issue is to simply avoid faunal communities suspected of representing several major habitat types or to divide them as much as possible according to specific habitats that can be well-defined within the overall area. Alternatively, a statistical approach can be used to identify species lists whose composition is significantly distinct and under- or over-representing particular species groups. Louys et al. (2009) suggest a taxonomic distinctness analysis based on a measure developed by Warwick and Clarke (1995, 2001) which can determine communities that have a higher than average taxonomic distinctness, the potential result of sampling multiple habitats across a large area. These can then be excluded from further community analysis as they are non-representative of a natural habitat-specific fauna.

### **Defining and Assessing Paleo-Species**

Although it is not necessary to taxonomically identify individual fossil specimens in a community, discrete taxonomic units still need to be identified. Often material is too fragmentary to allow this and it is common to see that fossil assemblages have fewer species

overall (see **Fossil Assemblage Suitability** above) when compared to modern communities; this may relate both to the under-sampling of fauna generally, but also to our inability to actually identify all of the unique paleo-taxa. It may be difficult to distinguish between multiple species in a particularly diverse family or tribe with numerous closely related taxa, such that faunal lists may be comprised of generic or tribal groups, i.e. “Alcelaphini A” and “Alcelaphini B” . These distinctions may be based on differences in body mass because the morphology is so similar. In these cases, as long as they can be recognized as distinct taxonomic units and we can assign ecovariation categories to them, they should be included in the analysis.

### **Taphonomy and Sampling**

There are many taphonomic factors that impact the composition of the final fossil assemblage in such a way that it no longer reflects the composition of the original living community. Most of them involve differential destruction and transportation of bones due to fluvial factors, carnivore and rodent activities, weathering, and trampling (for examples, see Voorhies, 1969a,b; Behrensmeyer, 1975, 1980; papers in Behrensmeyer & Hill, 1980; Blumenschine, 1988). Furthermore, collection strategies and/or sampling, regardless of how thorough, are probably not sampling the entirety of what was once the living community. For example, at Laetoli, all specimens identifiable by taxon or anatomical elements are collected (Harrison 2011). This is an intensive collecting strategy that ensures a representative fossil assemblage. However, even then, smaller taxa are significantly under-represented. Size-abundance analysis on herbivores larger than 1 kg indicate that species less than 10 kg are under-represented in the Laetoli fossil assemblage and that the assemblage is taphonomically biased by surface processes, such as carnivore ravaging, weathering, erosion, and trampling (Su 2005). It is important to note that because community structure analysis relies on faunal

lists, many of these taphonomic biases are ameliorated and we are not necessarily concerned with correcting these biases prior to conducting an analysis (Andrews et al. 1979a). It is, however, useful to identify the taphonomic biases present in the paleontological assemblage of interest for interpreting the results.

Another important taphonomic consideration is the general under-representation of small mammals in fossil assemblages. Although they are useful habitat indicators in their own right and can be subjected to their own community structure analysis to aid paleoecological interpretations (e.g., D. Reed 2007, Avery 2007, Stoetzel et al. 2011), they present unique challenges to paleoenvironmental inferences based on community structure. They are often accumulated under different taphonomic regimes from the large mammals in fossil assemblages (Andrews 1990, Reed 1997, D. Reed 2007) or are so poorly preserved that they are under-represented in the fossil community (Andrews et al. 1979a, Reed 1997, Su and Harrison 2007). Many researchers make the decision, based on these factors, to exclude them from the analysis, sometimes using 500 g, 1 kg, or 4 kg thresholds, to reduce the need to consider their taphonomic biases in interpreting the results (e.g., Reed 1997, 2008, Su and Harrison 2007, Louys et al. 2009, Su 2011, 2016). Where small species are clearly biased in an assemblage, a statistical solution may be to employ a sampling technique that modifies the comparative modern data such that the body mass profiles match that of the target paleontological community (Kovarovic et al 2013). However, eliminating smaller body mass categories would also eliminate important ecological information and the analysis would thus be based on a limited array of niches that may not discriminate between different habitats as well.

## **FURTHER CONSIDERATIONS AND FUTURE DIRECTIONS**

## **A dynamic biome**

In community structure analysis and other faunal-based methods of paleoenvironmental reconstruction, we are using ecological information derived from animals as a proxy for the vegetation structure that was present. We are, in effect, assuming that mammalian community structures in similar biomes are convergent regardless of their location in time and space and that we are able to distinguish between the biomes based on the fauna (and their adaptations). However, biomes are dynamic entities, changed by temporal, geographic and scale-dependant events, which also change the structure of the associated mammalian communities.

*Temporal factors.* Temporal changes in the climate and physical environment occur over ecological (days to millenia) and geological (orbital to tectonic) time frames and floral and faunal communities have to adjust to these changes (Overpeck et al., 2003). The circadian response of vegetation to daily differences in the amount of solar energy can be observed as the movement of leaves or the opening and closing of flowers, for example, but it is change that occurs on a longer time scale that can impact on palaeoenvironmental reconstruction.

Within decadal and centennial time frames, primary or secondary succession of vegetation alters plant species composition of the biome itself, and as a result, of the mammalian community structure. For example, after a disturbance to a forested area, whether by storm, wildfire or some other event, pioneering species invade the new area quite rapidly and begin a new succession of plants and animals. Often, at the beginning, the animals are r-selected species but as the succession continues, more K-selected animals eventually inhabit the forest (Pianka, 1970). Thus, within a relatively short time span, the faunal community structure can change in the same geographic location, which may impact paleoenvironmental reconstruction.

Climatic events, like ice ages, can cause millennial changes in vegetation (Hessler et al., 2010, Overpeck et al., 2003). Pollen records from both marine and terrestrial sediment cores demonstrate that during the last glacial period, as a result of global temperature change, cold- and warm-adapted plant species altered their distributions (Hessler et al., 2010, Overpeck et al., 2003). As plant composition alters and the biomes gradually change, mammalian communities respond to these changes by regional- or continental-scale migration, extinction, or by subspecies level selection (Overpeck et al., 2003), which likely results in a shift in community structure.

Larger changes in the biomes occur over geological time scales, both at the orbital and tectonic levels. The orbital level, which spans between 10,000 and 100,000 years, includes glacial-interglacial cycles and relates to the orbital forces described by Milankovitch (Ruddiman, 2007). These cycles have repeatedly formed and broken up biomes, causing extinction and speciation events in both plant and mammalian communities. There would have been biomes in the past that have no present-day equivalent. Biomes are also affected by plate tectonics and continental drift. These changes take millions of years and cause major changes in biomes by creating new mountain ranges, opening and closing of water courses, and altering global climatic conditions. At this scale, mammalian communities can change drastically in species composition and community structure as they are being shaped by major extinction and speciation events (Janis, 1993).

*Geographic factors.* Geography is significant for the evolution of biomes and their associated mammalian fauna. Both latitudinal and altitudinal change or the presence of a continental or island environment alter climatic variability and the rate of dispersion and speciation. The amount of received solar energy is directly related to latitude; low and mid-latitudes have high solar radiation and higher latitudes have low solar radiation (Ruddiman, 2007). Similar phenomena can be observed with patterns of species richness: it is higher in

the tropics and lower towards the poles. Several hypotheses have been proposed to explain this pattern, including (1) warm climates have faster rates of microevolution (Gillman et al., 2009) and (2) the tropics have more energy available in usable form, which results in more organisms, and hence, more species, the environment can support (=species richness-energy hypothesis; Brown, 1981). While these have not yet been sufficiently tested, it is clear that latitude and the amount of solar energy have important implications for climate, vegetation and mammalian communities. A similar pattern exists with altitude. Atmospheric temperature decreases by 6.5 °C/km as altitude increases (Coe, 2009), which has a significant impact on the vegetation and the associated mammalian community (see, for example, Geise et al., 2004).

Other geographical factors such as the size, shape and type of the observed area (continent or smaller island) also impact vegetation type and structure of the mammalian community. Studies of island biogeography have noted that islands closer to the continental area have higher species richness than those further away, as they are easier to reach by migrating populations (MacArthur & Wilson, 1967). In addition, larger islands have more resources than smaller ones, thus there is a higher probability of immigration of continental animals (MacArthur & Wilson, 1967).

*Geographic scale.* Changes in vegetation or mammalian communities can also be analyzed at different geographical scales, from the micro- to macro-level. At the macro-level, the analyzed scale is at the continental level where local changes are insignificant from an analytical perspective. For example, as the climate became increasingly arid during the Miocene, grass-dominated open habitats spread while closed forested areas decreased. This had a direct impact on mammalian herbivore tooth structure, which evolved from low-crowned (brachyodont) to medium (mesodont) or high-crowned (hypsodont) teeth in response to their changing trophic resources (Eronen et al., 2010a, Strömberg, 2011). At the

meso-level, changes in vegetation and mammalian communities are observed *within* continents. A change in climate may cause the disappearance of one biome and the appearance or increase in size of another, as was the case during the late Pleistocene and Holocene when the same general climatic events affected both South America and Africa. African mammalian communities remained almost unchanged while those of South America changed dramatically, leading to present-day faunal dissimilarities between the two continents (De Vivo & Carmignotto, 2004). At the micro-level, changes in vegetation and mammalian communities are observed in local areas such as a specific island, mountain, valley, or basin. For example, high levels of endemism exist not only in the Philippine islands as a whole, but within individual islands, as well. The repeated dropping and rising of sea levels during the Pleistocene joined and disconnected the Philippine islands numerous times, forming the so called Pleistocene island groups with unique sets of species (Catibog-Singa et al., 2006).

*Implications for community structure analysis:* Most of the above mentioned temporal, geographic and scale-dependant events relate to historical contingency, or the effect of the order and timing of past events on community assembly. Abiotic events like wildfires, floods, storms and earthquakes change species arrival history of a location or an area. These biotic events have different priority effects on the order in which species arrive to the primary or secondary succession area. Although quite often the succession may follow an expected order, the consequences of species arrival history can cause differences in the structure of communities, as effects are amplified over time and space via population growth and interactions (Fukami, 2015). The results of this are that even in relatively similar environments, faunal composition can be different due to their different histories. This suggests that historical contingency may be an important factor to consider when interpreting the results of community structure analyses (see **Moving Forward** for more discussion).



## Moving Forward

The more nuanced our understanding of the relationship between a biome and its associated mammalian community (and its ecological structure), the better we can infer past environmental and climatic conditions based on proxies. There is much that can be done to refine community structure analyses, thus guaranteeing ever more robust and detailed environmental reconstructions. We suggest that development in the following three areas would be most beneficial to improving the methodology:

(1) *Historical contingency*. As we previously discussed, paleoecological studies rely on our assumption of community convergence and, indeed, in the Tropical and northern (Palearctic and Nearctic) zones, there are general similarities in community structure between localities of similar latitudes on different continents. However, this apparently only applies to broad locomotor and dietary variables in the Tropics (Lintulaakso & Kovarovic, 2016) and between extreme environments that are similar, e.g. desert and steppe communities with high levels of water stress and low complexity in vegetation structure (Rodríguez et al., 2006). Historical effects are likely to be responsible for differences that have been shown to exist between communities in similar habitats (Losos 1996; Rodríguez et al., 2006; Graham 2008; Fukami, 2015; Lintulaasko and Kovarovic, 2016), but it is not yet clear how historical contingency impacts on the application of community structure analysis to palaeoenvironmental reconstruction. Thus, a better understanding of the effects of historical contingency should be considered as one of the next steps in improving the method.

(2) *Continuous variables*. Community structure analysis is an excellent way of reconstructing the spatial and trophic niches that were available to past communities. But, it can be a coarse approximation that forces habitats into discrete and broadly defined categories, glossing over the nuanced differences between them (regardless of the

mechanisms that are responsible for them). Steps are being taken to to move away from category-driven approaches, with greater emphasis placed on the continuous nature of many environmental variables, such as the amount of tree cover and how this can be inferred (Louys et al., 2015). The incorporation of continuous variables, rather than just the use of simplistic categories, will be essential for our ability to provide a more nuanced and accurate reconstruction of paleoenvironment and a better understanding of the relationship between the biome and their associated faunal communities.

(3) *Paleoclimate reconstructions*. The method as it is generally applied also lacks the ability to clearly identify paleoclimatic factors, focusing largely on a reconstruction of the physical environment rather than climate. However, the use of averages of ecometric traits (species traits which are easily measurable, whose structure is closely related to their function, and whose function interacts directly with local environment) in communities has been shown to be connected to precise measures of climate variables, such as the amount of precipitation or temperature (Eronen et al., 2010b, Fortelius et al., 2016). These types of data allow us to characterize physical habitats at the regional scale, linking them specifically to primary production (Fortelius et al., 2016).

Community structure analysis is rooted in the niche concept and the idea that faunal communities and the adaptations of their members can provide information about their environment. It is a powerful tool for reconstructing paleoenvironments. While it does not rely on a single mammalian taxon, it does incorporate ecometrics and taxon-specific analyses (dietary and locomotor reconstructions for specific taxa) to arrive at a holistic and community-level perspective on paleoenvironmental reconstruction. With further development and refinement, this methodology can help move us away from categorical descriptions of paleoenvironments and emphasize the continuous nature of ecological

variables. This will allow for more nuanced reconstructions of paleoenvironments and a better understanding of the relationship between biotic communities and their environments.

#### ACKNOWLEDGEMENTS

KK and KL gratefully acknowledge Denise Su, Darin Croft and Scott Simpson for the invitation to participate in the “Latest Methods in Reconstructing Cenozoic Terrestrial Environments and Ecological Communities” symposium in Cleveland 2015. KK sincerely thanks the organisers for support and assistance with childcare during the symposium and for useful discussions throughout. We also thank Peter Andrews for his continued advice and insight.

## REFERENCES

- Alexander, R.M. (1977). Allometry of the limbs of antelopes (Bovidae). *Journal of Zoology, London*, 183, 125-146.
- Alverson, K. D., Bradley, R. S., & Pedersen, T. F. (2003). *Paleoclimate, global change and the future*. Springer Science & Business Media.
- Anderson, M.J., & Willis, T.J. (2003). Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. *Ecology*, 84, 511-525.
- Andrews, P. (1989). Paleocology of Laetoli. *Journal of Human Evolution*, 18, 173–181.
- Andrews, P. (1990a). *Owls, Caves, and Fossils*. Chicago: University of Chicago Press.
- Andrews, P. (1990b). Small Mammal Taphonomy. In Lindsey, E.H., Fahlbusch, V., Mein, P. (Eds.) *European Neogene Mammal Chronology* (pp. 487-494). New York: Plenum Press.
- Andrews, P. (1996). Paleocology and hominoid palaeoenvironments. *Biological Reviews*, 71, 257–300.
- Andrews, P. & Humphrey, L. (1999). African Miocene environments and the transition to early hominines. In T.G.Bromage & F. Schrenk (Eds.), *African Biogeography. Climate Change & Human Evolution* (pp. 282-300). Oxford: Oxford University Press.
- Andrews, P., & Nesbit Evans, E. (1979). The environment of *Ramapithecus* in Africa. *Paleobiology*, 5, 22-30.
- Andrews, P., & Van Couvering, J. H. (1975). Paleoenvironments in the East African Miocene. In F. Szalay (Ed.), *Approaches to Primate Paelobiology* (pp. 62-103). Basel: Karger.
- Andrews, P., Lord, J.M. & Nesbit Evans, E.M. (1979). Patterns of ecological diversity in fossil and modern mammalian faunas. *Biological Journal of the Linnean Society*, 11, 177-205.
- Ashley, G.M., Tactikos, J.C. & Owen, R.B. (2009). Hominin use of springs and wetlands: paleoclimate and archaeological records from Olduvai Gorge (1.79-174 Ma). *Palaeogeography Palaeoclimatology Palaeoecology*, 272, 1-16.
- Avery, D.M. (2007). Micromammals as palaeoenvironmental indicators of the southern African Quaternary. *Transactions of the Royal Society of South Africa*, 62, 17-23.
- Badgley, C., Damuth, J., Theodor, J. & Werdelin, L. (2001). Global variation in ecological structure of mammalian faunas and paleoenvironmental inferences. *Journal of Vertebrate Paleontology*, 21 (suppl.), 30A.

- Bedaso, Z.K., Wynn, J.G., Alemseged, Z. & Geraads, D. (2013). Dietary and paleoenvironmental reconstruction using stable isotopes of herbivore tooth enamel from middle Pliocene Dikika, Ethiopia: implication for *Australopithecus afarensis* habitat and food resources. *Journal of Human Evolution*, 64, 21-38.
- Behrensmeyer, A.K. (1975). Taphonomy and paleoecology in the hominid fossil record. *Yearbook of Physical Anthropology*, 19, 36-50.
- Behrensmeyer, A.K. (1978). Taphonomic and ecologic information from bone weathering. *Paleobiology*, 4, 150-162.
- Behrensmeyer, A.K., & Hill, A.P. (1980). *Fossils in the making*. Chicago: University of Chicago Press.
- Behrensmeyer, A.K. (2015). Four million years of African herbivory. *Proceedings of the National Academy of Sciences of the United States of America*, 112, 11428–11429.
- Blumenschine, R.J. (1988). An experimental model of the timing of hominid and carnivore influence on archaeological bone assemblages. *Journal of Archaeological Science*, 15, 483-502.
- Blumenschine, R.J. & Peters, C.R. (1998). Archaeological predictions for hominid land use in the paleo-Olduvai Basin, Tanzania, during lowermost Bed II times. *Journal of Human Evolution*, 34, 565-607.
- Blumenschine, R.J., Peters, C.R., Masao, F.T., Clarke, R.L., Deino, A.L., Hay, R.L., Swisher, C.C., Stanistreet, I.G., Ashley, G.M., McHenry, L.J., Sikes, N.E., van der Merwe, N.J., Tactikos, J.C., Cushing, A.E., Deocampo, D.M., Njau, J.K. & Ebert, J.I. (2003). Late Pliocene Homo and hominid land use from western Olduvai Gorge, Tanzania. *Science*, 299, 1217-1221.
- Blumenschine, R.J., Masao, F.T., Stollhofen, H., Stanistreet, I.G., Bamford, M.K., Albert, R.M., Njau, J.K. & Prassack, K.A. (2012a). Landscape distribution of Oldowan stone artifact assemblages across the fault compartments of the eastern Olduvai Lake Basin during early lowermost Bed II times. *Journal of Human Evolution*, 63, 384-394.
- Blumenschine, R.J., Stanistreet, I.G. & Masao, F.T. (2012b). Olduvai Gorge and the Olduvai landscape paleoanthropology project. *Journal of Human Evolution*, 63, 247-250.
- Brown, J. H. (1981). Two decades of homage to Santa Rosalia: toward a general theory of diversity. *American Zoologist*, 21(4), 877-888.
- Campisano, C.J. & Feibel, C.S. (2007). Connecting local environmental sequences to global climate patterns: evidence from the hominin-bearing Hadar Formation, Ethiopia. *Journal of Human Evolution*, 53, 515-527.
- Catibog-Singa, C., Catibog, C. S., & Heaney, L. R. (2006). *Philippine biodiversity: principles and practice*. Haribon Foundation.

- Cerling, T.E., Andanje, S.A., Blumenthal, S.A., Brown, F.H., Chritz, K.L., Harris, J.M., Hart, J.A., Kirera, F.M., Kaleme, P., Leakey, L.N., Leakey, M.G., Levin, N.E., o Manthi, F.K., Passey, B.H. & Uno, K.T. (2015). Dietary changes of large herbivores in the Turkana Basin, Kenya from 4 to 1 Ma. *Proceedings of the National Academy of Sciences of the United States of America*, 112, 11467-11472.
- Cerling, T.E. & Hay, R.L. (1986). An isotopic study of paleosol carbonates from Olduvai Gorge. *Quaternary Research*, 25, 63-78.
- Clarke, K.R., & Warwick, R.M. (2001). A further biodiversity index applicable to species lists: variation in taxonomic distinctness. *Marine Ecology Progress Series*, 216, 265–278.
- Clarke, K.R. & Warwick, R.M. (1994). An approach to statistical analysis and interpretation. *Change in Marine Communities*, 2.
- Coe, H. (2009). Atmospheric energy and the structure of the atmosphere. In C.N. Hewitt & A.V. Jackson (Eds.), *Atmospheric science for environmental scientists* (pp. 54-82). John Wiley & Sons.
- Coe, M. (1980). The role of modern ecological studies in the reconstruction of palaeoenvironments in sub-Saharan Africa. In A.K. Behrnesmeyer & A.P. Hill (Eds.), *Fossils in the Making: Vertebrate Taphonomy and Paleocology* (pp. 55-67). Chicago: The University of Chicago Press.
- Croft, D. A., Flynn, J. J., & Wyss, A. R., 2008. The Tinguiririca Fauna of Chile and the early stages of “modernization” of South American mammal faunas. *Archivos Museu Nacional*, 66, 191-211.
- Cushing, A.E. (2002). The Landscape Zooarchaeology and Paleontology of Plio-Pleistocene Olduvai, Tanzania and Their Implications for Early Hominid Ecology. PhD Dissertation: Rutgers, the State University of New Jersey.
- Damuth, J., & McFadden, B.J. (1990). *Body Size in Mammalian Paleobiology: Estimation and Biological Implications*. New York: Cambridge University Press.
- Damuth, J., Fortelius, M., Andrews, P., Badgley, C., Hadly, E.A., Hixon, S., Janis, C., Madden, R.H., Reed, K., Smith, F.A., Theodor, J., Van Dam, J.A., Van Valkenburgh, B. & Werdelin, L. (2002). Reconstructing mean annual precipitation based on mammalian dental morphology and local species richness. *Journal of Vertebrate Paleontology*, 22 (suppl.), 48A.
- De Vivo, M., & Carmignotto, A. P. (2004). Holocene vegetation change and the mammal faunas of South America and Africa. *Journal of Biogeography*, 31(6), 943-957.
- Deino, A.L. (2012). <sup>40</sup>Ar/<sup>39</sup>Ar dating of Bed I, Olduvai Gorge, Tanzania, and the chronology of early Pleistocene climate change. *Journal of Human Evolution*, 63, 251-273.

- Eisenberg, J.F. (1981). *The mammalian radiations: an analysis of trends in evolution, adaptation, and behavior*. University of Chicago Press
- Eiten, G. (1992). How names are used for vegetation. *Journal of Vegetation Science*, 3, 419-424.
- Eronen, J. T., Puolamäki, K., Liu, L., Lintulaakso, K., Damuth, J., Janis, C., & Fortelius, M. (2010). Precipitation and large herbivorous mammals II: application to fossil data. *Evolutionary Ecology Research*, 12(2), 235-248.
- Eronen, J.T., Polly, P.D., Fred, M., Damuth, J., Frank, D.C., Mosbrugger, V., Scheidegger, C., Stenseth, N.C., Fortelius, M. (2010) Ecometrics: the traits that bind the past and present together. *Integrative zoology*. 5(2):88-101.
- Faith, J.T., Tryon, C.A., Peppe, D.J., Beverly, E.J. & Blegen, N. (2014). Biogeographic and evolutionary implications of an extinct late Pleistocene impala from the Lake Victoria Basin, Kenya. *Journal of Mammalian Evolution*, 21, 213-222.
- Fernandez-Jalvo, Y., Denys, C., Andrews, P., Williams, T., Dauphin, Y., Humphrey, L. (1998). Taphonomy and paleoecology of Olduvai Bed-I (Pleistocene, Tanzania). *Journal of Human Evolution*, 34, 137-172.
- Flagstad, Ø., Syversten, P.O., Stenseth, N.C. & Jakobsen, K.S. (2001). Environmental change and rates of evolution: the phylogeographic pattern within the hartebeest complex as related to climatic variation. *Proceedings of the Royal Society B*, 268, 667-677.
- Fleming, T. H. (1972). Numbers of mammal species in north and central American forest communities. *Ecology*, 54, 555-563.
- Flynn, J. J., Wyss, A. R., Croft, D. A., & Charrier, R., 2003. The Tinguiririca Fauna, Chile: biochronology, paleoecology, biogeography, and a new earliest Oligocene South American Land Mammal 'Age'. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 195, 229-259.
- Fortelius, M. (1990). Problems using fossil teeth to estimate body sizes of extinct mammals. In Damuth, J. & McFadden, B.J. (Eds.) *Body Size in Mammalian Paleobiology: Estimation and Biological Implications* (pp. 207-288). New York: Cambridge University Press.
- Fortelius, M., Žliobaitė, I., Kaya, F., Bibi, F., Bobe, R., Leakey, L., Leakey, M., Patterson, D., Rannikko, J., Werdelin, L. (in press). An ecometric analysis of the fossil mammal record of the Turkana Basin. *Philosophical Transactions B*.
- Fortelius, M., Eronen, J., Jernvall, J., Liu, L., Pushkina, D., Rinne, J., Tesakov, A., Vislobokova, I., Zhang, Z., & Zhou, L. (2002). Fossil mammals resolve regional patterns of Eurasian climate change over 20 million years. *Evolutionary Ecology Research*, 4, 1005-1016.

- Fukami, T. (2015). Historical contingency in community assembly: integrating niches, species pools, and priority effects. *Annual Review of Ecology, Evolution, and Systematics*, 46(1), 1.
- Geise, L., Pereira, L. G., Bossi, D. E. P., & Bergallo, H. G. (2004). Pattern of elevational distribution and richness of non volant mammals in Itatiaia National Park and its surroundings, in southeastern Brazil. *Brazilian Journal of Biology*, 64(3B), 599-612.
- Gentry, A.W. & Gentry, A. (1978). Fossil Bovidae (Mammalia) of Olduvai Gorge, Tanzania, Part II. *Bulletin of the British Museum of Natural History (Geology)*, 30, 1-83.
- Gillman, L. N., Keeling, D. J., Ross, H. A., & Wright, S. D. (2009). Latitude, elevation and the tempo of molecular evolution in mammals. *Proceedings of the Royal Society of London B: Biological Sciences*, 276(1671), 3353-3359.
- Gingerich, P.D., Smith, B.H., & Rosenberg, K. (1982). Allometric scaling in the dentition of primates and prediction of body weight from tooth size in fossils. *American Journal of Physical Anthropology*, 58, 81-100.
- Gould, S.J. (1975). On the scaling of tooth size in mammals. *American Zoologist*, 15, 351-362.
- Graham, C. H., & Fine, P. V. (2008). Phylogenetic beta diversity: linking ecological and evolutionary processes across space in time. *Ecology letters*, 11(12), 1265-1277.
- Greenacre, M.J. (1984). *Theory and applications of Correspondence Analysis*. New York: Academic Press.
- Greenacre, M.J. & Vrba, E.S. (1984). Graphical display and interpretation of antelope census data in African wildlife areas, using correspondence analysis. *Ecology*, 65, 984-997.
- Happold, D.C.D. (1987). *The mammals of Nigeria*. Oxford: Clarendon Press.
- Harrison, J. L. (1962). The distribution of feeding habits among animals in a tropical rain forest. *Journal of Animal Ecology*, 31, 53-63.
- Harrison, T. (2011). Laetoli Revisited: Renewed Paleontological and Geological Investigations at Localities on the Eyasi Plateau in Northern Tanzania. In T. Harrison (Ed.), *Paleontology and Geology of Laetoli: Human Evolution in Context. Volume 1: Geology, Geochronology, Paleoecology and Paleoenvironment* (pp. 1-15). Dordrecht: Springer Press.
- Hay, R.L., (1976). *The Geology of Olduvai Gorge: a Study of Sedimentation in a Semiarid Basin*. Berkeley: University of California Press.
- Hay, R.L., 1990. Olduvai Gorge: a case history in the interpretation of hominid paleoenvironments in East Africa. In: L.F. Laporte (Ed.), *Establishment of a Geological Framework for Paleoanthropology* (pp. 23-37). Geological Society of America Special Paper 242.



- Hessler I, Dupont L, Bonnefille R, Behling H, González C, Helmens K.F., Hooghiemstra H, Lebamba J, Ledru M.P., Lézine A.M., Maley J., 2010. Millennial-scale changes in vegetation records from tropical Africa and South America during the last glacial. *Quaternary Science Reviews*, 29(21), 2882-99.
- Hopkins, S.S.B. (2008). Reassessing the mass of exceptionally large rodents using toothrow length and area as proxies for body mass. *Journal of Mammalogy*, 89, 232–243.
- Hutchinson, G.E. (1957). The multivariate niche. *Cold Spring Harbor Symposium of Quantitative Biology*, 22, 415-421.
- Janis, C. M. (1993). Tertiary mammal evolution in the context of changing climates, vegetation, and tectonic events. *Annual Review of Ecology and Systematics*, 467-500.
- Kappelman, J. (1984). Plio-Pleistocene environments of Bed I and lower Bed II, Olduvai Gorge, Tanzania. *Palaeogeography Palaeoclimatology Palaeoecology*, 48, 171–196.
- Karr, J. R. (1971). Structure of avian communities in selected Panama and Illinois habitats. *Ecological Monographs*, 41, 207-233.
- Kovarovic, K., Aiello, L.C., Cardini, A. & Lockwood, C.A. (2011). Discriminant function analyses in archaeology: are classifications rates too good to be true? *Journal of Archaeological Science*, 38, 3006-3018.
- Kovarovic, K., Andrews, P. & Aiello, L.C (2002). The palaeoecology of the Upper Ndolanya Beds at Laetoli, Tanzania. *Journal of Human Evolution*, 43, 395–418.
- Kovarovic, K., Slepkov, R. & McNulty K.P. (2013). Ecological continuity between Lower and Upper Bed II, Olduvai Gorge, Tanzania. *Journal of Human Evolution*, 64, 538-555.
- Kurtén, B. (1952): The Chinese Hipparion fauna. *Commentationes Biologicae, Societas Scientiarum Fennica* 13:1–82
- Lawesson, J.E. (1994). Some comments on the classification of African vegetation. *Journal of Vegetation Science*, 5, 441-444.
- Leakey, M.D., 1971. *Olduvai Gorge: Excavations in Beds I and II, 1960-1963, vol. 3*. Cambridge: Cambridge University Press.
- Leakey, M.D. & Harris, J.M., Eds. (1987). *Laetoli: A Pliocene Site in Northern Tanzania*. Oxford: Clarendon Press.
- Legendre, S. (1986). Analysis of mammalian communities from the late Eocene and Oligocene of Southern France. *Paleovertebrata*, 16, 191-212.
- Legendre, S. (1989). Les communautés de mammifères du Paléogène (Eocène supérieur et Oligocène) d'Europe occidentale: structures, milieux et evolution. *Münchner Geowissenschaftliche Abhandlungen A*, 16, 1-110.

- Lintulaakso, K., & Kovarovic, K. (2016). Diet and locomotion, but not body size, differentiate mammal communities in worldwide tropical ecosystems. *Palaeogeography, Palaeoclimatology, Palaeoecology*, *454*, 20-29.
- Lomolino, M. V., Riddle, B. R., Brown, J. H., & Brown, J. H. (2006). *Biogeography*. Sunderland, MA: Sinauer Associates.
- Losos, J.B. (1996). Phylogenetic perspectives on community ecology. *Ecology*, 1344-1354.
- Louys, J. (2007). *Ecology And Extinction of Southeast Asia's Megafauna*. PhD thesis: University of New South Wales.
- Louys, J., Meloro, C., Elton, S., Ditchfield, P., & Bishop, L. C. (2015). Analytical framework for reconstructing heterogeneous environmental variables from mammal community structure. *Journal of human evolution*, *78*, 1-11.
- Louys, J., Meloro, C., Elton, S., Ditchfield, P. & Bishop, L.C. (2011). Mammal community structure correlates with arboreal heterogeneity in faunally and geographically diverse habitats: implications for community convergence. *Global Ecology and Biogeography*, *20*, 717-729.
- Louys, J., Travouillon, K.J., Bassarova, M. & Tong, H. (2009). The use of protected natural areas in palaeoecological analyses: assumptions, limitations and application. *Journal of Archaeological Science*, *36*, 2274–2288.
- Magurran, A.E. (1988). *Ecological Diversity and its Measurement*. London: Croom Helm.
- Meloro, C., & Kovarovic, K. (2013). Spatial and ecometric analyses of the Plio-Pleistocene large mammal communities of the Italian peninsula. *Journal of Biogeography*, *40*, 1451-1462.
- Mendoza, M., Goodwin, B. & Criado, C. (2004). Emergence of community structure in terrestrial mammal-dominated ecosystems. *Journal of Theoretical Biology*, *230*, 203-214.
- MacArthur, R. H., & Wilson, E. O. (1967). *Theory of Island Biogeography*. Princeton, NJ: Princeton University Press.
- Nesbit Evans, E. M., Van Couvering, J. A. H., & Andrews, P. (1981). Paleoecology of Miocene sites in western Kenya. *Journal of Human Evolution*, *10*, 99-116.
- Odum, E.P. (1959). *Fundamentals of Ecology*. 2<sup>nd</sup> ed. Philadelphia: W.B. Saunders Company.
- Olson, J.S., Watts, J.A. & Allison, L.J. (1983). *Carbon In Live Vegetation Of Major World Ecosystems*. Oak Ridge, TN: Oak Ridge National Laboratory.
- Olson, J.S., Watts, J.A. & Allison, L.J. (1985). *Major World Ecosystem Complexes Ranked by Carbon in Live Vegetation* (NDP-017).

- Overpeck, J., Whitlock, C., & Huntley, B. (2003). Terrestrial biosphere dynamics in the climate system: past and future. In *Paleoclimate, global change and the future* (pp. 81-103). Springer Berlin Heidelberg.
- Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007). *Updated world map of the Köppen-Geiger climate classification*. Hydrology and Earth System Sciences Discussions Discussions, 4(2), 439-473.
- Pianka, E.R. (1970). On r-and K-selection. *The American Naturalist*, 104, 592-597.
- Pimm, S.L., & Lawton, J.H. (1978). On feeding on more than one trophic level. *Nature*, 275, 542-544.
- Pineda-Munoz, S., Evans, A. R., & Alroy, J. (2016). The relationship between diet and body mass in terrestrial mammals. *Paleobiology*, 1–11.
- Potts, R. (1988). *Early Hominid Activities at Olduvai Gorge*. New York: Aldine de Gruyter.
- Reed, D.N. (2007). Serengeti micromammals and their implications for Olduvai paleoenvironments. In (Bobe, R., Alemseged, Z., & Behrensmeyer, A.K. (Eds) *Hominin environments in the East African Pliocene: An assessment of the faunal evidence* (pp. 217-255). Dordrecht: Springer.
- Reed, K.E. (1997). Early hominid evolution and ecological change through the African Plio-Pleistocene. *Journal of Human Evolution*, 32, 289-322.
- Reed, K.E. (1998). Using large mammal communities to examine ecological and taxonomic structure and predict vegetation in extant and extinct assemblages. *Paleobiology*, 24, 384-408.
- Reed, K.E. (2008). Paleoecological patterns at the Hadar hominin site, Afar Regional State, Ethiopia. *Journal of Human Evolution*, 54, 743-768.
- Rodríguez, J., Hortal, J., & Nieto, M. (2006). An evaluation of the influence of environment and biogeography on community structure: the case of Holarctic mammals. *Journal of Biogeography*, 33(2), 291-303.
- Ruddiman, W.F. (2007). *Earth's Climate: Past and Future*. Macmillan.
- Schick, R.S., Halpin, P.N., Read, A.J., Urban, D.L., Best, B.D., Good, C.P., Roberts, J.J, LaBrecque, E.A., Dunn, C., Garrison, L.P., Hyrenbach, K.D., McLellan, W.A., Pabst, D.A., Palka, D.L. & Stevick, P. (2011). Community structure in pelagic marine mammals at large spatial scales. *Marine Ecology Progress Series*, 434, 165-181.
- Schrenk, F., Bromage, T.G., Gorthner, A., & Sandrock, O. (1995). Paleoecology of the Malawi Rift: vertebrate and invertebrate faunal contexts of the Chiwondo Beds, northern Malawi. *Journal of Human Evolution*, 28, 59–70.

- Scott, K.M. (1990). Postcranial dimensions of ungulates as predictors of body mass. In Damuth, J. & McFadden, B.J. (Eds.) *Body Size in Mammalian Paleobiology: Estimation and Biological Implications* (pp. 301-355). New York: Cambridge University Press.
- Shipman, P. & Harris, J. (1988). Habitat preference and paleoecology of *Australopithecus boisei* in Eastern Africa. In F.E. Grine (Ed.), *Evolutionary History of the "Robust" Australopithecines* (pp. 343–381). New York: Aldine de Gruyter.
- Simpson, G.G. (1964). Species density of North American Recent Mammals. *Systematic Zoology*, 13, 57-73.
- Sokal, R.R., & Rohlf, F.J. (1995). *Biometry: the principles and practice of statistics in biological research*. New York: W.H. Freeman and Company.
- Sponheimer, M. & Lee-Thorp, J.A. (2003). Using carbon isotope data of fossil bovid communities for paleoenvironmental reconstruction. *South African Journal of Science*, 99, 273–275.
- Spurr, S.H. & Burton, V.B. *Forest ecology*. John Wiley & Sons Inc., 1980.
- Stanistreet, I.G. (2012). Fine resolution of early hominin time, Beds I and II, Olduvai Gorge, Tanzania. *Journal of Human Evolution*, 63, 300-308.
- Stoetzel, E., Marion, L., Nespoulet, R., El Hajraoui, M.A., & Denys, C. (2011). Taphonomy and paleoecology of the late Pleistocene to middle Holocene small mammal succession of El Harhoura 2 cave (Rabat-Témara, Morocco). *Journal of Human Evolution*, 60, 1-33.
- Strömberg, C. A. (2011). Evolution of grasses and grassland ecosystems. *Annual Review of Earth and Planetary Sciences*, 39, 517-544.
- Su, D.F. (2005). *The paleoecology of Laetoli, Tanzania: evidence from the mammalian fauna of the Upper Laetolil Beds*. Ph.D. thesis: New York University.
- Su, D.F. (2011). Large Mammal Evidence for the Paleoenvironment of the Upper Laetolil and Upper Ndolanya Beds of Laetoli, Tanzania. In T. Harrison (Ed.), *Paleontology and Geology of Laetoli: Human Evolution in Context. Volume 1: Geology, Geochronology, Paleoecology and Paleoenvironment* (pp. 381-392). Dordrecht: Springer.
- Su, D.F. (2016). The taphonomy and paleoecology of Korsi Dora Vertebrate Locality 1, Woranso-Mille Study Area, Ethiopia. In Y. Haile-Selassie & D.F Su (Eds.), *The Postcranial Anatomy of Australopithecus afarensis: New Insights from KSD-VP-1/1* (pp. 25-37). New York: Springer-Verlag.
- Su, D.F., & Haile-Selassie, Y. (in review). Mosaic habitats at Woranso-Mille (Ethiopia) during the Pliocene and implications for *Australopithecus* paleoecology. In S.

- Reynolds & R. Bobe (Eds.) *African Paleoecology and Human Evolution*. Cambridge: Cambridge University Press.
- Su, D.F., & Harrison, T., 2007. The paleoecology of the Upper Laetolil Beds at Laetoli. In: Bobe, R., Alemseged, Z., Behrensmeyer, A.K. (Eds.), *Hominin environments in the East African Pliocene: An assessment of the faunal evidence* (pp. 279-313). Dordrecht: Springer.
- Su D.F., Ambrose S.H., Degusta D., & Haile-Selassie Y. (2009). Paleoenvironment. In Y. Haile-Selassie & G. WoldeGabriel (Eds.), *Ardipithecus kadabba: late Miocene evidence from the Middle Awash, Ethiopia* (pp. 521-547). Berkeley: University of California Press.
- Thibault, K.M. & Brown, J.H. (2008). Impact of an extreme climatic event on community assembly. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 3410–3415.
- Thibault, K.M., White, E.P. & Morgan Ernest, S.K. (2004). Temporal dynamics in the structure and composition of a desert rodent community. *Ecology*, 85, 2649-2655.
- Van Couvering, J. A. H. (1980). Community evolution in East Africa during the late Cenozoic. In A. K. Behrensmeyer, & A. P. Hill (Eds.) *Fossils in the making* (pp. 272-298). Chicago: University of Chicago Press.
- van Dam, J.A., Andrews, P., Badgley, C., Damuth, J., Fortelius, M., Hadly, E.A., Hixson, S., Janis, C., Madden, R.H., Reed, K., Smith, F.A., Theodor, J., Van Valkenburgh, B., & Werdelin, L. (2001). Within-habitat mammal diversity and productivity and their Recent patterns across latitude. *Journal of Vertebrate Paleontology*, 21 (suppl), 43A.
- Voorhies, M.R. (1969a). Sampling difficulties in reconstructing late Tertiary mammalian communities. *Proceedings of the North American Paleontological Convention, Part E*, 454-468.
- Voorhies, M.R. (1969b). *Taphonomy and population dynamics of an early Pliocene vertebrate fauna, Knox County, Nebraska*. Laramie: University of Wyoming.
- Warton, D.I., & Hui, F.K. (2011). The arcsine is asinine: the analysis of proportions in ecology. *Ecology*, 92, 3-10.
- Warwick, R.M., & Clarke, K.R. (1995). New ‘biodiversity’ measures reveal a decrease in taxonomic distinctness with increasing stress. *Marine Ecology Progress Series*, 129, 301–305.
- Western, D., & Behrensmeyer, A.K. (2009). Bone assemblages track animal community structure over 40 years in an African savannah ecosystem. *Science*, 324, 1061-1064.
- White, F., & Unesco. (1983). *The Vegetation of Africa: A Descriptive Memoir to Accompany the Unesco/AETFAT/UNSO Vegetation Map Of Africa*. Unesco.

- White, T.D., Ambrose, S.H., Suwa, G., Su, D.F., DeGusta, D., Bernor, R.L., Boisserie, J., Brunet, M., Delson, E., Frost, S., Garcia, N., Giaourtsakis, I.X., Haile-Selassie, Y., Howell, F.C., Lehmann, T., Likius, A., Pehlevan, C., Saegusa, H., Semperebon, G., & Vrba, E. (2009). Macrovertebrate paleontology and the Pliocene habitat of *Ardipithecus ramidus*. *Science*, 326, 87-93.
- Williams, S.E., Marsh, H., & Winter, J. (2002). Spatial scale, species diversity, and habitat structure: small mammals in Australian tropical rain forest. *Ecology*, 83, 1317-1329.
- Wilson, V.J. (1975). Mammals of the Wankie National Park. *Museum Memoirs of the National Museums and Monuments of Rhodesia* 4, 1-147.
- Vrba, E.S. (1980). The significance of bovid remains as indicators of environment and predation patterns. In A.K. Behrnesmeyer & A.P. Hill (Eds.), *Fossils in the Making: Vertebrate Taphonomy and Paleoecology* (pp. 247-271). Chicago: The University of Chicago Press.
- Yodzis, P. (1984). How rare is omnivory? *Ecology*, 65, 321-323.
- Zuur, A.F., Ieno, E.N., & Smith, G.M. (2007). *Analysing Ecological Data*. New York: Springer.

## FIGURE CAPTIONS

**Figure 1.** Locomotor niche exploitation profiles from four habitat types in Africa, based on data from 23 modern mammalian communities. Redrawn from Andrews et al., 1979.

LGM=large ground mammals, SGM=small ground mammals, A=arboreal, S=scansorial, Aq=aquatic, Ae=aerial.

**Figure 2.** Locomotor niche exploitation profiles for 15 habitat types across the world, based on data from 44 modern mammalian communities. Redrawn from Andrews & Humphrey, 1999: pg 287. T=terrestrial, S-A=scansorial-arboreal, Ar=arboreal, S=scansorial, Aq=aquatic, Ae=aerial, F=fossorial.

**Figure 3.** Percentage of frugivorous and grazing species in modern ecosystems. More open habitats have higher proportions of grazing species and lower proportions of arboreal species, (ecosystems 10-15). More wooded habitats have the opposite pattern with higher proportions of frugivorous species and lower proportions of grazing species (ecosystems 1-9).

Ecosystems: 1=African tropical forest, 2=African montane forest, 3=African seasonal tropical forest, 4=Malayan tropical forest, 5=Indian tropical forest, 6=Central American tropical forest, 7=Chinese paratropical forest, 8=Burmese subtropical forest, 9= Indian subtropical forest, 10=Miombo Woodland, 11=South African summer-rainfall woodland, 12=African mixed wood and grassland, 13=African tropical woodland, 14=African tropical grassland, 16=African tropical arid bushland.

Figure 1.






Figure 2.




Figure 3.

Table 1. Twenty-two modern African localities with habitat classification. Modified from Kovarovic et al (2002), which is based on a habitat classification system first outlined by Andrews et al (1979).

<b>Locality</b>	<b>Country</b>	<b>Habitat type</b>	<b>Description</b>
Irangi	Zaire	tropical forest	lowland evergreen forest; primary and secondary-riverine cultivation types
Seredou	Guinea	tropical forest	lowland evergreen forest
Semliki a	Uganda	tropical forest	lowland semi-deciduous forest
Budongo	Uganda	tropical forest	lowland semi-deciduous forest; secondary and monotypic types
Semliki b	Uganda	montane forest	montane evergreen forest continuous with lowland
Mt. Kenya	Kenya	montane forest	montane evergreen forest; primary and secondary
Lemera	Zaire	montane forest	montane evergreen forest in river valley
Amani	Tanzania	seasonal tropical forest	intermediate semi-deciduous forest
Kakamega	Kenya	seasonal tropical forest	intermediate semi-deciduous forest, secondary with glades, hills with bush
Rwenzori Forest	Uganda	seasonal tropical forest	lowland semi-deciduous forest
Sokoike-Gedi	Kenya	seasonal tropical forest	lowland deciduous forest and woodland
Tana River	Kenya	mixed wood and grassland	flood plain grassland with patches of deciduous forest
Gabiro	Rwanda	mixed wood and grassland	wooded grassland next to lake flats
Legaja	Tanzania	mixed wood and grassland	lake in short grassland with woodland on the fringes
Rukwa	Tanzania	mixed wood and grassland	flood plain grassland with bordering woodland and swamps
Rwenzori	Uganda	tropical woodland	short grass plains-bush-woodland mosaic
Banagi	Tanzania	tropical woodland	deciduous woodland and bushland
Kafue	Zambia	tropical grassland	flood plain grassland with bordering woodland
Serengeti	Tanzania	tropical grassland	short grass plains with small patches of bush
Karamoja	Uganda	tropical arid bushland	arid bushland
Kapiti Plains	Kenya	tropical arid bushland	grassland-bushland mosaic
Tsavo	Kenya	tropical arid bushland	bushland and arid bush