



This is the author's version of a work that was accepted for publication:

Mitchell, J. A., Pumpa, K. L., Williams, K. J., & Pyne, D. B. (2016). Variable Changes in Body Composition, Strength and Lower-Body Power During an International Rugby Sevens Season. *Journal of Strength and Conditioning Research*, 30(4), 1127-1136. <https://doi.org/10.1519/JSC.0000000000001188>

This file was downloaded from:

<https://researchprofiles.canberra.edu.au/en/publications/variable-changes-in-body-composition-strength-and-lower-body-powe>

©2016 National Strength & Conditioning Association

Notice: This is the non-final version of an article that was accepted for publication in the *Journal of Strength and Conditioning Research*, which has been published at <http://doi.org/10.1519/JSC.0000000000001188>

Changes resulting from the publishing process may not be reflected in this document

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

ABSTRACT

This study determined whether body composition, strength and power changes that occur during pre-season can be maintained during an international rugby sevens season. Fourteen male international rugby sevens players (age 21.4 ± 2.2 y; mean \pm SD) were categorized as a forward ($n=7$) or back ($n=7$), and assessed for height, mass (M), skinfolds (S) ($\Sigma 7$), upper body (UB) strength, lower body (LB) strength and LB power. Bench press, back squat and a countermovement jump were used to measure strength and power at three time points: initial, early season and late season. Forwards were taller (185 ± 4 cm), heavier (95 ± 6 kg) and possessed a greater lean body mass (55.5 ± 4.0 M·S^{-0.14}) than backs (181 ± 8 cm, 88.5 ± 5.5 kg, 51.9 ± 3.4 M S^{-0.14}). Over the full season, small ($\sim 5\% \pm 5\%$; mean \pm 90% confidence limits) positive changes occurred in body composition. LB strength gained during the pre-season decayed in-season, whereas UB strength increased moderately ($\sim 10\% \pm 3\%$) across the season. Power showed inconsistencies between measured variables with a moderate positive change across the season in mean velocity and relative peak power. Forwards showed a small decrease in peak power (relative and absolute). Moderate changes were observed in mean power over the season, forwards decreasing ($\sim 6\% \pm 6\%$) and backs increasing ($\sim 8\% \pm 6\%$). Rugby sevens forwards in this study found it difficult to maintain and improve power qualities in-season. Training loads of forwards and backs should be differentiated to maximize strength and power in-season.

KEY WORDS

Pre-season, in-season, back squat, bench press, countermovement jump

INTRODUCTION

Rugby sevens is a collision-based, high intensity modified version of 15-a-side rugby union. The Sevens World Series (SWS) is regarded as the highest level of competition in which 15 core nations compete in nine tournaments played in nine different countries over a period of eight months. International rugby sevens players require highly-developed speed, mobility, power and endurance to tolerate the physical demands of competition (16, 23). Players demonstrate acceleration, speed, muscular power, and endurance qualities that are similar to, if not better than, professional rugby union players (17). Rugby sevens players are categorized into 2 distinct playing positions i.e. forward or back and typically commence their playing career in rugby union as a loose forward or back (25). Rugby sevens players cover on average 113 m per min, 45% greater distance per min than rugby union with a higher proportion of that distance at high velocity ($>6 \text{ m s}^{-1}$) highlighting the high running demands required (16). Game analyses indicate rugby sevens forwards spend 19% more time jogging and backs spend 8% more time running (24, 25), although other studies reported no clear delineation between playing positions in rugby sevens (17, 26). However, rugby sevens forwards perform exclusive skill sets such as scrummaging, line outs and restarts in training and competition. These exclusive skills require careful consideration when planning the strength and conditioning program to maximize the expression of lower body strength and power

To meet the demands of rugby sevens, players possess similar body mass and height to professional backs in rugby union, while being substantially leaner (17). Forwards in rugby sevens are typically taller and heavier than backs, although the magnitude of difference is less than in rugby union (28), while international players are taller and heavier than provincial level players (28). The anthropometric characteristics of international rugby sevens players correlate with estimates of work-rate such as work-to-rest ratio's, distance, intensity and time (23). As such, an investigation

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

into the effect body mass has on relative strength and power characteristics would be valuable to assist in determining appropriate training variables and in the selection of rugby sevens players.

Field-based contact team sports, such as rugby union and rugby league, are characterized by frequent high-intensity sprints and a high degree of physical contact (29). In rugby league, high levels of strength and power are required to effectively perform tackling, lifting, pushing and pulling tasks, and to tolerate the heavy tackles and collisions that occur during a match (15). High levels of muscular strength and power are thought to be critical to success in collision-based sports (25). Given the high collision nature of rugby sevens it follows that assessment of strength and power qualities would be advantageous in determining player readiness to participate in high collision activities.

Several authors have highlighted the importance of strength and power in rugby sevens (17, 25, 30). Other collision-based sports such as rugby union and rugby league have reported comparable ranges of lower and upper body strength in players (1, 4, 9). Forwards in rugby union are generally stronger than backs in the lower and upper body, likely due to requirements of strength in scrums and a higher frequency of tackles and rucks (21, 29, 30). More recently, rugby union players were physically assessed in the pre-season and then differentiated at the end of the season by the coach into high and low playing ability groups. Backs identified as having high playing ability by the coach demonstrated higher absolute lower body strength as measured by a 1RM back squat suggesting a relationship between lower body strength and playing performance (1). In rugby sevens, international players appear to be stronger in the upper body than provincial level players and forwards are stronger than backs (28). Although cross-sectional comparisons are useful, practitioners need more information on the patterns of within-and-between athlete changes, and differences in lower body and upper body strength that may occur across a full international rugby sevens season.

Power in rugby sevens, expressed through the skills of sprinting, jumping and tackling, is typically assessed in a countermovement jump using a linear position transducer to measure speed of movement and distance (25, 28). Peak power and relative peak power are the primary variables of interest when analyzing neuromuscular fatigue and performance in rugby sevens (28, 33). In testing completed one month prior to rugby sevens tournaments (one international and one provincial), international level players and forwards were more powerful (than backs) in both absolute and relative measures (28). Players reported as possessing greater peak power in countermovement jumps were more effective in both attacking and defensive rucks during game play (27). Further research is required in characterizing power output variables and the time course, direction and magnitude of changes that may occur over a full season of rugby sevens. This information is useful in determining any player positional differences that may occur as a result of specific training requirements and game demands associated with scrummaging, line outs and restarts.

It is apparent that concurrent strength, rugby sevens skills practice and energy system training (or endurance) across the rugby sevens season needs to be considered carefully to optimize the physical requirements of players. To date there has been limited research into the strength and power requirements of rugby sevens players. With the inclusion of rugby sevens as an Olympic sport, the strength and power characteristics of rugby sevens playing positions and how they change across an international SWS season becomes increasingly important when formulating a periodized annual plan.

METHODS

Experimental approach to the problem.

To establish whether changes occurred across the season, initial upper body and lower body strength and lower body power measurements were collected in bench press, back squat and an instrumented countermovement jump at the start of the pre-season (initial), start of the competition season (early season) and end of the competition season (late season). Changes within-subjects and differences between forwards and backs were assessed for pre-season (initial to early season) in-season (early to late season) and full season (initial to late season). Effects were quantified to establish whether there had been substantial changes between phases and/or the full season as a result of the training and game demands of playing international Rugby sevens.

Subjects

Fourteen male international rugby sevens squad players (seven backs and seven forwards) participated in this study (age 21.4 ± 2.2 y; mean \pm SD). These players had ~ 1.7 years of specific international rugby sevens training and playing experience (range 1-4 years). Each player was informed of the benefits and risks of this investigation prior to signing a University of Canberra informed consent document to participate in this study. This study was approved by the University of Canberra Committee for Ethics in Human Research (Project 12-68).

Procedures

The SWS season consisted of 40 weeks that were divided into the pre-season and in-season training periods. The pre-season training period consisted 12 weeks, one regional international tournament, and ~ 3 lifting sessions per week (n=13 upper body, n=12 lower body and n=11 combined upper/lower sessions) with an average weekly running volume of ~ 12 km. The in-season training

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

period consisted of 28 weeks, 9 SWS tournaments, ~2.5 lifting sessions per week (n=40 upper body, n=33 lower body sessions) with an average weekly running volume of ~15km. Players completed lifting programs that were tailored to the individual's training experience, musculo-skeletal concerns and rugby sevens requirements. Testing was conducted as part of regular strength and conditioning lifting sessions (see Table 1) in line with the rugby sevens periodized plan. Tables 2 and 3 provide examples of pre-season, in-season and tournament training weekly schedules. Tests were selected using recommendations for the assessment of rugby union players as previously documented (32). Players were instructed to consume their regular pre-training diet, and where possible, refrain from intense exercise at least 48 hours prior to any scheduled testing. Only data collected on players free of injury and illness at the time of testing are reported.

Insert Table 1-3 here

Body composition was assessed by measurement of body mass, height and skinfold thickness using standard laboratory techniques (32). The sum of seven sites (mm): biceps, triceps, subscapular, supriliac, abdomen, front thigh and medial calf skinfolds, is reported. The typical error of measurement for sum of seven skinfolds is 1.3 mm and <1% for body mass and height. A lean mass index (LMI) was calculated for each subject as $M \cdot S^{-0.14}$, where M is body mass (kg) and S is sum of skinfolds (mm) (32).

Upper body and lower strength was assessed using an estimated one repetition maximum (1RM) for the bench press and back squat exercises respectively (1, 32). The bench press required the player to lay in a supine position, and using a self-selected grip width to lower a free weight barbell to their chest and then return the barbell to full arm extension. Players were not permitted to bounce the bar off the chest, and the feet and hips had to remain in contact with the ground and bench respectively. The grip width was self-selected by the subject in accordance with previous protocols.

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

The back squat required the subject to support a free weight barbell on their upper back, and through movements of the lower body, descend to a depth whereby the anatomical hip crease had moved below the apex of the patella. Players were instructed to have a comfortable foot placement approximately shoulder width apart with feet abducted. No supportive equipment of any kind was permitted during the back squat. Testing loads were based on previous training loads as determined by the strength and conditioning coach (1). Subjects then completed the following preliminary loads 5 x 65%, 5 x 75% and 3 x 85% of their predicted 1RM. Players were required to lift 95% of their prescribed 1RM load to failure. An estimated 1RM was determined from the final number of completed repetitions at the 95% load using an established 1RM prediction equation (10). Data for the bench press and back squat is presented in absolute and relative terms, and scaled allometrically to the power of 0.62 as recommended in previous studies (11, 25).

Lower body muscular power assessment was assessed using a linear position transducer given acceptable validity, reliability and portability to estimate power similar to a force plate when measuring jump performance (12). Prior to the lower body muscular power assessment all players completed a 10 minute standardized warm-up using a space approximately 20m long x 3m wide consisting of a series of dynamic exercises (e.g. 60-70% jogging, lateral/carioca, low to high skipping, jumping/hopping, dynamic hamstring/quadriceps/low back movements) (33). The linear position transducer was anchored to the ground on the right side of the player and attached to the right hand side of a broomstick (45g). In the start position the subject was required to support the broomstick on their upper back [with their hands], and through movements of the lower body perform a countermovement jump. The foot position and depth of the countermovement jump was self-selected by the player with the instruction to jump as high as possible (5). Players completed six countermovement jumps with allowance to reset the feet to the start position between repetitions, with the mean value of all six jumps being reported. A displacement-time curve for each jump was obtained by attaching a digital optical encoder via a cable (GYMAWARE Power Tool. Kinetic

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

Performance Technologies, Canberra, Australia) to one side of the broomstick. This data was uploaded to the GYMAWARE website then exported to excel for analysis. This system recorded displacement-time data for reported measures (see table 6) with a mean coefficient of variation of 13% and intraclass correlation of 0.815.

Statistical Analyses

Descriptive data are reported as mean and standard deviation (SD). Inferences on changes in a body composition or strength/power test score from phase to phase of the rugby sevens season were made using a student's t-test for dependent samples. Precision of estimation was indicated with 90% confidence limits. Effect sizes were reported in standardized (Cohen) units as the change in the mean with 90% confidence limits (18). Criteria to assess the magnitude of observed changes were: 0.0 to 0.2 trivial; 0.20 to 0.60 small; 0.60 to 1.20 moderate; and > 1.20 Large. An effect was deemed unclear if its confidence interval spanned both substantial positive and substantial negative values ($\pm 0.20 \times$ between-subject SD) (18).

RESULTS

Rugby sevens forwards were substantially taller, heavier and possessed greater lean muscle mass than backs (see Table 4). The combined cohort of backs and forwards showed little change in body mass, skinfolds and lean muscle mass across the overall course of the season. The backs showed small ($\sim 5\% \pm 5\%$; mean \pm 90% CL) decreases in skinfolds over the full season, with a small increase in lean mass in-season.

Both forwards and backs showed small ($\sim 7\% \pm 4\%$) increases in lower body strength in the pre-season (see Table 5). Backs had a moderate ($\sim 10\% \pm 3\%$) increase in lower body strength (back squat) in the pre-season when expressed relative to body weight and allometrically scaled, whereas

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

changes in the forwards were unclear and trivial respectively. The increases in lower body strength gained during the pre-season were lost in-season with the backs having the same levels by the end of the season and the forwards decreasing ($\sim 4\% \pm 3\%$) below initial levels. Conversely, upper body strength (bench press) showed small (forwards) to moderate (backs) improvements over the full season. Backs exhibited the biggest increases in the pre-season ($\sim 9\% \pm 4\%$) plateauing in-season to show a moderate increase by the end of the season, while the forwards continued to have small increases in upper body strength throughout the season ($\sim 5\% \pm 3\%$).

Lower body power scores in the countermovement jump are presented in Table 6. The combined cohort showed moderate positive changes ($\sim 6\% \pm 4\%$) across the season in mean velocity. Backs had a small increase in peak velocity with moderate increases in mean velocity, mean power and mean power w.kg^{-1} ($\sim 8\% \pm 6\%$). Forwards showed small increases ($\sim 3\% \pm 3\%$) in peak velocity in the pre-season with moderate ($\sim 18\% \pm 7\%$) decreases in peak power both absolute and relative with no change in jump height. All power variables for forwards showed a marked decrease over the full season. In both backs and forwards moderate changes (positive and negative) were observed in mean power both relative and absolute for the full season.

Insert Tables 4-6 here

DISCUSSION

This is the first study to quantify the changes in body composition, strength and power as they relate to playing position over a full season of international rugby sevens. During the pre-season, early season and late season phases upper body strength showed small improvements for the full season, whereas lower body strength increased in the pre-season before falling away in-season. Lower body power increased in the backs and decreased in the forwards over the duration of the season. The upper body and lower body strength scores confirm preliminary reports that rugby sevens players

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

display similar physical qualities to outside backs in rugby union. Given the different time course of changes in strength and power between forwards and backs it is prudent for the strength and conditioning coach to prescribe position-specific training programs account for potential auxiliary lower limb loading on field during specific skill practice.

The rugby sevens players as a combined group were younger and heavier with similar stature to other reports (17, 28, 30). When differentiating between playing position, the backs were ~2% heavier and within the height ranges previously reported, whereas the forwards were ~2% heavier at the same height in one study (24) and ~2% lighter and ~1% shorter than players in two other studies (14, 28). In this study the weight and stature of rugby sevens forwards and backs compared closely to outside backs in rugby union (14). When players were differentiated by playing ability and position, both forwards and backs showed similar body composition to backs of high playing ability in rugby union and to rugby sevens players of international standard (1, 28). The combined player sum of seven skinfolds was consistent with previous rugby sevens research whereas lean mass was 3.3% higher across the full season (17). Rugby sevens skinfolds and lean mass presented over one season is substantially leaner than professional rugby union players reported over a period of two full seasons (13). Forwards had higher total skinfolds than backs, which is consistent with the two other studies that investigated skinfolds and differentiated between playing positions in rugby sevens (24, 28). The higher sum of seven skinfolds in forwards is likely related to the extra collisions in scrums, rucking, mauling and tackling (21, 29, 30). Across the full season the backs showed small decreases in skinfolds with a small increase in lean mass in-season as shown previously (20). Significantly, there was no negative body composition change for the full season suggesting that the training and nutritional program maintained player body composition for a full international rugby sevens season.

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

To date no other study has reported on the assessment of lower body strength in rugby sevens using a back squat. Lower body strength showed a small increase in absolute and relative values during the pre-season although notably these improvements were not maintained in-season. The early season squat testing showed lower absolute strength when compared to other rugby codes. In contrast, these results were higher or comparable with relative lower body strength scores reported in rugby union (1, 5, 9). When compared to backs in professional rugby union, rugby sevens forwards and backs have similar relative lower body strength values whereas rugby sevens backs are 15% lower in absolute terms (1). Previous research reported an 8.5% increase in lower body strength across a rugby union season (4). The inverse relationship in lower body strength between the pre-season gains and in-season losses may be attributed to a number of factors including changes in lower body stimulus scheduling in-season, international travel requirements, increased tournament scheduling, and the ability to recover from high running loads (~20% higher - unpublished research) in-season and impact collisions that are integral to rugby sevens (16, 26, 30). Although it appears that maximal strength and power can be maintained at pre-season levels for long in-season periods (6, 19). There are two other possible explanations for this decay in lower body strength. First, forwards need superior lower body strength to perform the skills of scrummaging, line outs and restarts of play in both training and games. Measurement of force during an instrumented individual player simulated scrum session showed values ~52% less than a reported box squat measures and ~50% less than back squat (22, 31). This additional lower body load experienced exclusively by forwards would contribute to overall lower body fatigue and should be considered when structuring the training week to optimize lower body strength gains. Secondly, a box squat replaced the back squat exercise in-season (see Table 1) as a quality control measure to ensure players were maintaining a minimum squat depth, this action may also account for the decrease in reported back squat strength. Biomechanical differences such as a narrower stance, lower force values and greater anterior displacement of the torso in a back squat when compared to a box squat could result in the

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

reduced ability to perform maximal back squat (due to the absence of back squat specificity) during the training phase (31).

Upper body strength gains in the combined cohort were small through the pre-season and in-season with an overall moderate increase across the full season similar to previous reports in rugby league (6). This outcome is contrary to a decay of 1.2% in rugby union players across the season which was attributed to professional athletes having a greater strength training background and thus reducing the likelihood of further strength improvements in-season (4). In the current study the players were ~3 years younger with a ~9% lower bench press, which may be due to an inherently shorter strength training background. The backs had a moderate increase in bench press strength for the pre-season similar to a previous study in rugby union (3). Compared to professional rugby union and rugby league players the combined cohort had up to 8% lower absolute bench press values; however, in relative terms they were much closer to reported measures of 1.2-1.5 w.kg⁻¹ (2, 8, 9). The group was relatively homogenous when upper body strength was differentiated for forwards and backs. A 5% difference between rugby sevens playing positions has been reported (combined international and provincial level players), whereas international rugby sevens players had a 20% greater bench press than provincial level players (combined backs and forwards) (28). While upper body strength continually improved over the duration of the rugby sevens playing season, the combined cohort bench press scores were 2.4% less than previously reported in international level players (28). It is important to note the increase in upper body strength across the season suggesting running and skill training loads had minimal impact on upper body strength development.

It is apparent that no single lower body power measure showed consistent changes in the combined cohort or when differentiating between forwards and backs. The reported variance in power variables may have been magnified by using a linear position transducer (in place of a force plate). Peak power and relative peak power can be useful in assessing rugby sevens players (28, 33).

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

Combined players showed peak power measures 16% higher than provincial and 13% lower than international level rugby sevens players, and peak power measures were 29% higher than a recent report on elite rugby sevens players (28, 33). The backs in the current study showed small to moderate increases over the full season. In a comparison of squatting techniques it was suggested that a box squat (due to the reported reduced forces and higher rate of force development) could be an effective exercise to develop explosive strength (31). The replacement of back squat with a box squat in-season (see table 1) may account for the moderate increases shown by the backs in the absence of scrummaging, line outs and restarts of play. The decrease in power identified in the forwards may be in part relate to the specific training loads associated with scrummaging, line outs and restarts of play possibly compounding neuromuscular fatigue, resulting in a decreased adaptation to power training. Additionally, in-season training schedule changes reduced the ratio of lower to upper body strength training stimulus by ~20% (see Table 2). Given the strong relationship between strength and power, the reduction in the strength training stimulus may have facilitated a similar reduction in power output over time (7).

A longitudinal study comprising of multiple seasons would be beneficial to address chronic adaptations to strength and power training. Future investigations could determine the effect of running loads and position-specific skills training on strength and power of positional groups during both the pre-season and in-season. Determining the optimal training week structure whereby strength and power gains in-season are maximized would be advantageous. It appears that neuromuscular function may be significantly reduced for greater than 120 hours post tournaments (34). Given that there were 9 tournaments and 28 in-season training weeks this research effectively excludes 32% of productive training time in-season.

Conclusion

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

This study systematically quantified the changes in strength and power across an international rugby sevens season. It appears that rugby sevens players have similar physique and physical qualities to professional level backs in rugby union. Upper body strength continually increased throughout an international rugby sevens season, whereas lower body strength gains made in preseason training but proved difficult to maintain in-season. The ability to maintain and improve power for forwards during the season may be affected by the additional set piece requirements of scrummaging, line outs and restarts. Training loads need to be carefully prescribed for forwards and backs to account for the potential decay of strength and power qualities.

Practical Applications

Consideration should be given to the lower body demands of rugby sevens forwards to ensure that lower body improvements made in the pre-season can be maintained throughout the in-season. These outcomes could be achieved by ensuring the ratio of lower body to upper body programming remains consistent through the season managing training time parity and recovery between positions. Use of the box squat in-season may be useful for maintaining and increasing lower body explosive power, however care should be exercised when assessing the back squat if the box squat has been the primary lower body strength modality.

Acknowledgements

The authors would like to acknowledge the support of the players and staff involved in this study.

References

1. Appleby B, Hori N, and Dorman C. The characteristics of professional rugby union players in strength, power and anthropometrics. *J Aust Strength Con* 17: 9-17, 2009.
2. Appleby B, Newton RU, and Cormie P. Changes in strength over a 2-year period in professional rugby union players. *J Strength Cond Res* 26: 2538-2546, 2012.

3. Argus CK, Gill N, Keogh J, Hopkins WG, and Beaven CM. Effects of a short-term pre-season training programme on the body composition and anaerobic performance of professional rugby union players. *J Sports Sci* 28: 679-686, 2010.
4. Argus CK, Gill ND, Keogh JW, Hopkins WG, and Beaven CM. Changes in strength, power, and steroid hormones during a professional rugby union competition. *J Strength Cond Res* 23: 1583-1592, 2009.
5. Argus CK, Gill ND, Keogh JWL, and Hopkins WG. Assessing Lower-Body Peak Power in Elite Rugby-Union Players. *J Strength Cond Res* 25: 1616, 2011.
6. Baker D. The effects of an in-season of concurrent training on the maintenance of maximal strength and power in professional and college-aged rugby league football players. *J Strength Cond Res* 15: 172-177, 2001.
7. Baker D, Nance S, and Moore M. The load that maximizes the average mechanical power output during jump squats in power-trained athletes. *J Strength Cond Res* 15: 92-97, 2001.
8. Baker DG and Newton RU. An analysis of the ratio and relationship between upper body pressing and pulling strength. *J Strength Cond Res* 18: 594-598, 2004.
9. Bevan HR, Bunce PJ, Owen NJ, Bennett MA, Cook CJ, Cunningham DJ, Newton RU, and Kilduff LP. Optimal loading for the development of peak power output in professional rugby players. *J Strength Cond Res* 24: 43, 2010.
10. Brzycki M. Strength testing—predicting a one-rep max from reps-to-fatigue. *J Phys Ed Rec Dance* 64: 88-90, 1993.
11. Crewther BT, McGuigan MR, and Gill ND. The ratio and allometric scaling of speed, power, and strength in elite male rugby union players. *J Strength Cond Res* 25: 1968-1975, 2011.
12. Cronin JB, Hing RD, and McNair PJ. Reliability and validity of a linear position transducer for measuring jump performance. *J Strength Cond Res* 18: 590-593, 2004.
13. Duthie G, Pyne D, Hopkins W, Livingstone S, and Hooper S. Anthropometry profiles of elite rugby players: quantifying changes in lean mass. *Br J Sports Med* 40: 202-207, 2006.
14. Fuller CW, Taylor A, and Molloy MG. Epidemiological study of injuries in international rugby sevens. *Clin J Sport Med* 20: 179, 2010.
15. Gabbett TJ. Influence of physiological characteristics on selection in a semi-professional first grade rugby league team: a case study. *J Sports Sci* 20: 399-405, 2002.
16. Higham DG, Pyne DB, Anson JM, and Eddy A. Movement patterns in rugby sevens: Effects of tournament level, fatigue and substitute players. *J Sci Med Sport* 15: 277-282, 2012.
17. Higham DG, Pyne DB, Anson JM, and Eddy A. Physiological, Anthropometric and Performance Characteristics of Rugby Sevens Players. *Int J Sports Physiol Perform* 8: 19-27, 2013.
18. Hopkins WG, Marshall SW, Batterham AM, and Hanin J. Progressive Statistics for Studies in Sports Medicine and Exercise Science. *Med Sci Sports Exerc* 41: 3-13, 2009.
19. Leveritt M, Abernethy PJ, Barry BK, and Logan PA. Concurrent strength and endurance training. *Sports Med* 28: 413-427, 1999.
20. Meir RA. Seasonal changes in estimates of body composition in professional rugby league players. *Sport Health* 11: 27-31, 1993.
21. Quarrie K, Handcock P, Toomey M, and Waller AE. The New Zealand rugby injury and performance project. IV. Anthropometric and physical performance comparisons between positional categories of senior A rugby players. *Br J Sports Med* 30: 53-56, 1996.
22. Quarrie K and Wilson B. Force production in the rugby union scrum. *J of sports sci* 18: 237-246, 2000.
23. Reilly T. Assessment of sports performance with particular reference to field games. *Eur J of Sport Sci* 1: 1-12, 2001.
24. Rienzi E, Reilly T, and Malkin C. Investigation of anthropometric and work-rate profiles of Rugby Sevens players. *J Sports Med Phys Fitness* 39: 160, 1999.

25. Ross A, Gill N, and Cronin J. Match Analysis and Player Characteristics in Rugby Sevens. *Sports Med* 44: 357-367, 2014.
26. Ross A, Gill N, and Cronin J. The match demands of international rugby sevens. *J Sports Sci* 33: 1035-1041, 2015.
27. Ross A, Gill N, Cronin J, and Malcata R. The relationship between physical characteristics and match performance in rugby sevens. *Eur J Sport Sci* Apr 13: 1-7, 2015.
28. Ross A, Gill ND, and Cronin JB. Comparison of the Anthropometric and Physical Characteristics of International and Provincial Rugby Sevens Players. *Int J Sports Physiol Perform* In print, 2014.
29. Smart DJ. Physical profiling of rugby union players: implications for talent development. Doctor of Philosophy. Auckland University of Technology, 2011.
30. Suarez-Arrones L, Arenas C, López G, Requena B, Terrill O, and Mendez-Villanueva A. Positional differences in match running performance and physical collisions in men rugby sevens. *Int J Sports Physiol Perform* 9: 316-323, 2014.
31. Swinton PA, Lloyd R, Keogh JW, Agouris I, and Stewart AD. A biomechanical comparison of the traditional squat, powerlifting squat, and box squat. *J Strength Cond Res* 26: 1805-1816, 2012.
32. Tanner R. *Physiological Tests for Elite Athletes, 2E*. Human Kinetics Publishers, 2012.
33. West DJ, Cook CJ, Beaven MC, and Kilduff LP. The influence of the time of day on core temperature and lower body power output in elite rugby union sevens players. *J Strength Cond Res* 28: 1524-1528, 2014.
34. West DJ, Cook CJ, Stokes KA, Atkinson P, Drawer S, Bracken RM, and Kilduff LP. Profiling the time-course changes in neuromuscular function and muscle damage over two consecutive tournament stages in elite rugby sevens players. *J Sci Med Sport* 17: 688-692, 2014.

Table 1. Key exercises performed during strength and conditioning lifting sessions.

| Upper body Lift | Lower body Lift | Combined Lift | In-tournament | In -tournament |
|-----------------|--------------------|----------------|-------------------|----------------|
| | | | Lift 1 - Strength | Lift 2 - Power |
| Jump Squats | Olympic Lift** | Olympic Lift** | Olympic Lift** | Jump Squats |
| Bench Press | Squat or Box Squat | Deadlift | Squats | Bench Press* |
| Chin Ups | Romanian Deadlift | Bench Pull | Bench Press | Push Press |
| Military Press | | | Chin Ups | |

*Explosive with bands for accommodative resistance. ** Power Clean or Snatch variations.

Table 2. Pre-season and in-season standard weekly training plans.

| Phase | Mon | Tues | Wed | Thurs | Fri | Sat | Sun |
|------------|--|--|--------------|--------------------------------|--|-----------------------|-----|
| Pre-Season | Combined Lift Conditioning Field skills training | Upper Lift Speed Skills | Conditioning | Speed Lower Lift | Field skills training | Recovery | Off |
| In Season | Lift* Speed Field skills training | Lift* Conditioning Field skills training | Recovery | Speed Field skills training | Lift* Conditioning Field skills training | Field skills training | Off |

*Lift is alternate day upper the lower body focus.

Table 3. Example of a SWS tournament week

| Game + 1* | Game + 2 | Game -3** | Game -2 | Game -1 | Game Day 1 | Game Day 2 |
|-----------------------|-----------------|-----------------------|--------------|-----------------------|------------|------------|
| Recovery | Mixed Lift 1 | Recovery | Mixed lift 2 | Field skills training | 3 games | 2/3 games |
| Field skills training | Game simulation | Field skills training | | | Recovery | Recovery |

* +1 representing one day post games/travel. ** -3 representing three days prior to game day 1

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

Table 4. Changes in anthropometric measures from the initial to early season (Pre-Season), early season to late season (In-Season), Initial to late season (Full-Season) phases. Mean change \pm 90% confidence limits (CL).

| | Time point | | | Standardized mean change (Cohen's) | | |
|-----------------------------------|-----------------------|----------------------------|---------------------------|-------------------------------------|------------------------------------|--------------------------------------|
| | Initial $x \pm SD$ | Early Season $x \pm SD$ | Late Season $x \pm SD$ | Pre-season $\Delta x \pm 90\%CL$ | In-season $\Delta x \pm 90\%CL$ | Full-season $\Delta x \pm 90\%CL$ |
| Body Composition | | | | | | |
| Players Combined n=14 | | | | | | |
| Height (cm) | 183.2 \pm 6.4 | 183.2 \pm 6.4 | 183.1 \pm 6.3 | | | |
| Mass (kg) | 92.1 \pm 7.1 | 91.5 \pm 6.3 | 92.2 \pm 6.5 | -0.07 \pm 0.11 | 0.09 \pm 0.09 | 0.02 \pm 0.15 |
| Skinfolds - Σ 7 sites (mm) | 54.5 \pm 11.6 | 53.59 \pm 10.9 | 51.65 \pm 12.5 | -0.07 \pm 0.25 | -0.16 \pm 0.18 | -0.23 \pm 0.34 * |
| LMI (M S ^{-0.14}) | 53.6 \pm 3.9 | 53.4 \pm 3.8 | 54.09 \pm 4.0 | -0.06 \pm 0.09 | 0.17 \pm 0.07 | 0.11 \pm 0.13 |
| Backs n=7 | | | | | | |
| Height (cm) | 181.1 \pm 7.8 | 181.1 \pm 7.8 | 181.1 \pm 7.8 | | | |
| Mass (kg) | 89.1 \pm 6.3 | 88.0 \pm 5.6 | 88.6 \pm 5.0 | -0.14 \pm 0.14 | 0.08 \pm 0.12 | -0.06 \pm 0.20 |
| Skinfolds - Σ 7 sites (mm) | 50.8 \pm 7.6 | 49.2 \pm 9.4 | 46.7 \pm 10.0 | -0.18 \pm 0.38 | -0.29 \pm 0.27 * | -0.47 \pm 0.55 * |
| LMI (M S ^{-0.14}) | 51.8 \pm 3.6 | 51.5 \pm 3.3 | 52.3 \pm 3.40 | -0.08 \pm 0.10 | 0.21 \pm 0.11 * | 0.13 \pm 0.17 |
| Forwards n=7 | | | | | | |
| Height (cm) | 185.2 \pm 4.2 | 185.3 \pm 4.1 | 185.1 \pm 4.2 | | | |
| Mass (kg) | 95.1 \pm 6.7 | 95.0 \pm 5.2 | 95.8 \pm 6.1 | -0.01 \pm 0.20 | 0.11 \pm 0.15 | 0.10 \pm 0.25 |
| Skinfolds - Σ 7 sites (mm) | 58.2 \pm 14.2 | 57.94 \pm 11.0 | 56.6 \pm 13.5 | -0.01 \pm 0.39 | -0.08 \pm 0.28 | -0.10 \pm .50 |
| LMI (M S ^{-0.14}) | 55.5 \pm 3.6 | 55.3 \pm 3.5 | 55.8 \pm 3.9 | -0.03 \pm 0.19 | 0.12 \pm 0.12 | 0.09 \pm 0.25 |

*0.2 -0.6 -Small ** 0.6-1.2 - Moderate

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

Table 5. Changes in lower and upper body strength measures from the initial to early season (Pre-Season), early season to late season (In-Season), Initial to late season (Full-Season) phases. Mean change \pm 90% confidence limits (CL).

| | Time point | | | Standardized mean change (Cohen's) | | |
|--------------------------------------|-----------------------|----------------------------|---------------------------|-------------------------------------|------------------------------------|--------------------------------------|
| | Initial $x \pm SD$ | Early Season $x \pm SD$ | Late Season $x \pm SD$ | Pre-season $\Delta x \pm 90\%CL$ | In-season $\Delta x \pm 90\%CL$ | Full-season $\Delta x \pm 90\%CL$ |
| Players Combined n=14 | | | | | | |
| Lower Body Strength | | | | | | |
| Squat Absolute (kg) | 143.4 \pm 23.3 | 152.10 \pm 22.3 | 142.4 \pm 17.9 | 0.35 \pm 0.24 * | -0.39 \pm 0.23 * | -0.04 \pm 0.17 |
| Squat Relative (w.kg ⁻¹) | 1.6 \pm 0.2 | 1.7 \pm 0.2 | 1.6 \pm 0.2 | 0.43 \pm 0.30 * | -0.48 \pm 0.27 * | -0.05 \pm 0.22 |
| Squat (w.kg ^{0.62}) | 8.6 \pm 1.0 | 9.2 \pm 1.1 | 8.6 \pm 1.0 | 0.59 \pm 0.34 ** | -0.59 \pm 0.34 ** | 0 \pm 0 |
| Upper Body Strength | | | | | | |
| Bench Absolute (kg) | 114.2 \pm 12.5 | 120 \pm 11.2 | 124.4 \pm 12.3 | 0.44 \pm 0.25 * | 0.33 \pm 0.17 * | 0.77 \pm 0.30 ** |
| Bench Relative (w.kg ⁻¹) | 1.2 \pm 0.1 | 1.3 \pm 0.2 | 1.4 \pm 0.2 | 0.45 \pm 0.26 * | 0.24 \pm 0.15 * | 0.70 \pm 0.27 ** |
| Bench (w.kg ^{0.62}) | 6.9 \pm 0.8 | 7.3 \pm 0.7 | 7.5 \pm 0.8 | 0.29 \pm 0.17 * | 0.29 \pm 0.17 * | 0.76 \pm 0.29 ** |
| Backs n=7 | | | | | | |
| Lower Body Strength | | | | | | |
| Squat Absolute (kg) | 133.7 \pm 22.0 | 145.6 \pm 20.7 | 138.3 \pm 18.7 | 0.47 \pm 0.34 * | -0.29 \pm 0.34 * | 0.18 \pm 0.19 |
| Squat Relative (w.kg ⁻¹) | 1.5 \pm 0.2 | 1.7 \pm 0.2 | 1.6 \pm 0.2 | 0.75 \pm 0.52 ** | -0.46 \pm 0.46 * | 0.29 \pm 0.22 * |
| Squat (w.kg ^{0.62}) | 8.3 \pm 1.1 | 9.1 \pm 1.0 | 8.6 \pm 0.9 | 0.64 \pm 0.45 ** | -0.39 \pm 0.41 * | 0.25 \pm 0.21 * |
| Upper Body Strength | | | | | | |
| Bench Absolute (kg) | 114.8 \pm 9.8 | 122.3 \pm 10.9 | 124.5 \pm 12.0 | 0.66 \pm 0.39 ** | 0.20 \pm 0.34 * | 0.86 \pm 0.47 ** |
| Bench Relative (w.kg ⁻¹) | 1.3 \pm 0.1 | 1.4 \pm 0.2 | 1.4 \pm 0.2 | 0.69 \pm 0.39 ** | 0.12 \pm 0.26 | 0.80 \pm 0.34 ** |
| Bench (w.kg ^{0.62}) | 7.1 \pm 0.6 | 7.6 \pm 0.7 | 7.7 \pm 0.8 | 0.72 \pm 0.41 ** | 0.16 \pm 0.30 | 0.88 \pm 0.41 ** |
| Forwards n=7 | | | | | | |

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

Lower Body Strength

| | | | | | | |
|--------------------------------------|--------------|--------------|--------------|---------------|----------------|----------------|
| Squat Absolute (kg) | 153.2 ± 21.6 | 158.6 ± 23.4 | 146.6 ± 17.5 | 0.22 ± 0.39 * | -0.48 ± 0.39 * | -0.26 ± 0.22 * |
| Squat Relative (w.kg ⁻¹) | 1.6 ± 0.3 | 1.7 ± 0.2 | 1.5 ± 0.2 | 0.16 ± 0.37 | -0.44 ± 0.37 * | -0.28 ± 0.26 |
| Squat (w.kg ^{0.62}) | 9.1 ± 1.4 | 9.4 ± 1.3 | 8.7 ± 1.1 | 0.19 ± 0.38 | -0.46 ± 0.39 * | -0.28 ± 0.24 |

Upper Body Strength

| | | | | | | |
|--------------------------------------|--------------|--------------|--------------|---------------|---------------|----------------|
| Bench Absolute (kg) | 113.7 ± 15.5 | 117.8 ± 11.7 | 124.4 ± 13.7 | 0.23 ± 0.33 * | 0.37 ± 0.15 * | 0.60 ± 0.41 ** |
| Bench Relative (w.kg ⁻¹) | 1.2 ± 0.2 | 1.2 ± 0.1 | 1.3 ± 0.1 | 0.23 ± 0.37 * | 0.32 ± 0.18 * | 0.55 ± 0.45 * |
| Bench (w.kg ^{0.62}) | 6.8 ± 0.88 | 7.0 ± 0.6 | 7.4 ± 0.7 | 0.24 ± 0.36 * | 0.35 ± 0.17 * | 0.58 ± 0.44 * |

*0.2 -0.6 -Small ** 0.6-1.2 - Moderate

Table 6. Changes in lower body power measures from the initial to early season (Pre-Season), early season to late season (In-Season), Initial to late season (Full-Season) phases. Mean change \pm 90% confidence limits (CL).

| | Time point | | | Standardized mean change (Cohen's) | | |
|----------------------------------|-----------------------|----------------------------|---------------------------|-------------------------------------|------------------------------------|--------------------------------------|
| | Initial $x \pm SD$ | Early Season $x \pm SD$ | Late Season $x \pm SD$ | Pre-season $\Delta x \pm 90\%CL$ | In-season $\Delta x \pm 90\%CL$ | Full-season $\Delta x \pm 90\%CL$ |
| Players Combined n=14 | | | | | | |
| Jump Height (m) | 0.49 \pm 0.07 | 0.50 \pm 0.06 | 0.50 \pm 0.07 | 0.21 \pm 0.41 | -0.01 \pm 0.23 | 0.20 \pm 0.38 * |
| Peak Velocity (m/sec)) | 3.8 \pm 0.3 | 3.9 \pm 0.3 | 3.8 \pm 0.4 | 0.30 \pm 0.29 * | -0.17 \pm 0.24 | 0.14 \pm 0.34 |
| Mean Velocity (m/sec) | 2.4 \pm 0.2 | 2.5 \pm 0.2 | 2.5 \pm 0.2 | 0.45 \pm 0.52 * | 0.23 \pm 0.44 | 0.69 \pm 0.56 ** |
| Peak Power (W) | 7764 \pm 1443 | 7643 \pm 1177 | 7194 \pm 1526 | -0.08 \pm 0.22 | -0.29 \pm 0.31 | -0.37 \pm .38 |
| Mean Power (W) | 4402 \pm 558 | 4502 \pm 442 | 4453 \pm 615 | 0.17 \pm 0.40 | -0.08 \pm 0.32 | 0.09 \pm 0.41 |
| Peak Power (w.kg ⁻¹) | 84.7 \pm 15.2 | 83.7 \pm 13.6 | 78.1 \pm 17.4 | -0.07 \pm 0.20 | -0.34 \pm 0.34 * | -0.41 \pm 0.39 * |
| Mean Power (w.kg ⁻¹) | 48.0 \pm 5.7 | 49.2 \pm 5.3 | 48.4 \pm 7.5 | 0.20 \pm 0.39* | -0.14 \pm 0.34 | 0.06 \pm 0.43 |
| Backs n=7 | | | | | | |
| Jump Height (m) | 0.52 \pm 0.88 | 0.54 \pm 0.06 | 0.55 \pm 0.07 | 0.30 \pm 0.62 | 0.03 \pm 0.36 | 0.33 \pm 0.57 |
| Peak Velocity (m/sec)) | 3.9 \pm 0.3 | 4.0 \pm 0.3 | 4.0 \pm 0.3 | 0.28 \pm 0.50 | 0.08 \pm 0.32 | 0.36 \pm 0.54 * |
| Mean Velocity (m/sec) | 2.4 \pm 0.2 | 2.5 \pm 0.2 | 2.6 \pm 0.1 | 0.49 \pm 0.75 | 0.46 \pm 0.61* | 0.96 \pm 0.56 ** |
| Peak Power (W) | 7752 \pm 1236 | 7755 \pm 1119 | 7991 \pm 1507 | 0.00 \pm 0.39 | 0.17 \pm 0.45 | 0.17 \pm 0.44 |
| Mean Power (W) | 4451 \pm 445 | 4640 \pm 492 | 4821 \pm 467 | 0.37 \pm 0.79 * | 0.35 \pm 0.59 * | 0.72 \pm 0.51 ** |
| Peak Power (w.kg ⁻¹) | 87.7 \pm 13.4 | 87.7 \pm 12.4 | 89.5 \pm 16.2 | 0.00 \pm 0.33 | 0.12 \pm 0.51 | 0.12 \pm 0.47 |
| Mean Power (w.kg ⁻¹) | 50.4 \pm 4.9 | 52.4 \pm 5.3 | 54.0 \pm 5.3 | 0.36 \pm 0.68 | 0.30 \pm 0.60 | 0.66 \pm 0.53 ** |
| Forwards n=7 | | | | | | |
| Jump Height (m) | 0.46 \pm 0.04 | 0.46 \pm 0.03 | 0.46 \pm 0.03 | 0.03 \pm 0.68 | -0.09 \pm 0.33 | -0.06 \pm 0.52 |
| Peak Velocity (m/sec)) | 3.6 \pm 0.3 | 3.7 \pm 0.3 | 3.6 \pm 0.3 | 0.34 \pm 0.42 * | -0.50 \pm 0.35 * | -0.16 \pm 0.43 |
| Mean Velocity (m/sec) | 2.4 \pm 0.1 | 2.4 \pm 0.1 | 2.4 \pm 0.1 | 0.29 \pm 0.69 | -0.18 \pm 0.45 | 0.11 \pm 0.90 |

Variable changes in body composition, strength and lower body power during an international rugby sevens season.

| | | | | | | |
|----------------------------------|-------------|-------------|-------------|--------------|-----------------|-----------------|
| Peak Power (W) | 7777 ± 1726 | 7531 ± 1312 | 6397 ± 1137 | -0.12 ± 0.26 | -0.57 ± 0.22 * | -0.70 ± 0.36 ** |
| Mean Power (W) | 4352 ± 686 | 4364 ± 369 | 4084 ± 535 | 0.02 ± 0.45 | -0.36 ± 0.22 * | -0.34 ± 0.43 * |
| Peak Power (w.kg ⁻¹) | 81.8 ± 17.4 | 79.6 ± 14.6 | 66.8 ± 9.7 | -0.11 ± 0.27 | -0.64 ± 0.27 ** | -0.75 ± 0.40 ** |
| Mean Power (w.kg ⁻¹) | 45.7 ± 5.9 | 46.0 ± 3.0 | 42.7 ± 4.2 | 0.05 ± 0.56 | -0.49 ± 0.25 * | -0.44 ± 0.52 * |

*0.2-0.6 -Small ** 0.6-1.2 - Moderate