

# Can kunzea oil (*Kunzea ambigua*) control head lice (*Pediculus humanus capitis*)?

C. R. WILLIAMS<sup>1</sup>, C. E. WEBB<sup>2</sup>, S. ORRE<sup>1</sup>, M. NAUNTON<sup>3</sup> and J. THOMAS<sup>3\*</sup>

<sup>1</sup> *Sansom Institute for Health Research, University of South Australia, Adelaide, SA 5001, Australia*

<sup>2</sup> *Department of Medical Entomology, University of Sydney and Pathology West – ICPMR Westmead, Westmead Hospital, NSW 2145, Australia*

<sup>3</sup> *Faculty of Health, University of Canberra, Bruce, Canberra ACT 2601, Australia*

(Received 8 June 2015; revised 28 July 2015; accepted 6 August 2015)

## SUMMARY

Head lice (*Pediculus humanus capitis*) infestations are a public health concern. The insecticidal properties of the Australian native plant *Kunzea ambigua* (commonly known as tick bush) have been documented. In this study, we tested activity of kunzea oil (KO) against head lice through *in vitro* bioassays. Head lice were exposed to filter paper treated with either KO, as either a 5 or 100% oil, or commercial formulations containing either permethrin or tea tree oil (TTO) for 120 min. Head lice exposure to KO, both as a 5 and 100% solution oil, resulted in 100% mortality within 120 min with a mean survival times of 17.1 and 34.8 min, respectively. There was no significant difference between the mean mortality of head lice exposed to 5% KO ( $17.1 \pm 1.0$ ; 95% CI: 115.2–19.0) and 5% TTO ( $21.2 \pm 1.9$ ; 95% CI: 17.4–25.1). This study revealed, for the first time, that KO holds great potential as an effective alternative to current active ingredients contained within commercial pediculicide formulations.

Key words: Kunzea oil, *Pediculus humanus capitis*, *Kunzea ambigua*, tea tree oil, head lice, pediculicide, resistance.

## INTRODUCTION

Pediculosis (infestation with head lice) has been reported as a chronic public health problem worldwide, particularly in children aged 3–13 (Speare and Buettner, 1999; Diamantis *et al.* 2009; Currie *et al.* 2010). Therapeutic management of pediculosis is challenging with increasing worldwide prevalence due to product failures resulting from resistance, incorrect application, formulation changes and misdiagnosis (Tolozza *et al.* 2008). With a limited number of effective pediculicides available, plant-derived compounds hold great potential due to demonstrated repellent, ovicidal, adulticidal and feeding inhibition activity against various arthropods including head lice (Audino *et al.* 2007).

Like tea tree oil (TTO), kunzea oil (KO) is a myrtaceous essential oil; although it differs in composition to TTO, comprising about 70% monoterpenes but with characteristic sesquiterpenes (Thomas *et al.* 2010). *Kunzea* spp. plant extracts have insecticidal properties (Khambay *et al.* 1999, 2002). KO was shown to be repellent to host-seeking mosquitoes in the laboratory testing with comparable efficacy to citronella (Thomas *et al.* 2009a, b). Given the demonstrated repellent activity against mosquitoes, it has been suggested that similar effectiveness may be achieved against other haematophagous

arthropods. The aim of this study was to investigate the ability of commercially available KO for its pediculicidal properties.

## EXPERIMENTAL METHODS

### *Collection of head lice for laboratory studies*

This work was approved by the University of South Australia's Human Research Ethics Committee (approvals 30059 and 32538). Head lice were obtained directly from the children (following written parental consent) at two primary schools in South Australia as well as a purpose run head lice clinic at the University of South Australia. Head lice were removed from individuals by combing dry hair from scalp to tip using a head lice comb (Barker and Altman, 2010). Head lice were removed and placed into a Petridish for holding (maximum 15 min) until the testing could commence.

### *Pediculicidal assay*

Each test Petridish contained a filter paper (7 cm diameter) that was uniformly covered in 1 mL treatment or control products. These amounts were pre-measured before trials and then applied immediately prior to testing against lice. KO (Therapeutic Goods Administration ARTG entry 72143; Ducane Estate, Waterhouse, Tasmania), both a 100 and 5% solution [in ethanol (90% of 70% v/v ethanol and 5% glycerol)], was tested along with two commercially

\* Corresponding author. Faculty of Health, University of Canberra, Building 12 Level D Room 36, Kirinari Street, Bruce, Canberra, ACT 2601, Australia. E-mail: [Jackson.Thomas@canberra.edu.au](mailto:Jackson.Thomas@canberra.edu.au)




Fig. 1. Pediculicidal properties of kunzea oil preparations compared with positive and negative controls.

available formulations, TTO head lice gel (Integria Healthcare Australia Pty Ltd, Eight Mile Plains, QLD; containing 5% TTO) and permethrin spray (Garrards Pesticides, Clovelly Park SA; containing 500 g kg<sup>-1</sup> permethrin). The permethrin was included here as a positive control insecticide. Dimethicone-free hair conditioner (Moogoo Skin Care Pty Ltd, Brisbane, QLD) was tested as a wet control that may possess a suffocant effect against lice but no toxicosis.

Each formulation was applied evenly onto the filter paper contained within a Petridish and allowed it to dry for approximately 1 h prior to testing. As only head lice removed from a volunteer within 15 min were used, for each replicate test, 5–10 head lice were used. Head lice were transferred to a dish using a small, soft, bristled paintbrush and placed within the centre of the treated filter paper. Head lice were observed every 15 min for up to 120 min and the number of dead individual lice recorded at each time point. The number of replicate tests for each formulation and control varied between 3 and 9. Moribund lice responsive to touch with a probe were considered alive, while death was determined to have occurred when there was no response to the probe. Lice classified as dead never recovered. Mortality of head lice was recorded as the mean number of dead head lice across the replicate tests. Kaplan–Meier survival analysis (STATA version 12, College Station, USA) was used to compare pediculicidal effects of

different treatments and to calculate mean survival times.

## RESULTS

Exposure to KO, both as a 5% and neat oil (100%, undiluted) resulted in 100% mortality within 120 min (Fig. 1) with the mean survival times of 17.1 min (5% solution) and 34.8 min (neat oil; 100%, undiluted) (Table 1). The mean survival time for the 5% KO was the shortest of any other formulation tested and has now been shown to be extremely effective at killing lice *in vitro* and demonstrated the similar effectiveness to TTO (Fig. 1; Table 1). Kaplan–Meier survival analysis showed no significant difference in the mortality rate of KO compared with the TTO-based formulation (log-rank test,  $P > 0.05$ ), which is also reflected in the mean survival times of lice exposed to the two oils (Table 1). Interestingly, the mean survival times of the permethrin-based formulation were substantially longer than both the KO and TTO-based formulations (Table 1). Over 40% mortality was recorded in head lice exposed to the plain hair conditioner after 90 min, a significantly lower rate (log-rank test,  $P < 0.001$ ) than any of the botanical or insecticide-based products. Less than 10% mortality was recorded in head lice in the dry control (nothing added to the filter paper) category after 120 min (Fig. 1). These differences in the mortality between KO/TTO and the dry and wet controls were

Table 1. Mean individual louse survival times and 95% confidence intervals (CI) when exposed to *in vitro* pediculicidal treatments, in descending order of pediculicidal activity

	Replicates	Mean survival time (min)	S.E.	95% CI
Kunzea oil (5%)	6	17.1	1.0	15.2–19.0
5% tea tree oil	4	21.2	1.9	17.4–25.1
Kunzea oil (100%)	9	34.8	3.5	27.8–41.7
Permethrin	4	56.4	7.9	41.0–71.8
Conditioner	3	157.5	25.9	106.8–208.2
Dry control	7	231.3	4.9	221.6–240.9

statistically significant (log-rank test,  $P < 0.001$ ). Survival analysis also revealed no significant differences in the mortality between KO and TTO.

#### DISCUSSION

This *in vitro* study demonstrates that the mean mortality of head lice exposed to KO does not differ appreciably to those exposed to currently registered formulations containing TTO. With the perception that formulations containing synthetic insecticides such as permethrin (Durand *et al.* 2012) may be driving the development of insecticide resistance, there is growing interest in novel pediculicides, particularly those of botanical origin, and a partial summary of the key candidates is summarized in Table 2. Botanical extracts avoid the problems arising from residual effects seen with synthetic pediculicides (Webb and Russell, 2011) and are very unlikely to yield resistance in lice because of multiple modes of action of each of their active principles (Carpinella *et al.* 2007). While commercial products containing botanical extracts are widely available and commonly used as both pediculicides and repellents against head lice, TTO-based formulations are most common. The results of this study demonstrate that KO as an active ingredient will provide comparable control of head lice to that of TTO.

Given the interest amongst the community for ‘natural’ solutions to head lice, it is important that appropriate testing be conducted to confirm the efficacy of any new pediculicide formulations (Heukelbach *et al.* 2008). Based on the results of this study, there is great potential in KO and that it has already demonstrated efficacy above the 80% mortality threshold suggested (Heukelbach *et al.* 2008). However, while *in vitro* testing shows great potential, it will be critical that controlled field trials and therapeutic potential of formulated products be assessed (Burkhart and Burkhart, 2001). The need for understanding the synergistic effect of non-active ingredients in commercial formulations was highlighted by the results of the testing presented here with 5% KO recording a lower mean survival time compared with the neat oil (100%, undiluted).

Many local health authorities continue to preference the ‘wet comb technique’ as a head lice management

strategy to reduce the possibility of overexposure to insecticides and minimizing the risk of insecticide resistance (Maxwell *et al.* 2014). However, the results of this study indicated that the use of conditioner does not provide the same level of pediculicidal action as that provided by KO. The dimethicone-free hair conditioner was tested but our data indicate that hair conditioner can kill only some adult lice, presumably through a suffocation effect. The use of hair conditioner without any active pediculicides present, physical suffocant properties of head lice will only assist in treatment if combined with a thorough combing and removal of ‘stunned’ head lice.

KO appears to be safe for topical application on the skin. KO has been tolerated well in clinical studies (human and animal) with concentrations ranging from 10 to 100% with no major documented solicited localized and/or systemic adverse drug reactions (Jacobson *et al.* 2009; Thomas *et al.* 2009a, b, 2015). This includes application of oil and/or formulations at skin sites including open wounds (animal trial), inflamed skin (e.g. scalp psoriasis) and nails infected with fungal infection (both human trials). However, more work is required to further establish the optimum safe concentration of the oil for topical application. Repeated analyses of aged KO (6–36 months, stored at room temperature) did not show any noticeable change in the chemical profile of the oil indicating little degradation of oil resulting from oxidation and polymerization (J. Thomas, unpublished data). KO has been recently examined for its activity against ectoparasites (scabies mites and head lice) and did not show any appreciable activity against scabies mites below 5% (J. Thomas, 2015, unpublished data). The 5% TTO has demonstrated remarkable activity against human scabies mites (Walton *et al.* 2004), and the commercial 5% TTO gel formulation has been widely used for the management of head lice in Australia. Therefore, a 5% KO formulation has been used in our study for a direct comparison with the proprietary 5% TTO head lice gel.

This study confirms the pediculicidal potential of KO and further investigation is required to determine formulations that can be used as effective alternatives to botanical or synthetic insecticides for the management of ectoparasitic infestations in

Table 2. A partial summary of various essential oil with documented activity against *Pediculus humanus capitis*

Essential oil	Species	Family
Eucalyptus oil	<i>Eucalyptus globules</i>	Myrtaceae
Marjoram oil	<i>Origanum majorana</i>	Lamiaceae
Pennyroyal oil	<i>Mentha pulegium</i>	Lamiaceae
Rosemary oil	<i>Rosmarinus officinalis</i>	Lamiaceae
Cade oil	<i>Juniperus oxycedrus</i>	Cupressaceae
Cardamome ceylon oil	<i>Elettaria cardamomum</i>	Zingiberaceae
Clove bud oil	<i>Eugenia caryophyllata</i>	Myrtaceae
Myrtle oil	<i>Myrtus communis</i>	Myrtaceae
Rosewood oil	<i>Aniba rosaeodora</i>	Lauraceae
Sage oil	<i>Salvia officinalis</i>	Lamiaceae
Tagetes oil	<i>Tagetes minuta</i>	Asteraceae
Cinnamon bark oil	<i>Cinnamomum zeylanicum</i>	Lauraceae
Nem oil	<i>Azadirachta indica</i>	Meliaceae
Clove oil	<i>Syzygium aromaticum</i>	Myrtaceae
Citronella oil	<i>Cymbopogon winteratus</i>	Poaceae
Lippia multiflora oil	<i>Lippia multiflora Moldenke</i>	Verbenaceae
Cardamom oil	<i>Elettaria cardamomum</i>	Zingiberaceae
Tea tree oil	<i>Melaleuca alternifolia</i>	Myrtaceae

Adopted from Rossini *et al.* (2008) and Yang *et al.* (2004).

humans and animals. Despite active promotion of the head lice treatments, studies have found carers vary greatly in the treatments used against pediculosis (Maxwell *et al.* 2014). Recommendations for pediculicide use should be regularly reviewed in light of developing resistance levels and new therapies. The ideal head lice treatment has been defined as being safe, effective, easy to use, inexpensive and available without a prescription and as a single dose (Frankowski and Bocchini, 2010). Regardless of the development of novel pediculicide formulations, community education will remain a critical component of head lice management strategies.

#### ACKNOWLEDGEMENTS

Louise Massie provided great assistance at the Sansom Institute for Health Research Clinical Trials Facility. The cooperation of primary schools and their Principals was pivotal to this study and greatly appreciated.

#### FINANCIAL SUPPORT

This study was funded by the Faculty of Health, University of Canberra internal grant awarded to Dr Jackson Thomas.

#### CONFLICT OF INTEREST

The authors have no competing interests.

#### ETHICAL STANDARDS

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki

Declaration of 1975, as revised in 2008. This work was approved by the University of South Australia's Human Research Ethics Committee (approvals 30059 and 32538).

#### REFERENCES

- Audino, P. G., Vassena, C., Zerba, E. and Picollo, M. (2007). Effectiveness of lotions based on essential oils from aromatic plants against permethrin resistant *Pediculus humanus capitis*. *Archives of Dermatological Research* **299**, 389–392.
- Barker, S. C. and Altman, P. M. (2010). A randomised, assessor blind, parallel group comparative efficacy trial of three products for the treatment of head lice in children-melaleuca oil and lavender oil, pyrethrins and piperonyl butoxide, and a "suffocation" product. *BMC Dermatology* **10**, 6.
- Burkhart, C. N. and Burkhart, C. G. (2001). Recommendation to standardize pediculicidal and ovicidal testing for head lice (Anoplura: Pediculidae). *Journal of Medical Entomology* **38**, 127–129.
- Carpinella, M. C., Miranda, M., Almiron, W. R., Ferrayoli, C. G., Almeida, F. L. and Palacios, S. M. (2007). *In vitro* pediculicidal and ovicidal activity of an extract and oil from fruits of *Melia azedarach* L. *Journal of the American Academy of Dermatology* **56**, 250–256.
- Currie, M. J., Reynolds, G. J., Glasgow, N. J. and Bowden, F. J. (2010). A pilot study of the use of oral ivermectin to treat head lice in primary school students in Australia. *Pediatric Dermatology* **27**, 595–599.
- Diamantis, S. A., Morrell, D. S. and Burkhart, C. N. (2009). Treatment of head lice. *Dermatologic Therapy* **22**, 273–278.
- Durand, R., Bouvresse, S., Berdjane, Z., Izri, A., Chosidow, O. and Clark, J. (2012). Insecticide resistance in head lice: clinical, parasitological and genetic aspects. *Clinical Microbiology and Infection* **18**, 338–344.
- Frankowski, B. L. and Bocchini, J. A. (2010). Head lice. *Pediatrics* **126**, 392–403.
- Heukelbach, J., Canyon, D., Oliveira, F., Muller, R. and Speare, R. (2008). *In vitro* efficacy of over-the-counter botanical pediculicides against the head louse *Pediculus humanus var capitis* based on a stringent standard for mortality assessment. *Medical and Veterinary Entomology* **22**, 264–272.
- Jacobson, G. A., Narkowicz, C., Thomas, J. and Peterson, G. M. (2009). *Clinical Trial of Kunzea Oil for Onychomycosis Treatment: a Randomised Controlled Trial*. Rural Industries Research & Development Corporation, Publication No. 09/098, pp. 4–5.
- Khambay, B. P., Beddie, D. G., Simmonds, M. S. and Green, P. W. (1999). A new insecticidal pyranocyclohexenedione from *Kunzea ericifolia*. *Journal of Natural Products* **62**, 1423–1424.

- Khambay, B. P., Beddie, D. G. and Simmonds, M. S.** (2002). An insecticidal mixture of tetramethylcyclohexenedione isomers from *Kunzea ambigua* and *Kunzea baxterii*. *Phytochemistry* **59**, 69–71.
- Maxwell, M., Crawford, B. and Rose, V.** (2014). Nitbusters: lessons from a school-based intervention study to reduce head lice in a disadvantaged community. *Health Promotion Journal of Australia* **25**, 67–68.
- Rossini, C., Castillo, L. and González, A.** (2008). Plant extracts and their components as potential control agents against human head lice. *Phytochemistry Reviews* **7**, 51–63.
- Speare, R. and Buettner, P. G.** (1999). Head lice in pupils of a primary school in Australia and implications for control. *International Journal of Dermatology* **38**, 285–290.
- Thomas, J., Narkowicz, C., Peterson, G. M., Jacobson, G. A. and Narayana, A.** (2009a). Treatment of pastern dermatitis with a formulation containing kunzea oil: a randomised controlled trial. *Veterinary Record* **20**, 619–623.
- Thomas, J., Webb, C., Narkowicz, C., Jacobson, G., Peterson, G., Davies, N. and Russell, R.** (2009b). Evaluation of repellent properties of volatile extracts from the Australian native plant *Kunzea ambigua* against *Aedes aegypti* (Diptera: Culicidae). *Journal of Medical Entomology* **46**, 1387–1391.
- Thomas, J., Narkowicz, C., Jacobson, G. and Davies, N.** (2010). An examination of the essential oils of Tasmanian *Kunzea ambigua*, other *Kunzea* spp. and commercial kunzea oil. *Journal of Essential Oil Research* **22**, 381–385.
- Thomas, J., Jacobson, G. A., Narkowicz, C. and Peterson, G. M.** (2015). Safety and efficacy of kunzea oil-containing formulations for the management of psoriasis: a randomized, controlled trial. *Journal of Clinical Pharmacy and Therapeutics* **40**, 566–572.
- Toloza, A. C., Lucia, A., Zerba, E., Masuh, H. and Picollo, M. I.** (2008). Interspecific hybridization of Eucalyptus as a potential tool to improve the bioactivity of essential oils against permethrin-resistant head lice from Argentina. *Bioresource Technology* **99**, 7341–7347.
- Walton, S. F., McKinnon, M., Pizzutto, S., Dougall, A., Williams, E. and Currie, B. J.** (2004). Acaricidal activity of *Melaleuca alternifolia* (tea tree) oil: *in vitro* sensitivity of *Sarcoptes scabiei* var. *hominis* to terpinen-4-ol. *Archives of Dermatology* **140**, 563–566.
- Webb, C. E. and Russell, R. C.** (2011). Advice to travellers on topical insect repellent use against dengue mosquitoes in far north Queensland, Australia. *Journal of Travel Medicine* **18**, 282–283.
- Yang, Y. C., Lee, H. S., Clark, J. M. and Ahn, Y. J.** (2004). Insecticidal activity of plant essential oils against *Pediculus humanus capitis* (Anoplura: Pediculidae). *Journal of Medical Entomology* **41**, 699–704.