

Gerlach, Tobias; Harant, Jochen:

A note on domination in bipartite graphs

URN: [urn:nbn:de:gbv:ilm1-2020200083](https://nbn-resolving.org/urn:nbn:de:gbv:ilm1-2020200083)

Original published in: Discussiones mathematicae. Graph theory / Uniwersytet Zielonogórski, Wydział Matematyki, Informatyki i Ekonometrii. - Warsaw : De Gruyter Open. - 22 (2002), 2, p. 229-231.

Original published: 2002

ISSN: 2083-5892

DOI: [10.7151/dmgt.1171](https://doi.org/10.7151/dmgt.1171)

[Visited: 2020-01-17]



This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/) license.
To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

A NOTE ON DOMINATION IN BIPARTITE GRAPHS

TOBIAS GERLACH AND JOCHEN HARANT

Department of Mathematics
Technical University of Ilmenau
D-98684 Ilmenau, Germany

Abstract

DOMINATING SET remains *NP*-complete even when instances are restricted to bipartite graphs, however, in this case VERTEX COVER is solvable in polynomial time. Consequences to VECTOR DOMINATING SET as a generalization of both are discussed.

Keywords: bipartite graph, domination.

2000 Mathematics Subject Classification: 05C35.

For terminology and notation not defined here we refer to [2, 3]. Given a finite, simple, and undirected graph G without isolated vertices, $V(G) = \{1, \dots, n\}$, $E(G)$, $G[V]$, and $d_i(G)$ denote its vertex set, its edge set, the subgraph of G induced by $V \subseteq V(G)$, and the degree of $i \in V(G)$ in G , respectively. Furthermore, let $\vec{1} = (1, \dots, 1) \in R^n$ and $\vec{d}(G) = (d_1(G), \dots, d_n(G))$. Given an integral vector $\vec{k} = (k_1, \dots, k_n)$ with $1 \leq k_i \leq d_i(G)$ for $i = 1, \dots, n$, a set $D \subseteq V(G)$ is called \vec{k} -dominating if $d_i(G[V(G) \setminus D]) \leq d_i(G) - k_i$ for $i \in V(G) \setminus D$. Consider the corresponding decision problem, which was investigated in [4]:

VECTOR DOMINATING SET

Instance: A graph G on $V(G) = \{1, \dots, n\}$, an integral vector $\vec{k} = (k_1, \dots, k_n)$ with $1 \leq k_i \leq d_i(G)$ for $i = 1, \dots, n$, and a positive integer l .

Question: Does G contain a \vec{k} -dominating set D with $|D| \leq l$?

The restriction of VECTOR DOMINATING SET to $\vec{k} = \vec{1}$ is the decision problem DOMINATING SET, remaining *NP*-complete even when instances are restricted to bipartite graphs [1]. It is easy to see that $D \subseteq V(G)$ is $\vec{d}(G)$ -dominating if and only if each edge of G has at least one endvertex in D .

The corresponding decision problem VERTEX COVER is NP -complete in general, however, is solvable in polynomial time for bipartite graphs [3]. The question arises in how many components \vec{k} may differ from $\vec{1}$ or from $\vec{d}(G)$ such that the restriction of VECTOR DOMINATING SET to bipartite graphs remains NP -complete or is solvable in polynomial time, respectively. Theorem 1 and Theorem 2 give partial answers to this question.

Theorem 1. *Given $0 < c < \frac{1}{2}$, the following restriction of VECTOR DOMINATING SET remains NP -complete.*

Instance: A bipartite graph G on $V(G) = \{1, \dots, n\}$, an integral vector $\vec{k} = (k_1, \dots, k_n)$ with $1 \leq k_i \leq d_i(G)$ for $i = 1, \dots, n$ and $|\{i \in V(G) \mid k_i > 1\}| = \lceil cn \rceil$, and a positive integer l .

Question: Does G contain a \vec{k} -dominating set D with $|D| \leq l$?

Theorem 2. *Given $c > 0$, the following restriction of VECTOR DOMINATING SET is solvable in polynomial time.*

Instance: A bipartite graph G on $V(G) = \{1, \dots, n\}$, an integral vector $\vec{k} = (k_1, \dots, k_n)$ with $1 \leq k_i \leq d_i(G)$ for $i = 1, \dots, n$ and $|E(G[\{i \in V(G) \mid k_i < d_i(G)\}])| \leq c \log_2 n$, and a positive integer l .

Question: Does G contain a \vec{k} -dominating set D with $|D| \leq l$?

Proof of Theorem 1. Given $0 < c < \frac{1}{2}$, we shall transform DOMINATING SET to the restriction of VERTEX DOMINATING SET of Theorem 1. Let a graph H on $V(H) = \{1, \dots, m\}$ and a positive integer l be an instance of DOMINATING SET, the positive integer r be chosen such that $\frac{2cm}{1-2c} \leq r < \frac{2cm}{1-2c} + 1$, hence, $c(2m + 2r) \leq r < c(2m + 2r) + (1 - 2c)$ implying $r = \lceil c(2m + 2r) \rceil \geq 1$, and G be constructed with $V(G) = \{1, \dots, 2m + 2r\}$ and $E(G) = \{(i, m + j), (j, m + i) \mid (i, j) \in E(H)\} \cup \{(i, m + i), (m + i, 2m + j), (2m + j, 2m + r + j) \mid i = 1, \dots, m, j = 1, \dots, r\}$. Following the ideas in [1], it is easy to see that G is bipartite and that for $D(H) \subseteq V(H)$, $D(G) = \{i + m \mid i \in D(H)\} \cup \{2m + 1, \dots, 2m + r\}$ is a $\vec{1}$ -dominating set of G if and only if $D(H)$ is a $\vec{1}$ -dominating set of H . Let $\vec{k} = (k_1, \dots, k_{2m+2r})$ be an arbitrary integral vector with $1 \leq k_p \leq d_p(G)$ for $p = 1, \dots, 2m + 2r$, and $k_p = 1$ if and only if $p \in V(G) \setminus \{2m + 1, \dots, 2m + r\}$. Then $|\{i \in V(G) \mid k_i > 1\}| = r = \lceil c|V(G)| \rceil$, and $D(G)$ is even a \vec{k} -dominating set of G if $D(H)$ is a $\vec{1}$ -dominating set of H . Hence, H contains a $\vec{1}$ -dominating set of cardinality l if and only if G contains a \vec{k} -dominating set of cardinality $l + r$. With $|V(G)| = 2m + 2r < 2m + 2c(2m + 2r) + 2(1 - 2c) = 2|V(H)| + 2c|V(G)| + 2(1 - 2c)$, hence, $|V(G)| < \frac{2}{1-2c}|V(H)| + 2$ we are done. ■

Proof of Theorem 2. For $D \subseteq V(G)$, let $E(D)$ be the set of edges having no endvertex in D , and $H(D)$ be the graph arising from G by deleting the edges of $E(D)$. It is easy to see that D is a \vec{k} -dominating set of G if and only if $E(D) \subseteq E(G[\{i \in V(G) \mid k_i < d_i(G)\}])$, each endvertex i of an edge in $E(D)$ is endvertex of at most $d_i(G) - k_i$ edges in $E(D)$, and D is a $\vec{d}(H(D))$ -dominating set of $H(D)$. Since VERTEX COVER is solvable in polynomial time and the number of sets $E(D)$ is at most $2^{|E(G[\{i \in V(G) \mid k_i < d_i(G)\}])|} \leq n^c$ we are done. ■

Remark. It remains open whether the bounds cn ($0 < c < \frac{1}{2}$) and $c \log_2 n$ ($c > 0$) of Theorem 1 and Theorem 2 can be made significantly greater such that VECTOR DOMINATING SET is still NP-complete or solvable in polynomial time, respectively.

References

- [1] G.J. Chang and G.L. Nemhauser, *The k -domination and k -stability problems in sun-free chordal graphs*, SIAM J. Algebraic Discrete Methods **5** (1984) 332–345.
- [2] R. Diestel, *Graph Theory* (Springer-Verlag, New York, 2000).
- [3] M.R. Garey and D.S. Johnson, *Computers and Intractability* (W.H. Freeman and Company, San Francisco, 1979).
- [4] J. Harant, A. Pruchnewski and M. Voigt, *On dominating sets and independent sets of graphs*, Combinatorics, Probability and Computing **8** (1999) 547–553.

Received 24 August 2000