Treatment of Eutrophic KotaSAS Lake by Phytoremediation using Macrophytes Species; Eichhornia *Crassipes* and *Pistia Stratiotes*

J Muhammad Haziq¹, I Farah Amalina¹, AR Abdul Syukor¹, S. Sulaiman¹ and A.H. Mohd Rashid²

 ¹Faculty of Civil Engineering and Earth Resources, Universiti Malaysia Pahang (UMP), Lebuhraya Tun Razak, 26300, Gambang, Kuantan, Pahang, Malaysia.
²Faculty of Industrial Management, Universiti Malaysia Pahang (UMP), Lebuhraya Tun Razak, 26300, Gambang, Kuantan, Pahang, Malaysia.

Corresponding Author: syukor@ump.edu.my; abdsyukor@gmail.com

Abstract. The wastewater treatment is known as a necessary attention for the process of retreatment towards the removal of suspended solids. Phytoremediation is a plant-based system which directly used of green plant in-situ to reduce pollutants in the lakes. This method is ecologically friendly and solar dependent clean-up technology. This study was undertaken ex situ where the aim of this study is to treat the eutrophic man-made lake at KotaSAS lakeside through following objectives; to identify the variation of physicochemical attributes of KotaSAS lake water through phytoremediation, to evaluate the potential of selected heavy metals and nutrients removal (nitrate and phosphorus) using Eichornia crassipes (water hyacinth) and Pistia stratiotes (water lettuce) and to determine the correlations between sampling points and physicochemical attributes using statistical analysis.. The method of analysis of lake water was referred by Standard Method for Examination of Water and Wastewater by APHA. Then, statistical notation was conducted on the results obtained to identify the accuracy and validity of data, which in this case, correlations and t-test statistical method was used. Referring to the statistical notation of (p<0.05), conclusion on the significance of the result and variables were made. It is justified statistically that the data obtained for each treatment using different types of plants are valid and concise. As a result, it is found that water lettuce and water hyacinth have different nutrients removal and heavy metals removal efficiency but, in all cases,, these phytoremediation agents exhibited nutrients removal efficiency from the range of 27% up to 58% followed by heavy metals removal efficiency from the range of 23% up to 60%. Water hyacinth exhibited great nitrate and phosphorus removal efficiency, 51.51% and 58.81% whereby as for water lettuce exhibited great heavy metals removal efficiency for manganese (Mn), iron (Fe) and copper (Cu), 60.68%, 58.2% and 26.4% respectively. With this achievement acquired in phytoremediation technology employing water hyacinth, it is of utmost important for this technology to be implemented in larger scales thereafter. Hence, this plant is suitable to be used in controlling eutrophic lake due to its hyper-accumulating ability.

- Mayo, A.W. & Hanai, E.E. 2016. Modeling phytoremediation of nitrogen-polluted water using water hyacinth (Eichhornia crassipes). Physics and Chemistry of the Earth, Parts A/B/C, 100, 170–180, 10.1016/j.pce.2016.10.016.
- Mthethwa, N.P., Nasr, M., Bux, F. & Kumari, S. 2018. Utilization of Pistia stratiotes (aquatic weed) for fermentative biohydrogen: Electron-equivalent balance, stoichiometry, and cost estimation. International Journal of Hydrogen Energy, 43, 8243–8255, 10.1016/j.ijhydene.2018.03.099.
- Ng, Y.S. & Chan, D.J.C. 2016. Wastewater phytoremediation by Salvinia molesta. Journal of Water Process Engineering, 15, 107–115, 10.1016/j.jwpe.2016.08.006.
- Nivetha, C., Subraja, S., Sowmya, R. & Induja, N.M. 2016. Water Lettuce for Removal of Nitrogen and Phosphate from Sewage. 13, 98–101, 10.9790/1684-13020198101.
- Qin, H., Zhang, Z., Liu, M., Liu, H., Wang, Y., Wen, X., Zhang, Y. & Yan, S. 2016. Site test of phytoremediation of an open pond contaminated with domestic sewage using water hyacinth and water lettuce. Ecological Engineering, 95, 753–762, 10.1016/j.ecoleng.2016.07.022.
- Rezania, S., Ponraj, M., Talaiekhozani, A., Mohamad, S.E., Md Din, M.F., Taib, S.M., Sabbagh, F. & Sairan, F.M. 2015. Perspectives of phytoremediation using water hyacinth for removal of heavy metals, organic and inorganic pollutants in wastewater. Journal of Environmental Management, 163, 125–133, 10.1016/j.jenvman.2015.08.018.
- Syukor, A.R.A., Sulaiman, S., Siddique, M.N.I., Zularisam, A.W. and Said, M.I.M. 2015. Integration of phytogreen for heavy metal removal from wastewater. Journal of Cleaner Production 112, pp. 3124–3131. Available at: http://linkinghub.elsevier.com/retrieve/pii/S0959652615015759.
- Suhendrayatna, Marwan, Andriani, R., Fajriana, Y. & Elvitriana. 2012. Removal of municipal wastewater BOD, COD, and TSS by phyto-reduction: a laboratory-scale comparison of aquatic plant at different species Typha latifolia and Saccharum spontaneum. International Journal of Engineering and Innovative Technology (IJEIT), 2, 333–337 Available at: http://www.ijeit.com/vol 2/Issue 6/IJEIT1412201212_59.pdf.
- Tanee, F.B.G. & Akonye, L.A. 2009. Effectiveness of Vigna Unguiculata as a Phytoremediation Plant in the remediation of Crude Oil polluted soil for Cassava (Manihot Esculenta; Crantz) Cultivation. J. Appl. Sci. Environ. Manage., 13, 43–47 Available at: www.bioline.org.br/ja.
- The Star. 2014. Ill effects of nitrogen overdose Available at: http://www.thestar.com.my/news/environment/2014/06/30/ill-effects-of-nitrogen-overdose/.
- Ugya A.Y., I.T.S. and T.S.M. 2016. The Role Of Phytoremediation In Remediation Of Industrial Waste, 10.20959/wjpr20161275.
- Yang, X., Wu, X., Hao, H. & He, Z. 2008. Mechanisms and assessment of water eutrophication. Journal of Zhejiang University. Science. B, 9, 197–209, 10.1631/jzus.B0710626.
- Zheng, J.C., Liu, H.Q., Feng, H.M., Li, W.W., Lam, M.H.W., Lam, P.K.S. & Yu, H.Q. 2016. Competitive sorption of heavy metals by water hyacinth roots. Environmental Pollution, 219, 837–845, 10.1016/j.envpol.2016.08.001.