

1 **CASE REPORT**

2

3 **Title:** Optimised visual outcome after asymmetrical multifocal IOL rotation

4

5 **Authors:** Eric E Pazo MD^{1, 2}, Olivier Richoz MD PhD³, Richard McNeely BSc^{1, 2},
6 Zachary A Millar⁴, Tara CB Moore PhD², Jonathan E Moore PhD FRCOphth^{1, 2}.

7

8 1 Cathedral Eye Clinic, University of Ulster, 89-91 Academy Street, Belfast, UK.

9 2 Biomedical Sciences Research Institute, University of Ulster, Coleraine, UK.

10 3 Department of Ophthalmology, Royal Victoria Hospital, Belfast, UK.

11 4 Faculty of Biology, University of Cambridge, Cambridgeshire, UK.

12

13 Prior Presentation: None

14

15 Financial Support: None.

16

17 The authors' have no financial or proprietary interest in a product, method, or
18 material, or lack thereof.

19

20 Name and address of author to receive reprint requests:

21 **Professor Jonathan E Moore, 89-91 Academy Street, Belfast, Co Antrim, BT1**

22 **2LS. Tel: +44 28 90322020 Fax: +44 28 90321808 Email:**

23 jmoorecathedral@gmail.com

24 **PRECIS**

25 Asymmetrical IOLs require active management to maximise the surface area of
26 either the distance or near-add, between the dominant and non-dominant eyes, to
27 ensure optimal visual performance.

28 **ABSTRACT**

29

30 **Purpose:** To report an improved visual outcome after rotation of an asymmetrical
31 multifocal intraocular lens.

32

33 **Methods:** A 58-year-old patient underwent bilateral phacoemulsification with
34 asymmetrical multifocal intraocular lens (MFIOL) implantation. Subsequently postop
35 the IOL was rotated to improve pupil centration.

36

37 Postop UDVA in both eyes was 0.0 logMAR, UNVA in both the eyes was 0.0
38 logMAR. QOV questionnaire scores for day and night were 5 and 7 respectively. The
39 centre of the MFIOL in the dominant eye was initially found to be 0.2 mm supero-
40 temporally displaced increasing the percentage area of 'near-add' compared to
41 'distance-add' within the physiological pupil. Rotation of this IOL 120° clockwise
42 greatly improved the IOL centration within the pupil centre and resulted in an
43 immediate improvement in UDVA to -0.1 logMAR and simultaneously QOV to 8 and
44 9 respectively.

45

46 **Conclusions:** Assessment of the centration of an asymmetrical MFIOL is important
47 particularly if there are dysphotopic or other visual complaints.

48 **Introduction**

49

50 The aim of multifocal intraocular lens (MFIOLs) use is to restore distance,
51 intermediate and near visual function post cataract surgery. The new generation of
52 refractive radially asymmetrical MFIOLs' aim to alleviate the occurrence of optical
53 side effects. The SBL-3 (Lenstec, Inc.) is a bi-aspheric asymmetrical refractive
54 MFIOL with a +3.00 D add in the inferior anterior optic (figure 1). A transition zone
55 separates the distance and the near-add sections of the lens and the near segment
56 occupies 42% of the total lens optic.

57

58 We report a case in which a patient underwent bilateral implantation of asymmetrical
59 MFIOL; the near-add in both eyes was positioned infero-nasally as per manufacturer
60 recommendation. During the 1-month post-cataract surgery assessment the patient
61 complained of problems driving during the day and problems with vision in
62 supermarket lighting. It was noted that under photopic conditions the exposure of the
63 near-add was maximised in the patient's dominant eye thereby resulting in a myopic
64 state of accommodation. To alleviate the patient's myopic state during photopic
65 conditions the MFIOL in the dominant eye was rotated 120° clock wise to a supero-
66 temporal position. One month after the MFIOL rotation (2-months post cataract
67 extraction) the patient was invited for an assessment and reported vision problems
68 with night-driving and in the supermarket were alleviated and quality of vision was
69 greatly improved. This improvement in the quality of vision was found to be stable
70 during the patient's 4-month post-operative (3 month post rotation) ophthalmic
71 assessment (Table 1). To our knowledge, this is the first report of decentration of a
72 new radially asymmetric MFIOL and alleviation of the decentration by rotation.

73 CASE REPORT

74

75 A 58-year-old man presented at the ophthalmology clinic with gradually decreased
76 visual acuity because of bilateral cataract. On presentation, unaided distance visual
77 acuity (UDVA) in the right and left eye was 0.5 logMAR and 0.6 logMAR. Unaided
78 near visual acuity (UNVA) for the right and left was 0.4 logMAR and 0.2 logMAR.
79 Using a validated questionnaire, the quality of vision (QOV) pre-operatively was
80 recorded to be 4 and 3 (on a scale of 0-10) for day and night respectively. The IOL-
81 Master (Zeiss) device was used to measure corneal curvature, anterior chamber
82 depth, axial length, and subsequent IOL calculation using the Hoffer Q formula.

83

84 Cataract extraction with asymmetrical multifocal intraocular lens implantation was
85 performed in both eyes. The near-add was placed infero-nasally in both eyes by an
86 experienced surgeon (JEM). Standard sutureless on-steep axis corneal
87 phacoemulsification (2.8 mm incision) was performed with a uniform capsulorhexis of
88 5.2 mm.

89

90 One month after this uneventful cataract surgery, the patient complained that he was
91 experiencing difficulty with vision while driving during the day and in supermarkets.
92 On assessment, the UDVA in both eyes was 0.0 logMAR. UNVA also improved in
93 both the eyes to 0.0 logMAR. QOV questionnaire scores for day and night were still
94 low at 5 and 7 respectively (pre cataract extraction:- 4 and 3 respectively). Slitlamp
95 anterior segment and fundus examinations were unremarkable and the near-add of
96 the MFIOOL was confirmed to be oriented infero-nasally in both eyes.

97

98 On assessing the pupil of the dominant eye under photopic conditions using a
99 slitlamp; it was observed that the near-add surface had high exposure in the
100 dominant eye. Digital retro-illumination photographs were taken of dilated and
101 undilated pupil. Adobe PS suite (Adobe Systems Inc, San Jose, California) was used
102 to determine surface area exposure, decentration, capsular dimension changes and
103 pupil shift. After the risk and benefits were explained to the patient, the infero-nasally
104 placed (near-add) asymmetrical MFIOL in the dominant right eye was rotated 120°
105 clockwise to a more supero-temporally positioned near-add (figure 2a) 1-month post
106 cataract operation, ensuring that a more normal degree of distance-add was now
107 present within the physiological pupil.

108

109 Three month following IOL rotation surgery the UDVA in both eyes was -0.1 logMAR.
110 UNVA in both the eyes was -0.1 logMAR (Table 1). The patient was happy with
111 driving and seeing in the supermarket during the day. This improvement in
112 satisfaction of visual performance was reflected in the QOV questionnaire scores
113 with day and night scores of 8 and 9 (1-month post rotation) and after 3-months post
114 rotation the QOV vision was 9 for QOV day and 9 for QOV night, which was
115 previously 5 and 7 for day and night respectively (prior to rotation). The final position
116 of the MFIOL near-add in the left eye was infero-nasal and supero-temporal in the
117 right eye (figure 2a and 2b).

118

119 **DISCUSSION**

120

121 SBL-3 multifocal IOL is a relatively new lens; the recent case series of bilateral
122 implantation on 53 eyes published by Venter *et al.*, reported a good range of

123 distance, intermediate and near visual acuity in patients.¹ The rotation of
124 asymmetrical MFIOL on its axis was compared before by Moore *et al*² and found
125 that the placement of the near-add in the superior or inferior position in Mplus
126 (Lentis, Inc.) had no significant overall difference in the mean subjective or objective
127 outcomes²

128

129 Decentration of any MFIOL can lead to decreased visual acuity and photopic
130 phenomenon, which has an adverse affect upon the quality of vision for the patient.³
131 The effect of decentration of a multifocal IOL upon visual quality can be further
132 compounded by a large angle kappa, resulting in central optical rays potentially
133 passing through the periphery of the MFIOL rather than its centre.⁴ The centration of
134 any MFIOL with respect to the physiological pupil centre can be difficult principally as
135 this is dictated generally by the position of capsular bag periphery.⁵ SBL-3 is radially
136 asymmetric and centration appears to play a crucial role for good quality of vision as
137 documented in this particular case report where the supero-temporal displacement of
138 approximately 0.2 mm in the dominant eye from the centre resulted in poor quality of
139 vision. Possible factors that may have influenced this decentration with respect to the
140 physiological pupil include capsular contraction, haptic movement, IOL rotation or
141 pupil shift.^{3,6,7}

142

143 Pupil shift refers to a slight change in reference to the pupil's central location
144 between mesopic, photopic, and pharmacologically dilated conditions⁸ and this
145 tendency of the pupil to shift makes it all the more difficult for a precise positioning of
146 the asymmetrical MFIOL. Closer examination of the photopic pupil of the patient's
147 dominant eye revealed that a photopic pupil shift occurred towards the infra-nasal

148 region⁹ and thereby maximised the light exposure to the inferiorly placed near-add of
149 the IOL; making distance vision during photopic conditions difficult for the patient.

150

151 In summary, postoperative rotation of asymmetrical MFIOL can be beneficial for
152 some patients experiencing dysphotopsia and poor quality of vision (figure 2a). It is
153 key to ensure that that the dominant eye is optimised for distance viewing by
154 maximising the area of distance add within the mesopic and photopic pupil.

155 Determining where the physiological pupil centre lies during surgery in an attempt to
156 centre the IOL within a pharmacologically dilated pupil is difficult. What we can
157 deduce however from this case report is that rotation of the IOL can result in different
158 final positions for the centre of the IOL. This is due to the asymmetric nature of the
159 capsular bag¹⁰ and the differences between the centre of the bag and the centre of
160 the pupil. Asymmetrical MFIOLs are not circular and nor is the capsular bag,
161 therefore one can actively alter the resultant centration of the IOL by rotating it into
162 different positions. Taking these factors into consideration the near-add positioning
163 should be assessed individually for optimal positioning of MFIOL and potentially
164 different positions utilised for the dominant and non-dominant eyes.

165 **References**

- 166 1. Blaylock J. Visual outcomes after bilateral or unilateral implantation of
167 nonrotational dual-optic bi-aspheric IOL. *2014 American Society of Cataract and*
168 *Refractive Surgery, conference abstract 2014*
- 169 2. de Wit DW, Diaz JM, Moore TC, Moore JE. Refractive lens exchange for a
170 multifocal intraocular lens with a surface-embedded near section in mild to moderate
171 anisometropic amblyopic patients. *J Cataract Refract Surg.* 2012;38(10):1796-1801.
- 172 3. Crnej A, Hirschall N, Nishi Y, et al. Impact of intraocular lens haptic design and
173 orientation on decentration and tilt. *J Cataract Refract Surg.* 2011;37(10):1768-1774.
- 174 4. Altmann GE, Nichamin LD, Lane SS, Pepose JS. Optical performance of 3
175 intraocular lens designs in the presence of decentration. *J Cataract Refract Surg.*
176 2005;31(3):574-585.
- 177 5. Sauer T, Mester U. Tilt and decentration of an intraocular lens implanted in the
178 ciliary sulcus after capsular bag defect during cataract surgery. *Graefes Arch Clin*
179 *Exp Ophthalmol.* 2013;251(1):89-93.
- 180 6. van der Linden JW, van der Meulen IJ, Mourits MP, Lapid-Gortzak R. In-the-bag
181 decentration of a hydrophilic radially asymmetric multifocal intraocular lens
182 secondary to capsule contraction. *J Cataract Refract Surg.* 2013;39(4):642-644.
- 183 7. Patel C, Ormonde S, Rosen PH, Bron AJ. Postoperative intraocular lens rotation:
184 A randomized comparison of plate and loop haptic implants. *Ophthalmology.*
185 1999;106(11):2190-2196.

- 186 8. Yang Y, Thompson K, Burns SA. Pupil location under mesopic, photopic, and
187 pharmacologically dilated conditions. *Invest Ophthalmol Vis Sci.* 2002;43(7):2508-
188 2512.
- 189 9. Porter J, Yoon G, Lozano D, et al. Aberrations induced in wavefront-guided laser
190 refractive surgery due to shifts between natural and dilated pupil center locations. *J*
191 *Cataract Refract Surg.* 2006;32(1):21-32.
- 192 10. Strenn K, Menapace R, Vass C. Capsular bag shrinkage after implantation of an
193 open-loop silicone lens and a poly (methyl methacrylate) capsule tension ring. *J*
194 *Cataract Refract Surg.* 1997;23(10):1543-1547.
- 195

196 **FIGURES LEGENDS**

197

198 **Figure 1.** Lenstec SBL-3, asymmetrical multifocal intraocular lens.

199

200 **Figure 2a.** Pupil (right eye) under pharmacological dilated conditions, post rotation of
201 MFIOL. Superio-temporal positioned near-add.

202

203 **Figure 2b.** Pupil (left eye) under pharmacological dilated conditions, infero-nasally
204 positioned near-add.

Table 1. Comparative Pre-op and post-operative data.

Biometry (preoperative)				
	OD		OS	
AL	23.65 mm		23.51 mm	
K1	41.46 D 10°		41.56 D 155°	
K2	41.87 D 100°		41.77 D 65°	
ACD	3.00 mm		3.05 mm	
Angle kappa	4.02°		3.75°	
Photopic pupil	3.2 mm		3.3 mm	
Mesopic pupil	4.1 mm		3.8 mm	
Sphere (D)	1.5		1.4	
Mean corneal astigmatism (D)	0.74		0.6	
	Pre-op	Post-op 1 month	Post rotation 1-month	Post rotation 3-months
Visual Acuity				
UDVA (OD)	0.5	0	-0.1	-0.1
UDVA (OS)	0.6	0	-0.1	-0.1
BCVA (OD)	0.2	-0.1	-0.1	-0.1
BCVA (OS)	0.2	0.04	0	-0.1
BCVA (Bino)	0.2	-0.1	-0.1	-0.1

UNVA (OD)	0.4	0	-0.1	-0.1
UNVA (OS)	0.2	-0.1	-0.1	-0.1
Questionnaire				
QOV (Day)	4	5	8	9
QOV (Night)	3	7	9	9
IOL surface area exposure of near-add in photopic pupil				
OD	NA	60%	20%	20%
OS	NA	40%	NA	NA
<p>AL=Axial length; ACD= Anterior chamber depth; QOV=Quality of vision; D=Dioptre; UDVA=Unaided Distance Visual Acuity; BCVA=Best Corrected Visual Acuity; UNVA=Unaided Near Visual Acuity; Bino=Binocular; IOL=Intraocular lens.</p>				





