

1

2

3 **Ursodeoxycholic acid and lithocholic acid exert anti-inflammatory actions in the colon**

4

5 Joseph B.J. Ward <sup>1</sup>, Natalia K. Lajczak<sup>1</sup>, Orlaith B. Kelly<sup>1</sup>, Aoife M. O'Dwyer<sup>1</sup>, Ashwini K.  
6 Giddam<sup>2</sup>, Joan N. Ni Gabhainn<sup>3</sup>, Placido Franco<sup>4</sup>, Murtaza M. Tambuwala<sup>5</sup>, Caroline A.  
7 Jefferies<sup>6</sup>, Simon Keely<sup>2</sup>, Aldo Roda<sup>4</sup>, and Stephen J. Keely<sup>1</sup>.

8 <sup>1</sup>Department of Molecular Medicine, Royal College of Surgeons in Ireland, Beaumont  
9 Hospital, Dublin 9, Ireland. <sup>2</sup>School of Biomedical Science and Pharmacy, University of  
10 Newcastle, Australia. <sup>3</sup>Department of Molecular and Cellular Therapeutics, Royal College of  
11 Surgeons in Ireland.<sup>4</sup>Department. of Chemistry, University of Bologna, Bologna, Italy.  
12 <sup>5</sup>School of Pharmacy and Pharmaceutical Sciences, Ulster University, Coleraine, Northern  
13 Ireland. <sup>6</sup>Department of Biomedical Sciences, Cedars Sinai Medical Centre, Los Angeles,  
14 USA.

15 **Corresponding Author:** Stephen J Keely, Department of Molecular Medicine, Royal  
16 College of Surgeons in Ireland, Education and Research Centre, Smurfit Building, Beaumont  
17 Hospital, Dublin 9, Ireland.

18 e-mail: [skeely@rcsi.ie](mailto:skeely@rcsi.ie)

19 Telephone: +3531 8093821

20 **Running Title:** UDCA and LCA prevent colonic inflammation

21

## 22 ABSTRACT

23 Inflammatory bowel diseases (IBD) are a group of common and debilitating chronic intestinal  
24 disorders for which currently-available therapies are often unsatisfactory. The naturally-  
25 occurring secondary bile acid, ursodeoxycholic acid (UDCA), has well-established anti-  
26 inflammatory and cytoprotective actions and may therefore be effective in treating IBD.  
27 Here, we aimed to investigate regulation of colonic inflammatory responses by UDCA and to  
28 determine the potential impact of bacterial metabolism on its therapeutic actions. The anti-  
29 inflammatory efficacy of UDCA, a non-metabolisable analogue, 6-methyl-UDCA (6-  
30 MUDCA), and its primary colonic metabolite, lithocholic acid (LCA), were assessed in the  
31 murine DSS model of mucosal injury. The effects of bile acids on cytokine release (TNF- $\alpha$ ,  
32 IL-6, IL-1 $\beta$ , IFN- $\gamma$ ) from cultured colonic epithelial cells and mouse colonic tissue *in vivo*  
33 were investigated. Luminal bile acids were measured by GC-MS. UDCA attenuated release  
34 of proinflammatory cytokines from colonic epithelial cells *in vitro* and was protective against  
35 the development of colonic inflammation *in vivo*. In contrast, although 6-MUDCA mimicked  
36 the effects of UDCA on epithelial cytokine release *in vitro*, it was ineffective in preventing  
37 inflammation in the DSS model. In UDCA-treated mice, LCA became the most common  
38 colonic bile acid. Finally, LCA treatment more potently inhibited epithelial cytokine release  
39 and protected against DSS-induced mucosal inflammation than did UDCA. These studies  
40 identify a new role for the primary metabolite of UDCA, LCA, in preventing colonic  
41 inflammation and suggest that microbial metabolism of UDCA is necessary for the full  
42 expression of its protective actions.

43

## 44 NEW AND NOTEWORTHY

45 Based on its cytoprotective and anti-inflammatory actions, the secondary bile acid,  
46 ursodeoxycholic acid (UDCA), has well-established uses in both traditional and Western  
47 medicine. Here, we identify a new role for the primary metabolite of UDCA, lithocholic acid,  
48 as a potent inhibitor of intestinal inflammatory responses and we present data to suggest that  
49 microbial metabolism of UDCA is necessary for the full expression of its protective effects  
50 against colonic inflammation.

51

52 **Keywords:** Bile acid; Epithelium; Inflammatory Bowel Disease; Cytokine, Barrier Function

## 53 INTRODUCTION

54 Inflammatory bowel diseases, such as ulcerative colitis (UC) and Crohn's disease (CD), are  
55 chronic, relapsing inflammatory disorders of the gastrointestinal tract affecting approximately  
56 1% of the adult population of Western countries. While the pathogenesis of inflammation  
57 associated with IBD is still not well-defined, it is widely accepted that a combination of  
58 genetic, environmental, and immunological factors are involved, which drive an  
59 inappropriate mucosal inflammatory response (17). With this in mind, current therapeutic  
60 options employ anti-inflammatory drugs, including glucocorticoids, immunosuppressants,  
61 aminosalicylates, and biologics to inhibit mucosal immune responses and production of  
62 proinflammatory cytokines (6). While each of these treatment approaches can be of benefit,  
63 they also have significant drawbacks in terms of the occurrence of side effects, lack of  
64 efficacy, and high cost (42). Thus, more effective, and safer, drugs to treat colitis are much  
65 needed.

66 Epithelial cells lining the colonic lumen play a key role in IBD pathogenesis (28, 36). One of  
67 the primary physiological roles of the epithelium is to act as an innate barrier against the  
68 uptake of luminal toxins and pathogens. There are several components to this barrier,  
69 including the physical barrier posed by the epithelium itself, along with numerous secreted  
70 factors, such as mucus and cytokines. A hallmark feature of IBD is dysregulation of epithelial  
71 barrier function with associated increases in permeability and induction of cytokine release  
72 (2, 30). Many endogenous and exogenous components of the luminal contents have been  
73 shown to have the capacity to promote epithelial cytokine release, including bacterial toxins  
74 and cell wall components, viral RNA, and bile acids, all of which are altered in the setting of  
75 gut inflammation (8, 26, 27). Thus, given its central role in the development of colitis, the  
76 epithelium is currently receiving a great deal of interest as a target for the development of  
77 new treatments (28, 42).

78 Ursodeoxycholic acid (UDCA) is a naturally-occurring secondary bile acid, produced in the  
79 colon by bacterial metabolism of the primary bile acid chenodeoxycholic acid (CDCA).  
80 UDCA is considered to be unique among bile acids as it has long been recognized to have  
81 broad-ranging protective actions. Indeed, UDCA is often referred to as the “therapeutic” bile  
82 acid as it has been used for centuries in Traditional Chinese Medicine, as a component of  
83 bear bile, to treat diverse maladies, such as failing eyesight, intestinal malaise, impotency,  
84 and fever (10). More recently, in Western medicine, UDCA has been used to treat liver  
85 inflammation and cholestasis (24, 47), and currently it is also under investigation for a  
86 number of conditions, including neurological, ocular, cardiovascular, and metabolic disorders  
87 (45). Importantly, unless it is used at high doses (9), UDCA is a safe drug with few side  
88 effects. While its mechanisms of action are not well-defined, it is believed that the therapeutic  
89 properties of UDCA are largely due to its anti-inflammatory and cytoprotective actions (5,  
90 45). The biological actions of UDCA have been mostly studied in the liver, where it has been  
91 shown to exert immunomodulatory and anti-apoptotic actions, and to prevent cytokine release  
92 (7, 33, 34, 37). In the current study, we hypothesised that by virtue of its anti-inflammatory  
93 and cytoprotective properties, UDCA is a represents a promising target for development of  
94 new treatments for diseases associated with intestinal inflammation. However, when  
95 considering UDCA as a potential therapeutic for intestinal disease, it is also important to  
96 consider that *in vivo*, it is extensively metabolised by the colonic microbiome and the effects  
97 that this has on its therapeutic activity are not known. Thus, in the current study we used *in*  
98 *vitro* and *in vivo* models to investigate the anti-inflammatory effects of UDCA in the colon  
99 and the potential consequences of bacterial metabolism on its therapeutic actions.

100

101

102 **MATERIALS AND METHODS:**

103 **Ethical Approval:** All experiments carried out on mice conformed to the Animal Research:  
104 Reporting of *In Vivo* Experiments (ARRIVE) guidelines and were approved by the RCSI  
105 Research Ethics Committee (REC739) and by the Irish Department of Health and Children  
106 (B100/4159).

107 **Animal Studies:** All experiments carried out on mice conformed to the Animal Research:  
108 Reporting of *In Vivo* Experiments (ARRIVE) guidelines and were approved by the RCSI  
109 Research Ethics Committee (REC739) and by the Irish Department of Health and Children  
110 (B100/4159). Male C57Bl/6 mice were used between 10 – 12 weeks of age. Colitis was  
111 induced in mice by addition of 2.5 % DSS (MP Biomedicals, Solon, OH) to their drinking  
112 water for 5 days. Disease activity index (DAI) was used as a measure of disease progression  
113 and was calculated by the addition of scores designated to body weight, faecal blood and  
114 stool consistency/diarrhoea, as previously described (39). Starting 24 hrs before  
115 administration of DSS, and once daily thereafter, animals received by intraperitoneal  
116 injection, either endotoxin-free PBS as vehicle control, Na<sup>+</sup>-UDCA (30 or 100 mg/kg), Na<sup>+</sup>-  
117 6-MUDCA or Na<sup>+</sup>-LCA (30 mg/kg) dissolved in PBS. Mice were sacrificed on day 6, the  
118 length of their colons was recorded, caecal contents were kept for analysis, and colonic tissue  
119 was processed for H&E staining, or for analysis of cytokine expression. For histological  
120 scoring, approximately 1 cm sections of colonic tissue were fixed in 10% paraformaldehyde  
121 (pH 7.4; PBS buffered) and embedded in paraffin. Sections (4 µm) were cut and stained with  
122 H&E. All sections were examined in a blinded fashion independently by 2 observers and  
123 histologic scoring was carried out, as previously described (39). Blood was collected at time  
124 of sacrifice by cardiac puncture. Serum was obtained by centrifugation (2,000 x g for 10  
125 minutes, 4°C), aliquoted, and stored at -80°C until use. Serum creatinine and ALT were

126 measured using the RXL Dimension Autoanalyser platform (Siemens Healthcare  
127 Diagnostics, Munich, Germany).

128 **Cytokine Measurements:** T<sub>84</sub> or HT29C119A cells were cultured on 96-well plates until they  
129 reached approximately 80% confluence. Cells were serum-starved for 1hr prior to stimulation  
130 with polyinosinic:polycytidylic acid (poly I:C) (25 µg/ml) or TNF-α (10 ng/ml) in the  
131 presence or absence of UDCA or LCA (24 hr, 37°C). Mouse colons were homogenised in  
132 liquid N<sub>2</sub> on dry ice, re-suspended by vortexing in lysis buffer (1% Nonidet P-40, 150 m  
133 NaCl, 50 mmol/L Tris Base, 1 x Complete mini EDTA free protease inhibitor tablet, 0.1  
134 mg/1mL PMSF, 1 mmol/L Na<sub>3</sub>VO<sub>4</sub>) in a m/v ratio of 1:5, lysed (45 minutes on ice),  
135 sonicated (3 x 10s pulses), centrifuged (15,294 x g, 20 mins, 4°C) and supernatants were  
136 retained for analysis. For measurements of TNF-α, IL-1β, IL-6, IFN-γ, IL-12p70, and GM-  
137 CSF cell culture supernatants or colonic lysates were then added to a pre-coated V-Plex  
138 Multi-array and Multi-sport Human Cytokine Assay plates (Catalogue #: K15007B-1) and  
139 assayed as per the manufacturer's protocol (Meso Scale Diagnostics; Rockville, MD).  
140 Measurements of IL-8 release from T<sub>84</sub> cells were carried out by ELISA (Beckton Dickinson,  
141 San Diego, CA).

142 **Caecal bile acid analysis:** Caecal contents were collected from treated and control animals  
143 and stored in isopropanol at -20°C. Caecal bile acid levels were measured by HPLC-ES-  
144 MS/MS, as previously described (38).

145 **Acid Phosphatase Assay:** T<sub>84</sub> cells grown to confluency on 96-well plates were serum-  
146 starved for 1hr prior to treatment with LCA. Cells were then washed in warm PBS, incubated  
147 in sodium acetate buffer (0.1M C<sub>2</sub>H<sub>3</sub>NaO<sub>2</sub>, pH 5.5, 0.1% Triton x-100) protected from light  
148 at 37°C for 30 mins, following which absorbance was recorded at 404 nm.

149 **Statistical Analysis:** Results are expressed as mean  $\pm$  SEM for a series of  $n$  experiments.  
150 Data were assumed to be normally distributed and statistical analyses were carried out using  
151 GraphPad Instat software (GraphPad, San Diego, CA). Paired t-test were used for  
152 comparisons of paired treatments between 2 groups, unpaired t-tests for comparisons of  
153 unpaired treatments between 2 groups, and one way ANOVA using Tukey multiple  
154 comparisons test for treatments of 3 groups or more.  $p$  values  $\leq 0.05$  were considered to be  
155 significant.

156

## 157 **RESULTS**

158 **UDCA inhibits pro-inflammatory cytokine release from colonic epithelial cells:** First, we  
159 investigated the effects of UDCA on release of pro-inflammatory cytokines from T<sub>84</sub> colonic  
160 epithelial cells. For these studies, we used the TLR-3 agonist, poly I:C (25  $\mu$ g/ml), as a  
161 stimulus and cytokines released into the bathing media were analysed using validated  
162 multiplex arrays. We found that Poly I:C induced secretion of TNF- $\alpha$  from T<sub>84</sub> cells and that  
163 UDCA significantly attenuated this response in a concentration-dependent manner, with a  
164 maximal effect occurring at 200  $\mu$ M (Figure 1A). UDCA (200  $\mu$ M) also attenuated Poly I:C-  
165 induced secretion of IL-1 $\beta$ , and IL-6 (Figures 1B and C). In contrast, UDCA did not alter  
166 Poly I:C-stimulated IFN- $\gamma$  release (Figure 1D), or that of IL-12p70 and GM-CSF (data not  
167 shown).

168 **UDCA exerts protective effects in the DSS model of mucosal inflammation:** Next, we  
169 went on to examine the effects of UDCA in the DSS mouse model of mucosal inflammation.  
170 The DSS model is considered to be a particularly good model for studying mucosal  
171 inflammation occurring as a consequence of disrupted epithelial barrier function (31, 48).  
172 Inclusion of 2.5% DSS in the drinking water of C57/BL6 mice led to a reduction in body

173 weight and increased DAI over the 5 day experimental period. Both effects were significantly  
174 attenuated by daily treatment with UDCA (30 mg/kg) (Figures 2A-B). UDCA at a higher  
175 dose of 100 mg/kg (Day 5 DAI =  $5.8 \pm 0.5$ ) did not confer additional protection when  
176 compared to its effects at 30 mg/kg (Day 5 DAI =  $6.8 \pm 0.9$ ; n = 6). Mice treated with DSS  
177 also had significantly shorter colons ( $60.8 \pm 2.1$  mm) and lack of faecal pellet formation  
178 compared to controls ( $87.2 \pm 2.1$  mm, n = 6 – 12,  $p \leq 0.001$ ), whereas treatment with UDCA  
179 (30 mg/kg) prevented shortening of the colon ( $69.0 \pm 1.5$  mm, n = 6 – 12,  $p \leq 0.05$ ) and  
180 restored faecal pellet formation (Figures 2 C). Histological studies revealed that UDCA  
181 reduced inflammatory cell infiltration and prevented epithelial damage, leading to a reduction  
182 in overall inflammation score (Figure 2D-E). As shown in Figure 3, UDCA also tended to  
183 reduce levels of TNF- $\alpha$ , IL-1 $\beta$ , and IL-6, although none of these effects achieved statistical  
184 significance. Similar to its effects in T<sub>84</sub> cells, UDCA did not attenuate IFN- $\gamma$  levels and, in  
185 fact, tended to enhance DSS-induced release of this cytokine.

186 **6-MUDCA is not protective against DSS-induced colonic inflammation:** In humans,  
187 UDCA is known to be metabolised to LCA in the colon and GC-MS analysis of the caecal  
188 contents revealed that this is also the case in mice (Figure 4A). Thus, we hypothesised that  
189 bacterial metabolism of UDCA likely limits its therapeutic effects. To test this, we employed  
190 a 6-methylated derivative of UDCA, 6 $\alpha$ -methyl-UDCA (6-MUDCA), which cannot be  
191 metabolised by bacteria to LCA or other metabolites (32). We have previously shown 6-  
192 MUDCA not to be metabolised to LCA in mice, but to retain the biological activity of UDCA  
193 *in vitro* (16). Here, we confirmed that 6-MUDCA also retains the activity of UDCA in  
194 preventing poly I:C-induced TNF- $\alpha$  release from T<sub>84</sub> cell monolayers (Figure 4B). 6-  
195 MUDCA was also active in HT29C119A cells, reducing Poly I:C (25  $\mu$ g/ml)-induced TNF- $\alpha$   
196 release from  $378 \pm 108$  pg/ml in controls to  $236 \pm 59$  pg/ml (n = 3;  $p \leq 0.01$ ), indicating its  
197 effects are not cell line-specific. However, despite its capacity to prevent colonic epithelial



198 cytokine secretion *in vitro*, in contrast to UDCA, 6-MUDCA was not protective against DSS-  
199 induced mucosal inflammation *in vivo*, as assessed by DAI measurements (Figure 4C).  
200 Similarly, 6-MUDCA did not prevent weight loss or colon shortening in response to DSS  
201 treatment. Body weight was reduced to  $94.8 \pm 0.5\%$  of controls in response to DSS-treatment,  
202 compared to  $90.0 \pm 2.1\%$  in 6-MUDCA-treated mice, whereas colon length in DSS-treated  
203 mice was  $59.3 \pm 1.5$  mm compared to  $57.7 \pm 1.5$  mm in those co-treated with 6-MUDCA.  
204 This lack of efficacy of 6-MUDCA was contrary to our original hypothesis, and suggest that  
205 bacterial metabolism of UDCA is necessary for it to exert its protective effects *in vivo*.

206 **LCA inhibits pro-inflammatory cytokine release from colonic epithelial cells:** Since  
207 metabolism of UDCA appears to be required for it to exert protective actions, we went on to  
208 investigate the effects of its major colonic metabolite, LCA, in regulating colonic  
209 inflammatory responses. First, we examined LCA effects on cytokine release from colonic  
210 epithelial cells *in vitro*. T<sub>84</sub> cells were treated with poly I:C, either in the absence or presence  
211 of LCA (0.1 – 10  $\mu$ M) and TNF- $\alpha$  secretion into the bathing medium was measured.  
212 Interestingly, we found that LCA treatment was considerably more effective than UDCA,  
213 practically abolishing poly I:C-induced TNF- $\alpha$  release (Figure 5A and c.f. Figure 1A).  
214 Furthermore, the effects of LCA were not specific to TLR3 activation by Poly I:C, since the  
215 bile acid also inhibited IL-8 cytokine secretion in response to another pro-inflammatory  
216 stimulus, TNF- $\alpha$  (Figure 5B). Use of the acid phosphatase activity assay, as a direct index of  
217 the number of cells present, revealed only a slight reduction associated with this effect of the  
218 bile acid (Figure 5C). To further assess potential LCA toxicity on colonic epithelial cells, we  
219 examined its effects on transepithelial resistance (TER), a sensitive index of epithelial  
220 monolayer integrity. After 24 hrs treatment, the TER of LCA (10  $\mu$ M)-treated T<sub>84</sub> cells was  
221  $94 \pm 2.6\%$  (n = 5) of that in controls, indicating that, at concentrations which abolish cytokine  
222 secretion, LCA does not alter monolayer integrity.

223 **LCA is protective against DSS-induced colonic inflammation and cytokine release:** We  
224 next examined the effects of LCA on DSS-induced colonic inflammation *in vivo*. Daily  
225 treatment with LCA (30 mg/kg; IP) significantly increased caecal LCA levels from  $6.1 \pm 0.5$   
226 to  $15.7 \pm 3.1$   $\mu\text{M}$  in controls and from  $2.0 \pm 0.3$  to  $11.5 \pm 2.1$   $\mu\text{M}$  in DSS-treated mice ( $n = 5$ ,  
227  $p \leq 0.05$ ). We noted that treatment with LCA alone induced a significant loss of body weight  
228 by day 5 to  $89.3 \pm 1.0$  % of that before LCA treatment (Figure 6A), consequently causing a  
229 slight, non-significant, increase in DAI (Figure 6B). Interestingly, LCA almost completely  
230 prevented the onset of inflammation, as measured by DAI, which in DSS-treated animals was  
231  $11.2 \pm 0.9$  compared to  $5.2 \pm 0.6$  in LCA-treated mice ( $n = 5$ ,  $p \leq 0.001$ ) (Figure 6B). LCA  
232 alone caused a slight shortening of the colon but prevented that caused by DSS treatment and  
233 restored the appearance of normal stool pellets (Figures 6C). Furthermore, LCA completely  
234 reversed DSS-induced changes in mucosal histology and increases in inflammation score  
235 (Figure 6D-E). An analysis of the effects of LCA on levels of proinflammatory cytokines  
236 revealed that it was even more effective than UDCA in reducing mucosal levels of TNF- $\alpha$ ,  
237 IL-6, and IL-1 $\beta$  in DSS-treated mice (Figure 7). Interestingly, in contrast to UDCA,  
238 administration of LCA also inhibited Poly I:C-induced increases in IFN- $\gamma$ . Mice treated with  
239 LCA actions were not associated with any apparent signs of systemic toxicity, as determined  
240 by measurements of serum creatinine and ALT. Serum creatinine levels were  $35.7 \pm 1.2$ ,  $29.0$   
241  $\pm 2.0$  and  $31.3 \pm 1.8$  mM/L in control, DSS, and DSS + LCA-treated mice, respectively ( $n =$   
242 3), while ALT levels were determined to be  $< 6$  U/L in all treatment groups.

243

244

**245 DISCUSSION**

246 By virtue of its potent anti-inflammatory and cytoprotective properties, UDCA is recognised  
247 as a drug with great therapeutic potential (45), and our current studies add to a growing body  
248 of evidence that suggest it may also be useful in treatment of intestinal inflammation. Our  
249 studies also show that the protective effects of UDCA are likely to be due, at least in part, to  
250 inhibition of epithelial cytokine production and point to an important role for bacterial  
251 metabolism in determining its efficacy *in vivo*.

252 An early step in intestinal inflammatory responses is the production of cytokines from the  
253 epithelium in response to various luminal factors, such as bacteria and their toxins and  
254 metabolites. Viruses are also present and their importance in IBD pathogenesis has recently  
255 been highlighted (27). Viruses promote cytokine secretion through the release of double-  
256 stranded RNA which activates epithelial Toll-like receptors (TLRs), in particular TLR3 (1,  
257 11), and here we found that such responses are inhibited by UDCA treatment. These findings  
258 are particularly interesting in the context of recently published data, where the effects of the  
259 conjugated derivative of UDCA, tauro-UDCA (TUDCA) were investigated in the DSS model  
260 (21). Although, significantly higher doses were required, similar to UDCA, TUDCA  
261 prevented the development of mucosal inflammation, an effect that was closely associated  
262 with inhibition of epithelial apoptosis. Also similar to our own studies, UDCA was found to  
263 prevent colonic inflammation in TNBS-treated rats, a model of intestinal inflammation  
264 distinct to that used in the current studies (25). Thus, UDCA has the capacity to prevent both  
265 the elevated cytokine levels and increased epithelial permeability associated with intestinal  
266 inflammation, suggesting it should be of therapeutic benefit in patients with IBD.

267 However, when considering the use of UDCA for treatment of colonic disease, it important to  
268 consider the potential impact of the colonic microbiota on its actions. Bile acids entering the

269 colon undergo rapid metabolism by resident bacteria by deconjugation, dehydroxylation and  
270 epimerisation and therefore, the fate of UDCA in the colon is determined by the relative  
271 expression of bacterial hydrolases, dehydratases, and epimerases (20, 22). How UDCA  
272 administration changes the makeup of the colonic bile acid pool is not well-defined but  
273 studies in humans show that after UDCA treatment, LCA becomes the most prominent  
274 colonic bile acid (44). This is supported by our current studies which showed extensive  
275 metabolism of UDCA to LCA in the cecum of normal mice. It was also interesting to note  
276 that in DSS-treated mice, despite the fact that it prevented inflammation, levels of UDCA in  
277 the colon did not increase appreciably after administration of the bile acid, while those of  
278 LCA increased approximately 4-fold. Also notable in these studies was the effect of DSS  
279 treatment in reducing cecal levels of UDCA and LCA. These data are in line with a previous  
280 study demonstrating fecal LCA levels to be decreased in DSS-treated mice (3), and a more  
281 recent study demonstrating that levels of both UDCA and LCA are reduced in this model of  
282 colonic inflammation. Furthermore, such changes were found to be associated with  
283 significant alterations in the colonic microbiota and were partially restored by UDCA  
284 treatment (43). Further studies to more precisely determine how changes in the microbiota  
285 and related alterations in the colonic bile acid signature contribute to the onset of  
286 inflammation and how UDCA administration influences such processes warrants further  
287 investigation.

288 LCA is the most lipophilic of the secondary colonic bile acids and is classically considered  
289 to be relatively toxic, particularly in the liver (15). Increased levels of hepatic LCA, which  
290 occur in conditions of cholestasis, are thought to contribute to liver damage through induction  
291 of apoptotic cell death. Indeed, several studies have demonstrated that supraphysiological  
292 levels of LCA, cause oxidative stress, DNA damage and induce apoptosis in both hepatocytes  
293 and colonic epithelial cells (4). Thus, since UDCA is normally metabolised to LCA in the

294 colon, we hypothesised that this may be a factor that limits its therapeutic actions. To test this  
295 hypothesis we used 6-MUDCA, a non-metabolizable derivative of UDCA, which we have  
296 previously shown to not be metabolised to LCA either in mouse colon or by the human fecal  
297 microbiota (16, 32). To our surprise we found that, even though, similar to UDCA, it inhibits  
298 epithelial cytokine production *in vitro*, 6-MUDCA did not confer protection in the DSS  
299 model. These findings were contrary to our hypothesis and suggest that, rather than limiting  
300 its therapeutic actions, bacterial metabolism of UDCA is actually required for it to fully exert  
301 its protective effects.

302 While most previous studies have focussed on the cytotoxic actions of LCA at high  
303 concentrations, few have investigated whether it might also have more physiological roles to  
304 play. Interestingly, one recent study showed that administration of LCA to mice by enema  
305 can prevent colonic epithelial apoptosis, and therefore presumably promote barrier function  
306 (18). In the current studies, we found that even at concentrations as low as 10  $\mu$ M, which  
307 approximates its normal physiological range in the colon (13), LCA was even more effective  
308 than UDCA in preventing TNF- $\alpha$  release from colonic epithelial cells *in vitro*. Even more  
309 remarkably, we found that when administered to mice, LCA was also more effective than  
310 UDCA in preventing DSS-induced inflammation. Further analysis showed that cytokine  
311 release from mucosal tissues was practically abolished in LCA-treated mice, compared to the  
312 partial inhibition observed with UDCA treatment. Notably, while UDCA tended to increase  
313 mucosal levels of IFN- $\gamma$  in DSS-treated mice, LCA inhibited accumulation of this cytokine.  
314 While we were concerned that the effects of LCA might be due to toxicity, this does not  
315 appear to be the case, as indicated by a lack of effect of the bile acid on TER across epithelial  
316 monolayers and only a modest effect on cell number at concentrations that abolish cytokine  
317 release. Furthermore, no overt toxicity was apparent in histological sections of colonic tissue  
318 from LCA-treated mice, nor were serum levels of creatinine or ALT altered by the bile acid.

319 However, it was notable that LCA treatment significantly reduced body weight over the  
320 course of the experiment. Given the lack of apparent local or systemic toxicity, we speculate  
321 that this could either be due to reduced food intake in the LCA-treated mice, or alternatively,  
322 might reflect effects of the bile acid on energy expenditure and fat metabolism. This latter  
323 hypothesis seems possible since previous studies have shown that bile acids prevent weight  
324 gain in mice on a high fat diet (46), and that this effect is mimicked by the TGR5-selective  
325 agonist, INT-777 (19, 41). TGR5 is now accepted to play an important role in regulating  
326 metabolism (23), suggesting that LCA, as a natural agonist of the receptor, could be an  
327 endogenous regulator of metabolism, energy expenditure and body weight. Separating such  
328 dual actions on metabolism and inflammation is an important issue to consider when  
329 developing bile acids, or synthetic agonists, as therapeutics for IBD. However, it is notable  
330 that studies by Harach and co-workers indicate that agonists of TGR5 influence metabolism  
331 only when they are present in the systemic circulation, suggesting that colonic or rectal  
332 delivery of such drugs may be the optimal approach for their use in treating colitis, while  
333 minimising effects on weight (14).

334 Although UDCA shows excellent potential for therapeutic development in treating intestinal  
335 inflammation, there is still much work to be done to elucidate mechanisms underlying its  
336 effects. While our current studies suggest that its metabolism to LCA may be important, it is  
337 also possible that other metabolites may be involved. For example, 7-keto-LCA, formed by  
338 the action of 7 $\beta$ -hydroxysteroid dehydrogenase, is the major metabolic intermediate of  
339 UDCA and LCA and its actions on colonic epithelial physiology are not yet known.  
340 Similarly, how sulfation of UDCA and LCA alter their physiological/pathophysiological  
341 actions remains to be determined. It is also important to develop our understanding of the role  
342 of the microbiota in modulating bile acid actions on colonic epithelial barrier function. This is  
343 particularly important in the setting of inflammation, where the microbiome is known to be

344 significantly altered (8). Such alterations would undoubtedly influence metabolism of UDCA,  
345 the generation of its metabolites, and consequently, its therapeutic actions. Finally, the  
346 molecular pathways underlying the anti-inflammatory effects of UDCA and its metabolites  
347 and their differential effects on epithelial cytokine secretion remain to be fully elucidated. In  
348 this regard, several bile acid receptors are expressed in the colonic epithelium, including  
349 TGR5 and the nuclear receptors, farnesoid x receptor, pregnane x receptor, and vitamin D  
350 receptor, each of which has been shown to protect against colonic inflammation in animal  
351 models (12, 29, 35, 40). Although structurally similar, UDCA and LCA have very different  
352 actions at these receptors, likely underlying different responses to the bile acids. Future work  
353 should aim to elucidate how expression of these receptors is altered in conditions of colonic  
354 inflammation and how this impacts the effects of UDCA and its metabolites on epithelial  
355 function.

356 In conclusion, our studies support the hypothesis that UDCA may be useful as a new therapy  
357 for alleviating or preventing chronic intestinal inflammation but that bacterial metabolism of  
358 the bile acid is necessary for its full therapeutic benefit to be apparent. We also demonstrate a  
359 new anti-inflammatory role for the primary UDCA metabolite, LCA, in the colon, which  
360 suggests it may be an important mediator of UDCA effects. Further studies are necessary to  
361 more completely understand how the colonic microbiome and bile acids interact in order to  
362 regulate epithelial barrier function in health and disease.

363

364 **FIGURE LEGENDS**

365 **Figure 1. UDCA attenuates proinflammatory cytokine release from colonic epithelial**  
366 **cells.** T<sub>84</sub> cells grown on 96-well plates were serum-starved for 1hr prior to stimulation with  
367 poly I:C (25 µg/ml) in the presence or absence of UDCA. After 24 hrs, supernatants were  
368 collected and analysed for **A)** TNF-α (n = 6), **B)** IL-6, **C)** IL-1β, and **D)** IFN-γ (n = 4). \*p <  
369 0.05, \*\*p < 0.01, \*\*\*p < 0.001 compared to control cells; #p < 0.05, ##p < 0.01 compared to  
370 cells treated with poly I:C alone.

371 **Figure 2. UDCA exerts protective effects in the DSS model of mucosal inflammation.**  
372 Starting 24 hrs prior to administration of DSS (2.5% in the drinking water), and daily  
373 thereafter, separate groups of male C57BL6 mice received either endotoxin-free PBS or Na<sup>+</sup>-  
374 UDCA (30 mg/kg or 100 mg/kg, dissolved in PBS) by IP injection. **A)** Disease activity index  
375 (DAI) and **B)** body weight were assessed daily to monitor disease progression (n = 6 - 12  
376 throughout). **C)** Mice were sacrificed on day 6 and their colons were removed and measured.  
377 **D)** Sections of colon from control, DSS-treated, UDCA-treated, and DSS+UDCA-treated  
378 C57BL6 mice were taken and processed for H&E staining. Sections were visualised by light  
379 microscopy under 10x magnification. **E)** Inflammation score was assessed as described in  
380 Materials and Methods. \*\*\* p < 0.001 compared to controls (no DSS treatment); # p < 0.05,  
381 ## p < 0.01, ### p < 0.001 compared to DSS-treated mice.

382 **Figure 3. UDCA modulates expression of pro-inflammatory cytokines in the DSS model**  
383 **of mucosal inflammation.** Sections of colon from control, DSS-treated, UDCA-treated, and  
384 DSS+UDCA-treated C57BL6 mice were homogenised in lysis buffer and were analysed by  
385 MSD assay for **A)** TNF-α, **B)** IL-6, **C)** IL-1β, and **D)** IFN-γ. n = 6 – 12; \*\*p < 0.01, \*\*\*p <  
386 0.001 compared to controls (no DSS treatment). n.s. = not significant.



387 **Figure 4. A metabolically stable analogue of UDCA, 6-MUDCA, is not protective**  
388 **against DSS-induced colonic inflammation:** **A)** Caecal contents were collected from treated  
389 and control mice and bile acid levels were measured by HPLC-ES-MS/MS. **B)** T<sub>84</sub> cells were  
390 stimulated with poly I:C (25 µg/ml) in the presence or absence of UDCA or 6-methyl-UDCA  
391 (200 µM; bilateral). After 24 hrs, supernatants were collected and analysed for TNF-α. Data  
392 are expressed as fold change with respect to cells treated with poly I:C alone (n = 5; \*\*\*p <  
393 0.001). **C)** Starting 24hrs prior to administration of DSS (2.5% in the drinking water), and  
394 daily thereafter, separate groups of male C57BL6 mice received either endotoxin-free PBS or  
395 Na<sup>+</sup>-6-MUDCA (30 mg/kg) by IP injection. DAI was assessed daily to monitor disease  
396 progression. (n = 3 - 9).

397 **Figure 5. LCA exerts anti-inflammatory effects *in vitro*.** **A)** T<sub>84</sub> cells were stimulated with  
398 poly I:C (25 µg/ml) in the presence or absence of LCA (1 nM - 10 µM). After 24 hrs,  
399 supernatants were collected and analysed for TNF-α. Data are expressed as fold change with  
400 respect to cells treated with poly I:C alone (n = 7; \*\*\*p < 0.001). **B)** T<sub>84</sub> cells were treated  
401 with TNF-α (10 ng/ml) and LCA (10 µM) alone or in combination. After 24 hrs apical media  
402 were collected and analysed for IL-8 levels by ELISA. Data are expressed as fold change  
403 with respect to cells treated with TNF-α alone (n = 4; \*\*\*p < 0.001). **C)** T<sub>84</sub> cells grown on  
404 96 well plates were serum starved for 1 hr prior to treatment with LCA (1 nM to 1 mM) for  
405 24 hrs (n = 4), after which acid phosphatase activity was measured (\*\*p < 0.01, \*\*\*p < 0.001  
406 compared to untreated cells).

407 **Figure 6. LCA exerts protective effects in the DSS model of mucosal inflammation.**  
408 Starting 24 hrs prior to administration of DSS in the drinking water, and daily thereafter,  
409 separate groups of male C57BL6 mice received either endotoxin-free PBS or Na<sup>+</sup>-LCA (30  
410 mg/kg) by IP injection. **A)** Body weight and **B)** disease activity index (DAI) were assessed

411 daily to monitor disease progression (n = 5). **C)** Mice were sacrificed on day 6 and their  
412 colons were removed and measured (image is representative of n = 5). **D)** Sections of colon  
413 from control, DSS-treated, LCA (30 mg/kg)-treated, and DSS+LCA-treated C57BL6 mice  
414 were taken and processed for H&E staining. Sections were visualised by light microscopy  
415 under 10x magnification. **E)** Inflammation score was assessed as described in Materials and  
416 Methods (n = 3 – 5). \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001 compared to controls (no DSS  
417 treatment); #p < 0.05, ##p < 0.01, compared to DSS-treated mice

418 **Figure 7. LCA modulates the expression of pro-inflammatory cytokines in murine**  
419 **colon.** Sections of colon from control, DSS-treated, LCA (30 mg/kg)-treated, and DSS+LCA-  
420 treated C57BL6 mice were homogenised in lysis buffer and were analysed for **A)** TNF- $\alpha$ , **B)**  
421 IL-6, **C)** IL-1 $\beta$ , and **D)** IFN- $\gamma$ . n = 3 – 10; \*\*p < 0.01, \*\*\*p < 0.001 compared to controls (no  
422 DSS treatment).

423

424

## 425 REFERENCES

- 426 1. **Alexopoulou L, Holt AC, Medzhitov R, and Flavell RA.** Recognition of double-  
427 stranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature* 413: 732-738,  
428 2001.
- 429
- 430 2. **Antoni L, Nuding S, Wehkamp J, and Stange EF.** Intestinal barrier in  
431 inflammatory bowel disease. *World J Gastroenterol* 20: 1165-1179, 2014.
- 432
- 433 3. **Araki Y, Andoh A, Tsujikawa T, Fujiyama Y, and Bamba T.** Alterations in  
434 intestinal microflora, faecal bile acids and short chain fatty acids in dextran sulphate sodium-  
435 induced experimental acute colitis in rats. *Eur J Gastroenterol Hepatol* 13: 107-112, 2001.
- 436
- 437 4. **Barrasa JI, Olmo N, Lizarbe MA, and Turnay J.** Bile acids in the colon, from  
438 healthy to cytotoxic molecules. *Toxicol In Vitro* 27: 964-977, 2013.
- 439
- 440 5. **Bellentani S.** Immunomodulating and anti-apoptotic action of ursodeoxycholic acid:  
441 where are we and where should we go? *Eur J Gastroenterol Hepatol* 17: 137-140, 2005.
- 442
- 443 6. **Bezzio C, Furfaro F, de Franchis R, Maconi G, Asthana AK, and Ardizzone S.**  
444 Ulcerative colitis: current pharmacotherapy and future directions. *Expert Opin Pharmacother*  
445 15: 1659-1670, 2014.
- 446
- 447 7. **Dosa PI, Ward T, Castro RE, Rodrigues CM, and Steer CJ.** Synthesis and  
448 evaluation of water-soluble prodrugs of ursodeoxycholic acid (UDCA), an anti-apoptotic bile  
449 acid. *ChemMedChem* 8: 1002-1011, 2013.
- 450
- 451 8. **Duboc H, Rajca S, Rainteau D, Benarous D, Maubert MA, Quervain E, Thomas**  
452 **G, Barbu V, Humbert L, Despras G, Bridonneau C, Dumetz F, Grill JP, Masliah J,**  
453 **Beaugerie L, Cosnes J, Chazouilleres O, Poupon R, Wolf C, Mallet JM, Langella P,**  
454 **Trugnan G, Sokol H, and Seksik P.** Connecting dysbiosis, bile-acid dysmetabolism and gut  
455 inflammation in inflammatory bowel diseases. *Gut* 62: 531-539, 2013.
- 456
- 457 9. **Eaton JE, Silveira MG, Pardi DS, Sinakos E, Kowdley KV, Luketic VA,**  
458 **Harrison ME, McCashland T, Befeler AS, Harnois D, Jorgensen R, Petz J, and Lindor**  
459 **KD.** High-dose ursodeoxycholic acid is associated with the development of colorectal  
460 neoplasia in patients with ulcerative colitis and primary sclerosing cholangitis. *Am J*  
461 *Gastroenterol* 106: 1638-1645, 2011.
- 462
- 463 10. **Feng Y, Siu K, Wang N, Ng KM, Tsao SW, Nagamatsu T, and Tong Y.** Bear bile:  
464 dilemma of traditional medicinal use and animal protection. *J Ethnobiol Ethnomed* 5: 2,  
465 2009.
- 466 11. **Furrie E, Macfarlane S, Thomson G, Macfarlane GT, Microbiology, Gut Biology**  
467 **G, Tayside T, and Tumour B.** Toll-like receptors-2, -3 and -4 expression patterns on human  
468 colon and their regulation by mucosal-associated bacteria. *Immunology* 115: 565-574, 2005.
- 469
- 470 12. **Gadaleta RM, van Erpecum KJ, Oldenburg B, Willemsen EC, Renooij W,**  
471 **Murzilli S, Klomp LW, Siersema PD, Schipper ME, Danese S, Penna G, Laverny G,**  
472 **Adorini L, Moschetta A, and van Mil SW.** Farnesoid X receptor activation inhibits

- 473 inflammation and preserves the intestinal barrier in inflammatory bowel disease. *Gut* 60: 463-  
474 472, 2011.
- 475
- 476 13. **Hamilton JP, Xie G, Raufman JP, Hogan S, Griffin TL, Packard CA, Chatfield**  
477 **DA, Hagey LR, Steinbach JH, and Hofmann AF.** Human cecal bile acids: concentration  
478 and spectrum. *Am J Physiol Gastrointest Liver Physiol* 293: G256-263, 2007.
- 479
- 480 14. **Harach T, Pols TW, Nomura M, Maida A, Watanabe M, Auwerx J, and**  
481 **Schoonjans K.** TGR5 potentiates GLP-1 secretion in response to anionic exchange resins.  
482 *Sci Rep* 2: 430, 2012.
- 483
- 484 15. **Hofmann AF.** Detoxification of lithocholic acid, a toxic bile acid: relevance to drug  
485 hepatotoxicity. *Drug Metab Rev* 36: 703-722, 2004.
- 486
- 487 16. **Kelly OB, Mroz MS, Ward JB, Colliva C, Scharl M, Pellicciari R, Gilmer JF,**  
488 **Fallon PG, Hofmann AF, Roda A, Murray FE, and Keely SJ.** Ursodeoxycholic acid  
489 attenuates colonic epithelial secretory function. *J Physiol* 591: 2307-2318, 2013.
- 490
- 491 17. **Knights D, Lassen KG, and Xavier RJ.** Advances in inflammatory bowel disease  
492 pathogenesis: linking host genetics and the microbiome. *Gut* 62: 1505-1510, 2013.
- 493
- 494 18. **Kozoni V, Tsioulis G, Shiff S, and Rigas B.** The effect of lithocholic acid on  
495 proliferation and apoptosis during the early stages of colon carcinogenesis: differential effect  
496 on apoptosis in the presence of a colon carcinogen. *Carcinogenesis* 21: 999-1005, 2000.
- 497
- 498 19. **Kumar DP, Asgharpour A, Mirshahi F, Park SH, Liu S, Imai Y, Nadler JL,**  
499 **Grider JR, Murthy KS, and Sanyal AJ.** Activation of Transmembrane Bile Acid Receptor  
500 TGR5 Modulates Pancreatic Islet alpha Cells to Promote Glucose Homeostasis. *J Biol Chem*  
501 291: 6626-6640, 2016.
- 502
- 503 20. **Kuo SM, Merhige PM, and Hagey LR.** The effect of dietary prebiotics and  
504 probiotics on body weight, large intestine indices, and fecal bile acid profile in wild type and  
505 IL10<sup>-/-</sup> mice. *PLoS One* 8: e60270, 2013.
- 506
- 507 21. **Laukens D, Devisscher L, Van den Bossche L, Hindryckx P, Vandenbroucke RE,**  
508 **Vandewynckel YP, Cuvelier C, Brinkman BM, Libert C, Vandenabeele P, and De Vos**  
509 **M.** Tauroursodeoxycholic acid inhibits experimental colitis by preventing early intestinal  
510 epithelial cell death. *Lab Invest* 94: 1419-1430, 2014.
- 511
- 512 22. **Lee JY, Arai H, Nakamura Y, Fukiya S, Wada M, and Yokota A.** Contribution of  
513 the 7beta-hydroxysteroid dehydrogenase from *Ruminococcus gnavus* N53 to ursodeoxycholic  
514 acid formation in the human colon. *J Lipid Res* 54: 3062-3069, 2013.
- 515
- 516 23. **Li T, and Chiang JY.** Bile acids as metabolic regulators. *Curr Opin Gastroenterol*  
517 31: 159-165, 2015.
- 518
- 519 24. **Maillette de Buy Wenniger L, and Beuers U.** Bile salts and cholestasis. *Dig Liver*  
520 *Dis* 42: 409-418, 2010.
- 521

- 522 25. **Martinez-Moya P, Romero-Calvo I, Requena P, Hernandez-Chirlaque C,**  
523 **Aranda CJ, Gonzalez R, Zarzuelo A, Suarez MD, Martinez-Augustin O, Marin JJ, and**  
524 **de Medina FS.** Dose-dependent antiinflammatory effect of ursodeoxycholic acid in  
525 experimental colitis. *Int Immunopharmacol* 15: 372-380, 2013.  
526
- 527 26. **Merga Y, Campbell BJ, and Rhodes JM.** Mucosal barrier, bacteria and  
528 inflammatory bowel disease: possibilities for therapy. *Dig Dis* 32: 475-483, 2014.  
529
- 530 27. **Norman JM, Handley SA, Baldrige MT, Droit L, Liu CY, Keller BC, Kambal**  
531 **A, Monaco CL, Zhao G, Fleshner P, Stappenbeck TS, McGovern DP, Keshavarzian A,**  
532 **Mutlu EA, Sauk J, Gevers D, Xavier RJ, Wang D, Parkes M, and Virgin HW.** Disease-  
533 specific alterations in the enteric virome in inflammatory bowel disease. *Cell* 160: 447-460,  
534 2015.  
535
- 536 28. **Odenwald MA, and Turner JR.** Intestinal permeability defects: is it time to treat?  
537 *Clin Gastroenterol Hepatol* 11: 1075-1083, 2013.  
538
- 539 29. **Ooi JH, Li Y, Rogers CJ, and Cantorna MT.** Vitamin D regulates the gut  
540 microbiome and protects mice from dextran sodium sulfate-induced colitis. *J Nutr* 143: 1679-  
541 1686, 2013.  
542
- 543 30. **Pastorelli L, De Salvo C, Mercado JR, Vecchi M, and Pizarro TT.** Central role of  
544 the gut epithelial barrier in the pathogenesis of chronic intestinal inflammation: lessons  
545 learned from animal models and human genetics. *Front Immunol* 4: 280, 2013.  
546
- 547 31. **Perse M, and Cerar A.** Dextran sodium sulphate colitis mouse model: traps and  
548 tricks. *J Biomed Biotechnol* 2012: 718617, 2012.  
549
- 550 32. **Roda A, Pellicciari R, Cerre C, Polimeni C, Sadeghpour B, Marinozzi M, Forti**  
551 **GC, and Sapigni E.** New 6-substituted bile acids: physico-chemical and biological  
552 properties of 6 alpha-methyl ursodeoxycholic acid and 6 alpha-methyl-7-epicholic acid. *J*  
553 *Lipid Res* 35: 2268-2279, 1994.  
554
- 555 33. **Rodrigues CM, Fan G, Wong PY, Kren BT, and Steer CJ.** Ursodeoxycholic acid  
556 may inhibit deoxycholic acid-induced apoptosis by modulating mitochondrial transmembrane  
557 potential and reactive oxygen species production. *Mol Med* 4: 165-178, 1998.  
558
- 559 34. **Roma MG, Toledo FD, Boaglio AC, Basiglio CL, Crocenzi FA, and Sanchez**  
560 **Pozzi EJ.** Ursodeoxycholic acid in cholestasis: linking action mechanisms to therapeutic  
561 applications. *Clin Sci (Lond)* 121: 523-544, 2011.  
562
- 563 35. **Sakanaka T, Inoue T, Yorifuji N, Iguchi M, Fujiwara K, Narabayashi K,**  
564 **Kakimoto K, Nouda S, Okada T, Kuramoto T, Ishida K, Abe Y, Takeuchi T, Umegaki**  
565 **E, Akiba Y, Kaunitz JD, and Higuchi K.** The effects of a TGR5 agonist and a dipeptidyl  
566 peptidase IV inhibitor on dextran sulfate sodium-induced colitis in mice. *J Gastroenterol*  
567 *Hepatol* 30 Suppl 1: 60-65, 2015.  
568
- 569 36. **Salim SY, and Soderholm JD.** Importance of disrupted intestinal barrier in  
570 inflammatory bowel diseases. *Inflamm Bowel Dis* 17: 362-381, 2011.

- 571 37. **Sola S, Aranha MM, Steer CJ, and Rodrigues CM.** Game and players:  
572 mitochondrial apoptosis and the therapeutic potential of ursodeoxycholic acid. *Curr Issues*  
573 *Mol Biol* 9: 123-138, 2007.  
574
- 575 38. **Tadano T, Kanoh M, Matsumoto M, Sakamoto K, and Kamano T.** Studies of  
576 serum and feces bile acids determination by gas chromatography-mass spectrometry. *Rinsho*  
577 *Byori* 54: 103-110, 2006.  
578
- 579 39. **Tambuwala MM, Manresa MC, Cummins EP, Aversa V, Coulter IS, and Taylor**  
580 **CT.** Targeted delivery of the hydroxylase inhibitor DMOG provides enhanced efficacy with  
581 reduced systemic exposure in a murine model of colitis. *J Control Release* 217: 221-227,  
582 2015.  
583
- 584 40. **Terc J, Hansen A, Alston L, and Hirota SA.** Pregnane X receptor agonists enhance  
585 intestinal epithelial wound healing and repair of the intestinal barrier following the induction  
586 of experimental colitis. *Eur J Pharm Sci* 55: 12-19, 2014.  
587
- 588 41. **Thomas C, Gioiello A, Noriega L, Strehle A, Oury J, Rizzo G, Macchiarulo A,**  
589 **Yamamoto H, Mataki C, Pruzanski M, Pellicciari R, Auwerx J, and Schoonjans K.**  
590 TGR5-mediated bile acid sensing controls glucose homeostasis. *Cell Metab* 10: 167-177,  
591 2009.  
592
- 593 42. **Torres J, Danese S, and Colombel JF.** New therapeutic avenues in ulcerative colitis:  
594 thinking out of the box. *Gut* 62: 1642-1652, 2013.  
595
- 596 43. **Van den Bossche L, Hindryckx P, Devisscher L, Devriese S, Van Welden S,**  
597 **Holvoet T, Vilchez-Vargas R, Vital M, Pieper DH, Vanden Bussche J, Vanhaecke L,**  
598 **Van de Wiele T, De Vos M, and Laukens D.** Ursodeoxycholic acid and its taurine/glycine  
599 conjugated species reduce colitogenic dysbiosis and equally suppress experimental colitis in  
600 mice. *Appl Environ Microbiol* 2017.  
601
- 602 44. **van Gorkom BA, van der Meer R, Boersma-van Ek W, Termont DS, de Vries**  
603 **EG, and Kleibeuker JH.** Changes in bile acid composition and effect on cytolytic activity of  
604 fecal water by ursodeoxycholic acid administration: a placebo-controlled cross-over  
605 intervention trial in healthy volunteers. *Scand J Gastroenterol* 37: 965-971, 2002.  
606
- 607 45. **Vang S, Longley K, Steer CJ, and Low WC.** The Unexpected Uses of Urso- and  
608 Tauroursodeoxycholic Acid in the Treatment of Non-liver Diseases. *Glob Adv Health Med* 3:  
609 58-69, 2014.  
610
- 611 46. **Watanabe M, Houten SM, Mataki C, Christoffolete MA, Kim BW, Sato H,**  
612 **Messaddeq N, Harney JW, Ezaki O, Kodama T, Schoonjans K, Bianco AC, and Auwerx**  
613 **J.** Bile acids induce energy expenditure by promoting intracellular thyroid hormone  
614 activation. *Nature* 439: 484-489, 2006.  
615
- 616 47. **Williamson KD, and Chapman RW.** Primary sclerosing cholangitis. *Dig Dis* 32:  
617 438-445, 2014.  
618
- 619 48. **Wirtz S, Neufert C, Weigmann B, and Neurath MF.** Chemically induced mouse  
620 models of intestinal inflammation. *Nat Protoc* 2: 541-546, 2007.

621 **COMPETING INTERESTS:** The authors have no competing interests to report.


622

623 **ACKNOWLEDGEMENTS:** This work was supported by a Science Foundation Ireland  
624 Principal Investigator Award (10/IN.1/B2983) and a Crohn's and Colitis Foundation of  
625 America Senior Investigator Award to SJK. The authors would like to thank Prof. Alan F.  
626 Hofmann (University of California, San Diego) for his helpful discussion during the  
627 preparation of this manuscript.


628

Figure 1


A




B



C



D





**Figure 2**










Figure 3


A



B



C



D






Figure 4

A



B



C






Figure 5


A



B




C




**Figure 6**


**A**




**B**



**C**



**D**



**E**




Figure 7

