

# Osteoporosis International

## Greater yogurt consumption is associated with increased bone mineral density and physical function in older adults

--Manuscript Draft--

<b>Manuscript Number:</b>	OSIN-D-16-00895R1	
<b>Full Title:</b>	Greater yogurt consumption is associated with increased bone mineral density and physical function in older adults	
<b>Article Type:</b>	Original Article	
<b>Funding Information:</b>	Irish Department of Agriculture, Food and the Marine (07FHRIUCD1)	Not applicable
	Irish Department of Agriculture, Food and the Marine (13F407)	Not applicable
	Northern Ireland Department for Employment and Learning (DEL)	Not applicable
	National Dairy Council Ireland	Not applicable
<b>Abstract:</b>	<p><b>Summary</b> In this cohort of community dwelling older adults (&gt;60 yrs), we observed significant positive associations between the frequency of yogurt intake and bone health.</p> <p><b>Introduction</b> The associations of yogurt intakes with bone health and frailty in older adults are not well documented. The aim was to investigate the association of yogurt intakes with bone mineral density [BMD], bone biomarkers and physical function in 4,310 Irish adults from the Trinity, Ulster, Department of Agriculture aging cohort study (TUDA).</p> <p><b>Methods</b> Bone measures included total hip, femoral neck and vertebral BMD with bone biochemical markers. Physical function measures included Timed Up and Go (TUG), Instrumental Activities of Daily Living Scale and Physical Self-Maintenance Scale.</p> <p><b>Results</b> Total hip and femoral neck BMD in females were 3.1 - 3.9 % higher among those with the highest yogurt intakes (n= 970) compared to the lowest (n= 1,109; P &lt;0.05) as were the TUG scores (-6.7%; P = 0.020). In males, tartrate-resistant acid phosphatase (TRAP 5b) concentrations were significantly lower in those with the highest yogurt intakes (-9.5%; P &lt;0.0001). In females, yogurt intake was a significant positive predictor of BMD at all regions. Each unit increase in yogurt intake in females was associated with a 29% lower risk of osteopenia (OR 0.71; 95% CI 0.51 - 1.01; P=0.037) and a 37% lower risk of osteoporosis (OR 0.63; 95% CI 0.44 - 0.91; P=0.014) and in males, a 51% lower risk of osteoporosis (OR 0.49; 95% CI 0.25 - 0.94; P=0.032).</p> <p><b>Conclusion</b> In this cohort, higher yogurt intake was associated with increased BMD and physical function scores. These results suggest that improving yogurt intakes could be a valuable and cost-effective health strategy for maintaining bone health in older adults.</p>	
<b>Corresponding Author:</b>	Eamon Laird, Ph.D. University of Dublin Trinity College IRELAND	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	University of Dublin Trinity College	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Eamon Laird, Ph.D.	
<b>First Author Secondary Information:</b>		
<b>Order of Authors:</b>	Eamon Laird, Ph.D.	
	Anne Molloy	

	Mary Ward
	Kevin McCarroll
	Leane Hoey
	Catherine F Hughes
	Conal Cunningham
	JJ Strain
	Helene McNulty
	Miriam Casey
<b>Order of Authors Secondary Information:</b>	
<b>Author Comments:</b>	<p>Drs. J. Kanis; F. Cosman  Editors-in Chief Osteoporosis International</p> <p>27th August 2016</p> <p>Dear Co-Editors,</p> <p>We would be very grateful if you would consider our manuscript "Greater yogurt consumption is associated with increased bone mineral density (BMD) and physical function in older adults" for publication in Osteoporosis International</p> <p>Osteoporosis is an increasingly common, chronic condition estimated to affect over 200 million individuals worldwide. Maintaining an optimal nutritional status is a key preventative measure, particularly for older adults (&gt;60 years). Of the major food groups, dairy foods (including milk, yogurt and cheese) are one of the richest sources of the macro and micro nutrients that contribute to bone health. There is however, little information on the associations of yogurt intake, with BMD and bone health bio-markers in addition to concomitant measurements of physical function in this older population.</p> <p>In this study we investigate the associations of daily dairy intakes (in particular yogurt) and the associations with BMD, bone biomarkers and measures of physical function in 4,310 older adults (60-102 years). The participants were originally recruited to the Trinity Ulster Department of Agriculture (TUDA) aging cohort study, a large study of older Community dwelling Irish adults designed to investigate nutritional factors and gene-nutrient interactions in the development of chronic diseases of aging. Data for the current study relate to &gt;4,000 participants, making it the largest study to-date examining yogurt intakes and associations with bone health and frailty. Bone biomarkers included serum osteocalcin, Bone specific alkaline phosphatase, C-terminal telopeptides of type I collagen and Tartrate-resistant acid phosphatase (TRAP 5b). BMD was measured at the total hip, femoral neck and vertebral spine. Physical function measures included Timed get Up and Go and the Instrumental Activities of Daily living scale. Total hip and femoral BMD in women were 3.1 - 3.9 % higher amongst those with the highest yogurt intakes, as were the TUG scores (-6.7%). TRAP 5b concentrations were significantly lower in men with the highest yogurt intakes (-9.5%). In a regression model examining predictors of osteoporosis, each unit increase in daily yogurt intake was associated with a 29% lower risk of osteopenia and a 37% lower risk of osteoporosis in women ( what about osteopenia in men) and a 51% lower risk of osteoporosis in men</p> <p>Our findings provide evidence that an increase in the frequency of yogurt intake is positively associated bone health status and measurements of physical function in a cohort of older adults. We believe the findings of the current study will contribute significantly to the growing evidence supporting a beneficial role for yogurt in health.</p> <p>Thank you for taking the time to consider this manuscript.</p> <p>Yours sincerely,</p> <p>Dr Eamon Laird</p>

Research Fellow, Trinity College Dublin, Ireland

**Response to Reviewers:**

Reviewer 1

This is a well written observational cross-sectional study carried out in a large cohort of community dwelling older (>60 years) females and male subjects.

Response: We thank reviewer 1 for their helpful comments regarding the manuscript. The clarifications suggested by the reviewer are welcomed and the specific replies to each comment are shown below.

1. My main concern is the excess of reliance on statistical analysis without considering the pathophysiological relevance of the significance as computed by using various tests including hierarchical regression models with adjustment for about 15 variables. For instance, what is the biological meaning of reporting that an increase in serum 25OHD is associated with greater (Odd ratio: 1.0, 95%CI 1.00-1.02 !) risk of osteoporosis with a probability level of <0.0001 (Table 5) ? This kind of significance test amalgamates precision with effect size, thus muddling two essential aspects of data interpretation. A weak association between an extremely mild increase in serum 25OHD and the risk of osteoporosis is incorrectly interpreted as "important" because it is statistically significant. In order to control Type I error rate (i.e. to conclude there is a difference though none exists) when multiple hypothesis are tested, a correction has to be applied. Among several statistical correction tools, the Bonferroni correction is often used, making it more difficult to reject the null hypothesis.

Response: We agree with the reviewer that this is an unexpected result and that we should have mentioned this in the text (we did not discuss this in the results or discussion). We feel this unexpected result could be due to the fact we did not include those on vitamin D supplements as a variable in the multinomial regression model. We have now included this for both women and men (Tables 5 and 6). Serum 25(OH)D is no longer a significant predictor of bone health for women or men. However, yes to vitamin D supplements is now a significant predictor of bone health. For women, those on vitamin D supplements had a significantly reduced risk of osteopenia (OR 0.51; 95% CI 0.34 – 0.76; P=0.001) and a significantly reduced risk of osteoporosis (OR 0.41; 95% CI 0.26 – 0.64; P<0.0001). For men, those on vitamin D supplements also had a significantly reduced risk of osteoporosis (OR 0.40; 95% CI 0.22 – 0.72; P=0.003). These results have been added both to Tables 5 and 6. With the addition of the vitamin D supplement variable to the model, the positive effect of daily yogurt intakes on reducing the risk of impaired bone health has also increased (lines 246-248).

In regards to the Bonferroni correction, the authors feel that it is statistically inappropriate for the multinomial regression model as each variable included is controlling for the other. However, for the other analysis (Tables 2-3 and Supplemental Tables 2-3) we have used the Bonferroni correction. We apologise that this was not made clear in the statistical methods and we have now added a sentence to this effect and a footnote to each of the appropriate Tables. It is important to note however, that in doing the Bonferroni correction, the correction directly targets the Type 1 error problem, but it does so at the expense of Type 2 error. By changing the p value needed to reject the null (or equivalently widening the uncertainty intervals) the number of claims of rejected null hypotheses will indeed decrease on average. Although this reduces the number of false rejections, it also increases the number of instances that the null is not rejected when in fact it should have been. Thus, the Bonferroni correction can severely reduce our power to detect an important effect (Gelman et al 2012).

2. According to DXA-BMD measurements (presumably by using either Total Hip or Femoral Neck BMD? This referenced skeletal site has to be clearly indicated), odd ratios were calculated for 15 variables in each gender (Tables 5 and 6). In females, 3 and 5 variables were found to be significantly associated with the risk of osteopenia (+Age, -BMI, -Daily Yogurt Serving) and osteoporosis (+Age, -BMI, +25OHD, -Daily Yogurt Serving), respectively. In males, 3 variables were found to be significantly associated with the risk of osteopenia (-BMI, +Daily Cheese Serving, -Daily Meat Serving) and osteoporosis (-BMI, -Daily Yogurt Serving, +Daily Cheese), respectively. Some of these associations appear to be significant statistically, but without any biological significance. Thus, in females mean serum 25OHD varies very little among the 3 groups categorized according to the frequency of yogurt intakes (Table 2). Similarly minimal variations of serum 25OHD were recorded in females distributed

according to the frequency of milk (Supplemental Table 2) and cheese (Supplemental Table 4) intakes.

Response: We have now inserted a sentence explaining the referenced skeletal sites for Tables 5 and 6 as a footnote on each. In response to the results of the multinomial regression model, some of the results that were observed were expected and have been reported extensively in the literature (e.g. increasing age, and lower BMI associated with increased risk of osteopenia/osteoporosis). We feel it is interesting some of the dietary components were also significant as few studies often examine these as predictors of bone health and thus little is known on the effects of diet on bone health. As stated in response to query 1, the multinomial regression model is a robust statistical model with each variable acting as a control/adjustment for another. In regards to Table 2, supplemental Tables 2 and 4 it is actually a positive that 25(OH)D concentrations did not differ between the groups as it can then be ruled out as a potential likely driver of any effects on bone health that we observed.

3. To determine the predictors of bone fragility (osteopenia and osteoporosis) a multinomial logistic regression model was used (with normal bone health as the reference category) with relevant co-predictors including serum 25OHD as a nominal variable (P.8, lines 184-190). As presented in Table 5, in females the risk of osteoporosis would be positively increased by 1.01 (95% CI 1.00-1.02,  $P < 0.0001$ ) by an augmentation of serum 25OHD. This is an example of highly statistical significance without any pathophysiological significance. This kind of statistic test mixes effect size and precision, thus muddling two essential aspects of data interpretation. Confidence intervals provide both an estimate of the effect size and the precision of the measurement.

Response: We have amended these results as explained in the response to queries 1, 2 and 6.

4. P.7, lines 168-169. References should be given for the functionality measures by PSM and IADL.

Response: A reference has now been inserted (line 176).

5. P.8, line 183-184. BMD unit is in gram of hydroxyapatite equivalent divided by the DXA-scanned surface ( $\text{g}/\text{cm}^2$ ) of the region of interest. Therefore it does not correspond to a "concentration". This wrong unit designation should be corrected.

Response: We have now removed any mention of 'concentration' to BMD measures or results in the manuscript.

6. Table 5. Females statistical analysis. Increasing age significantly augments the Odd ratio (augments the risk) for both osteopenia (1.03 |1.00-1.06|  $P = 0.043$ ) and osteoporosis (1.04 |1.01-1.08|  $P = 0.008$ ) (Table 5). Increasing BMI (0.92 |1.00-1.06|  $P < 0.0001$ ). Physical activity and Yogurt consumption reduces the Odd ratio (reduces the risk) of osteoporosis (Table 5). These findings are consistent with previous knowledge on the risk factors of bone fragility in later age. A discordant finding concerns bone fragility (osteopenia, osteoporosis) related to the vitamin D status. It is reported that an increase in serum 25OHD rises the risk of osteoporosis (Table 5). This is inconsistent with most published studies providing evidence that increasing vitamin D status, as assessed by measuring the serum level of 25OHD, reduces bone turnover in menopausal women and is associated with a lower risk of fragility fractures with aging.

Response: We have agree with the reviewer and have amended these results as explained in the response to queries 1, 2 and 3.

7. -Table 6. Males statistical analysis. Increased BMI significantly reduces the Odd ratio for osteoporosis. Increase in both yogurt and meat consumption also reduces the risk of osteopenia and osteoporosis. These findings corroborate several published results. In contrast, in this epidemiological study an increased consumption of cheese would augment the risk of developing either osteopenia or osteoporosis. The authors recognize that this statistical finding is in contradiction with recent reports indicating that an increase of cheese consumption exerts a rather bone-protective effect. They

mention the hypothesis that a high sodium content of certain varieties of cheese would increase the urinary excretion of calcium and, thereby, would negatively influence bone mineral balance. However, the sodium-induced calciuria hypothesis is not supported by any long term observations. To mention an untenable mechanistic hypothesis to account for a statistically unexpected and contradictable result lowers the scientific quality of the discussion.

Response: We acknowledge the reviewers comments the sodium-induced calciuria hypothesis is not supported by any long term observations (e.g. the recent Women's Health initiative Study) and we have mentioned this in the text. However we also recognize that this could offer a potential hypothesis to explain our observed results and given that it has not been extensively examined in the literature, in particular for the effects on men, we feel we must mention it as a possibility.

8. P.8, lines 195 - P9, line 196. The sentence "The majority of participants were females, who were significantly older, lighter, and contained a higher proportion of individuals receiving bone, vitamin D or calcium supplements in comparison with males." is not clear. The word "bone" should be deleted.

Response: We have deleted this from the sentence.

9. P.9, lines 220 – P10, lines 221-223. The differences highlighted in terms of milk consumption, for Time up and Go, IADL and PSM (Supplemental Table 2 and 3, not Suppl.Tables 1 and 2) are, though statistically significant, of trivial biological significance. A more stringent statistical analysis using a multiple-comparison procedure (e.g. Bonferroni's method) that increases the critical F or t is needed for declaring the comparison to be significant.

Response: As stated in response to query 1, these results are adjusted for Bonferroni. We again apologize for not making this clear in the manuscript and have added this as a footnote to the Tables of interest.

10. P.10, line 224. Please substitute Supplemental Table 3 and 4 by Supplemental Table 4 and 5.

Response: We have now switched these Supplemental Tables.

11. -Table 2 and Table 3. Reference range of the listed biochemical markers (CTX, OC, BAP; TRAP 5b, 25OHD and PTH) should be indicated in the legend to these two tables.

Response: We have now added these reference ranges in the methods section and in the legends to Table 2 and 3.

12. P.10, lines 244-245 and P.11, lines 246-247. In men, Table 6 does not corroborate the statement that in men increased yogurt consumption was associated with a significant decrease in the Odd ratio of osteopenia (Table 6, 0.85, 95% CI 0.58-1.24, P =0.410).

Response: We apologize for this error and have now amended the sentence.

13. P.11, lines 256-259. The reference #22 (Hochberg et al., 2002 J Clin Endocrinol Metab) is a meta-analysis of randomized placebo-controlled trials in postmenopausal women who were all diagnosed at baseline as being osteoporotic. The effects of antiresorptive pharmaceutical agents (particularly the bisphosphonates alendronate and risedronate) were analyzed regarding the later incidence of nonvertebral fractures in relation with early changes in BMD and biochemical markers (BCM). This important meta-analysis estimated that a 3% increase in hip BMD reduced nonvertebral fracture risk by about 46%, and a 70% decrease in resorption BCM reduced nonvertebral fracture risk by about 40 %. In contrast, the submitted observational cross-sectional study was carried out in less than 60 % of osteoporotic women (360/626=57.5%, Table 5). Therefore, the hypothesis that the effects of increased yogurt consumption on BMD and resorption biochemical markers would cause in the long term such a large reduction in nonvertebral fracture risk should be expressed with some caution.

Response: We thank the reviewer for this comment and have inserted this information into the text and we have also stated that such a reduction is expressed with caution 'However it is important to note that the meta-analysis [27] was conducted in women all diagnosed with osteoporosis whereas in the current study only 60% of the women were osteoporotic and thus the potential for yogurt to reduce fractures at the same rate should be viewed with caution' (lines 273-276).

14. The punctuation should be checked throughout the manuscript, particularly some full stops are missing e.g.: P13, line 302.

Response: We apologize for this error and have now checked and amended the manuscript throughout.

#### References

Gelman, Andrew, Jennifer Hill, and Masanao Yajima. "Why we (usually) don't have to worry about multiple comparisons." *Journal of Research on Educational Effectiveness* 5.2 (2012): 189-211.

#### Reviewer 2:

In this cross-sectional study conducted in a large cohort of healthy men and women aged 60 years and older, an association between bone health and physical function, and yogurt consumption was assessed. The results indicate some positive association, and suggest the possibility of yogurt based preventive regimens for maintaining bone health. This paper raises the following comments.

Response: We thank reviewer 2 for their valuable comments regarding the manuscript. The clarifications suggested by the reviewer are again highly appreciated and the specific replies to each comment are shown below.

Major points:

1. The level of physical activity in yogurt consumers and non-consumers should be presented.

Response: We have now added the sentence 'In yogurt consumers, the proportion who answered yes to physical activity was 80.9% while in non-consumers it was 74.7%.' (lines 207-208).

2. The size of yogurt servings as well as whether yogurts are usually enriched in milk powder in the region should be specified, in other words, whether 120 g of yogurt are providing more calcium and protein than 120 g of milk.

Response: We have no data in relation to the serving size of each dairy type or the particular product brand (lines 136 - 137) and thus we cannot state whether there is a difference in the calcium or protein content. We have mentioned this as a limitation of the study (lines 355-358). However, we have also mentioned the average serving intake size of milk, yogurt and cheese in a representative study of older Irish adults (>65 yrs) recruited at the same time period (as part of the National Adult Nutrition Study (NANs) (lines 137 - 140).

3. The results should also be adjusted for calcium and protein of dairy origin, to address the issue of yogurt specificity.

Response: As stated in our response to major point 2, we have no data in relation to the serving size of each dairy type or the particular product brand and thus we cannot adjust for the exact protein or calcium amount of dairy origin. However, through all of the analysis of the associations of yogurt with BMD and bone health we have adjusted for the frequency of intakes for other dairy products such as milk and cheese and have adjusted for the frequency of intakes for non-dairy products including red and white meat (total meat), oily and white fish (total fish) and egg intakes.



4. What would be the results when adjusted for weight and height, instead of BMI?

Response: We have repeated the analysis with adjustment for weight and height instead of BMI and the results are very similar to what has been reported in the manuscript. For women, total hip BMD was 3.1% higher ( $P=0.004$ ) and the femoral neck BMD was 3.9% higher ( $P<0.0001$ ) in those with the highest yogurt intakes compared to the lowest. All other results for women were also similar. For men, the concentrations of the bone biomarker TRAP were 9.6% lower in those with the highest yogurt intakes compared to the lowest ( $P=0.004$ ).

5. The associations differ between men and women. What kind of explanation can be provided?

Response: The results differ significantly between genders and we suggested that given significantly fewer men than women were high yogurt consumers, it is possible that there were only subtle bone turnover changes in men as not enough yogurt was being consumed to affect BMD but enough to affect bone turnover (lines 283 - 286). However, it is beyond the scope of the current study to detect the reason behind the gender differences.

5. Did T-score differ between groups as well, without or with adjustment?

Response: For women only (without adjustment), total hip T-scores were significantly higher ( $P=0.006$ ) as were femoral neck T-scores ( $P<0.0001$ ) in those with the highest yogurt intakes compared to the lowest/none yogurt intake group. For women only (with adjustment), total hip T-scores were significantly higher ( $P=0.004$ ) as were femoral neck T-scores ( $P<0.0001$ ) in those with the highest yogurt intakes compared to the lowest/none yogurt intake group. There was no difference in vertebral T-scores with or without adjustment.

For men only, vertebral T-scores were significantly higher in the low yogurt consumer group compared with the lowest/none yogurt intake group with adjustment ( $P=0.028$ ) and without adjustment ( $P=0.040$ ). There was no differences in total hip or femoral neck T-scores with or without adjustment.

Minor points

1. Abstract: cost-effectiveness of yogurt consumption has not been evaluated nor discussed in the paper.

Response: This statement has been removed from the abstract

2. Introduction: the paper showing reduced fracture risk and mortality with fermented dairy products may be quoted.

Response: This reference has now also been mentioned in the introduction (lines 92 - 93).

3. Did the subjects without BMD measurements differ from those included in the present study?

Response: The TUDA study was based on three diseased cohorts: hypertension, cognitive dysfunction and osteoporosis. Those who were on current bone treatment were removed from the analysis. Subjects in the cognitive cohort (this cohort did not have BMD measures) were recruited from geriatric clinics and a day hospital service and cognitive impairment was assessed based on testing with the RBANS (Repeatable Battery for the Assessment of Neuropsychological Status). This cohort was significantly older and their ability to accurately recall dietary intakes may be compromised.

4. Which database was used to calculate T-score in men? The same as for women?

Response: The database used was from NHANES III for the femoral neck and total hip and the manufacturer's own database for the vertebral.

5. Prevalent fracture should be reported

Response: A very small proportion of the cohort self-reported either a hip fracture (n = 54) or self-reported a vertebral fracture (n = 128) and thus we could not examine the associations with dairy intakes as the study was not powered to detect or examine prevalent fracture rates. Additionally, we cannot ensure the accuracy of these fractures rates as they have been self-reported.

6. Phosphate supplements are very unusual. What was the reason for these 6%?

Response: We agree with the reviewer that phosphate supplements on their own is quite unusual. However, the TUDA study collected information on all supplements/medications consumed and thus an exhaustive search was conducted for any preparation, medication or supplement that contained any phosphate to be as thorough as possible.

7. P 11: the paper by Hochberg et al cannot be used in the present discussion since the changes in BMD were achieved with pharmacological agents. There is no argument that the relationship may be similar with nutritional intervention.

Response: We acknowledge the reviewers comments and although the changes in BMD were achieved with pharmacological agents in the paper by Hochberg et al., there is still a possibility that there is a similar relationship with nutritional intervention (dairy products) as this has not been investigated before. However we have modified this part of the discussion in line with suggestions with reviewer 1 and have stated that these possibilities should be treated with caution until further evidence is available.

8. The possibility that yogurt consumption may be a marker of a healthy lifestyle should be discussed.

Response: We agree with the reviewer that yogurt consumption may be a marker of a healthy lifestyle and have discussed this in the manuscript (lines 315 - 316).



[Click here to view linked References](#)

1 **Greater yogurt consumption is associated with increased bone mineral density and**  
2 **physical function in older adults**

3  
4  
5  
6  
7 4 E. Laird<sup>1</sup>, A. M. Molloy<sup>1</sup>, H.McNulty<sup>2</sup>, M.Ward<sup>2</sup>, K.McCarroll<sup>3</sup>, L.Hoey<sup>2</sup>, C.Hughes<sup>2</sup>,  
8  
9 5 C.Cunningham<sup>3</sup>, J.J. Strain<sup>2</sup>, M. C. Casey<sup>3</sup>.

10  
11  
12  
13  
14 7 <sup>1</sup>School of Medicine, Trinity College Dublin, Ireland; <sup>2</sup>Northern Ireland Centre for Food and  
15  
16 8 Health, University of Ulster, Coleraine, United Kingdom; <sup>3</sup>The Mercers Institute for Research  
17  
18 9 on Ageing, St James's Hospital, Dublin, Ireland.

19  
20  
21  
22 10  
23  
24 11 **Key words:** Yogurt; BMD; Physical function; Ageing; Frailty

25  
26 12 **Word count:** 3,276 words

27  
28  
29 13 Corresponding Author/Reprint requests: Dr Eamon Laird

30  
31  
32 14 Corresponding Author address:

33  
34  
35 15 School of Medicine

36  
37 16 Trinity College Dublin, Dublin 2, Ireland

38  
39 17 Email: lairdea@tcd.ie

40  
41  
42 18 **Conflict of interest**

43  
44  
45  
46 19 Eamon Laird, Anne Molloy, Helene McNulty, Mary Ward, Kevin McCarroll, Leane Hoey,  
47  
48 20 Catherine Hughes, Conal Cunningham, JJ Strain and Miriam Casey declare that they have no  
49  
50  
51  
52 21 conflict of interest

53  
54  
55 22

56  
57  
58 23

24 **Mini-Abstract**

1  
2 25 In this cohort of community dwelling older adults (>60 yrs), we observed significant positive  
3  
4 26 associations between the frequency of yogurt intake with measures of bone density, bone  
5  
6  
7 27 biomarkers and indicators of physical function. Improving yogurt intakes could be a valuable  
8  
9  
10 28 health strategy for maintaining bone health in older adults.

11  
12 29  
13  
14 30 **Abstract**

15  
16  
17 31 *Introduction* The associations of yogurt intakes with bone health and frailty in older adults are  
18  
19 32 not well documented. The aim was to investigate the association of yogurt intakes with bone  
20  
21  
22 33 mineral density [BMD], bone biomarkers and physical function in 4,310 Irish adults from the  
23  
24 34 Trinity, Ulster, Department of Agriculture aging cohort study (TUDA).

25  
26 35 *Methods* Bone measures included total hip, femoral neck and vertebral BMD with bone  
27  
28  
29 36 biochemical markers. Physical function measures included Timed Up and Go (TUG),  
30  
31  
32 37 Instrumental Activities of Daily Living Scale and Physical Self-Maintenance Scale.

33  
34 38 *Results* Total hip and femoral neck BMD in females were 3.1 - 3.9 % higher among those with  
35  
36 39 the highest yogurt intakes ( $n= 970$ ) compared to the lowest ( $n= 1,109$ ;  $P <0.05$ ) as were the  
37  
38  
39 40 TUG scores ( $-6.7\%$ ;  $P = .013$ ). In males, tartrate-resistant acid phosphatase (TRAP 5b)  
40  
41  
41 41 concentrations were significantly lower in those with the highest yogurt intakes ( $-9.5\%$ ;  $P$   
42  
42  
42 42  $<0.0001$ ). In females, yogurt intake was a significant positive predictor of BMD at all regions.

43  
43 43 Each unit increase in yogurt intake in females was associated with a 31% lower risk of  
44  
44 44 osteopenia (OR 0.69; 95% CI 0.49 – 0.96;  $P=0.032$ ) and a 39% lower risk of osteoporosis (OR  
45  
45 45 0.61; 95% CI 0.42 - 0.89;  $P=0.012$ ) and in males, a 52% lower risk of osteoporosis (OR 0.48;  
46  
46 46 95% CI 0.24 - 0.96;  $P=0.038$ ).

47 *Conclusion* In this cohort, higher yogurt intake was associated with increased BMD and  
48 physical function scores. These results suggest that improving yogurt intakes could be a  
49 **valuable public health strategy** for maintaining bone health in older adults.

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72 **Introduction**

1  
2 73 Osteoporosis is an increasingly common, chronic condition estimated to affect over 200 million  
3  
4 74 individuals worldwide [1] with 6% of men and 21% of women aged 50-84yrs affected in the  
5  
6  
7 75 EU alone [2]. It is characterized by decreased bone mineral density (BMD) with a significantly  
8  
9  
10 76 increased risk of fracture and subsequently, morbidity and mortality [3]. The condition has  
11  
12 77 been estimated to cause over 8.9 million fractures annually, with osteoporotic fractures  
13  
14 78 accounting for 0.8% of the global burden of non-communicable disease and the loss of over  
15  
16  
17 79 5.8 million disability-adjusted life years (DALYs) [4-6].  
18

19 80

20  
21  
22 81 The development of osteoporosis can be influenced by a range of both demographic and  
23  
24 82 lifestyle factors [1,7,8]. However, maintaining an optimal nutritional status is also a key  
25  
26  
27 83 preventative measure, particularly for older adults (>50 years) [9]. Of the major food groups,  
28  
29 84 dairy foods are one of the richest sources of the macro and micro nutrients that contribute to  
30  
31  
32 85 bone health such as protein, calcium, magnesium and the B-vitamins [10-14]. For example,  
33  
34 86 dairy products are the primary source of calcium across most industrialized countries in Europe  
35  
36  
37 87 and the USA [11,12]. Previous data from observational studies and randomized controlled trials  
38  
39 88 (RCTs) have reported significant positive associations between dairy intakes and bone health  
40  
41  
42 89 as reviewed in recent commentaries and Government reports [15,16]. In one 12-yr follow-up  
43  
44 90 analysis of the Framingham Offspring Study ( $n= 2,506$ ; mean age 55 yrs), yogurt intake alone  
45  
46  
47 91 was positively associated with hip trochanter BMD and had a weak protective trend with hip  
48  
49 92 fracture reduction [17]. Furthermore, fermented milk products have been associated with a  
50  
51 93 lower fracture incidence and mortality [18]. There is however, little information on the  
52  
53  
54 94 associations of yogurt intake with bone health bio-markers and with measures of functionality.  
55  
56 95 In the current study, we examined the association of yogurt intakes with BMD, biochemical  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 96 markers of bone health and physical function measures in a large cohort of free-living older  
2 97 adults ( $n= 4,310$ , age range 60-102 yrs).  
3  
4

5 98

## 7 99 **Subjects and Methods**

8  
9  
10 100 Data analyzed for the current study originated from the Trinity Ulster Department of Agriculture  
11 101 (TUDA) ageing cohort study, a large study of older Irish adults (>60 yrs) designed to investigate  
12 101 (TUDA) ageing cohort study, a large study of older Irish adults (>60 yrs) designed to investigate  
13  
14 102 nutritional factors, related gene-nutrient interactions and a range of health and lifestyle factors  
15  
16 102 nutritional factors, related gene-nutrient interactions and a range of health and lifestyle factors  
17 103 in the development of chronic diseases of aging. Further details of sampling and recruitment  
18  
19 104 have been described previously [19-22]. Of the 5,186 participants recruited, 3 with severe frailty  
20  
21 104 have been described previously [19-22]. Of the 5,186 participants recruited, 3 with severe frailty  
22 105 (replied no or had a missing answer to the self-feeding question in the Physical Self-  
23  
24 106 Maintenance questionnaire (PSM)), 866 with cognitive impairment (Mini-Mental State  
25  
26 106 Maintenance questionnaire (PSM)), 866 with cognitive impairment (Mini-Mental State  
27 107 Examination (MMSE) score <25) and those with a missing response to the yogurt intake  
28  
29 108 question ( $n= 7$ ) were excluded from the physical function analysis leaving a total of 4,310  
30  
31 108 question ( $n= 7$ ) were excluded from the physical function analysis leaving a total of 4,310  
32 109 participants (Supplemental Figure 1). Approximately 1,699 participants did not have BMD  
33  
34 110 measures taken. In addition to these exclusions, participants who reported receiving medications  
35  
36 111 that could affect bone mineral metabolism (Bisphosphonates; Aromatase Inhibitors;  
37  
38 111 that could affect bone mineral metabolism (Bisphosphonates; Aromatase Inhibitors;  
39 112 Gonadotropin releasing hormone analogues or Luteinizing hormone releasing agonists; Anti-  
40  
41 112 Gonadotropin releasing hormone analogues or Luteinizing hormone releasing agonists; Anti-  
42 113 androgen medication; Parathyroid hormone (PTH) treatment; Strontium treatment; Anti-  
43  
44 113 androgen medication; Parathyroid hormone (PTH) treatment; Strontium treatment; Anti-  
45 114 epileptic medications; Paget's disease treatment) were also excluded from the BMD and bone  
46  
47 115 biomarker analysis (Supplemental Figure 1). Ethical approval was granted by the relevant  
48  
49 116 authorities in each jurisdiction: the Research Ethics Committee of St. James's Hospital and The  
50  
51 117 Adelaide and Meath Hospital, Dublin, and the Office for Research Ethics Committees Northern  
52  
53 117 Adelaide and Meath Hospital, Dublin, and the Office for Research Ethics Committees Northern  
54 118 Ireland (ORECNI; reference 08/NI/RO3113) with corresponding approvals from the Northern  
55  
56 119 and Western Health and Social Care Trusts, Northern Ireland.  
57

58 120  
59  
60  
61  
62  
63  
64  
65

121 **Lifestyle and medications**

1  
2 122 Data associated with lifestyle factors were obtained by questionnaire. Information included  
3  
4  
5 123 gender, age, ethnicity, physical activity (reported as yes/no in the last two weeks), smoking  
6  
7 124 status and alcohol intake. Full details of dietary supplement and vitamin use including dose,  
8  
9  
10 125 frequency and duration was confirmed from packaging or prescription information. A small  
11  
12 126 number of supplements listed were unidentifiable or contained unidentifiable ingredients;  
13  
14  
15 127 individuals consuming such supplements were excluded from analysis.  
16

17 128

19 129 **Dietary dairy intake servings**

21  
22 130 Participants were given a modified food frequency questionnaire (FFQ) which asked if the  
23  
24 131 participant consumed yogurt, milk (approximated as glasses of milk per day) and/or cheese and  
25  
26  
27 132 if yes, how often for each dairy type. The total frequency of the daily intake serving for yogurt,  
28  
29 133 milk and cheese was calculated from the FFQ responses (Supplemental Table 1). These values  
30  
31  
32 134 were then separated into tertiles of non-consumers, low consumers and high consumers for  
33  
34 135 each dairy type. Information was also recorded for the frequency of consumption of red meat  
35  
36 136 and poultry (total meat), oily and white fish (total fish) and egg intakes. No product brand name  
37  
38  
39 137 or serving size information was available for the dairy intakes, however the average serving  
40  
41 138 intake size in a representative study of older Irish adults (>65 yrs) recruited at the same time  
42  
43  
44 139 period (as part of the National Adult Nutrition Study (NANs)) was 114 grams (g) for yogurt,  
45  
46 140 123g for milk and 35g for cheese [23].  
47

48  
49 141

51 142 **Biochemical analysis**

52  
53 143 A non-fasting blood sample (50ml) was collected by venipuncture into an evacuated clotting  
54  
55  
56 144 tube (Sarstedt; Numbrecht, Germany) by a trained phlebotomist. Samples were kept chilled  
57  
58 145 and centrifuged (3000 rpm for 15 minutes) within 3 hours of collection and serum aliquots  
59  
60



146 were labeled and stored at -80°C until required for analysis. Serum bone bio-markers were  
147 measured in duplicate using an automated enzyme immunoassay method (EIA) following the  
148 manufacturer's instructions (Triturus®, Immunodiagnosics (IDS) limited, Boldon, Tyne &  
149 Wear, UK). Inter-assay CVs were as follows: Serum osteocalcin (OC) <4.5% (reference range  
150 for males of 9.6-40.8 ng/ml and for postmenopausal women 12.8-55.0 ng/ml), C-terminal  
151 telopeptides of type I collagen (CTX) <3.1% (reference range is 0.020 ng/mL to 3.380 ng/ml),  
152 Bone specific alkaline phosphatase (BAP) <1.5 % (reference range for males of 5.7-32.9 µg/ml  
153 and for postmenopausal women 5.5-27.1 µg/ml) and Tartrate-resistant acid phosphatase  
154 (TRAP 5b) <1.6% (reference range for males of 55-79 ng/ml and for postmenopausal women  
155 41-81 ng/ml). Intact PTH was measured at St. James's Hospital, Dublin using an  
156 electrochemiluminescence immunoassay (ECLIA) (Modular E170, Roche Diagnostics,  
157 Dublin, Ireland) with an inter-assay CV of <2.9% and an assay measurement range of 1.2 –  
158 5000 pg/ml. Vitamin D (25-hydroxyvitamin D (25(OH)D)) concentrations were quantified  
159 using LC-MS/MS (API 4000; AB SCIEX; Chromsystems GmbH) with an inter-assay CV of  
160 <5.7% (detection range 7.5 – 624 nmol/L) [19,20]. Renal function tests (creatinine) were  
161 analyzed using a Roche Cobas c701 (Roche 8000 modular system) with an inter-assay CV  
162 <5%. Glomerular filtration rate (GFR) was estimated by use of the Cockcroft-Gault equation.

#### 164 **BMD and Physical function measures**

165 BMD was measured by dual energy X-ray absorptiometry (DXA) (Lunar iDXA™, UK)  
166 performed at the hip, femoral neck and the vertebral column by a fully trained operator  
167 according to ionizing radiation medical exposure regulations (IRMER) and scans were  
168 subsequently interpreted with the assistance of a radiographer. Results were expressed as grams  
169 of BMD per square centimeter (g/cm<sup>2</sup>) and as T-scores using the manufacturer's reference  
170 database. Osteopenia was defined as a BMD T-score between -1.0 and -2.5 at any site and

171 osteoporosis defined as a BMD T score  $> -2.5$  at any site (below the young adult mean) [24].  
172 Physical function was primarily assessed by use of the Timed Up and Go test (TUG) which  
173 measured the time it took a participant to rise from a chair, walk three meters, turn around,  
174 walk back to the chair and sit down again. A score of 12 seconds or more has been reported as  
175 an indication of reduced mobility [25]. Additional functionality measures included the PSM  
176 and the Instrumental Activities of Daily living scale (IADL) [26].

177

### 178 **Statistical analysis**

179 The statistical analysis was performed using the Statistical Package for Social Sciences  
180 (Version 23.0; SPSS UK Ltd; Chersey, UK). Data were assessed for normality and where  
181 necessary, data were log-transformed for normalization purposes. Data within tables are  
182 primarily expressed as adjusted means with 95% confidence intervals. Estimated marginal  
183 means were adjusted for age, gender, BMI, GFR, calcium and vitamin D supplement usage  
184 (yes/no). Where appropriate, an independent Student's T-test, one-way ANOVA or ANCOVA  
185 with pair-wise comparisons were applied to determine statistical differences between groups  
186 ( $P < 0.05$ ). Data were corrected for multiple comparison using the Bonferroni correction.

187 Categorical variables were assessed by chi-square analysis. Hierarchical multiple regression  
188 models with adjustment for age, gender, education, BMI, smoking status, alcohol consumption,  
189 physical activity (in past two weeks), vitamin D and calcium supplement usage, 25(OH)D  
190 concentration, daily milk, yogurt, cheese, total meat (red meat and poultry), total fish (oily and  
191 white) and daily egg servings were applied to determine significant predictors of BMD  
192 concentrations and physical function measure scores. To determine the predictors of bone  
193 health (osteopenia or osteoporosis), a multinomial logistic regression model was used (with  
194 normal bone health as the reference category) with relevant co-predictors including the nominal  
195 variables: age, BMI, education, 25(OH)D concentration, PTH concentration, GFR, frequency

196 of daily servings of milk, yogurt, cheese, total meat, total fish and eggs and the categorical  
197 variables: gender (reference male), vitamin D supplement user (reference non-vitamin D  
198 supplement user), non-smoker (reference smoker), non-alcohol consumer (reference alcohol  
199 consumer), physical activity: yes (reference physical activity: no).

## 201 Results

202 General characteristics of participants in the TUDA cohort as defined by gender are shown in  
203 Table 1. The majority of participants were female (67.4%), who were significantly older  
204 ( $P=0.004$ ), lighter ( $P<0.0001$ ), and contained a higher proportion of individuals receiving  
205 vitamin D or calcium supplements ( $P<0.0001$ ) in comparison with males. A higher percentage  
206 of females were yogurt consumers with mean daily yogurt servings significantly higher than  
207 males (0.42/d vs. 0.32/d respectively) ( $P<0.0001$ ). In yogurt consumers, the proportion who  
208 answered yes to physical activity was 80.9% while in non-consumers it was 74.7%. In  
209 participants who had measures of BMD performed, 41.3% had osteopenia while 27% had  
210 osteoporosis which was more common in females than males (35.6% vs 14.8% respectively;  
211  $P<0.0001$ ). Data for BMD, the bone biomarker concentrations and physical function measures  
212 across the frequency of daily yogurt intakes (split by gender) are presented in Table 2 and Table  
213 3. In females, after adjustment for covariates (and exclusion of those receiving medications  
214 that may affect BMD), total hip BMD was 3.1% higher ( $P=0.005$ ) and femoral neck BMD was  
215 3.9% higher ( $P<0.0001$ ) in the high yogurt consumers (>once per day serving) compared to the  
216 non-consumers (<once per week serving/never). In males, vertebral BMD was 4.1% higher in  
217 low yogurt consumers compared with non-consumers ( $P=0.028$ ). Similarly, mean vitamin D  
218 concentrations (after exclusion of those receiving vitamin D supplements) were 12.9% higher  
219 ( $P=0.006$ ) and mean TRAP 5b concentrations were 9.5% lower ( $P=0.003$ ) in the male high  
220 yogurt consumers compared to the non-consumers. No significant change in concentration

221 across yogurt consumption was observed for PTH or the remaining bone biomarkers in either  
222 gender.

223  
224 For physical function measures in females, non-consumers of yogurt were 0.9 seconds (6.7%)  
225 slower than the high consumers (13.8 vs 12.9 seconds;  $P=0.020$ ). Similarly, PSM and IADL  
226 scores were significantly higher in the yogurt high consumers compared to the non-consumers  
227 ( $P=0.010$  &  $P=0.003$  respectively). No significant difference was observed for males. This  
228 analysis was then repeated to examine BMD, bone biomarker and physical function measures  
229 across frequency of milk and cheese intakes (Supplemental Tables 2-5). No significant  
230 difference was observed across milk intake frequencies for BMD or bone biomarker  
231 concentrations. However, TUG scores were significantly lower in the non-milk consumers  
232 compared to the high milk consumers in both men and women ( $P<0.05$ ). In addition there were  
233 slight increases in PSM and IADL scores across milk intakes in both genders while no  
234 significant difference was observed across cheese intake frequencies for any of the measures.

235  
236 In a hierarchical multiple regression model (Table 3) examining predictors of BMD, bone  
237 markers and physical function measures, increasing yogurt intake was a significant positive  
238 predictor for BMD in females at all three sites after adjustment for relevant covariates. For  
239 instance, with each unit increase in yogurt intake (i.e., an increase of one serving per week)  
240 total hip BMD increased by  $0.015 \text{ g/cm}^2$  ( $P=0.002$ ), vertebral BMD by  $0.026 \text{ g/cm}^2$  ( $P=0.005$ )  
241 and femoral neck BMD increased by  $0.023 \text{ g/cm}^2$  ( $P<0.0001$ ). Furthermore, with each yogurt  
242 unit increase, TUG scores decreased by 0.59 seconds ( $P=0.021$ ). In men only, with each unit  
243 increase in yogurt intake, concentrations of TRAP 5b decreased by  $0.118 \text{ } \mu\text{g/L}$  ( $P<0.0001$ ).  
244 Significant predictors of bone health status are outlined in Table 4. Daily yogurt intake was a  
245 significant predictor of bone health with each unit increase in yogurt intake associated with a

246 31% lower risk of having osteopenia (OR 0.69; 95% CI 0.49 – 0.96;  $P=0.032$ ) and a 39% lower  
247 risk of being characterized as osteoporotic (OR 0.61; 95% CI 0.42 - 0.89;  $P=0.012$ ) in females  
248 and in males a 52% lower risk of osteoporosis (OR 0.48; 95% CI 0.24 - 0.96;  $P=0.038$ ). For  
249 females, those on vitamin D supplements had a significantly reduced risk of osteopenia (OR  
250 0.51; 95% CI 0.34 – 0.76;  $P=0.001$ ) and a significantly reduced risk of osteoporosis (OR 0.41;  
251 95% CI 0.26 – 0.64;  $P<0.0001$ ). For males, those on vitamin D supplements also had a  
252 significantly reduced risk of osteoporosis (OR 0.40; 95% CI 0.22 – 0.72;  $P=0.003$ ).

## 254 Discussion

255 In this study we observed significant positive associations of increased frequency of yogurt  
256 intakes with bone health and measures of physical function in a cohort of older adults. Females  
257 with the highest yogurt intakes had significantly higher BMD and better physical function  
258 scores compared to individuals with the lowest intakes. Furthermore, we show for the first time  
259 that, after adjustment for covariate predictors, each unit increase in yogurt intake significantly  
260 decreased the odds of being characterized as osteopenic or osteoporotic in women and as  
261 osteoporotic in men.

262  
263 The significant positive associations of yogurt with BMD within this large study are consistent  
264 with previous observations from the Framingham Offspring observational study [17] In 2,733  
265 adults (26-85 yrs), higher yogurt intake was positively associated with trochanteric BMD over  
266 a 12-year follow-up with a weak protective trend of yogurt (but not other dairies) on the risk  
267 of hip fracture. In a cohort of 61,000 Swedish women (aged 39-74 yrs), fermented milk  
268 products (yogurt) were associated with a significant decrease in fracture incidence and  
269 mortality over a mean follow-up of 20 years. With each increase in fermented dairy intakes,  
270 hip fractures were reduced by 10-15% [18]. Although the current data-set did not have data on

271 fracture incidence, the effect of increased yogurt intake seen in this cohort has the potential to  
272 reduce non-vertebral fractures by up to 46% in women, as fracture risk reduction has been  
273 modelled as 46% decrease for 3% hip BMD increase [27]. However it is important to note that  
274 the meta-analysis [27] was conducted in women all diagnosed with osteoporosis whereas in  
275 the current study only 60% of the women were osteoporotic and thus the potential for yogurt  
276 to reduce fractures at the same rate should be viewed with caution. The potential protective  
277 effects of yogurt on bone health are also supported by the positive associations of yogurt with  
278 the bone biomarker Trap 5b, the concentrations of which were 9.5% lower in those with the  
279 highest yogurt intake compared to the lowest, though only in men. Trap 5b is a direct marker  
280 of osteoclast number and bone resorption (indicating positive bone balance), with better  
281 sensitivity than CTX (a by-product of collagen breakdown) [28] and has been described as one  
282 of the most sensitive markers to monitor the response of diet intervention on bone resorption  
283 [29]. Significantly fewer men than women were high yogurt consumers, and it is possible this  
284 marker was detecting subtle bone turnover changes in men only as not enough yogurt was being  
285 consumed to affect BMD but enough to affect bone turnover, though this hypothesis needs to  
286 be tested. If the results from the current study are confirmed, there is the potential that increased  
287 yogurt intakes may add an inexpensive and relatively low-risk strategy to improve bone health  
288 in conjunction with bone treatment. However, future research and randomized controlled trials  
289 are needed to explore this approach.

290

291 Notably, this study also observed that greater consumption of yogurt was associated with a  
292 significantly lower TUG score (6.7% difference lowest vs highest yogurt intakes) in women  
293 only. TUG has been described as a composite measure of functional mobility with worse scores  
294 associated with poorer muscle strength and balance, both of which are risk factors for falling  
295 in older adults [30]. Our results are in agreement with Lana et al. who observed that higher



296 consumption of yogurt (and milk) was associated with a lower risk of frailty and a lower risk  
297 of a slow walking speed in 1,871 community dwelling older adults [31]. Furthermore, in a  
298 cross-sectional study of elderly Australian women (n 1,456), higher dairy intake was associated  
299 with increased grip strength and decreased likelihood of a lower TUG score [32].

300  
301 A number of potential mechanisms may explain the observed positive associations. Yogurt  
302 naturally contains significant concentrations of bone promoting minerals and vitamins [10-14]  
303 which have also been associated with improved frailty measures [33,34]. In data from the  
304 Framingham Heart Study offspring cohort, yogurt consumers were 47% and 55% less likely to  
305 have inadequate intakes of vitamins B2 and B12 (respectively) [35], while in 2,797 Italian  
306 adults (aged 18-97 yrs), yogurt consumers were more likely to have adequate intakes of  
307 vitamins and minerals compared to non-consumers [36]. Yogurt also contains significant  
308 quantities of protein, bio-active peptides and bio-cultures which have been associated with  
309 bone health and immunological benefits [37-43]. For example, yogurt (and other dairy  
310 products) contain branched chain amino acids (BCAAs) which are potent stimulators of muscle  
311 protein synthesis [44,45]. Furthermore, in a recent review it was suggested that the modifiable  
312 nature of the gut microbiome could provide a potential therapeutic target to intervene in  
313 musculoskeletal conditions of aging [46]. It is perhaps this unique combination of macro and  
314 micronutrients with bio-active compounds within yogurt that confers bone promotion and  
315 improved physical function. It is also possible that increased yogurt intakes could also be a  
316 reflection of a long term dietary habit of an overall healthy eating pattern and lifestyle [36],  
317 though diet quality (including vitamin D and calcium) was adjusted for in the current analysis.  
318 Yogurt has been the target of some criticism, especially with the renewed concerns regarding  
319 excess sugar intakes and associations with obesity [47] given that some processed 'sweetened'  
320 yogurts can contain substantial quantities of sugar [48]. Yet not all yogurts have a high sugar

1 321 content [49] and further exploration is required to identify the types of yogurts and the  
2 322 individual components within that may exhibit health benefits.  
3

4  
5 323

6  
7 324 We also examined the associations of the other dairy products (milk and cheese) with BMD  
8  
9 325 and functionality. We observed no significant difference in BMD across milk intakes, in line  
10  
11 326 with inconsistent data from previous observational studies. For example, some studies have  
12  
13 327 observed strong associations between childhood and adolescent milk consumption with BMD  
14  
15 328 [50]. For older adults (>60 yrs), studies have observed no associations or a negative association  
16  
17 329 of milk intakes with fracture risk [50]. The majority of positive randomized trials with milk  
18  
19 330 which have observed significant decreases in the concentrations of bone biomarkers and  
20  
21 331 improvements in bone metabolism have all utilized fortified milk [51]. The milk intakes in the  
22  
23 332 current study were not heavily fortified at this time period (2008-2012) and could account for  
24  
25 333 the lack of any such association. Furthermore, we have previously reported that in this  
26  
27 334 population, with increasing age, milk intakes increased while yogurt intakes significantly  
28  
29 335 decreased [22]. This could help explain why some of the physical function measures became  
30  
31 336 poorer with increased milk intakes. As milk intakes increased, we suggest that there was a loss  
32  
33 337 of a particular protective component within the yogurt that enhanced bone health/improved  
34  
35 338 physical function, though this hypothesis requires verification. Interestingly, we observed no  
36  
37 339 significant associations of cheese intakes with BMD though male participants with  
38  
39 340 osteoporosis were more likely to have a higher frequency of cheese consumption. Cheese  
40  
41 341 products generally have a different nutritional profile in comparison with yogurts and we  
42  
43 342 previously observed in TUDA that cheese intakes had no significant effect on the  
44  
45 343 concentrations of vitamin D, folate, vitamin's B12, B6 or B2 [22]. Furthermore, it has been  
46  
47 344 suggested that the high sodium content of certain cheeses could be less beneficial for bone  
48  
49 345 health by negatively altering calcium metabolism, though few studies have examined this issue  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 346 [52] and the sodium-induced calciuria hypothesis has not been supported by any long term

2 347 observations. Moreover, fortified cheese products have been positively associated with bone

3 348 metabolism [53]. Further research is needed to identify the relationship between cheese intakes

4 349 and BMD in men.

5 350

6 351 Our study has several limitations. The data are observational and cross-sectional, and such

7 352 observed associations between yogurt intakes and bone health do not necessarily indicate a

8 353 causal relationship. However, one of the major strengths of this study was the size, as to the

9 354 best of our knowledge it is the largest observational study conducted to date investigating such

10 355 associations. Potential weaknesses of this study also include our reliance on self-reported

11 356 intakes and we were unable to quantify the dairy or yogurt intakes through food dairies or other

12 357 more quantitative dietary collection (and thus did not have information on serving sizes or

13 358 product types). However, although we could not adjust for total energy intake, we did adjust

14 359 for frequency of intake of other important dietary components including meat, fish, egg and

15 360 other dairy constituents which can give a proxy measure of diet quality. Furthermore, those

16 361 with severe cognitive impairment or frailty were removed from the analysis to increase recall

17 362 accuracy.

18 363

19 364 In conclusion, to our knowledge, this is the largest study to demonstrate an association between

20 365 the frequency of yogurt intakes, BMD, bone biomarkers and measures of physical function

21 366 exclusively within free-living, older adults (>60y). The findings provide evidence that lower

22 367 frequency of yogurt intake is significantly associated with a lower BMD and that improving

23 368 yogurt intakes could be a valuable and cost-effective health measure for maintaining bone

24 369 health and in reducing frailty in older adults. Future RCT trials are required to assess and

25 370 investigate the efficacy of such approaches.

371 **Acknowledgments**

1  
2 372 This research was funded by the Irish Department of Agriculture, Food and the Marine through  
3  
4  
5 373 the grants 07FHRIUCD1 ("JINGO" 2007-2013) and 13F407 ("JINGO-JPI"/"ENPADASI"  
6  
7 374 2014-2016) and from the Northern Ireland Department for Employment and Learning (DEL),  
8  
9  
10 375 Cross-Border Research and Development Programme: "Strengthening the all Island Research  
11  
12 376 Base". Funding was also provided from The National Dairy Council (NDC) Ireland. The  
13  
14 377 funding organizations had no role in the design and conduct, collection, management, analysis,  
15  
16  
17 378 and interpretation of the data, or in the preparation, review, or approval of this manuscript. The  
18  
19 379 authors also acknowledge Drs Martin Healy, Adrian McCann and Liadhan McAnena for bone  
20  
21  
22 380 biochemical analysis. Authors' roles: Conception and study design: AM, MC, HM, MW, CC,  
23  
24 381 and JJS. Data analysis: EL and AM. Input on statistical analysis: EL, AM, MC, HM. Drafting  
25  
26  
27 382 manuscript: EL, AM and MC. Revising manuscript content, approval of final version: All  
28  
29 383 authors. EL and AM take responsibility for the integrity of the data analysis.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 1.** Demographic and health characteristics of the TUDA cohort study by gender<sup>a</sup>

Variable	Total ( <i>n</i> = 4,310)	Male ( <i>n</i> = 1,405)	Female ( <i>n</i> = 2,905)	<i>P</i> value
Age, <sup>b</sup> y	73.1 (7.9)	72.6 (7.8)	73.3 (8.0)	0.004
60-69, <sup>c</sup> y <i>n</i> (%)	1690 (39.2)	567 (40.4)	1123 (38.7)	0.284
70-79, <sup>c</sup> y <i>n</i> (%)	1697 (39.4)	568 (40.4)	1129 (38.9)	0.325
>80, <sup>c</sup> <i>n</i> (%)	923 (21.4)	270 (19.2)	653 (22.5)	0.014
Age Finished Education, <sup>b</sup> (yrs)	16.2 (3.0)	16.2 (3.2)	16.2 (2.9)	0.564
<b>Health &amp; Lifestyle</b>				
BMI, <sup>b</sup> kg/m <sup>2</sup>	28.0 (5.3)	28.6 (4.4)	27.7 (5.7)	<0.0001
GFR, <sup>b</sup> ml/min	69.4 (24.4)	77.1 (25.4)	65.6 (22.9)	<0.0001
Current smoker, <sup>c</sup> <i>n</i> (%)	515 (12.0)	156 (11.1)	359 (12.4)	0.231
Current alcohol consumer, <sup>c</sup> <i>n</i> (%)	2551 (59.2)	919 (65.5)	1632 (56.2)	<0.0001
Physical activity in last two weeks, <sup>c</sup> <i>n</i> (%)	3403 (79.0)	1094 (77.9)	2309 (79.5)	0.214
Receives Bone medications, <sup>c</sup> <i>n</i> (%)	1484 (34.4)	244 (17.4)	1240 (42.7)	<0.0001
Yogurt consumer, <sup>c</sup> <i>n</i> (%)	2658 (61.7)	725 (51.6)	1933 (66.5)	<0.0001
Milk (as a drink) consumer, <sup>c</sup> <i>n</i> (%)	1806 (42.9)	640 (46.4)	1166 (41.1)	0.001
Cheese consumer, <sup>c</sup> <i>n</i> (%)	3651 (84.7)	1209 (86.0)	2442 (84.1)	0.089
<b>Supplement use,<sup>c</sup> <i>n</i> (%)</b>				
Vitamin D supplement user	2042 (47.8)	447 (33.2)	1595 (58.4)	<0.0001
Calcium supplement user	1742 (40.4)	308 (21.9)	434 (49.4)	<0.0001
Phosphate supplement user	262 (6.1)	82 (5.8)	180 (6.2)	0.643

<sup>a</sup> Values are means ( $\pm$ SD) for continuous variables.

<sup>b</sup> Student's independent *t* test was used to test differences between log-transformed continuous variables.

<sup>c</sup> Chi-square tests were used to test differences between categorical variables.

**Table 2.** Comparison of bone mineral density (BMD), biomarkers of bone health and mean frailty measures across frequencies of daily yogurt intake in females in the TUDA cohort study<sup>a</sup>

Variable	Tertile of daily yogurt intakes								
	Non-consumer			Low consumer			High consumer		
	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI
		Mean yogurt frequency (0.0 daily / <once per week/never) <i>n</i> = 970			Mean yogurt frequency (0.34 daily / 2-3 times per week) <i>n</i> = 826			Mean yogurt frequency (1.03 daily / >once per day) <i>n</i> = 1109	
BMD region, <sup>b</sup> g/cm <sup>2</sup>									
Total Hip	320	0.890	0.877 – 0.903	331	0.904	0.891 – 0.917	406	0.918*	0.907 – 0.930
Femoral Neck	319	0.824	0.812 – 0.836	331	0.843	0.831 – 0.855	405	0.857**	0.846 – 0.867
Vertebral	260	1.005	0.984 – 1.025	251	1.027	1.006 – 1.047	330	1.036	1.018 – 1.054
Bone health biomarkers									
CTX, <sup>b</sup> ng/ml	226	0.34	0.32 – 0.36	258	0.33	0.31 – 0.35	278	0.32	0.30 – 0.34
OC, <sup>b</sup> ng/ml	225	19.2	18.0 – 20.5	252	18.7	17.5 – 19.8	278	18.8	17.7 – 19.9
BAP, <sup>b</sup> µg/L	226	17.8	16.9 – 18.7	252	18.0	17.1 – 18.8	278	17.7	16.9 – 18.5
TRAP 5b, <sup>b</sup> µg/L	226	3.30	3.17 – 3.43	252	3.29	3.17 – 3.41	279	3.24	3.12 – 3.35
25(OH)D, <sup>c</sup> nmol/L	412	41.0	38.9 – 43.2	347	43.4	41.1 – 45.7	361	43.5	41.2 – 45.8
PTH, <sup>b</sup> pg/ml	505	50.0	47.5 – 52.5	446	47.0	44.4 – 49.7	578	46.3	43.9 – 48.6
Physical function measures <sup>c</sup>									
Timed up and Go, (sec)	868	13.8	13.3 – 14.2	740	13.0	12.5 – 13.5	1,016	12.9*	12.5 – 13.3
IADL	906	24.4	24.2 – 24.6	751	24.5	24.3 – 24.8	1,014	24.8*	24.6 – 25.0
PSM	913	22.9	22.8 – 23.0	767	22.9	22.8 – 23.0	1,035	23.1*	23.0 – 23.2

<sup>a</sup> Values are estimated marginal means (95% CI) adjusted for multiple covariates. Differences in means were assessed by pair-wise comparisons **and adjusted for multiple comparisons: Bonferroni correction.** \*,\*\*Different from the lowest yogurt intake tertile: \**P*<0.05, \*\**P*<0.0001. Non-consumer frequency range (0 – 0.07 units / <once per week/never); low consumer frequency range (>0.07- 0.50 units / >once per week to 3-4 times per week); high consumer frequency range (>0.50 – 2.00 units / >3-4 times per week to twice per day).

<sup>b</sup>Adjusted for age, education, BMI, GFR, physical activity, total daily serving milk (glass only), total daily serving of cheese, calcium or vitamin D supplements (Participants receiving medications that could affect bone metabolism were removed from the analysis).

<sup>c</sup>Adjusted for age, BMI, total daily serving milk (glass only), total daily serving of cheese; (participants receiving vitamin D supplements were removed from the 25(OH)D analysis). Abbreviations: BAP, Bone specific alkaline phosphatase; CTX, C-terminal telopeptides of type I collagen; IADL, Instrumental Activities of Daily Living Scale; OC, Osteocalcin; PSM, Physical Self-Maintenance Scale; PTH, Parathyroid hormone; TRAP 5b, Tartrate-resistant acid phosphatase 5b; /d, per day; 25(OH)D, 25-hydroxyvitamin D. **OC reference range for males of 9.6-40.8 ng/ml and for postmenopausal women 12.8-55.0 ng/ml; CTX reference range is 0.020 ng/mL to 3.380 ng/ml; BAP reference range**



15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

for males of 5.7-32.9 µg/ml and for postmenopausal women 5.5-27.1 µg/ml; TRAP 5b reference range for males of 55-79 ng/ml and for postmenopausal women 41-81 ng/ml; Intact PTH measurement range of 1.2 – 5000 pg/ml. and 25(OH)D detection range 7.5-624 nmol/L.

**Table 3.** Comparison of bone mineral density (BMD), biomarkers of bone health and mean frailty measures across frequencies of daily yogurt intake in males in the TUDA cohort study<sup>a</sup>

Variable	Tertile of daily yogurt intakes								
	Non-consumer			Low consumer			High consumer		
	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI
		(0.0 daily / <once per week/never) <i>n</i> = 680			(0.29 daily / 2-3 times per week) <i>n</i> = 392			(1.00 daily / >once per day) <i>n</i> = 333	
<b>BMD region,<sup>b</sup> g/cm<sup>2</sup></b>									
Total Hip	339	1.041	1.026 – 1.056	239	1.057	1.039 – 1.075	185	1.058	1.038 – 1.079
Femoral Neck	339	0.925	0.911 – 0.939	239	0.945	0.928 – 0.962	185	0.947	0.928 – 0.966
Vertebral	271	1.207	1.183 – 1.231	180	1.258*	1.229 – 1.287	134	1.235	1.201 – 1.269
<b>Bone health biomarkers</b>									
CTX, <sup>b</sup> ng/ml	306	0.27	0.25 – 0.29	231	0.27	0.25 – 0.29	169	0.27	0.25 – 0.29
OC, <sup>b</sup> ng/ml	305	14.7	13.9 – 15.7	231	14.9	13.9 – 15.8	169	15.0	13.8 – 16.2
BAP, <sup>b</sup> µg/L	305	16.2	15.3 – 17.1	231	15.7	14.6 – 16.7	168	16.2	15.0 – 17.5
TRAP 5b, <sup>b</sup> µg/L	306	2.96	2.87 – 3.06	231	2.89	2.78 – 3.00	169	2.69**	2.56 – 2.82
25(OH)D, <sup>c</sup> nmol/L	443	41.8	39.7 – 44.0	243	49.3**	46.3 – 52.2	205	47.6*	44.4 – 50.8
PTH, <sup>b</sup> pg/ml	527	45.2	42.7 – 47.8	310	43.7	40.4 – 47.0	258	47.9	44.3 – 51.5
<b>Physical function measures<sup>c</sup></b>									
Timed up and Go, (sec)	627	13.3	12.6 – 13.9	361	12.0*	11.1 – 12.8	302	12.7	11.8 – 13.6
IADL	636	24.6	24.3 – 24.9	372	25.4*	25.0 – 25.8	309	24.7	24.4 – 25.2
PSM	647	23.2	23.1 – 23.4	374	23.4	23.2 – 23.5	315	23.2	23.0 – 23.4

<sup>a</sup> Values are estimated marginal means (95% CI) adjusted for multiple covariates. Differences in means were assessed by pair-wise comparisons **and adjusted for multiple comparisons: Bonferroni correction.** \*,\*\*Different from the lowest yogurt intake tertile: \**P*<0.05, \*\**P*<0.0001. Non-consumer frequency range (0 – 0.07 units / <once per week/never); low consumer frequency range (>0.07- 0.50 units / >once per week to 3-4 times per week); high consumer frequency range (>0.50 – 2.00 units / >3-4 times per week to twice per day).

<sup>b</sup>Adjusted for age, education, BMI, GFR, physical activity, total daily serving milk (glass only), total daily serving of cheese, calcium or vitamin D supplements (Participants receiving medications that could affect bone metabolism were removed from the analysis).

<sup>c</sup>Adjusted for age, BMI, total daily serving milk (glass only), total daily serving of cheese; (participants receiving vitamin D supplements were removed from the 25(OH)D analysis). Abbreviations: BAP, Bone specific alkaline phosphatase; CTX, C-terminal telopeptides of type I collagen; IADL, Instrumental Activities of Daily Living Scale; OC, Osteocalcin; PSM, Physical Self-Maintenance Scale; PTH, Parathyroid hormone; TRAP 5b, Tartrate-resistant acid phosphatase 5b; /d, per day; 25(OH)D, 25-hydroxyvitamin D. **OC reference range for males of 9.6-40.8 ng/ml and for postmenopausal women 12.8-55.0 ng/ml; CTX reference range is 0.020 ng/mL to 3.380 ng/ml; BAP reference range**

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

for males of 5.7-32.9 µg/ml and for postmenopausal women 5.5-27.1 µg/ml; TRAP 5b reference range for males of 55-79 ng/ml and for postmenopausal women 41-81 ng/ml; Intact PTH measurement range of 1.2 – 5000 pg/ml. and 25(OH)D detection range 7.5-624 nmol/L.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 4.** Yogurt consumption as a predictor of markers of bone health and physical function in the TUDA cohort study<sup>a</sup>

	Total Hip BMD, <sup>b</sup>	Femoral Neck BMD, <sup>b</sup>	Vertebral BMD, <sup>b</sup>	TRAP 5b, <sup>b</sup>	Timed up and go, <sup>c</sup>	IADL <sup>c</sup>	PSM <sup>c</sup>
	g/cm <sup>2</sup>	g/cm <sup>2</sup>	g/cm <sup>2</sup>	µg/L	sec		
<b>Variable</b>	<b>β</b>	<b>β</b>	<b>β</b>	<b>β</b>	<b>β</b>	<b>β</b>	<b>β</b>
Total sample	0.015 (0.007)	0.023 (0.006)	0.026 (0.011)	-0.118 (0.055)	-0.599 (0.252)	0.220 (0.115)	0.084 (0.054)
P-value	0.015	<0.0001	0.016	0.032	0.018	0.056	0.121
Female only	0.024 (-0.008)	0.031 (0.007)	0.034 (0.012)	0.015 (-0.077)	-0.641 (0.277)	0.272 (0.126)	0.157 (0.065)
P-value	0.002	<0.0001	0.005	0.847	0.021	0.031	0.016
Male only	0.004 (-0.013)	0.009 (-0.012)	0.012 (-0.021)	-0.292 (0.080)	-0.496 (-0.543)	0.06 (-0.248)	-0.141 (-0.096)
P-value	0.761	0.431	0.557	<0.0001	0.361	0.809	0.144

<sup>a</sup>Values are unstandardized Beta (β) coefficients (standard error) derived from a hierarchical multiple regression analysis.

<sup>b</sup>Adjustment for age, gender (total sample only), education, BMI, smoking status, alcohol consumption, vitamin D or calcium supplement use, 25(OH)D, GFR, physical activity, total daily serving milk (glass only), total daily serving of cheese, total daily serving of meat (red meat and poultry), total daily serving of fish (white and oily) and total daily serving of eggs. (Participants receiving medications that could affect bone metabolism were removed from the analysis).

<sup>c</sup>Adjustment for age, gender (total sample only), BMI, total daily serving milk (glass only), total daily serving of cheese, total daily serving of meat (red meat and poultry), total daily serving of fish (white and oily) and total daily serving of eggs.

Abbreviations: IADL, Instrumental Activities of Daily Living Scale; PSM, Physical Self-Maintenance Scale; TRAP 5b, Tartrate-resistant acid phosphatase 5b; 25(OH)D, 25-hydroxyvitamin D.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 5.** The predictors of bone health status of females within the TUDA cohort study<sup>a</sup>

Variable	Odds ratio [95% CI]	P Value	Odds ratio [95% CI]	P Value
	<i>Osteopenia vs Normal</i>		<i>Osteoporosis vs Normal</i>	
	(n = 411 vs 266)		(n = 360 vs 266)	
Age (y)	1.02 [0.99-1.06]	0.073	1.04 [1.00-1.08]	0.018
BMI (kg/m <sup>2</sup> )	0.93 [0.89-0.96]	<0.0001	0.79 [0.76-0.83]	<0.0001
25(OH)D (nmol/L)	0.99 [0.99-1.00]	0.869	1.00 [0.99-1.01]	0.148
Vitamin D supplement user	0.51 [0.34-0.76]	0.001	0.41 [0.26-0.64]	<0.0001
PTH (pg/mL)	1.00 [0.99-1.01]	0.322	1.00 [0.99-1.01]	0.068
Education (yrs)	0.97 [0.91-1.03]	0.360	0.95 [0.88-1.02]	0.181
Non-Smoker <sup>b</sup>	0.89 [0.50-1.58]	0.710	0.61 [0.34-1.12]	0.116
Non-Alcohol user <sup>c</sup>	0.85 [0.60-1.20]	0.372	1.06 [0.72-1.57]	0.740
Physical activity: Yes <sup>d</sup>	0.89 [0.56-1.43]	0.645	0.61 [0.35-1.56]	0.078
GFR (ml/min)	0.99 [0.98-1.00]	0.810	1.00 [0.99-1.01]	0.601
Daily yogurt serving	0.69 [0.49-0.96]	0.032	0.61 [0.42-0.89]	0.012
Daily milk serving	0.77 [0.53-1.11]	0.167	0.75 [0.49-1.13]	0.175
Daily cheese serving	1.05 [0.65-1.69]	0.831	1.18 [0.70-2.00]	0.516
Daily meat serving	0.91 [0.58-1.41]	0.676	1.16 [0.70-1.92]	0.546
Daily fish serving	0.94 [0.39-2.26]	0.892	0.48 [0.17-1.35]	0.168
Daily egg serving	1.24 [0.63-2.46]	0.528	1.66 [0.76-3.61]	0.201

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

<sup>a</sup>Values are odds ratios (95 % CI lower and upper) derived from a multi-nominal logistic regression analysis. Overall reference category is normal bone health based on the WHO definition of osteopenia and osteoporosis [24] using the combination of total hip, femoral neck or vertebral BMD where available. Participants receiving medications that could affect bone metabolism were removed from the analysis.

<sup>b</sup>Reference is smoker

<sup>c</sup>Reference is user.

<sup>d</sup>Reference is physical activity: No in the last two weeks

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 6.** The predictors of bone health status of males within the TUDA cohort study<sup>1</sup>

Variable	Odds ratio [95% CI]	P Value	Odds ratio [95% CI]	P Value
	<i>Osteopenia vs Normal</i>		<i>Osteoporosis vs Normal</i>	
	<i>(n = 332 vs 315)</i>		<i>(n = 104 vs 315)</i>	
Age (y)	0.98 [0.95-1.01]	0.318	1.01 [0.96-1.06]	0.643
BMI (kg/m <sup>2</sup> )	0.96 [0.91-1.01]	0.128	0.80 [0.73-0.87]	<0.0001
25(OH)D (nmol/L)	1.00 [0.99-1.00]	0.703	0.99 [0.98-1.00]	0.213
Vitamin D supplement user	0.79 [0.52-1.21]	0.286	0.40 [0.22-0.72]	0.003
PTH (pg/mL)	1.00 [0.99-1.01]	0.238	1.01 [1.00-1.01]	0.026
Education (yrs)	0.95 [0.91-1.00]	0.106	0.96 [0.88-1.04]	0.389
Non-Smoker <sup>3</sup>	0.58 [0.32-1.07]	0.085	0.39 [0.18-0.85]	0.019
Non-Alcohol user <sup>2</sup>	0.77 [0.53-1.11]	0.170	0.94 [0.54-1.64]	0.846
Physical activity: Yes <sup>4</sup>	0.59 [0.37-0.92]	0.022	1.09 [0.51-2.35]	0.812
GFR (ml/min)	0.98 [0.97-0.99]	0.016	0.98 [0.97-1.00]	0.128
Daily yogurt serving	0.88 [0.59-1.30]	0.537	0.48 [0.24-0.96]	0.038
Daily milk serving	0.84 [0.58-1.19]	0.338	1.58 [0.99-2.52]	0.055
Daily cheese serving	1.79 [1.11-2.88]	0.016	2.36 [1.16-4.82]	0.018
Daily meat serving	0.64 [0.42-0.99]	0.045	1.10 [0.58-2.07]	0.756
Daily fish serving	0.74 [0.30-1.81]	0.522	0.69 [0.17-2.85]	0.617
Daily egg serving	1.27 [0.69-2.32]	0.439	1.16 [0.47-2.86]	0.740

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

<sup>1</sup>Values are odds ratios (95 % CI lower and upper) derived from a multi-nominal logistic regression analysis. Overall reference category is normal bone health based on the WHO definition of osteopenia and osteoporosis [24] using the combination of total hip, femoral neck or vertebral BMD where available. Participants receiving medications that could affect bone metabolism were removed from the analysis.

<sup>2</sup>Reference is user.

<sup>3</sup>Reference is smoker.

<sup>4</sup>Reference is physical activity: No in the last two weeks



1 **References**

- 2
- 3 1. Cooper C (1999) Epidemiology of osteoporosis. *Osteoporos Int* 9(8):2-8.
- 4
- 5
- 6
- 7 4 2. Ström O, Borgström F, Kanis AJ, Compston J, Cooper C, McCloskey EV, Jönsson B (2011)
- 8
- 9 5 Osteoporosis: Burden, healthcare provision and opportunities in the EU. *Arch Osteoporos*
- 10
- 11 6 6(1):59-155.
- 12
- 13
- 14
- 15 7 3. Curtis JR, Blume SW (2011) Medical costs of osteoporosis in the elderly Medicare
- 16
- 17 8 population. *Osteoporos Int* 22(6):1835-1844.
- 18
- 19
- 20
- 21 9 4. Holroyd C, Cooper C, Dennison E (2008) Epidemiology of osteoporosis. *Best Pract Res Clin*
- 22
- 23 10 *Endoc Meta* 22(5):671-685.
- 24
- 25
- 26
- 27 11 5. Hernlund E, Svedbom A, Ivergård M, Compston J, Cooper C, Stenmark J, McCloskey EV,
- 28
- 29 12 Jönsson B, Kanis JA (2013) Osteoporosis in the European Union: medical management,
- 30
- 31 13 epidemiology and economic burden. *Arch Osteoporos* 8(1-2):1-15.
- 32
- 33
- 34 14 6. Becker DJ, Kilgore ML, Morrisey MA (2010) The societal burden of osteoporosis. *Curr*
- 35
- 36 15 *Rheumatol Rep* 2(3):186-191.
- 37
- 38
- 39
- 40 16 7. Mitchell BD, Yerges-Armstrong LM (2011) The genetics of bone loss: challenges and
- 41
- 42 17 prospects. *J Clin Endocrinol Metab* 96(5):1258-1268.
- 43
- 44
- 45
- 46 18 8. Greco EA, Fornari R, Rossi F et al (2010) Is obesity protective for osteoporosis? Evaluation
- 47
- 48 19 of bone mineral density in individuals with high body mass index. *Int J Clin Pract* 64(6):817-
- 49
- 50 20 820.
- 51
- 52
- 53 21 9. Rizzoli R (2008) Nutrition: its role in bone health. *Best Pract Res Clin Endocrinol Metab*
- 54
- 55 22 22(5):813-829.
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65


- 23 10. Prentice AM (2014) Dairy products in global public health. *Am J Clin Nutr* 99(5):1212-  
24 1216.
- 25 11. Rice BH, Quann EE, Miller GD (2013) Meeting and exceeding dairy recommendations:  
26 effects of dairy consumption on nutrient intakes and risk of chronic disease. *Nutr Rev* 71(4):209-  
27 223.
- 28 12. Heaney, RP (2013) Dairy intake, dietary adequacy and lactose intolerance. *Adv Nutr*  
29 4(2):151-156.
- 30 13. Clarke M, Ward M, Strain JJ, Hoey L, Dickey W, McNulty H (2014) B-vitamins and bone  
31 in health and disease: the current evidence. *Proc Nutr Soc* 73(2):330-339.
- 32 14. Rizzoli R, Abraham C, Brandi ML (2014) Nutrition and bone health: turning knowledge  
33 and beliefs into healthy behaviour. *Curr Med Res Opin* 30(1):131-141.
- 34 15. Heaney RP. Dairy and bone health (2009) *J Am Coll of Nutr* 28(Sup 1):82-90.
- 35 16. Dietary guidelines for Americans. Dietary Guidelines Advisory Committee Report. 6. U.S.  
36 Department of Health and Human Services and U.S. Department of Agriculture, U.S.  
37 Government; Washington DC: 2005.
- 38 17. Sahni S, Tucker KL, Kiel DP, Quach L, Casey VA, Hannan MT (2013) Milk and yogurt  
39 consumption are linked with higher bone mineral density but not with hip fracture: the  
40 Framingham Offspring Study. *Arch Osteoporos* 8(1-2):1-9.
- 41 19. Laird E, McNulty H, Ward M et al (2014) Vitamin D deficiency is associated with  
42 inflammation in older Irish adults. *J Clin Endocrinol Metab* 99(5):1807-1815.
- 43 20. McCarroll K, Beirne A, Casey M et al (2015) Determinants of 25-hydroxyvitamin D in  
44 older Irish adults. *Age Ageing* 44(5):847-853.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
21. Molloy AM, Pangilinan F, Mills JL et al (2016) A Common Polymorphism in HIBCH Influences Methylmalonic Acid Concentrations in Blood Independently of Cobalamin. *Am J Hum Genet* 98(5):869-882.
22. Laird E, Casey MC, Ward M et al (2016) Dairy intakes in older Irish adults and effects on vitamin micronutrient status: Data from the TUDA study. *J Nutr Health Aging*. In Press
23. Irish Universities Nutrition Alliance. The Irish food portion sizes database (1<sup>st</sup> edition) 2013.
24. World Health Organization. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis: report of a WHO study group (meeting held in Rome from 22 to 25 June 1992).
25. Bischoff HA, Stähelin HB, Monsch AU et al (2003) Identifying a cut-off point for normal mobility: a comparison of the timed 'up and go' test in community-dwelling and institutionalised elderly women. *Age Ageing* 32(3):315-320.
26. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist* 1969;9(3):179-186.
27. Hochberg MC, Greenspan S, Wasnich RD, Miller P, Thompson DE, Ross PD (2002) Changes in bone density and turnover explain the reductions in incidence of non-vertebral fractures that occur during treatment with anti-resorptive agents. *J Clin Endocrinol Metab* 87(4):1586-1592.
28. Henriksen K, Tanko LB, Qvist P, Delmas PD, Christiansen C, Karsdal MA (2007) Assessment of osteoclast number and function: application in the development of new and improved treatment modalities for bone diseases. *Osteoporos Int* 18(5):681-685
29. Bonjour JP, Benoit V, Payen F, Kraenzlin M (2013) Consumption of yogurts fortified in vitamin D and calcium reduces serum parathyroid hormone and markers of bone resorption: a

- 69 double-blind randomized controlled trial in institutionalized elderly women. *J Clin Endocrinol*  
70 *Metab* 98(7):2915-2921.
- 71 30. Schoene D, Wu SM, Mikolaizak AS, Menant JC, Smith ST, Delbaere K, Lord SR (2013)  
72 Discriminative Ability and Predictive Validity of the Timed Up and Go Test in Identifying  
73 Older People Who Fall: Systematic Review and Meta-Analysis. *J Am Geriatr Soc* 61(2):202-  
74 208.
- 75 31. Lana A, Rodriguez-Artalejo F, Lopez-Garcia E (2015) Dairy consumption and risk of  
76 frailty in older adults: A prospective cohort study. *J Am Geriatr Soc* 63(9):1852-1860.
- 77 32. Radavelli-Bagatini S, Zhu K, Lewis JR, Dhaliwal SS, Prince RL (2013) Association of  
78 dairy intake with body composition and physical function in older community-dwelling  
79 women. *J Acad Nutr Diet* 113(12):1669-1674.
- 80 33. Matteini AM, Walston JD, Fallin MD et al (2008) Markers of B-vitamin deficiency and  
81 frailty in older women. *J Nutr Health Aging* 12(5):303-308.
- 82 34. Shardell M, Hicks GE, Miller RR, Kritchevsky S, Andersen D, Bandinelli S, Cherubini A,  
83 Ferrucci L (2009) Association of low vitamin D levels with the frailty syndrome in men and  
84 women. *J Gerontol A Biol Sci Med Sci* 64(1):69-75.
- 85 35. Wang H, Livingston KA, Fox CS, Meigs JB, Jacques PF (2013) Yogurt consumption is  
86 associated with better diet quality and metabolic profile in American men and women. *Nutr*  
87 *Res* 33(1):18-26.
- 88
- 89 36. Mistura L, D'Addezio L, Sette S, Piccinelli R, Turrini A (2016) Diet quality of Italian  
90 yogurt consumers: an application of the probability of adequate nutrient intake score  
91 (PANDiet). *Int J Food Sci Nut* 67(3):232-238.
- 92
- 93 37. Parvaneh K, Jamaluddin R, Karimi G, Erfani R (2014) Effect of probiotics supplementation  
94 on bone mineral content and bone mass density. *ScientificWorldJournal* 22:595962.

95 38. Parvaneh K, Ebrahimi M, Sabran MR, Karimi G, Hwei AN, Abdul-Majeed S, Ahmad Z5,  
1 96 Ibrahim Z, Jamaluddin R (2015) Probiotics (*Bifidobacterium longum*) Increase Bone Mass  
2  
3 97 Density and Upregulate Sparc and Bmp-2 Genes in Rats with Bone Loss Resulting from  
4  
5 98 Ovariectomy. *BioMed Res Int* 897639  
6  
7  
8  
9 99 39. Korhonen H, Pihlanto A (2006) Bioactive peptides: production and functionality. *Int Dairy*  
10  
11 100 *J* 16(9):945-960.  
12  
13  
14  
15 101 40. Rizzoli R, Bonjour JP (2004) Dietary protein and bone health. *J Bone Miner Res* 19(4):527-  
16  
17 102 531.  
18  
19  
20  
21 103 41. Darling AL, Millward DJ, Torgerson DJ, Hewitt CE, Lanham-New SA (2009) Dietary  
22  
23 104 protein and bone health: a systematic review and meta-analysis. *Am J Clin Nutr* 90(6):1674-  
24  
25 105 1692.  
26  
27  
28 106 42. Narva M, Collin M, Lamberg-Allardt C, Kärkkäinen M, Poussa T, Vapaatalo H, Korpela  
29  
30 107 R (2004) Effects of long-term intervention with *Lactobacillus helveticus*-fermented milk on  
31  
32 108 bone mineral density and bone mineral content in growing rats. *Ann Nutr Metab* 48(4):228-  
33  
34 109 234.  
35  
36  
37  
38 110 43. Tu MY, Chen HL, Tung YT, Kao CC, Hu FC, Chen CM (2015) Short-Term Effects of  
39  
40 111 Kefir-Fermented Milk Consumption on Bone Mineral Density and Bone Metabolism in a  
41  
42 112 Randomized Clinical Trial of Osteoporotic Patients. *PLoS one* 10(12):e0144231.  
43  
44  
45 113 44. Gilbert JA, Bendsen NT, Tremblay A, Astrup A (2011) Effect of proteins from different  
46  
47 114 sources on body composition. *Nutr Metab and Cardiovasc Dis* 21:B16-31.  
48  
49  
50  
51 115 45. Moreno LA, Bel-Serrat S, Santaliestra-Pasías A, Bueno G (2015) Dairy products, yogurt  
52  
53 116 consumption, and cardiometabolic risk in children and adolescents. *Nutr Rev* 73(suppl 1):8-  
54  
55 117 14.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 118 46. Steves CJ, Bird S, Williams FM, Spector TD (2016) The Microbiome and Musculoskeletal  
1 119 Conditions of Aging: A Review of Evidence for Impact and Potential Therapeutics. *J Bone*  
2 120 *Miner Res* 1;31(2):261-9.  
3  
4  
5  
6  
7 121 47. Stanhope KL (2016) Sugar consumption, metabolic disease and obesity: The state of the  
8 122 controversy. *Crit Rev Clin Lab Sci* 53(1):52-67.  
9  
10  
11  
12  
13 123 48. United States Department of Agriculture (USDA). National Nutrient Database for Standard  
14 124 Reference Release 8. Yogurt, fruit, low fat, 9 grams protein per 8 ounce, Basic Report 01120.  
15 125 Accessed 4<sup>th</sup> July 2016.  
16  
17  
18  
19  
20  
21 126 49. United States Department of Agriculture (USDA). National Nutrient Database for Standard  
22 127 Reference Release 8. Yogurt, Greek, plain, low fat, Basic Report 01287. Accessed 4<sup>th</sup> July  
23 128 2016.  
24  
25  
26  
27  
28 129 50. Rizzoli R (2014) Dairy products, yogurts, and bone health. *Am J Clin Nutr* 99(5):1256S-  
29 130 1262S.  
30  
31  
32  
33  
34 131 51. Rozenberg S, Body JJ, Bruyère O et al (2016) Effects of Dairy Products Consumption on  
35 132 Health: Benefits and Beliefs - A Commentary from the Belgian Bone Club and the European  
36 133 Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and  
37 134 Musculoskeletal Diseases. *Calcif Tissue Int* 98(1):1-7.  
38  
39  
40  
41  
42  
43 135 52. Teucher B, Dainty JR, Spinks CA et al (2008) Sodium and bone health: Impact of  
44 136 moderately high and low salt intakes on calcium metabolism in postmenopausal women. *J*  
45 137 *Bone Miner Res* 23(9):1477-1485.  
46  
47  
48  
49  
50  
51 138 53. Bonjour JP, Benoit V, Rousseau B, Souberbielle JC (2012) Consumption of vitamin D-and  
52 139 calcium-fortified soft white cheese lowers the biochemical marker of bone resorption TRAP  
53 140 5b in postmenopausal women at moderate risk of osteoporosis fracture. *J Nutr* 142(4):698-703.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



**Supplemental Figure 1.** Study Design for yogurt intake frequency analysis within the TUDA cohort study

Abbreviations: BAP, Bone specific alkaline phosphatase; CTX, C-terminal telopeptides of type I collagen; OC, Osteocalcin; Parathyroid hormone; TRAP 5b, Tartrate-resistant acid phosphatase 5b; 25(OH)D, 25-hydroxyvitamin D.

**Supplemental Table 1.** Calculation of daily yogurt and dairy intakes in the TUDA cohort study<sup>1</sup>

Questionnaire Dairy Intake Frequency	Daily Yogurt intake frequency	Daily Milk intake frequency	Daily Cheese intake frequency
Twice per day	2	2	2
Once per day	1	1	1
5-6 times per week	0.785	0.785	0.785
3-4 times per week	0.5	0.5	0.5
1-2 times per week	0.21	0.21	0.21
<Once per week	0.07	0.07	0.07

<sup>1</sup>Values are self-reported intakes derived from a modified food frequency questionnaire (FFQ)



**Supplemental Table 2.** Comparison of bone mineral density (BMD), biomarkers of bone health and mean frailty measures across frequencies of daily milk intake in females in the TUDA cohort study<sup>a</sup>

Variable	Tertile of daily milk intakes								
	Non-consumer			Low consumer			High consumer		
	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI
		Mean milk frequency (0.0 daily / <once per week/never) <i>n</i> = 1,746			Mean milk frequency (0.18 daily / 1-2 times per week) <i>n</i> = 305			Mean milk frequency (1.00 daily / >once per day) <i>n</i> = 854	
BMD region, <sup>b</sup> g/cm <sup>2</sup>									
Total Hip	673	0.907	0.898 – 0.916	133	0.903	0.883 – 0.923	252	0.902	0.888 – 0.917
Femoral Neck	672	0.845	0.837 – 0.853	133	0.838	0.820 – 0.857	250	0.838	0.824 – 0.852
Vertebral	545	1.029	1.015 – 1.043	99	1.026	0.993 – 1.059	197	1.008	0.985 – 1.032
Bone health biomarkers									
CTX, <sup>b</sup> ng/ml	465	0.33	0.32 – 0.35	107	0.34	0.31 – 0.37	184	0.3`	0.29 – 0.34
OC, <sup>b</sup> ng/ml	464	19.2	18.3 – 20.0	107	18.5	16.7 – 20.2	184	18.4	17.1 – 19.8
BAP, <sup>b</sup> µg/L	466	17.8	17.1 – 18.4	107	17.3	16.0 – 18.6	183	18.3	17.3 – 19.3
TRAP 5b, <sup>b</sup> µg/L	566	3.31	3.22 – 3.40	107	3.22	3.04 – 3.41	184	3.21	3.07 – 3.35
25(OH)D, <sup>c</sup> nmol/L	691	43.2	41.5 – 44.8	129	41.8	38.0 – 45.7	301	41.5	39.0 – 44.0
PTH, <sup>b</sup> pg/ml	942	46.7	44.9 – 48.6	167	49.0	44.6 – 53.3	420	49.4	46.7 – 52.1
Physical function measures <sup>c</sup>									
Timed up and Go, (sec)	1,688	12.8	12.5 – 13.1	295	12.5	11.8 – 13.3	808	13.7*	13.2 – 14.2
IADL	1,706	24.9	24.7 – 25.0	298	24.8	24.5 – 25.2	831	24.2*	24.0 – 24.4
PSM	1,738	23.1	23.0 – 23.2	302	22.9	22.8 – 23.1	847	22.9*	22.8 – 23.0

<sup>a</sup>Values are estimated marginal means (95% CI) adjusted for multiple covariates. Differences in means were assessed by pair-wise comparisons and adjusted for multiple comparisons: Bonferroni correction. \*,\*\*Different from the lowest milk intake tertile: \**P*<0.05, \*\**P*<0.0001. Non-consumer frequency range (0 – 0.07 units / <once per week/never); low consumer frequency range (>0.07- 0.50 units / >once per week to 1-2 times per week); high consumer frequency range (0.50 – 2.00 units / >3-4 times per week to twice per day).

<sup>b</sup>Adjusted for age, education, BMI, GFR, physical activity, total daily serving of yogurt, total daily serving of cheese, calcium or vitamin D supplements (Participants receiving medications that could affect bone metabolism were removed from the analysis).

<sup>c</sup>Adjusted for age, BMI, total daily serving yogurt, total daily serving of cheese; (participants receiving vitamin D supplements were removed from the 25(OH)D analysis). Abbreviations: BAP, Bone specific alkaline phosphatase; CTX, C-terminal telopeptides of type I collagen; IADL, Instrumental Activities of Daily Living Scale; OC, Osteocalcin; PSM, Physical Self-Maintenance Scale; PTH, Parathyroid hormone; TRAP 5b, Tartrate-resistant acid phosphatase 5b; /d, per day; 25(OH)D, 25-hydroxyvitamin D. OC reference range for males of 9.6-40.8 ng/ml and for postmenopausal women 12.8-55.0 ng/ml; CTX reference range is 0.020 ng/mL to 3.380 ng/mL; BAP reference range

for males of 5.7-32.9 µg/ml and for postmenopausal women 5.5-27.1 µg/ml; TRAP 5b reference range for males of 55-79 ng/ml and for postmenopausal women 41-81 ng/ml; Intact PTH measurement range of 1.2 – 5000 pg/ml. and 25(OH)D detection range 7.5-624 nmol/L.

**Supplemental Table 3.** Comparison of bone mineral density (BMD), biomarkers of bone health and mean frailty measures across frequencies of daily cheese intake in females in the TUDA cohort study<sup>a</sup>

Variable	Tertile of daily cheese intakes								
	Non-consumer			Low consumer			High consumer		
	Mean cheese frequency (0.0 daily / <once per week/never)			Mean cheese frequency (0.33 daily / 2-3 times per week)			Mean cheese frequency (0.95 daily / >once per day)		
	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI
		<i>n</i> = 602			<i>n</i> = 1,673			<i>n</i> = 631	
BMD region, <sup>b</sup> g/cm <sup>2</sup>									
Total Hip	241	0.903	0.888 – 0.918	614	0.908	0.899 – 0.918	202	0.898	0.882 – 0.914
Femoral Neck	241	0.840	0.826 – 0.854	612	0.846	0.838 – 0.855	202	0.833	0.818 – 0.849
Vertebral	187	1.015	0.991 – 1.039	489	1.026	1.011 – 1.041	165	1.026	1.001 – 1.052
Bone health biomarkers									
CTX, <sup>b</sup> ng/ml	192	0.32	0.30 – 0.34	434	0.33	0.31 – 0.35	130	0.34	0.31 – 0.37
OC, <sup>b</sup> ng/ml	192	19.0	17.7 – 20.4	438	18.7	17.8 – 19.6	130	19.4	17.8 – 21.0
BAP, <sup>b</sup> µg/L	192	17.9	16.9 – 18.8	434	18.0	17.3 – 18.6	130	17.2	16.0 – 18.4
TRAP 5b, <sup>b</sup> µg/L	192	3.25	3.11 – 3.39	435	3.28	3.19 – 3.37	130	3.29	3.13 – 3.46
25(OH)D, <sup>c</sup> nmol/L	221	40.6	37.7 – 43.5	532	43.4	41.5 – 45.3	163	39.9	36.5 – 43.3
PTH, <sup>b</sup> pg/ml	335	47.0	44.0 – 50.1	893	46.9	45.1 – 48.8	301	50.7	47.5 – 54.0
Physical function measures <sup>c</sup>									
Timed up and Go, (sec)	577	13.3	12.8 – 13.9	1,603	12.9	12.6 – 13.2	611	13.2	12.6 – 13.7
IADL	583	24.5	24.3 – 24.7	1,634	24.8	24.6 – 24.9	618	24.6	24.4 – 24.9
PSM	596	23.0	22.9 – 23.2	1,662	23.0	23.0 – 23.1	629	23.0	22.9 – 23.1

<sup>1</sup>Values are estimated marginal means (95% CI) adjusted for multiple covariates. Differences in means were assessed by pair-wise comparisons and adjusted for multiple comparisons: Bonferroni correction. No significant difference observed from the lowest cheese intake tertile to the highest. Non-consumer frequency range (0 – 0.07 units / <once per week/never); low consumer frequency range (>0.07- 0.50 units / >once per week to 3-4 times per week); high consumer frequency range (>0.50 – 2.00 units / >3-4 times per week to twice per day).

<sup>2</sup>Adjusted for age, education, BMI, GFR, physical activity, total daily serving milk (glass only), total daily serving of yogurt, calcium or vitamin D supplements (Participants receiving medications that could affect bone metabolism were removed from the analysis).

<sup>3</sup>Adjusted for age, BMI, total daily serving milk (glass only), total daily serving of cheese; (participants receiving vitamin D supplements were removed from the 25(OH)D analysis). Abbreviations: BAP, Bone specific alkaline phosphatase; CTX, C-terminal telopeptides of type I collagen; IADL, Instrumental Activities of Daily Living Scale; OC, Osteocalcin; PSM, Physical Self-Maintenance Scale; PTH, Parathyroid hormone; TRAP 5b, Tartrate-resistant acid phosphatase 5b; /d, per day; 25(OH)D, 25-hydroxyvitamin D. OC reference range for males of 9.6-40.8 ng/ml and for postmenopausal women 12.8-55.0 ng/ml; CTX reference range is 0.020 ng/mL to 3.380 ng/mL; BAP reference range

for males of 5.7-32.9 µg/ml and for postmenopausal women 5.5-27.1 µg/ml; TRAP 5b reference range for males of 55-79 ng/ml and for postmenopausal women 41-81 ng/ml; Intact PTH measurement range of 1.2 – 5000 pg/ml. and 25(OH)D detection range 7.5-624 nmol/L.

**Supplemental Table 4.** Comparison of bone mineral density (BMD), biomarkers of bone health and mean frailty measures across frequencies of daily cheese intake in males in the TUDA cohort study<sup>a</sup>

Variable	Tertile of daily cheese intakes								
	Non-consumer			Low consumer			High consumer		
	Mean cheese frequency (0.0 daily / <once per week/never)			Mean cheese frequency (0.34 daily / 2-3 times per week)			Mean cheese frequency (0.95 daily / >once per day)		
	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI
		<i>n</i> = 269			<i>n</i> = 835			<i>n</i> = 304	
BMD region, <sup>b</sup> g/cm <sup>2</sup>									
Total Hip	176	1.064	1.043 – 1.088	447	1.048	1.035 – 1.061	140	1.041	1.048 – 1.064
Femoral Neck	176	0.953	0.934 – 0.972	447	0.932	0.920 – 0.944	140	0.929	0.907 – 0.951
Vertebral	129	1.246	1.211 – 1.280	342	1.221	1.200 – 1.242	114	1.234	1.197 – 1.271
Bone health biomarkers									
CTX, <sup>b</sup> ng/ml	173	0.27	0.25 – 0.30	413	0.27	0.25 – 0.28	120	0.27	0.24 – 0.30
OC, <sup>b</sup> ng/ml	173	15.1	13.9 – 16.3	413	15.0	14.3 – 15.8	121	13.8	12.4 – 15.2
BAP, <sup>b</sup> µg/L	172	16.4	15.2 – 17.6	412	15.8	15.0 – 16.6	120	16.3	14.8 – 17.7
TRAP 5b, <sup>b</sup> µg/L	173	2.94	2.81 – 3.06	413	2.86	2.78 – 2.95	120	2.81	2.66 – 2.97
25(OH)D, <sup>c</sup> nmol/L	163	45.7	42.0 – 49.3	473	45.1	43.0 – 47.2	172	46.3	42.8 – 49.3
PTH, <sup>b</sup> pg/ml	221	44.0	40.0 – 47.9	647	46.3	44.0 – 48.6	227	44.2	40.4 – 48.1
Physical function measures <sup>c</sup>									
Timed up and Go, (sec)	255	12.1	11.1 – 13.1	804	12.8	12.2 – 13.3	288	13.2	12.3 – 14.2
IADL	261	25.1	24.6 – 25.5	815	25.0	24.7 – 25.2	298	24.8	24.3 – 25.2
PSM	265	23.4	23.2 – 23.6	828	23.2	23.1 – 23.3	301	23.3	23.1 – 23.5

<sup>a</sup>Values are estimated marginal means (95% CI) adjusted for multiple covariates. Differences in means were assessed by pair-wise comparisons and adjusted for multiple comparisons: Bonferroni correction. No significant difference observed from the lowest cheese intake tertile to the highest. Non-consumer frequency range (0 – 0.07 units / <once per week/never); low consumer frequency range (>0.07- 0.50 units / >once per week to 3-4 times per week); high consumer frequency range (>0.50 – 2.00 units / >3-4 times per week to twice per day).

<sup>b</sup>Adjusted for age, education, BMI, GFR, physical activity, total daily serving milk (glass only), total daily serving of yogurt, calcium or vitamin D supplements (Participants receiving medications that could affect bone metabolism were removed from the analysis).

<sup>c</sup>Adjusted for age, BMI, total daily serving milk (glass only), total daily serving of cheese; (participants receiving vitamin D supplements were removed from the 25(OH)D analysis). Abbreviations: BAP, Bone specific alkaline phosphatase; CTX, C-terminal telopeptides of type I collagen; IADL, Instrumental Activities of Daily Living Scale; OC, Osteocalcin; PSM, Physical Self-Maintenance Scale; PTH, Parathyroid hormone; TRAP 5b, Tartrate-resistant acid phosphatase 5b; /d, per day; 25(OH)D, 25-hydroxyvitamin D. OC reference range for males of 9.6-40.8 ng/ml and for postmenopausal women 12.8-55.0 ng/ml; CTX reference range is 0.020 ng/mL to 3.380 ng/mL; BAP reference range

for males of 5.7-32.9 µg/ml and for postmenopausal women 5.5-27.1 µg/ml; TRAP 5b reference range for males of 55-79 ng/ml and for postmenopausal women 41-81 ng/ml; Intact PTH measurement range of 1.2 – 5000 pg/ml. and 25(OH)D detection range 7.5-624 nmol/L.

**Supplemental Table 5.** Comparison of bone mineral density (BMD), biomarkers of bone health and mean frailty measures across frequencies of daily milk intake in males in the TUDA cohort study<sup>a</sup>

Variable	Tertile of daily milk intakes								
	Non-consumer			Low consumer			High consumer		
	Mean milk frequency (0.0 daily / <once per week/never)			Mean milk frequency (0.18 daily / 2-3 times per week)			Mean milk frequency (0.99 daily / >once per day)		
	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI	<i>n</i>	Mean	95% CI
		<i>n</i> = 768			<i>n</i> = 201			<i>n</i> = 438	
<b>BMD region,<sup>b</sup> g/cm<sup>2</sup></b>									
Total Hip	424	1.050	1.036 – 1.063	132	1.050	1.026 – 1.074	207	1.051	1.032 – 1.071
Femoral Neck	424	0.939	0.926 – 0.951	132	0.936	0.914 – 0.958	207	0.932	0.915 – 0.950
Vertebral	331	1.234	1.213 – 1.256	98	1.216	1.176 – 1.256	156	1.226	1.195 – 1.257
<b>Bone health biomarkers</b>									
CTX, <sup>b</sup> ng/ml	392	0.26	0.25 – 0.28	125	0.27	0.25 – 0.30	189	0.28	0.26 – 0.30
OC, <sup>b</sup> ng/ml	391	15.0	14.3 – 15.8	125	14.3	12.9 – 15.6	189	14.8	13.7 – 15.9
BAP, <sup>b</sup> µg/L	391	16.1	15.2 – 16.9	125	15.9	14.5 – 17.4	188	16.1	14.9 – 17.2
TRAP 5b, <sup>b</sup> µg/L	392	2.85	2.76 – 2.93	125	2.95	2.80 – 3.10	189	2.87	2.75 – 2.99
25(OH)D, <sup>c</sup> nmol/L	486	44.7	42.6 – 46.8	127	46.5	42.4 – 50.6	279	45.5	42.8 – 48.3
PTH, <sup>b</sup> pg/ml	591	45.7	43.3 – 48.1	168	42.0	37.5 – 46.5	336	46.6	43.5 – 49.8
<b>Physical function measures<sup>c</sup></b>									
Timed up and Go, (sec)	739	12.4	11.8 – 13.0	193	11.9	10.8 – 13.0	415	13.7*	12.9 – 14.5
IADL	753	25.2	25.0 – 25.5	197	25.2	24.7 – 25.7	424	24.3**	24.0 – 24.7
PSM	762	23.3	23.2 – 23.4	200	23.4	23.2 – 23.6	432	23.1*	23.0 – 23.2

<sup>a</sup>Values are estimated marginal means (95% CI) adjusted for multiple covariates. Differences in means were assessed by pair-wise comparisons and adjusted for multiple comparisons: Bonferroni correction. \*,\*\*Different from the lowest milk intake tertile: \**P*<0.05, \*\**P*<0.0001. Non-consumer frequency range (0 – 0.07 units / <once per week/never); low consumer frequency range (>0.07- 0.50 units / >once per week to 3-4 times per week); high consumer frequency range (>0.50 – 2.00 units / >3-4 times per week to twice per day).

<sup>b</sup>Adjusted for age, education, BMI, GFR, physical activity, total daily of yogurt, total daily serving of cheese, calcium or vitamin D supplements (Participants receiving medications that could affect bone metabolism were removed from the analysis).

<sup>c</sup>Adjusted for age, BMI, total daily serving milk (glass only), total daily serving of cheese; (participants receiving vitamin D supplements were removed from the 25(OH)D analysis). Abbreviations: BAP, Bone specific alkaline phosphatase; CTX, C-terminal telopeptides of type I collagen; IADL, Instrumental Activities of Daily Living Scale; OC, Osteocalcin; PSM, Physical Self-Maintenance Scale; PTH, Parathyroid hormone; TRAP 5b, Tartrate-resistant acid phosphatase 5b; /d, per day; 25(OH)D, 25-hydroxyvitamin D. OC reference range for males of 9.6-40.8 ng/ml and for postmenopausal women 12.8-55.0 ng/ml; CTX reference range is 0.020 ng/mL to 3.380 ng/mL; BAP reference range

for males of 5.7-32.9 µg/ml and for postmenopausal women 5.5-27.1 µg/ml; TRAP 5b reference range for males of 55-79 ng/ml and for postmenopausal women 41-81 ng/ml; Intact PTH measurement range of 1.2 – 5000 pg/ml. and 25(OH)D detection range 7.5-624 nmol/L.





Click here to access/download  
**Authorship and Disclosure Form**  
OS signature page.pdf

