

Title:

Investigation of the contraction ratio of transversus abdominis and internal oblique muscles during lumbopelvic stability test

Names of the authors:

Patraporn Sitolertpisan¹, Leonard Joseph², Aatit Paungmali¹, Ubon Pirunsan¹, Titikorn Chunchai³

Affiliations of the authors:

¹Department of Physical Therapy, Faculty of Associated Medical Sciences, Chiang Mai University, Chiangmai, Thailand

²School of Health Science, University of Brighton, East Sussex, United Kingdom

³Department of Physiology, Faculty of Medicine, Chiang Mai University, Chiangmai, Thailand

Corresponding author:

Dr Patraporn Sitolertpisan
Assistant Professor in Physical Therapy
Department of Physical Therapy
Faculty of Associated Medical Sciences
Chiang Mai University
Chiang Mai, Thailand
Phone: +66 0-5394-9243
Email: patraporn.s@cmu.ac.th

Abstract:

Background: The clinicians administer lumbopelvic stability to test core stability of lumbar spine and TrA muscle function. While the reduced thickness and contraction ratio of the TrA is well documented among people with nonspecific low back pain (NSLBP), the direct morphological changes of TrA during the lumbopelvic test has not been reported so far.

Purpose: To determine and compare the contraction ratio of deep abdominal muscles (TrA and IO) during the 7 progressive stages of the lumbopelvic stability test (LPST) in healthy individuals.

Methods: Thirty healthy volunteers (15 males and 15 females) aged 21.83 ± 0.46 years participated in an experimental study. The TrA and IO thickness was assessed by ultrasound

imaging (USI) TOSHIBA, Famio8, SSA-530A) at the right side in mid axillary line between 12th rib and iliac crease. Images were taken for 2 trials during each level of LPST (7 levels) as measured by the pressure biofeedback unit (PBU). The thickness of the TrA and IO muscles were measured by Image J program (Image J®, NIH, USA) and the contraction ratio was calculated. A one way repeated measure ANOVA with post hoc analysis using the Bonferroni test at $p < 0.05$ was used to analyse the data.

Results: The contraction ratio of TrA showed a decreasing trend as the LPST was progressed to the 7 levels. Post hoc analysis showed a significant reduction in the contraction ratio of the TrA muscle in the 7th level of the LPST ($F=14.53$, $p=0.001$). However, the contraction ratio of IO remains unchanged during all the 7 levels of LPST.

Conclusion: The TrA muscle responds to the lumbopelvic stability testing by reduced contraction ratio across the 7 levels of the test procedure as measured by the PBU and USI. Further studies are warranted to compare the adaptation of the TrA muscle across 7 levels of the testing among patients with NSLBP.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Title page - Original article

2 Investigation of the contraction ratio of transversus abdominis and internal oblique muscles
3 during lumbopelvic stability test

9 Running title - Contraction ratio of transversus abdominis and internal oblique muscles during
10 lumbopelvic stability test

1 **Abstract**

2 **Background:** The clinicians administer lumbopelvic stability test (LPST) to evaluate core
3 stability of lumbar spine and TrA muscle function. While the reduced thickness and contraction
4 ratio of the TrA was well documented among people with nonspecific low back pain (NSLBP),
5 the direct morphological changes of TrA during the LPST had not been reported so far.

6 **Purpose:** To determine and compare the contraction ratio of deep abdominal muscles (TrA
7 and IO) during the 7 progressive stages of the LPST in healthy individuals.

8 **Methods:** Thirty healthy volunteers (15 males and 15 females) aged 21.83 ± 0.46 years
9 participated in an experimental study. The TrA and IO thickness was assessed by ultrasound
10 imaging (USI) TOSHIBA, Famio8, SSA-530A) at the right side in mid axillary line between
11 12th rib and iliac crease. Images were taken for 2 trials during each level of LPST (7 levels) as
12 measured by the pressure biofeedback unit (PBU). The thickness of the TrA and IO muscles
13 were measured by Image J program (Image J®, NIH, USA) and the contraction ratio was
14 calculated. A one way repeated measure ANOVA with post hoc analysis using the Bonferroni
15 test at $p < 0.05$ was used to analyse the data.

16 **Results:** The contraction ratio of TrA showed a decreasing trend as the LPST was progressed
17 to the 7 levels. Post hoc analysis showed a significant reduction in the contraction ratio of the
18 TrA muscle in the 7th level of the LPST ($F=14.53$, $p=0.001$). However, the contraction ratio
19
20 of IO remains unchanged during all the 7 levels of LPST.

21 **Conclusion:** The TrA muscle responded by reduced contraction ratio across the 7 levels of the
22 LPST as measured by the PBU and USI. Further studies are warranted to compare the
23 adaptation of the TrA muscle across 7 levels of the LPST among patients with NSLBP.

24 **Level of Evidence – 2b**

1 **Key words:** low back pain, musculoskeletal, physiotherapy, rehabilitation, transversus
2 abdominis, ultrasound imaging
3
4
5 **3**
6
7 **4**
8
9
10 **5**
11
12 **6**
13
14 **7**
15
16
17 **8**
18
19 **9**
20
21
22 **10**
23
24 **11**
25
26
27 **12**
28
29 **13**
30
31 **14**
32
33
34 **15**
35
36 **16**
37
38
39 **17**
40
41 **18**
42
43
44 **19**
45
46 **20**
47
48
49 **21**
50
51 **22**
52
53 **23**
54
55
56 **24**
57
58 **25**
59
60
61
62
63
64
65

1 **INTRODUCTION:**

2 Approximately 90% of the pain originating from the lumbar spine are classified as non-specific
3 low back pain (NSLBP) (1). The mechanical overuse and dysfunction of the spine and the
4 surrounding structures such as core muscles were proposed as one of the contributing factors
5 for NSLBP (1). An impaired function of the deep core muscles such as transversus abdominis
6 (TrA) and internal oblique (IO) was reported to affect stability, robustness and movement
7 control of the lumbar spine leading to recurrent low back pain (2). TrA and IO as deep
8 abdominal muscles particularly had received greater attention due to their close proximities
9 resulting in pre-activation prior to the limb movement and thus contributing to the core stability
10 of the lumbopelvic region (3). For example, 20.9% of reduced thickness and the delayed
11 anticipatory onset of TrA muscle during limb loading had been reported among patients with
12 NSLBP (4,5). Recently, the measure of TrA morphometry had been suggested to be a treatment
13 effect modifier for NSLBP patients (6). On the other hand, the thickness of TrA had been
14 studied in terms of gender difference, between rest and contraction, hand dominance and side
15 to side differences among healthy participants to understand the normative function of TrA
16 muscle (7,8). Therefore, many clinicians might consider the assessment and treatment of the
17 core muscles as one of the factors towards the management of NSLBP (9). Although several
18 pieces of evidence were available on TrA muscle anatomy and function, the changes in TrA
19 muscle during the different levels of the LPST through an abdominal draw-in maneuver had
20 never been studied before.

21 The function of the TrA was evaluated by clinicians through an abdominal drawing-in
22 manoeuvre using a pressure biofeedback unit (PBU) (10,11). In theory, the PBU **was** placed
23 under the lumbar spine in crook lying position and the patients **were** instructed to tuck in the
24 abdomen. Any pressure fluctuations greater than 4 mmHg during a standard test using
25 40mmHg **was** considered as poor stability of the lumbar spine. If no fluctuations, the

1 participants performed a unilateral leg lift in the sagittal plane followed by the seven
2 progressive levels of LPST (12). An ability to maintain the registered pressure at 40 mmHg
3 during the testing movement was marked as a successful performance. When the participants
4 were not able to hold 40 ± 4 mmHg in the PBU, the progression of the LPST would be stopped.
5 Although testing the TrA muscle using the PBU was reported to have good reliability of the
6 TrA function (11), the concurrent validity was reported to be poor against the
7 electromyography (13). A recent systematic review concluded that the ability of the test to
8 directly assess the function of TrA had not been answered (14). Perhaps, this was due to the
9 fact that the test did not measure directly the changes in the TrA, instead, it measured the
10 changes in the pressure of the PBU as an indirect evaluation of the TrA muscle contractility.
11 Till now, no studies had investigated the changes in the TrA function during different levels of
12 the LPST as indicated by the different upper and lower limb movements. Furthermore, an
13 absence of any normative understanding of the TrA muscle function among healthy individuals
14 remains a challenge for clinicians to interpret its outcome among low back pain patients.
15 Therefore, the main aim of the study is to investigate the changes in the contraction ratio of the
16 TrA muscle during different levels of the LPST among healthy participants. The study
17 hypothesizes that the contraction ratio of TrA and IO differs significantly during different levels
18 of the LPST. The current study is the first study to report on the contraction ratio of the TrA
19 and IO during the different levels of the LPST. It is
20 important to establish normative data in the healthy population as it becomes a benchmark for
21 any conditions such as NSLBP.

22 **Study design**

23 A cross-sectional study with an experimental design was conducted to investigate the
24 differences in TrA and IO during different levels of the LPST. The details of the study were
25 advertised through posters around a community setting. The participants for the study were

1 recruited through pre-defined study criteria from a community population. The study was
2 carried out in the pain and neuromusculoskeletal research laboratory in a university setting.
3 The study was conducted according to the international ethical standards recommended in the
4 clinical and field science research (15). A university institutional ethics board approved the
5 ethics of the study according to the Declaration of Helsinki.

6 **Participant characteristics**

7 A total of among a total of 30 healthy individuals (15 males and 15 females) participated in the
8 study. Any participants who did not have body aches or pain over the last 3 months were invited
9 to participate in the study. The participants were excluded if they had referred pain or
10 neurological involvement in lower limbs, had any history of past surgery and had any history
11 of injury in the last 3 months. The purpose of the study and the study procedure were explained
12 to all the participants and written informed consent was obtained prior to their participation in
13 the study.

14 **Procedure**

15 **Lumbopelvic stability testing**

16 A pressure biofeedback stabilizer unit (PBU) (Stabilizer Pressure Biofeedback-Chattanooga
17 Group, USA) was used during the test. PBU was reported to be a reliable device to test the
18 lumbopelvic stability. (16). The LPST was performed in supine crook lying position with the
19 participant hip and knee flexed to 70° and 90° respectively through an established protocol (12).

20 Before the test, the PBU was calibrated and pre-tested through an established protocol by
21 loading the unit for 24 hours with 4 kg of weight (17). The PBU unit was considered for use
22 only if the device lost no more than 0.4 mm Hg during the testing period and the same unit was
23 used throughout the study period. After calibration, the PBU was placed beneath the lumbar
24 spine between the levels of the second lumbar spine and the first sacral spine. The participants
25 were instructed to tuck in their naval by drawing in the abdomen. The pressure in the pressure

1 gauge was maintained at 40 mm Hg and the participants were instructed not to exceed more
2 than 4 mm Hg during the abdominal drawing-in manoeuvre. Once the participants were able
3 to maintain the pressure, the tests were progressed into 7 levels in the following order (12);
4 core with alternate hip abduction, core with alternate knee raise, core with both arms adduction,
5 core with both arms extension, core with alternate arm lift, and core with alternate leg lift and
6 finally core with alternate leg and arm lift. The test was stopped when the participants exceeded
7 the pressure more than 40 ± 4 mm Hg and the level of the test was noted as completed.

8 **Real-time ultrasound measurement of TrA**

9 The real-time ultrasound imaging (USI) measurement of the TrA muscle was carried using a
10 B-mode real-time ultrasound imaging unit (TOSHIBA, Fami08, SSA-530A) through an
11 established protocol (18). The ultrasound measurement was carried out simultaneously when
12 the participants performed the different levels of the LPST. The participants were positioned
13 in crook lying with their head and knee supported by pillows. Ultrasonic gel was applied to a
14 5-MHz curvilinear transducer and the transducer was placed in the transverse plane over the
15 midaxillary line at a midpoint between the lower rib and the iliac crest. An experienced
16 physiotherapist who was trained to conduct USI measurement carried out all the measurements.
17 A surface marking was made on the skin using a surface marker to ensure the transducer was
18 placed over the same location for every measurement during the seven levels of the test.

19 The images of TrA thickness were captured and measured using NIH (Bethesda, MD) Image
20 J Software (Version 1.8). The total resting muscle thickness was defined as the distance
21 between the superior border and the inferior border of TrA muscle. Then, the patients were
22 instructed to draw in their abdomen and TrA muscle thickness during contraction was measured
23 during the 7 levels of the test. **Figure 1** shows the USI measurement of the muscle thickness
24 during the LPST. All the measurements were taken at the end of the expiration. The mean
25 thickness of the three measurements on each side was calculated for analysis. The contraction

1 ratio of the TrA muscle was calculated for all the 7 levels of the test using an established
2 formula (19).

$$3 \quad \text{Contraction ratio} = \frac{\text{Muscle thickness during contraction}}{\text{resting muscle thickness}}$$

4 **Statistical analysis**

5 The sample size for the study was calculated using the G*Power software program by selecting
6 2-tails comparison between groups with an effect size of 0.71 for the primary outcome TrA
7 with a power of 95%, alpha error probability of 0.05 which suggested a sample size of at least
8 15 per gender. The statistical software package for social sciences (SPSS windows version 22.0)
9 was used for data analysis (20). The measured values were normally distributed according to the
10 Kolmogorov-Smirnov test. The differences in the contraction ratio of TrA and IO muscles
11 between the 7 levels of testing were analysed using One-way Repeated measures ANOVA.
12 The level of significance was set as 0.05. In terms of any significant differences, a Post hoc
13 was analysed using Bonferroni at $p < 0.05$.

14 **RESULTS:**

15 A total of 30 healthy individuals (15 males and 15 females) with a mean age of 21.3 years
16 participated and completed the study. The descriptive characteristics of the study participants
17 were given in **Table I**. The mean (SD) value of the resting thickness and the thickness during
18 contraction of TrA and IO for all the 7 levels of the LPST were shown in **Table II**. The result
19 showed that the values of the contraction ratio of the TrA and IO decreased as the tests were
20 progressed into the different 7 levels of the testing. The results of the post hoc analysis showed
21 that the contraction ratio of TrA significantly reduced in the 7th levels of the testing ($F=14.53$,
22 $p=0.001$). However, the contraction ratio of IO remains unchanged in all the 7 levels of LPST.

23 **DISCUSSION:**

1 This was the first study that investigated the changes in the contraction ratio of TrA and IO
2 muscles during the 7 different progressive stages of the LPST. The findings of the study
3 suggested that TrA muscle possibly were showing signs of fatigue as the muscle was repeatedly
4 made to contract during the 7 progressive levels of the LPST. In clinical practice, one might
5 argue that this was the characteristic pattern of the LPST which might explain that the difficulty
6 levels of the 7 progressive stages of the test might demand an increasing workload on TrA
7 muscle. It was worth noting that a past study challenged the notion that the PBU could clinically
8 assess TrA contraction during the abdominal drawing-in manoeuvre (10). However, the
9 findings on TrA muscle were proposed based on a population of NSLBP and the TrA muscle
10 was not tested in different levels of LPST. In contrast, the current study might suggest that the
11 abdominal drawing-in task through PBU had engaged TrA and the TrA muscle responded with
12 decreasing values of contraction thickness across the 7 levels of the LPST. While the past
13 studies used PBU to measure the lumbopelvic stability as an indirect measurement of TrA
14 muscle function (10,11) none of the past studies directly measured how the TrA muscle adapts
15 and functions to the 7 progressive stages of the LPST. Therefore, the current study findings
16 might provide a direct measurement of the TrA with the normative value which could provide
17 clinicians an understanding of how TrA muscle responds to the LPST on healthy individuals.
18 An understanding of the normative values and function of TrA muscle might help clinicians to
19 interpret the test results among NSLBP.

20 The average values of the contraction ratio of TrA muscle in the current study were similar to
21 other established evidence. For instance, the range of TrA contraction ratio measured in our
22 study ranged between 1.5 mm Hg and 1.8 mm Hg which fell within the range of 1.5 mm Hg
23 up to 2.2 mm Hg (21,22). It could be suggested that the norms of measurements were acceptable
24 within the practice standards. The contraction ratio of IO had been reported to be
25 the thinnest during rest and thickest during the abdominal drawing-in manoeuvre (23). Also,

1 irrespective of the measurements of the cross-section of TrA and IO either at rest or during the
2 testing procedure, it was reported that IO and TrA muscles remained the thickest and the
3 thinnest, respectively (23). However, the cross-section ratio of TrA was the thickest ranging
4 from 1.5-1.8 mm Hg, while the cross-section ratio of the IO was the thinnest ranging from 1.1-
5 1.3 mm Hg. Perhaps, the difference could be because the previous study tested the contraction
6 ratio of TrA and IO during the abdominal drawing-in task, while the current study used a
7 tougher testing procedure of testing the muscles across 7 levels of the LPST. Also, the past
8 study reported values from the low back pain population, while the findings of the current study
9 were from healthy individuals. Furthermore, the current findings were in line with a recently
10 reported evidence among healthy participants that the activation levels of the TrA were between
11 1.5-2.0 mm Hg (24,25). Therefore, it could be possible that the cross-section ratio of TrA and
12 IO muscles might adapt and respond differently between healthy individuals and NSLBP
13 population. Therefore, further studies were warranted to test the adaptation of the cross-section
14 ratio of TrA and IO muscles across the 7 levels of LPST between healthy individuals and
15 NSLBP.

16 **Clinical Implication**

17 The current study might have some practice implications. Reference values of the clinical tests
18 among healthy individuals are necessary, as these results may contribute to clinical practice
19 and for setting up training targets (26). For instance, an understanding of certain baseline
20 normative data on the contraction ratio of TrA and IO among the healthy individuals during
21 the LPST might provide clinicians a reference point and comparison standard for NSLBP. A
22 past study raised a clinical question about the measurement property of PBU to engage TrA
23 (15). Also, the engagement of TrA with PBU was questioned due to poor correlation with
24 electromyography (27). In the above context, the current study findings supported that TrA and
25 IO were engaged and activated using a PBU and USI supporting the notion that the contraction

1 ratio of TrA muscle could be evaluated using the 7 levels of the LPST. The small sample size
2 is one of the limitation for the study. Although a sample size of 15 participants were identified
3 as sufficient for a gender in the study, one might suggest that the study consisted of a small
4 number of research participants and hence, the findings might be more prone to variability.
5 Also, another limitation of this study might be mixing the results of men and women, as the
6 thickness of the muscles might be associated with gender.

7 **Conclusion:**

8 A significant reduction in the contraction ratio of TrA was observed in the 7th level of the LPST
9 as measured by the PBU and USI. However, the contraction ratio of IO remains unchanged
10 during all the 7 levels of LPST. As the levels of the LPST increased, the contraction ratio of
11 deep stabilizing muscles were insufficient to stabilize the lumbopelvic region. Further studies
12 are warranted to compare the adaptation of TrA and IO muscles across 7 levels of the testing
13 among patients with NSLBP.

14
15 **Conflict of Interest** – The authors declare that they have no conflict of interests (15).
16
17
18
19
20
21
22
23
24
25

1
2
3
4
5 **References:**
6
7

- 8 1. M.M. Panjabi. A hypothesis of chronic back pain: ligament sub failure injuries lead to
9 muscle control dysfunction. *Eur Spine J*, 2006, 15 (5): p. 668-676.
10
11 2. P.W. Hodges, G.L. Moseley. Pain and motor control of the lumbopelvic region: effect
12 and possible mechanisms. *J Electromyogr Kinesiol*, 2003. 13 (4): p. 361-370.
13
14 3. Hodges PW. Spinal segmental stabilization training in: rehabilitation of the spine.
15 Lippincott: Williams & Wilkins; 2007.
16
17 4. Teyhen DS1, Bluemle LN, Dolbeer JA, Baker SE, Molloy JM, Whittaker J, ChildsJD.
18 Changes in lateral abdominal muscle thickness during the abdominal drawing-in
19 maneuver in those with lumbopelvic pain. *J Orthop Sports Phys Ther*, 2009, 39(11): p.
20 791-8.
21
22 5. MacDonald D, Moseley GL, Hodges PW. Why do some patients keep hurting their
23 back? Evidence of ongoing back muscle dysfunction during remission from recurrent
24 back pain. *PAIN®*, 2009, 142: p. 183-188.
25
26 6. Unsgaard-Tøndel M, Nilsen TIL, Magnussen J, Vasseljen O. Is activation of
27 transversus abdominis and obliquus internus abdominis associated with long-term
28 changes in chronic low back pain? A prospective study with 1-year follow-up. *BMJ*
29 *Sports Med*, 2012, 46: p. 729-734.
30
31 7. Springer BA1, Mielcarek BJ, Nesfield TK, Teyhen DS. Relationships among lateral
32 abdominal muscles, gender, body mass index, and hand dominance. *J Orthop Sports*
33 *Phys Ther*, 2006, 36(5): p. 289-97.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1 8. Rho M, Spitznagle T, Van Dillen L, Maheswari V, Oza S, Prather H. Gender
2 differences on ultrasound imaging of lateral abdominal muscle thickness in
3 asymptomatic adults: a pilot study. *PM R*, 2013, 5(5): p. 374-80.
- 4 9. Şahin N, Karahan AY, Albayrak I. Effectiveness of physical therapy and exercise on
5 pain and functional status in patients with chronic low back pain: a randomized
6 controlled trial. *Turk J Phys Med Rehab* 2018;64(1):52-58.
- 7 10. Grooms DR, Grindstaff TL, Croy T, Hart JM, Saliba SA. Clinimetric analysis of
8 pressure biofeedback and transversus abdominis function in individuals with
9 stabilization classification low back pain. *J Orthop Sports Phys Ther*, 2013, 43(3): p.
10 184-93.
- 11 11. Lima PO, de Oliveira RR, de Moura Filho AG, Raposo MC, Costa LO, Laurentino GE.
12 Reproducibility of the pressure biofeedback unit in measuring transversus abdominis
13 muscle activity in patients with chronic nonspecific low back pain. *J Bodyw Mov Ther*,
14 2012, 16(2): p. 251-7.
- 15 12. Paungmali A, Henry LJ, Sitalertpisan P, Pirunsan U, Uthaikhup S. Improvements in
16 tissue blood flow and lumbopelvic stability after lumbopelvic core stabilization training
17 in patients with chronic non-specific low back pain. *J Phys Ther Sci*, 2016, 28(2): p.
18 635-40.
- 19 13. Lima PO, Oliveira RR, Moura Filho AG, Raposo MC, Costa LO, Laurentino GE.
20 Concurrent validity of the pressure biofeedback unit and surface electromyography in
21 measuring transversus abdominis muscle activity in patients with chronic nonspecific
22 low back pain. *Rev Bras Fisioter*, 2012, 16(5): p. 389-95.
- 23 14. De Paula Lima PO, de Oliveira RR, Costa LO, Laurentino GE. Measurement properties
24 of the pressure biofeedback unit in the evaluation of transversus abdominis muscle
25 activity: a systematic review. *Physiotherapy*, 2011, 97(2): p. 100-6.

- 1 15. Padulo J., Oliva F., Frizziero A., Maffulli N. *Muscles, Ligaments and Tendons Journal*
2 – Basic principles and recommendations in clinical and field Science Research: 2018
3 update. *MLTJ* 2018; 8(3): 305 – 307.
4
5 16. Azevedo DC, Lauria AC, Pereira ARS, Andrade GT, Ferreira ML, Ferreira PH, Van
6 Dillen L. Intraexaminer and interexaminer reliability of pressure biofeedback unit for
7 assessing lumbopelvic stability during 6 lower limb movement test. *J Manipulative*
8 *Physiol Ther* 2013;36:33-43.
9
10 17. K von Garnier, K Koveker, B Rackwitz, U Kober, S Wilke, T Ewert, *et al.* Reliability
11 of a test measuring transversus abdominis muscle recruitment with a pressure
12 biofeedback unit. *Physiotherapy*, 2009, 95: p. 8-14.
13
14 18. Sitalertpisan P, Pirunsan U, Puangmali A, Ratanapinunchai J, Kiatwattanacharoen S,
15 Neamin H, Laskin JJ. Comparison of lateral abdominal muscle thickness between
16 weightlifters and matched controls. *Phys Ther Sport*, 2011, 12: p. 171-4.
17
18 19. Pulkovski N, Mannion AF, Caporaso F, Toma V, Gubler D, Helbling D, Sprott H.
19 Ultrasound assessment of transversus abdominis muscle contraction ratio during
20 abdominal hollowing: a useful tool to distinguish between patients with chronic low
21 back pain and healthy controls? *Eur Spine J.* 2012;21 (Suppl 6):S750–S759.
22
23 20. IBM Corp. Released 2011. *IBM SPSS Statistics for Windows, Version 22.0.* Armonk,
24 NY: IBM Corp.
25
26 21. Linek P, Saulicz E, Wolny T, Myśliwiec A, Kokosz M. Lateral abdominal muscle size
27 at rest and during abdominal drawing-in manoeuvre in healthy adolescents. *Man Ther*,
28 2015, 20(1): p. 117-23.
29
30 22. F.D. Manshadi, M. Parnianpour, J. Sarrafzadeh, M.R. Azghani, A. Kazemnejad.
31 Abdominal hollowing and lateral abdominal wall muscles' activity in both healthy men
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

& women: an ultrasonic assessment in supine and standing positions. *J Bodyw Mov Ther*, 2011, 15: p. 108-113.

23. Mannion et al., 2008b, A.F. Mannion, N. Pulkovski, V. Toma, H. Sprott. Abdominal muscle size and symmetry at rest and during abdominal hollowing exercises in healthy control subjects. *J Anat*, 2008, 213: p. 173-182.

24. Teyhen DS, Rieger JL, Westrick RB, Miller AC, Molloy JM, Childs JD. Changes in deep abdominal muscle thickness during common trunk-strengthening exercises using ultrasound imaging. *J Orthop Sports Phys Ther*, 2008, 38: p. 596– 605.

25. Gorbet N, Selkow NM, Hart JM, Saliba S. No Difference in Transverse Abdominis Activation Ratio between Healthy and Asymptomatic Low Back Pain Patients during Therapeutic Exercise. *Rehabil Res Pract*. 2010;2010:459738.

26. Oliveira IO, Pilz B, Santos RLG Junior, Vasconcelos RA, Mello W, Grossi DB. Reference values and reliability for lumbopelvic strength and endurance in asymptomatic subjects. *Braz J Phys Ther*, 2018, 22(1): p. 33-41.

27. O.M. Giggins, U.M. Persson, B. Caulfield. Biofeedback in rehabilitation. *J Neuroeng Rehabil*, 2013, 10: p. 60.

Figure I: Ultrasound imaging of the internal oblique and transversus abdominis




Table I. Characteristics of the study participants

Variables	Mean±SD	Min	Max
age (years)	21.83±0,46	20.00	22.00
weight(kg)	55.48±11.6	41.00	100.00
height(cm)	163.87±8.14	150.00	181.00
BMI (Kg/m ²)	20.541±3.28	16.61	35.43

Table II. Transversus abdominis and internal oblique muscles thickness and contraction ratio during 7 levels of lumbopelvic stability test (LPST)

Variables	Muscle	LPST level							
		resting	1 st	2 nd	3 rd	4 th	5 th	6 th	7 th
Muscle Thickness (mm)	TrA	2.23 ± 0.5	4.07 ± 0.9	4.21 ± 0.9	4.10 ± 1.1	3.76 ± 1.1	3.85 ± 1.2	3.67 ± 1.2	3.17 ± 1.2
	IO	5.38 ± 1.9	6.11 ± 2.7	6.10 ± 2.4	6.41 ± 2.6	7.09 ± 5.4	6.21 ± 2.3	6.24 ± 2.5	6.22 ± 2.2
Contraction Ratio	TrA	-	1.89±0.4	1.95±0.4	1.88±0.4	1.73±0.4	1.76±0.5	1.70±0.5	1.46±0.5*
	IO	-	1.13±0.3	1.14±0.2	1.20±0.3	1.28±0.6	1.17±0.2	1.17±0.2	1.18±0.2

*significantly difference at $p < 0.001$ when compared to 1st level of LPST