

Washington University in St. Louis

Washington University Open Scholarship

Spring 2017

Washington University
Senior Honors Thesis Abstracts

Spring 2017

Characterizing the Physiological Impact of Carbon Starvation on the Model Bacterium *Escherichia coli*

Jesse Kao

Washington University in St. Louis

Follow this and additional works at: https://openscholarship.wustl.edu/wushta_spr2017

Recommended Citation

Kao, Jesse, "Characterizing the Physiological Impact of Carbon Starvation on the Model Bacterium *Escherichia coli*" (2017). *Spring 2017*. 59.

https://openscholarship.wustl.edu/wushta_spr2017/59

This Abstract for College of Arts & Sciences is brought to you for free and open access by the Washington University

Senior Honors Thesis Abstracts at Washington University Open Scholarship. It has been accepted for inclusion in Spring 2017 by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

CHARACTERIZING THE PHYSIOLOGICAL IMPACT OF CARBON STARVATION ON THE MODEL BACTERIUM *ESCHERICHIA COLI*

Jesse Kao

Mentors: Petra Levin and Corey Westfall

As single celled organisms, bacteria constantly experience stressful changes in their environment. For an enteric organism such as *Escherichia coli*, the sudden dilution of nutrients as they travel through the host's colon and into the environment creates significant challenges that must be dealt with through modifications in growth rate, metabolic flux, and cellular composition.

To understand the impact of sudden changes in nutrient availability on *E. coli* physiology, I have analyzed the impact of rapid depletion of carbon on *E. coli* growth and morphology. In previous experiments, scientists have determined that the slow depletion of carbon as bacteria enter stationary phase results in the formation of short, rounded cells. In contrast, my data indicate that rapidly shifting *E. coli* from carbon rich to carbon free conditions does not lead to stationary-like rounded cells, instead resulting in the surprising detachment of the inner plasma membrane from the cell wall and outer membrane. Further experiments indicate that the detachment phenotype persists in the absence of protein, RNA and lipid synthesis, and is independent of the so-called "stringent response" that is typically seen in *E. coli* when subjected to nutrient stress. Membrane detachment is also independent of osmotic conditions and appears to be specific to carbon starvation; rapid depletion of phosphate and nitrogen starvation does not detectably impair the cell envelope structure.

The re-addition of carbon results in inner membrane reattachment to the cell wall. The rate of reattachment differs based on the carbon source introduced into the system. These add back experiments provide a way to visualize the phenomenon in real time. Characterizing this novel carbon starvation phenotype may provide insight into new mechanisms *E. coli* utilize under carbon depleted environments that may simulate depleted conditions met during bacterial host transmission.