Washington University in St. Louis Washington University Open Scholarship

Volume 13

Washington University Undergraduate Research Digest

Spring 2018

Employing Human Induced Pluripotent Stem Cells (hiPSCs) to Model NF1-Associated Low Grade Gliomas

Kelly Hartigan Washington University in St. Louis

Follow this and additional works at: https://openscholarship.wustl.edu/wuurd_vol13

Recommended Citation

Hartigan, Kelly, "Employing Human Induced Pluripotent Stem Cells (hiPSCs) to Model NF1-Associated Low Grade Gliomas" (2018). *Volume 13*. 79. https://openscholarship.wustl.edu/wuurd_vol13/79

This Abstracts A-I is brought to you for free and open access by the Washington University Undergraduate Research Digest at Washington University Open Scholarship. It has been accepted for inclusion in Volume 13 by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

Employing Human Induced Pluripotent Stem Cells (hiPSCs) to Model NF1-Associated Low Grade Gliomas

Kelly Hartigan

Mentor: David Gutmann

Neurofibromatosis type 1 (NF1) is a clinically heterogeneous cancer predisposition syndrome caused by a germline mutation in one of the two alleles of the NF1 gene. Low grade gliomas (LGGs), particularly in the optic pathway, are common in patients with NF1 (15-20%). These tumors arise when the remaining allele is somatically inactivated in glial progenitor cells. Nf1 genetically-engineered mouse (GEM) models have been employed to study NF1-associated LGGs; however, there are still a limited number of therapeutic targets available. This lack of clinical translation could reflect species-specific differences between rodents and humans or the use of "knockout" mutations in the mouse models, whereas patients with NF1 harbor a diverse number of germline NF1 gene mutations. To study the importance of the germline mutation to LGG development and progression in human cells, the Gutmann Laboratory has created a collection of human induced pluripotent stem cells (hiPSCs) with NF1 patient mutations and hiPSC-derived cerebral organoids. Using this resource, we have generated mixed cerebral organoids comprised of hiPSCs homozygous for patient-specific NF1 mutations and hiPSCs heterozygous for distinct NF1 mutations spanning the NF1 gene. Our preliminary results demonstrate that these mixed organoid cultures exhibit histological characteristics of NF1-LGGs. Future experiments employing this model will evaluate the differential effects of NF1 mutations on LGG formation and growth.