

Washington University in St. Louis

## Washington University Open Scholarship

---

Volume 12

Washington University  
Undergraduate Research Digest

---

Spring 2017

### Genetic Variation Linked to Neuroticism Is Associated with Amygdala Function

William W. Pan

*Washington University in St. Louis*

Follow this and additional works at: [https://openscholarship.wustl.edu/wuurd\\_vol12](https://openscholarship.wustl.edu/wuurd_vol12)

---

#### Recommended Citation

Pan, William W., "Genetic Variation Linked to Neuroticism Is Associated with Amygdala Function" (2017).

*Volume 12*. 152.

[https://openscholarship.wustl.edu/wuurd\\_vol12/152](https://openscholarship.wustl.edu/wuurd_vol12/152)

This Abstracts J-R is brought to you for free and open access by the Washington University Undergraduate Research Digest at Washington University Open Scholarship. It has been accepted for inclusion in Volume 12 by an authorized administrator of Washington University Open Scholarship. For more information, please contact [digital@wumail.wustl.edu](mailto:digital@wumail.wustl.edu).

# GENETIC VARIATION LINKED TO NEUROTICISM IS ASSOCIATED WITH AMYGDALA FUNCTION

*William W. Pan*

*Mentor: Ryan Bogdan*

Neuroticism is a heritable personality trait characterized by emotional instability and psychological stress that places individuals at risk for psychopathology. The amygdala is a brain region that plays a critical role in behavioral vigilance and assigning emotional significance to stimuli that may contribute to the expression of neuroticism. A recent GWAS of 180,911 individuals identified common genetic variation associated with neuroticism. Here, we explored whether single nucleotide polymorphisms (SNPs) that were associated with neuroticism at genomewide levels of significance are associated with threat-related amygdala function.

Genomic, neuroimaging, and self-report data were available for 448 non-Hispanic European-American participants who completed the ongoing Duke Neurogenetics Study. Threat-related amygdala reactivity was assayed using an emotional face-matching task while functional magnetic resonance imaging data were acquired. Neuroticism was assessed with self-report. We tested whether 11 genome-wide significant single nucleotide polymorphisms (SNPs) were associated with neuroticism through GWAS. Covariates in analyses included sex and ancestrally-informative principal components.

We found that the risk alleles of three (*TYRP1* rs10809559, *SBF2* rs13923776, *PAFAH1B1* rs12938775) SNPs were associated with elevated amygdala reactivity (all  $\beta > 0.032$ , all  $p < 0.04$ ). The *PAFAH1B1* rs12938775 allele associated with neuroticism in the GWAS was also associated with neuroticism in our dataset ( $\beta = 3.31$ ,  $p < 0.016$ ), however, neither *TYRP1* rs10809559 nor *SBF2* rs13923776 were (both  $p > 0.68$ ). We conclude that common genetic risk for neuroticism is associated with elevated threat-related amygdala reactivity. Increased threat-related amygdala response may be a genetically influenced neural mechanism conferring neuroticism and risk for psychopathology.