
Content lists available at www.jacsonline.org/D176750082-LROR/ JACSONline GP, DOI:10.22341

Journal of Applied Chemical Science

Journal homepage: www.jacsonline.org/journals/jacson/

Physiological Status of Fattening Bali Cattle Feeding a Concentrate Containing *Gliricidia sepium* Leaves Meal Fortified with Vitamin B-Complex and Vermicide

Sukawaty Fattah*¹, Yohanis U. L. Sobang*¹, Marthen Yunus*¹, F. D. Samba*², and Erna Hartati*¹¹Lecturer at the Faculty of Animal Husbandry, Nusa Cendana University, ² Student of Animal Sciences Post Graduate Program, Nusa Cendana University, INDONESIA

Article history: Received in revised form 15-07-2018

Accepted 29-08-2018

Available online September 28, 2018

Cite this article as: Fattah S, Sobang YUL, Yunus M, Samba FD, and Hartati E. *Physiological Status of Fattening Bali Cattle Feeding a Concentrate Containing Gliricidia Sepium Leaves Meal Fortified with Vitamin B-Complex and Vermicide*. J Applied Chem. Sci. 2018, 5(2): 464-468 DOI: <https://dx.doi.org/10.22341/jacs.on.00501p464> p-ISSN: 2089-6328, e-ISSN: 2580-1953 © 2018 JACSONline GP. All right served



The JACSONline Group Publisher publishes the work of Jacson-Journal of Applied Chemical Science eISSN: 2580-1953/pISSN: 2089-6328 under the licensing of a [Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License](https://creativecommons.org/licenses/by-nc-sa/3.0/). Authors retain the copyright to their work. Users may read, copy, and distribute the work in any medium provided the authors and the journal are appropriately credited. The users may not use the material for [commercial purposes](https://creativecommons.org/licenses/by-nc-sa/3.0/).

ABSTRACTS

This research has been conducted at Oeletsala village, for 10 weeks from 23 November 2015 to 23 January 2016, comprised of 2 weeks preliminary and 8 weeks for data collection. The aim of the research was to study the effect of feeding a *Gliricidia sepium* leaves meal concentrate fortified with B. complex vitamin and vermicide on rectal temperature, respiratory frequency, and heart rate of fattening Bali cattle. Experimental animals used were nine young male Bali cattle of 1.5-2 years old, with an initial body weight ranging from 82 to 124 kg (KV=15.114%) in average of 98.7±14.93 kg. The experimental design used was Randomized Block Design with three treatments and three replications. Those treatments were R0 = leaves of *Leucaena leucocephala* and *Ceiba pentandra*+ native grass *ad libitum* (as commonly practiced by local farmers), R1 = R0 + 2 kg concentrate, and R2 = R1 + B. complex vitamin, and vermicide. Statistical analysis showed that there was no significant effect of the treatments on rectal temperature, respiratory frequency, and heart rate of the fattening Bali cattle ($P \leq 0.05$). In conclusion, there was no significant effect of *Gliricidia sepium* leaves meal concentrate, vitamin B. complex, and vermicide on rectal temperature, respiratory frequency, and heart rate of the fattening Bali cattle.

Keywords: concentrate, B complex vitamin, worm vermicide, rectal temperature, respiratory frequency, and heart rate

* Corresponding author: sukawatifattah@yahoo.com

1. Introduction

East Nusa Tenggara (ENT) has been well known as a source of livestock in Indonesia and played an important role in fulfillment of the national meat demand. Although beef cattle's farming was the most leader commodity in ENT, but still facing some handicaps --which cause low productivity of beef cattle especially fattening cattle-- such as not continuity of feed availability, still lay on traditional culture, unused available local feed sources, which in turn impact to the low of animal productivity. One of the factors causes the low productivity of animal in farmer's level was the low quality of feed offered led to increasing of digest tract performance resulted in increasing of stress and in turn disturb thermo regulation process of animal.

Some researcher reported that the stresses on animal were caused by both metrological factors, temperature, humidity, and radiation (Genswein *et al.*, 2012) and non metrological factors such as animal density, quantity, and quality of feed offered, and drinking water availability (Aradom, 2013; Bulitta *et al.*, 2015). In addition, Ake *et al.* (2013) stated that physiological change was also caused by transportation. Some factors above can stimulate changes in physiology status of animal which bring negative impacts on animal such as the increasing of body temperature, heart rate,

and abnormal in respiratory frequency (Eniolorunda *et al.*, 2009). It also impacts to the changes of hematological performance (Sporer *et al.*, 2014), which cause decrease of body weight and when it happens in long term will cause to decrease immune function which in turn can cause death (Knowles, 1999). Decreasing of immune function up to the death of animal in a relatively large number can cause highly disadvantages to the farmers on village condition that strongly dependence their life upon agriculture and farming. Therefore, strategy to improve feeding management by offering a balance ratio between protein and energy in the diet is needed by feeding on feed that more degradable in the rumen led to which press the activity of microbial rumen and performance of digest tract in digesting low quality feed and press physiological status of animal in order to adapted to feed of low quality and environment by feeding concentrate comprised of protein source feed such as *Gliricidia sepium* leaves

On local farmers kept level, the feeding of this *Gliricidia sepium* resulted in not optimum yet, therefore supplementation of vitamin B-complex and vermicide are needed. Vitamin B-complex is functions to improve body resistant and appetite, meanwhile vermicide is functions to kill internal parasites, so the feed offered could be better digested and in turn, the growth of animal became optimum. This rese-

arch was expected would able to eliminate physiological status disturbance on Bali cattle fattened as commonly practiced by local farmers which in turn could improve animal productivity. The aim of this research was to study the effect of the supplementation of *Gliricidia sepium* leaves meal concentrate, with the addition of vitamin B-complex and vermicide on rectal temperature, respiratory frequency, and heart rate of Bali cattle fattened as commonly practiced by local

2. Materials and Methods

This experiment was carried out at Oeletsala village, Taebenu sub-district, the regency of Kupang for 10 weeks, from 23 November 2015 up to 29 January 2016, comprised of 2 weeks preliminary and 8 weeks collecting data. Nine young male Bali cattle of 1.5 – 2 years old with an initial body weight ranging from 82 to 124 kg and coefficient variation (CV) 17.10 % were employed in the experiment. Those experimental animals were placed in 9 individual pens of 2.10 m x 1.20 m each, cement floor, roofs from coconut leaves, and equipped with drinking and feeding manger. Feed offered were concentrate and feed ingredients commonly used by local farmers, also 5 cc vitamin B-complex by injected and 20 g vermicide per 100 kg body weight. Drinking water was offered *ad libitum*. Equipment used were: bucket, shovel, electrical scale *Excellent Scale* with a sensitivity of 0.5 kg and capacity of 1000 kg for weighing experimental animal, static balance *Kondo* with a sensitivity of 0.05 kg and capacity of 5 kg, thermometer, stethoscope, and stopwatch. The experimental design used was Randomized Block Design, consisted of 3 treatments and 3 replications. Those three treatments were:

R₀ : Feed commonly used by farmers (*Leucaena leucocephala* leaves, *Ceiba pentandra* leaves, and grass)

R₁ : R₀ + 2 kg concentrate

R₂ : R₁ + vitamin B-complex and vermicide

Variables measured as indicator of the effect of treatment applied in this research were:

Table 1. Concentrate ingredients for fattening male Bali cattle (on dry matter basis)

No.	Ingredients	Percentage (%)
1.	Rice-bran	50
2.	Grinding maize	25
3.	<i>Gliricidia sepium</i> leaves meal	16
4.	Fish meal	5
5.	Urea	3
6.	Mineral mix	0.5
7.	<i>Starbio</i>	0.5
Total		100

Table 2. Chemical composition of feed ingredients (based on % DM)

Ingredients	DM	OM	CP	Crude Fat	CF	CHO	NEE	Energy	
								MJ/kg DM	Kcal/kg DM
Grass	16.12	92.48	8.55	1.43	25.34	82.50	57.16	16.83	4,007.85
<i>Leucaena leucocephala</i>	24.23	90.35	23.04	3.33	13.39	63.98	50.59	17.79	4,235.30
<i>Ceiba pentandra</i>	26.11	86.46	16.78	1.95	11.95	67.73	55.78	16.45	3,916.81
Concentrate	90.30	90.22	20.77	6.04	6.30	63.41	57.12	18.07	4,301.25

1. Rectal temperature (in °C). By pushing in thermometer as depth as 5 cm in the rectum.
2. Respiratory frequency. By measuring the up-down movement of stomach-rib surface and approaching hand palm on nose of the experimental animal by using stethoscope for 1 minute.
3. Heart rate. By measuring the heart beating by using stethoscope for 1 minute on both sides of front left foot of the experimental animal.

Experimental Procedure

- 1) Preparation phase. Starting with prepare concentrate and nine heads young male Bali cattle of 1.5 – 2 years old. This phase was prior to research conducted.
- 2) Randomizing of animal experimental:
 - a) Before the research carried out, the experimental animals were weighed to know their initial body weight, and they were numbered 1 to 9.
 - b) Experimental pens were numbered regularly.
 - c) Randomizing experimental animals by using lottery number, to be placed in the pen.
 - d) Randomizing feeds to be treated by using lottery number.
- 3) Experimental phase. This phase was divided into two periods: period-1 the experimental animals were adapted for 14 days to feed treated in the pen environment. Treatment feed was offered little by little to ensure they were familiar.
- 4) On day-14 the experimental animals were weighed. On period-2 feed intake data was collected and also for the rest feed every day.
- 5) Feed offering. Feed was offered twice a day, at 7 am and 4 pm, respectively.
- 6) Feeding. Concentrate was first offered then followed by feeds commonly used by the local farmers. Before offered, feed was weighed first and also for the rest feed on the next morning.
- 7) Environment temperature. It can be read from dry and wet thermometer which hung on the pen's wall.
- 8) Procedure of preparing concentrate:
 - a) Equipment used in the experiment was: scale with a capacity of 300 kg, bucket, and shovel.
 - b) Ingredients for preparing concentrate were: rice-bran, grinding maize, *Gliricidia sepium* leaves meal, fish meal, urea, mineral mix, and *starbio*.
 - c) Supplement mixing procedure: ingredients for supplement and feed were homogenized mixed, started with the least part of the ingredients.

Table 3. Mean of rectal temperature, heart rate, and respiratory frequency

Parameters	Treatments			MS	P-Value
	R ₀	R ₁	R ₂		
Rectal temperature (°C).	38 ^a ±0.20	38 ^a ±0.06	37 ^a ±0.32	0.747	0.530
Heart rate (times/minute)	69 ^a ±1.52	69 ^a ±0.57	66 ^a ±1.02	2.111	0.334
Respiratory frequency (times/minute)	25 ^a ±1.52	25 ^a ±2.3	22 ^a ±1.00	9.333	0.163

Data Analysis

Data collected was tabulated and subjected to Analysis of Variance (ANOVA) to explore effects of treatment on parameters observed. The statistical significant of the data value was considered at $P \leq 0.05$.

3. Results and Discussion

3.1. Feed and Environment Temperature

In keeping their animals including cattle, the local farmers strongly dependent on forages available surrounding them, without consider on nutritive value of the forages. Although the forages were quiet high protein content, but low energy content. Surroundings temperature has a direct effect on animal physiology which impact to the thermoregulation of animal body. Low quality feed and surroundings temperature was altogether resulted in stress on cattle which affected the animal physiology and in turn decrease animal production. The higher temperature and humidity than comfort zone will cause the higher attempting of the animal to keep normal their body temperature through thermoregulation. The increasing of body heat was caused by the combination of air temperature, air humidity, air movement, and sun radiation. This increasing of body heat and respiratory frequency led to less of feed intake and production (Astuti et al.,2015). Physiology status of Bali Cattle Fattened as commonly practiced by Local Farmers are shown on Table 3. The same superscripts in the row indicated no significant difference ($P \leq 0.05$)

3.2. Effect of Treatments on Rectal Temperature

Table 3 showed that the highest mean of rectal temperature was found on treatments R₀ and R₁ 38.0 °C then followed by treatment R₂ (37.8 °C). Forages offered for each treatment were consisted of *Leucaena leucocephala* leaves, *Ceiba pentandra* leaves, and native grasses, as commonly practiced by the local farmers, and added with 2 kg concentrate, vitamin B-complex and vermicide, was able to decrease rectal temperature of the experimental animals. Based on statistical analysis (ANOVA), there was no significant effect of all of the three treatments on rectal temperature. This was because of the low quality of diet commonly fed by the local farmers, produces metabolism heat although *Gliricidia sepium* leaves meal was added. The high feed digestion activity will increase body heat, therefore the relatively the same quality of feed offered and feed intake resulted in the same body heat produced. This result was in ranging of normal body temperature for mammalian and in accordance with the finding of Astuti et al. (2015) by feeding forages and concentrate in different time i.e. 38.57±0.25, 38.89±0.19, and 38.77±0.14, for treatments P₁, P₂, and P₃, respectively. Meanwhile Astuti and Sudarman (2015) found that by adding

Ca-saponified lemuru oil coated with herbs on lambs resulted in average of body temperature were 38.90, 38.83, 38.73, 38.73 and 38.83 for R₁, R₂, R₃, R₄, and R₅, respectively. Sporer et al. (2014) also reported that cattle transported increased 0.13 °C their body temperature during the transportation, from 38.73 ± 0.06 °C to 38.86 ± 0.05 °C. According to Farooq et al. (2010) effort of the animal to keep their body temperature in normal condition, is by increasing the dissipation of their body heat through increasing of their respiratory frequency and heart rate.

The high body temperature on treatment R₀ was caused by low quality of nutrients and high content of crude fiber in that treatment led to increase the activity of animal in masticating feed consumed which in turn released by/through conduction and radiation via respiratory tract which impact to high body temperature. According to Astuti et al., (2015^a), the heat releasing by conduction, radiation, and evaporation was to keep the body temperature in normal range because animal needs a balance between heat production and heat released by their body. Meanwhile, according to Gaughan et al.(2004) and Kendall et al. (2007) heat releasing by conduction and convection; because when animal body was watered, there will a heat transfer process from animal body to water media which has a cooler temperature on the thin layer around skin.

3.3. Effect of Treatments On heart Rate

Table 3 showed that the highest mean of heart rate was the treatments R₀ and R₁ which was 69 times/minute, and the lowest was R₂ (66 times/minute). Forages offered for each treatment were consisted of *Leucaena leucocephala* leaves, *Ceiba pentandra* leaves, and native grasses, as commonly practiced by the local farmers, and added with 2 kg concentrate, vitamin B-complex and vermicide.

Treatments R₀ and R₁ have the same effect on heart rate, therefore animal would always try to keep normal their heart rate through physiological, metabolic, and thermoregulation activities. Based on statistical analysis, there was no significant effect of treatments on heart rate. This was analog with the result on respiratory frequency of this experiment. According to Astuti et al. (2015), that the increasing of respiratory frequency in ruminant was related to heart rate and rectal temperature due to daily activity and surroundings temperature stress. Results of this experiment was lower than finding of Astuti et al. (2015) who feeding forages and concentrate in different time, found that the heart rate were 87.06±2.17, 90.69±2.18, and 88.13±2.17, for treatments P₁, P₂, and P₃, respectively. Meanwhile, Astuti and Sudarman (2015) found that by adding Ca-saponified lemuru oil coated with herbs on lambs resulted in average of heart rate were 104.6, 99.6, 93.6,

102.6 and 98.3 for R₁, R₂, R₃, R₄, and R₅, respectively. According to Eniolorunda *et al.* (2009) that the increasing of heart rate was physiologically related to the increasing of respiratory frequency which cause increasing of respiratory muscles activity resulted in fasting heat distribution to the periphery skin to be released to surrounding to keep balance of body heat.

The high heart rate on treatments R₀ and R₁ was suspected due to increasing of muscles activity as the resulted of high cell wall content of forages of diet commonly practiced by local farmers and also high level of concentrate offered which cause difficult to be consumed and digested in term of took long time to be masticated by animal and needed more energy. According to West (2003) that the higher level of feed offered the higher energy consumed, and this led to increasing of heat produced from the body as the result of metabolic process. Heat produced by animal body was derived from metabolic activity and heat from the surroundings, would be released by conduction, radiation, and evaporation via skin and respiratory tract (Ewing dan Borell, 1999).

3.5. Effect of Treatments On respiratory Frequency

Table 3 showed that the highest mean of respiratory frequency were the treatments R₀ and R₁ 25 times/minute, and the lowest was R₂ (22 times/minute). Forages offered for each treatment were consisted of *Leucaena leucocephala* leaves, *Ceiba pentandra* leaves, and native grasses, as commonly practiced by the local farmers, and added with 2 kg concentrate, vitamin B-complex and vermicide. Both treatments R₀ and R₁ indicated to have the same effect on respiratory frequency. Based on statistical analysis, there was no significant effect of treatments on respiratory frequency. This no significant difference among the treatments was suspected due to low quality of feed and/or high content of crude fiber, resulted in disturbing of physiology condition of animal and impact to increasing of respiratory frequency. This was because of high crude fiber consumed could produce a high metabolic heat. The high feed digestion activity lead to high body heat, therefore the relatively the same feed quality and feed consumed of all the three treatments resulted in also the same on body heat produced. High crude fiber of feed affected the physiologic condition of animal and also respiratory frequency, where high quality of feed will increase respiratory frequency as the result of feed metabolic activity.

The respiratory frequency found in this experiment were far lower than the findings of Astuti *et al.* (2015) who feeding forages and concentrate in different time, found that the respiratory frequency were 22.66±2.27, 26.81±2.21, and 25.00±0.94, for treatments P₁, P₂, and P₃, respectively. Meanwhile, Astuti and Sudarman (2015) found that by adding Ca-saponified lemuru oil coated with herbs on lambs resulted in average of respiratory frequency were 43.33, 48.66, 49.00, 44.33, 45.66, for treatments R₁, R₂, R₃, R₄, and R₅, respectively. The high respiratory frequency on treatments R₀

and R₁ was suspected due to high animal activity in masticating low quality feed which in turn digestion in rumen need longer time and this will push the diaphragm and also the lung, resulted in short in respiration lead to decreasing of tidal volume or inward air (inspiration) and outward air (expiration) in respiratory tract. According to Swenson dan Reece (1993), factors that affected there spiratory frequency were body size, age, muscle activity, surroundings temperature, pregnancy, and fully of digestive tract. The high surroundings temperature will also determine the high and the low of respiratory frequency (Ambius *et al.*, 2016).

4. Conclusions

Based on the results, it is concluded that the feeding of supplement contained *Gliricidia sepium* leaves meal with the addition of vitamin B-complex and vermicide was not affected significantly (P ≤ 0.05) the rectal temperature, respiratory frequency, and heart rate of Bali cattle fattened as commonly practiced by local farmers in the Oeletsala village.

References

- Ambius A, Lalu Muhammad Kasip, Lalu Wirapribadi, SulaimanNgonguDepamede, A. Rai Somaning Asih. 2016. *Physiological Status and Body Weight Changes of Bali Cattle Heifers Transported from Lombok Island to West Kalimantan*)JurnalIlmudanTeknologiPeternakan Indonesia, 2 (1): 86 – 95
- Ayu A, Erwanto, Santosa PE. The Effect of Providing Forage-Concentrate on Physiological Response and Performance of Simmental Cross Beef Cattle. *Jurnal Ilmiah Peternakan Terpadu*. 2015; 3 (4) : 201-207.
- Astuti A, and Sudarman A. Physiological Status, Blood Profile and Body Composition of Sheep Fed With Ca-Saponified Lemuru Oil Coated by Herbs. *Buletin Peternakan*. 2015; 39 (2):116-122. DOI: <https://doi.org/10.21059/buletinpeternak.v39i2.6716>
- Ake AS, Ayo JO, and Aluwong T. Effects of transportation and thermal stress on donkeys in the Northern Guinea Savannah zone of Nigeria: A review. *Journal of Cell and Animal Biology*. 2013; 7 (8): 92-101. DOI: <https://doi.org/10.5897/JCAB2013.0370>
- Aradom S. Animal Transport and Welfare with Special Emphasis on Transport Time and Vibration. Doctoral Thesis: Faculty of Natural Resources and Agricultural Sciences, Department of Energy and Technology Uppsala. Swedish University of Agricultural Sciences. 2013; Uppsala.
- Bulitta FS, Aradom S, and Gebresenbet G. Effect of transport time of up to 12 hours on welfare of cows and bulls. *Journal of Service Science and Management*. 2015; 8: 161-182. DOI: <https://doi.org/10.4236/jssm.2015.82019>
- Eniolorunda E. Fashina, and Aro. Adaptive physiological response to load time stress during transportation of cattle in Nigeria. *Journal of Archive Zootechnology*. 2009;58 (222): 223-230.
- Ewing SA, Lay DCJR, and Borell EV. Farm animal well being stress physiology animal behavior and environmental design. Prentice- Hall.Inc. New Jersey. 1999.
- Farooq U, Samad HA, Shehzad F, and Qayyum A. Physiological responses of cattle to heat stress. *Journal of World Applied Sciences*, (Special Issue of Biotechnology and Genetic Engineering); 2010. 8: 38-43.
- Gaughan JB, Davis MS, and Mader TL. Wetting and the physiological responses of grain-fed cattle in a heated environment. *Aust. J. Agric. Res.* 2004; 55: 253-260. DOI: <https://doi.org/10.1071/AR03110>

- Genswein KS, Faucitano L, Dadgar S, Shand P, González LA, and Crowe TG. Road transport of cattle, swine and poultry in North America and its impact on animal welfare, carcass and meat quality: a review. *Journal of Meat Sciences*. 2012. 92 (3): 227-43. **DOI: <https://doi.org/10.1016/j.meatsci.2012.04.010>**
- Kendall PE, Verkerk GA, Webster JR, and Tucker CB . Sprinklers and shade cool cows and reduce insect-avoidance behavior in pasture-based dairy systems. *J. Dairy Sci*. 2007; 90: 3671-3680. **DOI: <https://doi.org/10.3168/jds.2006-766>**
- Knowles TG. Review of the Road Transport of Cattle. *Veterinary Record*. 1999; 144: 8: 197-201. **DOI: <https://doi.org/10.1136/vr.144.8.197>**
- Sporer KRB, Weber PSD, Burton JL, Earley B, and Crowe MA.. Transportation of young beef bulls alters circulating physiological parameters that may be effective biomarkers of stress. *Journal of Animal Science*. 2014; 86:1325-1334. **DOI: <https://doi.org/10.2527/jas.2007-0762>**
- Swenson M.J and Reece WO. *Duke's Physiology of Domestic Animals*. 11th Edn. Comstock Publishing Associates. 1993.
- West JW. Effect of heat-stress on production in dairy cattle. *J Dairy Sci*. 2003; 6: 2131. **DOI: [https://doi.org/10.3168/jds.S0022-0302\(03\)73803-X](https://doi.org/10.3168/jds.S0022-0302(03)73803-X)**

Conflict of interest: Non declare