



Estimation of Growth, Mortality, and Exploitation.....from The New Calabar River, Nigeria (Olopade, O.A., et al)



Available online at: <http://ejournal-balitbang.kkp.go.id/index.php/ifrj>
e-mail: ifrj.puslitbangkan@gmail.com
INDONESIAN FISHERIES RESEARCH JOURNAL
Volume 25 Nomor 2 December 2019
p-ISSN: 0853-8980
e-ISSN: 2502-6569
Accreditation Number RISTEKDIKTI: 21/E/KPT/2018



ESTIMATION OF GROWTH, MORTALITY, AND EXPLOITATION STATUS OF NURSE TETRA (*Brycinus nurse*) AND TRUE BIG SCALE TETRA (*Brycinus macrolepidotus*) (FAMILY: ALESTIDAE) FROM THE NEW CALABAR RIVER, NIGERIA

Olaniyi Alaba Olopade*¹, Henry Eyina Dienne¹ and Uzoma C Nworgu¹

¹Department of Fisheries, Faculty of Agriculture, University of Port Harcourt, Nigeria

Received; April 20-2019 Received in revised from December 02-2018; Accepted December 11-2019

ABSTRACT

This study aimed to determine the growth patterns, mortality, and exploitation status of two species of Alestidae in the New Calabar River, Nigeria. For this purpose, fish samples were collected monthly from three landing sites from the local fishermen using gill nets (mesh sizes: 15-25mm), beach seine (mesh sizes: 2.3-10mm), and cast nets (mesh sizes: 15 -25mm). The length-weight relationship revealed exponent "b" value for *Brycinus nurse* was 3.54 and 3.21 for *Brycinus macrolepidotus* while the condition factors were 1.08 and 1.02 for *Brycinus nurse* and *Brycinus macrolepidotus* respectively. The growth parameters of *Brycinus nurse* asymptotic length (L_{∞}) and growth coefficient (K) were 24.46 cm and 0.52 yr⁻¹, respectively, while those for *Brycinus macrolepidotus* L_{∞} was 28.88 cm and K was 0.22 yr⁻¹. The reproductive load (L_{50}/L_{∞}) ratio was found to be 0.59 and 0.61 for *B. nurse* and *B. macrolepidotus*, respectively. Exploitation rate (E) for *B. nurse* was 0.26 and 0.11 for *B. macrolepidotus* while length-at-first capture (L_c) was 14.49 cm for *B. nurse* and 17.64 cm for *B. macrolepidotus*. The natural mortality was greater than the fishing mortality for both species and Logistic regression of the probability of capture routine values recorded for *B. nurse* were higher than that of *B. macrolepidotus*. Maximum exploitation rate (E_{max}) was less than 0.5 for both *B. nurse* (0.41) and *B. macrolepidotus* (0.42). These values were close to the maximum allowable limit; therefore, the species may be unsustainable when fishery intensifies in the future. To ensure sustainable exploitation of the two Alestid species in the area, fishing effort should be regulated.

Keywords: Alestid species; length-weight relationship; reproductive load; demographic structure; Nigeria

INTRODUCTION

Alestidae has been described as the most speciose of all African characiform families (Paugy & Schaefer, 2007; Arroyave & Stiassny, 2011). About 118 valid species from family Alestidae greatly vary in body and fin sizes, shapes, and occupied ecological niches (Froese & Pauly, 2016). According to Froese & Pauly (2017), about 21 species spread across eight genera (*Alestes*, *Alestopetersius*, *Anordichthys*, *Brycinus*, *Hydrocinus*, *Bryconaethiops*, *Micralestes*, and *Rhabdalestes*) have been found in Nigerian waters. Off the eight genera, the genus *Brycinus* is the most prevalent in Nigerian freshwaters with ten species. The genus *Brycinus* is characterized by presence of rudimentary adipose eyelids, and by having two rows of pluricuspid teeth on the upper jaw (Paugy, 2003). The exploitation history of Characidae from Nigerian waters, according to the NBS (2015), revealed that

the annual catch of the characid species from 2010 to 2015 ranged between 14, 784 to 23, 124 metric tons (mt), where 23,124 mt was the highest landing occurred in 2014 then decreased to 18,356 mt in 2015.

Brycinus macrolepidotus (Valenciennes, 1850) and *Brycinus nurse* (Rüppell, 1832) are native to freshwater systems in Africa thriving well in both lacustrine and riverine conditions (Boulenger, 2002), particularly in the Niger Delta, Nigeria. *Brycinus macrolepidotus* is easily identified by red marks on pectoral, ventral, and anal fins, the tip of adipose fins is reddish, the tail fin of *B. nurse* is bright red, other fins are tinged with red, and there is black patch on the tail peduncle (Adesulu & Sydenham, 2007). These species are said to be of commercial importance due to the fact that they are widely consumed locally and have food value (Reed *et al.*, 1967; Saliu & Fagade, 2004). Like other commercially important fish species in Nigerian inland

correspondence author:

e-mail: olaniyi.olopade@uniport.edu.ng

DOI: <http://dx.doi.org/10.15578/ifrj.25.2.2019.113-122>

waters, this fish family is currently subjected to intense fishing pressure due to lack of proper management policy. Since unregulated exploitation is still on there are urgent needs to monitor and assess the status of the stocks that are being fished.

Fish stock change depends on recruitment, natural mortality, individual growth, and harvesting. The growth parameters and the mortality rate are important tools to assess the exploitation level of the pelagic species (Wang & Liu, 2006). Length and weight data of fish can be used for the estimation of the length and age structures growth and mortality rates of the fish (Kohler *et al.*, 1995) and the condition factor that may help determine whether somatic growth is isometric or allometric (Ricker, 1975). Relationship between length and weight is required for setting up yield equation (Beverton & Hold 1957; Ricker 1968).

The biological information on fish is of paramount importance for the development of effective strategies for their management and conservation. Information on population parameters, i.e., growth, reproduction, recruitment, as well as mortality of fish, is vital to the implementation of sustainable management strategies for their better conservation (Hossain *et al.*, 2009). The present study was thus undertaken to estimate the key biological information on two most dominant Alestid species caught in the New Calabar River, Nigeria.

MATERIALS AND METHODS

The Niger Delta is crossed by many distributaries. Off these distributaries is the New Calabar River in the Niger River Delta, Nigeria. It is a partially mixed estuary river lies between latitude 4°25'2" N and longitude 7°16'2" E (Olopade *et al.*, 2018). It runs through the most densely populated areas in the hinterland and empties into the Atlantic Ocean at the southern tip of Bonny in the south. The river has high water volume during rainy season due to high runoff. The main rainy season is from April to September with the annual rainfall between 2000 and 3000 mm (Abowei & Hart, 2009). Rainfall is generally negligible in December and February.

Three non-overlapping landing sites were selected along the New Calabar River, namely Choba, Aluu, and Ogbogoro (Figure 1) to provide a representative overview of the fisheries in the river. The fish samples were also collected monthly from three landing sites from the local fishermen using gill nets (mesh sizes: 15-25 mm), beach seine (mesh sizes: 2.3-10 mm), and cast nets (mesh sizes: 15 -25 mm). The fish species were identified using fish identification keys (Paugy, 2003; Adesulu & Sydenham, 2007). Total length, standard length, and weight of the fish species were measured in centimeters and grams, respectively.




Figure 1. Map Showing New Calabar River, Rivers State Nigeria.

Data Analysis

The length-weight relationship is expressed by the equation $W = aL^b$, where W = Body weight (g), and L = Total length (cm), Ricker (1973) as follows:

$\log W = a + b \log L$ (1)

The value of the growth exponent was used to calculate the condition factor using the formula:

$K = 100 W / L^b$ (2)

Where;
 K = condition factor

W = total body weight (g)
 L = Total length (cm) and
 b = growth exponent

$$E = F/Z \dots\dots\dots (4)$$

The length-frequency data for each species were collected monthly from the different sampling sites and subsequently grouped into class intervals for analysis. The data was analyzed using FISAT II (FAO-ICLARM Stock Assessment Tools) as explained in detail by (Gayaniilo *et al.*, 2005).

Relative yield per recruit (Y/R) was estimated using the model of Beverton & Holt (Beverton & Holt, 1966) as modified by Pauly & Soriano (1986) and incorporated in the FISAT software.

The von Bertalanffy growth parameters (Pauly, 1980), L_{∞} and annual growth coefficient K were computed by ELEFAN I (Electronic Length Frequency Analysis) (Beverton & Holt, 1966). The total mortality rate (Z) was estimated by length-converted catch curve (Pauly, 1984). The natural mortality rate (M) was also calculated by using Pauly's empirical formula (Pauly, 1980). The fishing mortality rate (F) was calculated by the difference between (Z) and (M) or

Resource status was evaluated by comparing estimates of the fishing mortality rate with a target (F_{opt}) and limit (F_{limit}) biological reference points (BRP) which were defined as $F_{opt} = 0.5M$ and $F_{limit} = 2/3M$ (Patterson, 1992).

$$F = Z - M \dots\dots\dots (3)$$

Probability of capture against mid-length a resultant curve was used to compute the length at first capture ($Lc50$). The length at first maturity ($Lm50$) was estimated as $Lm50 = (2 * L_{\infty}) / 3$ (Hoggarth *et al.*, 2006)

The rate of exploitation (E) was calculated by the quotient between fishing and total mortality (Pauly, 1984):

RESULTS AND DISCUSSION

Results

A total of 196 individuals Alestid species belong to two species, namely *B. macrolepidotus* and *B. nurse*, were analyzed during this study. *Brycinus nurse* occurred in considerable number (64.3%) more than *B. macrolepidotus* with 35.7% (Table 1).

Table 1. Species composition by number from all types of gears

Species	Beach seine	Cast net	Gillnet	N	Percentage (%)
<i>B. macrolepidotus</i>	29	23	18	70	35.7
<i>B. nurse</i>	53	27	46	126	64.3
Total	82	50	64	196	100

The mean total lengths were estimated to be 15.08 ± 0.37 and 14.58 ± 0.25 cm for *B. macrolepidotus* and *B. nurse*, respectively. The mean weight for *B. macrolepidotus* was estimated as 40.00 ± 3.57 g, whereas it was 39.44 ± 2.73 g for *B. nurse* (Table 2). The length-weight relationship results revealed exponent of (b) values of 3.2103 and 3.5405 for *B.*

macrolepidotus and *B. nurse*, respectively (Table 2, Fig. 2 and 3). Also the correlation value (r) for the two species were estimated as 0.8868 and 0.8853, respectively. The mean value for condition factor (K) recorded for *B. macrolepidotus* was 1.02 ± 0.03 , and *B. nurse* had the value of 1.08 ± 0.02 (Table 3).

Table 2. Total length, standard length and Mean weight of Alestid species from New Calabar River landed by all type of fishing gears

Species	TL (cm)		SL(cm)		WEIGHT (g)		GIRTH (cm)	
	Mean±SE	Range	Mean±SE	Range	Mean±SE	Range	Mean±SE	Range
<i>B. macrolepidotus</i>	5.08 ± 0.37	11.50 - 27.00	11.88 ± 0.30	8.70 - 22.30	40.00 ± 3.57	10.00 - 160.00	8.80 ± 0.25	6.00 - 15.30
<i>Brycinus nurse</i>	14.58 ± 0.25	7.30 - 23.00	11.49 ± 0.20	5.50 - 18.60	39.44 ± 2.73	5.00 - 210.00	8.80 ± 0.19	2.50 - 16.50




Figure 2. Length-weight relationship of *B. macrolepidotus* in the New Calabar River.




Figure 3. Length-weight relationship of *B. nurse* in the New Calabar River.

Table 3. Condition factors and growth patterns of two Alestid species in the New Calabar River

Species	K	a	b	r ²	
	Mean	Range			
<i>B. macrolepidotus</i>	1.02±0.03	0.61 - 1.56	-5.1712	3.2103	0.8868
<i>B. nurse</i>	1.08±0.02	0.49 - 1.73	-5.9999	3.5405	0.8853

The growth parameters estimated for *B. nurse* using the length frequency data in ELEFAN I program revealed the best fit for $L_{\infty} = 24.46$ cm and $k = 0.52$ per year while for *B. macrolepidotus*, the best fit were for $L_{\infty} = 28.88$ cm and $k = 0.22$ per year (Table 4). The total mortality (Z) of *B. nurse* estimated by the length converted catch curve was 1.88 for *B. nurse* while it was 0.77 for *B. macrolepidotus*. The natural mortality (M / year) as per Pauly's empirical formula was found to be 1.05 for *B. nurse* while it was 0.69

for *B. macrolepidotus*. The estimated fishing mortality ($Z-M=F$) stood at 0.83 and 0.77 while the exploitation ratio (E) was found to be 0.26 and 0.11 for *B. nurse* and *B. macrolepidotus*, respectively (Table 4).

The logistic regression of the probability of capture routine values recorded for *B. nurse* was lower than that of *B. macrolepidotus* (Table 5, Fig 4 and 5). The estimated L_{50} was 14.49 cm for *B. nurse* and 17.64 cm for *B. macrolepidotus*. The L_{25} was calculated as

12.88 cm and 15.13 cm while L_{75} was found to be 16.10 cm and 20.16 cm for both species, respectively. In this study, the reproductive load (L_{50}/L_{∞}) ratio was found to be 0.59 and 0.61 for *B. nurse* and *B. macrolepidotus*, respectively.

Table 4. Population parameters of *B. nurse* and *B. macrolepidotus*

Von Bertalanffy's Growth parameters	<i>B. nurse</i>	<i>B. macrolepidotus</i>
Asymptotic length (L_{∞})	24.46	28.88
Growth coefficient (k)	0.52	0.22
t_0 (yr)	0.04	0.07
Total mortality (Z)	0.83	0.77
Fishing mortality (F)	0.22	0.08
Natural mortality (M)	1.05	0.69
Exploitation rate (E)	0.26	0.11

Table 5. Length at capture of *B. nurse* and *B. macrolepidotus*

Length at capture	<i>B. nurse</i>	<i>B. macrolepidotus</i>
L_{25}	12.88 cm	15.13 cm
L_{50}	14.49 cm	17.64 cm
L_{75}	16.10 cm	20.16 cm




Figure 4. Length converted catch curves of *B. nurse*.




Figure 5. Length converted catch curves of *B. macrolepidotus*.




Figure 6. Length at first capture (L_{50}) for *B. nurse*.




Figure 7. Length at first capture (L_{50}) for *B. macrolepidotus*.

The Beverton-Holt relative yield per recruit (Y'/R) and relative biomass per recruit (B'/R) were estimated using selective Ogive procedure of FiSAT. The analysis indicated that the exploitation rate, which maximises

yield per recruit, produced values of $E_{max} = 0.41, 0.42$; $E_{10} = 0.36, 0.36$ and $E_{50} = 0.28, 0.29$ for *B. nurse* and *B. macrolepidotus*, respectively (Figs 7 & 8).




Figure 8. Stock status of *B. nurse* using Beverton and Holt's relative Y/R analysis.




Figure 9. Stock status of *B. macrolepidotus* using Beverten and Holt's relative Y/R analysis.

Discussion

The calculated 'b' values of the LWRs, which were 3.2103 for *B. macrolepidotus* and 3.5405 for *B. nurse*, indicate positive allometric growth. The values fall within the acceptable range of 2.5 and 3.5, which is typical for tropical fish stocks (Carlander, 1969; Froese, 2006). Gopakumar *et al.* (1991) opined that higher b values imply a relatively productive environment. Positive allometric growth implies that the fish grows faster in weight than in length. A similar observation was also reported in *B. nurse* (3.0737) from the White Volta River in Ghana (Abobi & Ekau, 2013). Echi & Ezenwaji (2016) reported contrasting findings for *B. macrolepidotus* from the Anambra River, Nigeria with the males and females showed negative allometric growth (2.309 and 2.734, respectively); in the same vein, the other Alestid species reported in the study, such as *Alestes baremoze*, *B. leuciscus*, and *Hydrocynus vittatus*, showed negative allometric growth for their respective male and female. The slight difference in the exponent 'b' values recorded could be associated with several factors, such as temperature, salinity, adequate food, seasonal changes, environmental conditions, sampling procedure, and sexual dimorphism.

The condition factor "K" of a fish can be defined as a measurement of the general health condition of the fish as calculated by the ratio of body weight to body length. This factor implies that whatever is able to affect the weight of fish will definitely affect the K values. Adesulu & Sydenham (2007) reported that condition factor in fish is mainly affected by the amount of food in stomach and stage of egg development that will affect the weight of the body and then the weight of fish is a function of condition factor in fish. The "K" values of the two Alestid species in this study indicate that the population is in moderate growth condition as "K" values are slightly greater than 1. This factor

implies that the fishes thrive well in the habitat (Komolafe & Arawomo, 2011). The condition factor for *B. longipinnis* was reported to range 1.94-2.80 in Ikomi & Sikoki (2003).

The asymptotic length (L_{∞}) is the largest theoretical mean length that a fish could attain in its natural habitat, assuming the fish grows throughout its life (Abobi & Ekau, 2013); while the growth curvature (K) is the rate at which it grows towards this final size (Etim *et al.*, 1999). In the present study, *B. nurse* had an asymptotic length (L_{∞}) of 24.46 cm, growth curvature (k) of 0.52 per year, and t_0 (per year) of 0.04 while *B. macrolepidotus* had L_{∞} of 28.88 cm, k of 0.22 per year, and t_0 (per year) of 0.07 (Table 4). These results indicate that both species are fast-growing.

The total mortality (Z), natural mortality (M / year), fishing mortality (F), and exploitation rate (E) of *B. nurse* were found to be 1.88, 1.05, 0.83 and 0.26, respectively; while for *B. macrolepidotus* it was 0.77, 0.69, 0.08 and 0.11, respectively. Values of Z, M, and F were higher for *B. nurse* than for *B. macrolepidotus*; however, since natural mortality (M) exceeded fishing mortality (F) for both species, the stock is not over-exploited. The exploitation rate (E) of *B. nurse* was 0.26. Based on the assumption that a stock is optimally exploited when $F=M$ or $E=0.5$ (Gulland, 1971). The results of the present study indicate that the current exploitation rate is close to the maximum level of 0.5. The low value of E in the two species in this finding suggests that levels of exploitation may be increased in order to reduce waste of stock through natural mortality; however, caution must be taken to avoid what can be called a skewed exploitation situation (Chukwu & Deekae, 2010). Ahmad *et al.* (2018) reported lower values of Z (1.31), M (1.0), F (0.31), and E (0.24) for *B. nurse* from the Challawa Gorge dam, Kano. Higher values of the fore mentioned parameters were observed for *B. nurse* by Kwarfo-

Apegyah *et al.* (2009) and Uneke & Nwani (2013), respectively; $Z = 2.54$ and 3.34 ; $M=1.3$ and 1.80 ; $F=1.24$ and 1.54 ; and $E=0.49$ and 0.46 . However, the relatively lower values of E in these studies indicate that the stocks were also not over-exploited.

The result of this study showed the estimated length at first capture (L_c) of both *Brycinus* species to be 14.49 and 17.64 cm for *B. nurse* and *B. macrolepidotus*, respectively. The L_c value recorded in this study falls within the range of the total length 7.30 to 23.00 cm for *B. nurse* and 11.50 - 27.00 cm for *B. macrolepidotus*. A lower value of L_c (7.58) was recorded for *B. nurse* by Ahmad *et al.* (2018). Uneke and Nwani (2013) found the L_c for *B. nurse* in Cross River to be 8.83 while Kwarfo-Apegyah (2008) found the L_c for *B. nurse* in Bontanga Reservoir (Northern Region of Ghana) to be 10.94 .

The predicted E_{max} of the Selective Ogive procedure for *B. nurse* and *B. macrolepidotus* (0.41 and 0.42 , respectively) were higher than their respective current exploitation rates E (0.28 and 0.29) showing that both *Brycinus* species were not over-exploited. Uneke & Nwani (2013) and Kwarfo-Apegyah *et al.* (2009) reported the E of *B. nurse* to be 0.46 and 0.49 , respectively; both values are higher than the value of E estimated for the two Alestid species in this study. Values of E_{max} , E_{50} , and E for *B. nurse* in the Challawa Gorge dam, Kano, were reported by Ahmad *et al.* (2018) as 0.65 , 0.2 , and 0.24 , respectively.

CONCLUSIONS

The present study provides some important baseline information on the growth and population of two commercially important fish species from the family Alestidae that will help in the development of efficient management strategies. The length-at-first capture (L_c) of *B. macrolepidotus* estimated in this study showed that this species was caught at very small sizes concerning the use of small mesh size nets, such as 2.3 mm and 15 mm. Natural mortality was greater than the fishing mortality for both *B. nurse* and *B. macrolepidotus* while exploitation rate (E) was less than 0.5 for both species. Even though the difference between biological reference point and fishing mortality is small, it would be better for the well-being of fishery to introduce some fishery regulation measures.

REFERENCES

Abobi, S. M., & Ekau, W. (2013). Growth, Mortalities and Exploitation Rates of *Alestes baremoze*(Joannis, 1835), *Brycinus nurse*

(Rüppell, 1832) and *Schilbe intermedius* (Rüppell 1832) from the lower reaches of the White Volta River (Yapei), Ghana. *Journal of Agriculture and Biodiversity Research*, 2(1), 1-10.

Abowei, J. F. N., & Hart, A. I. (2009). Some morphometric parameters of ten finfish species from the lower Nun River, Niger Delta, Nigeria. *Research Journal of Biological Sciences* 4(3), 282-288.

Adesulu, E. A., & Sydenham, D. H. J. (2007). The freshwater and fisheries of Nigeria. Macmillan Nigeria Publishers, Lagos, Nigeria. 397 pp.

Ahmad, I. M., Yola, I.D., & Suleiman, N. (2018). Mortality and Exploitation Rates of Challawa Gorge Dam Fishes, Kano State, Nigeria. *Journal of Fisheries and Livestock Production*, 6: 262, DOI: 10.4172/2332-2608.1000262.

Arroyave, J., & Stiassny, M. L. J. (2011). Phylogenetic relationships and the temporal context for the diversification of African characins of the family Alestidae (Ostariophysi: Characiformes): Evidence from DNA sequence data. *Molecular Phylogenetics and Evolution*, 60, 385–397.

Beverton, R. J. H., & Holt, S. J. (1966). Manual of methods for fish stock assessment. Part II. Tables of yield function. *FAO Fisheries Technical Paper*, 38(1), 67p.

Beverton, R.J.H. & Holt, S.J. (1957) On the Dynamics of Exploited Fish Populations. Gt Britain Fish Invest. Ser. 2, Vol. 19, 1-533.

Boulenger, D. P. (2002). The biology of *Brycinus* species from Lekki Lagoon, Lagos State, Nigeria *Journal of Science*, 12, 73-84.

Carlander, K. D. (1969). *Handbook of freshwater fishery biology*. The Iowa State University Press, Ames, IA. 1:752p.

Chukwu, K. O., & Deekae, S. N. (2010). Growth, mortality, and recruitment of *Periophthalmus barbarus* (Linnaeus, 1766) in New Calabar River, Nigeria. *Agriculture and Biology Journal of North America*, 2(7), 1066-1068.

Echi, P. C. & Ezenwaji, H. M. G. (2016). Length-Weight relationships and food and feeding habits of some Characids (Osteichthyes: Characidae) from Anambra River basin, Nigeria. *Animal Research International*, 13(1), 2316-2320.

- Etim, L., Lebo, P. E., & King, R. P. (1999). The dynamics of an exploited population of a siluroid catfish (*Schilbe intermedius*, Reupell 1832) in the Cross River, Nigerian. *Fisheries Research*, 40, 295–307.
- Froese, R., & Pauly, D. (eds.) (2017). List of Freshwater Fishes reported from Nigeria. FishBase. World Wide Web electronic publication. Retrieved from www.fishbase.org. Accessed 1 August 2018.
- Froese, R., & Pauly, D. (Editors) (2016). FishBase. World Wide Web electronic publication. [Online]. www.fishbase.org. Accessed 14 September, 2018.
- Froese, R. (2006). Cube law, condition factor, and length-weight relationships: History, meta-analysis, and recommendations. *Journal of Applied Ichthyology*, 22, 241-253.
- Gayanilo, F. C., Sparre, P., & Pauly, D. (2005). FAO-ICLARM Stock Assessment Tools II (FiSAT II). User's guide. *FAO Computerized Information Series (Fisheries)*. No. 8, revised version, FAO, Rome, 168 pp.
- Gopakumar, G., Pillai, P. P., & Koya, K. P. (1991). Population characteristics of tuna live baits in Lakshadweep. *Journal of Marine Biological Association India*, 33(1-2), 255-277.
- Gulland, J. A. (1971). The Fish Resources of the Ocean. 1st Edn. Fishing News Books, London, pp. 255.
- Hoggarth, D. D., Abeyasekera, S., Arthur, R., Beddington, J. R., Burn, R. W., Halls, A. S., Kirkwood, G. P., McAllister, M., Medley, P., Mees, C. C., Pilling, G. M., Wakeford, R. & Welcomme, R. L. (2006). Stock assessment and fishery management - A framework guide to the FMSP stock assessment tools", FAO Fisheries Technical Paper No. 487, Rome, Italy, 261 pp.17
- Hossain, M. Y., Ohtomi, J., & Ahmed, Z. F. (2009). Morphometric, meristic characteristic and conservation of the threatened fish, *Puntius sarana* (Hamilton 1822) (Cyprinidae) in the Ganges River, northwestern Bangladesh. *Turkish Journal Fisheries and Aquatic Sciences*, 9, 223-225.
- Ikomi, R. B., & Sikoki, F. D. (2003). Studies on the ecology of the African longfin tetra, *Brycinus longipinnis* (Gunther, 1864) in the Jamieson River (Niger Delta, Nigeria) *Acta Ichthyologica Piscatoria*, 33(1), 17–36.
- Kohler, N., Casey, J. & Turner, P. (1995). Length-weight Relationship for 13 Species of Sharks from Western North Atlantic. *Fisheries Bulletin*, 93, 412-414.
- Komolafe, O. O., & Arawomo, G. A. O. (2011). Observations in the composition, physiological condition, and fisheries in Erinle Lake, Osun state, Nigeria. *West African Journal of Applied Ecology*, 18, 71-78.
- Kwarfo-Apegyah, K. (2008). Ecology and stock assessment of major fish species of Bontanga reservoir for sustainable management. (Ph.D. Thesis.) Department of Oceanography and Fisheries, University of Ghana, Legon, 216 pp.
- Kwarfo-Apegyah, K., Ofori-Danson, P. K., & Nunoo, F. K. E. (2009). Exploitation rates and management implications for the fisheries of Bontanga Reservoir in the Northern region of Ghana. *West African Journal of Applied Ecology*, 14, 1-7. <http://dx.doi.org/10.4314/wajae.v14i1.44710>
- National Bureau of Statistics (NBS) (2017). Nigerian's fish production 2010–2015. Data is supplied, verified and validated by the National Bureau of Statistics, Nigeria (NBS). <https://www.nigerianstat.gov.ng/elibrary>. Accessed on 17 June 2018
- Olopade, O. A., Dienye, H. E., & Eyekpegba, A. (2018). Length frequency distribution, length-weight relationship and condition factor of cichlid fishes (Teleostei: Cichlidae) from the New Calabar River, Nigeria. *Iranian Journal of Ichthyology* 5(1), 74-80.
- Patterson, K. (1992). Fisheries for small pelagic species: an empirical approach to management targets. *Reviews in Fish Biology and Fisheries*, 2, 321-338. <https://doi.org/10.1007/BF00043521>
- Pauly, D. & Soriano, M. L. (1986). Some practical extensions to Beverton and Holt's relative yield-per-recruit mode. In: Maclean, J.L. Dizon, L.B., Hosillos. L.V. (Eds.) The First Asian Fisheries Forum, Asian Fisheries Society, Manila, Philippines, pp. 491-496.
- Pauly, D. (1984). Fish population dynamics in tropical waters: a manual for use with programmable calculators. *ICLARM Stud. Rev.*, (8), 325.

- Pauly, D. (1980). On the interrelationships between natural mortality, growth performance and mean environmental temperature in 175 fish stock. *Journal of Marine Sciences*, 39(2), 175-192.
- Paugy, D., & Schaefer, S. A. (2007). Alestidae. p. 347-411. In: Stiassny, M.L.J., Teugels, G.G. and C.D. Hopkins (Editors), *Poissons d'eaux douces et saumâtres de basse Guinée, ouest de l'Afrique centrale/The fresh and brackish water fishes of Lower Guinea, west-central Africa*. Vol. 1. Coll. Faune et Flore tropicales 42. Institut de recherche pour le développement, Paris, France, Muséum nationale d'histoire naturelle, Paris, France and Musée royale de l'Afrique centrale, Tervuren, Belgique. 800 p.
- Paugy, D. (2003). Alestidae. p. 236-282. In: Paugy, D., Lévêque, C. and Teugels, G. G. (Editors), *The fresh and brackish water fishes of West Africa* Volume 1. Coll. faune et flore tropicales 40. Institut de recherche de développement, Paris, France, Muséum national d'histoire naturelle, Paris, France and Musée royal de l'Afrique Central, Tervuren, Belgium, 457 pp.
- Reed, W. J., Buchard, A. J., Hopson, J., Jennes, J., & Yaro, I. (1967). *Fish and fisheries of Northern Nigeria*. Ministry of Agriculture, Northern Nigeria. 226 pp.
- Ricker, W. E. (1975). Computations and interpretation of biological statistics of fish populations. *Fisheries Research Board Canada Bulletin*, 191: 382.
- Ricker, W. E. (1973). Linear regressions in fishery research. *Journal Fisheries Research Board Can.* 30, 409–434.
- Ricker, W. E. (1968). *Methods for Assessment of Fish Production in Freshwaters*. IBP Handbook No. 3. Blackwell Scientific Publications, Oxford and Edinburgh, UK. 313 pp.
- Saliu, J. K., & Fagade, S. O. (2004). The Structural and Functional Morphology of the Hard Parts of *Brycinus nurse* (Pisces: Cypriniformes, Characidae), From Asa Reservoir, Ilorin, Nigeria. *Turkish Journal of Fisheries and Aquatic Sciences*, 4: 23-31
- Uneke, B. I., & Nwani, C. D. (2013). Stock assessment of *Brycinus nurse* (Characiformes: Alestidae) in a tropical flood river basin. *Continental Journal of Fisheries and Aquatic Science*, 7(3), 22-33.
- Wang, Y. & Liu, L. (2006). Estimation of natural mortality using statistical analysis of fisheries catch-at age data. *Fisheries Resources*, 78, 342-351.