

FOREST SOIL HETEROGENEITY AND SOIL SAMPLING PROTOCOLS ON LIMESTONE OUTCROPS: EXAMPLE FROM SW CHINA

PROTOKOL VZORČENJA IN HETEROGENOST GOZDNE PRSTI NA APNENČASTIH IZDANKIH: PRIMER IZ JZ KITAJSKJE

ZHOU Yunchao¹, WANG Shi-jie^{1*}, LU Hong-mei^{1,2}, XIE Liping^{1,2} & XIAO De'an^{1,2}

Abstract

UDC 630*114:551.44(510)

Zhou Yunchao, Wang Shi-jie, Lu Hong-mei, Xie Liping & Xiao De'an: Forest Soil Heterogeneity and Soil Sampling Protocols on limestone outcrops: Example from SW China

Forest soil heterogeneity of outcrop karst was studied by testing soil organic carbon at a selected 20 m × 30 m plot set up in Maolan primitive karst forest area, southern Guizhou Province, China. 90 microhabitats in the plot are roughly classified into 7 distinct microhabitat. There was great heterogeneity in the plot not only for soil distribution but for great difference of soil organic carbon content. Soil organic carbon (SOC) contents were much higher, ranging from 40.1 g/kg to 203.5 g/kg. The CV (coefficient of variation) value of SOC from 38 soil sampling points, 22 sampling microhabitats, each microhabitat type, and individual microhabitat was 43%, 41%, 2.2% to 42%, 14% to 57.3% respectively. This showed high soil heterogeneity in terms of soil distributing pattern, size and soil organic carbon SOC content in outcropped karst. Different methods were applied to calculate SOC content in the plot. Small discrepancies suggested that the mixed soil samples collected from the major microhabitat types covering 95% soil distributing area were capable of representing soil nutrient status of the whole plot.

Keywords: karst, prst, organic carbon, sampling protocol, Southwestern China.

Izveček

UDK 630*114:551.44(510)

Zhou Yunchao, Wang Shi-jie, Lu Hong-mei, Xie Liping & Xiao De'an: Protokol vzorčenja in heterogenost gozdne prsti na apnenčastih izdankih: primer iz JZ Kitajske

Z analizo organskega ogljika smo raziskovali heterogenost gozdne prsti na testni površini velikosti 20 m x 30 m. Raziskave so potekale v gozdnem območju Maolan v južnem delu province Guizhou, Kitajska. Na površini smo našli 90 mikrohabitats in jih razvrstili v 7 različnih mikrohabitatsnih tipov glede na položaj vzorčnega mesta. Tudi količina organskega ogljika (SOC, soil organic content) v prsti kaže na veliko heterogenost prsti. Ta zavzema vrednosti med 40,1 g/l in 203,5 g/l. Koeficient variacije SOC na 28 vzorčevalnih mestih vzorčenj je 43%, med 22 mikrohabitatsi 41%, znotraj mikrohabitatsnega tipa 2.2% do 42% in znotraj posameznega mikrohabitata med 14% in 57,3%. Pri izračunu SOC smo uporabili različne metode, majhna odstopanja pa kažejo, da mešani vzorci prsti iz glavnih mikrohabitatsnih tipov, ki pokrivajo 95% testne površine, reprezentativno predstavljajo vsebnost hranil na celotni testni površini.

Ključne besede: kras, prst, organski ogljik, vzorčenje prsti, južna Kitajska.

¹ State Key Laboratory of Environment Geochemistry, Institute of Geochemistry Chinese Academy of Sciences, Guiyang, e-mail: fc.yczhou@gzu.edu.cn, wangshijie@vip.gyig.ac.cn;

² Graduate School of Chinese Academy of Sciences, Beijing, e-mail: hongmeilu0826@126.com, Xieliping@163.com, xiaodean@hotmail.com

Received/Prejeto: 28.8.2009

INTRODUCTION

Soil heterogeneity has been world-wide interested recently (Boerner *et al.* 1998; Wang *et al.* 1998; Park & Vlek 2002; Yang *et al.* 2002; Jhorar *et al.* 2004). Researchers either focused on assessing properties of heterogeneous soils based on valid sampling protocols or on identifying the aims to study the validities of sampling protocols (Theocharopoulos *et al.* 2001; Barahona & Iriarte 2001; Fernando *et al.* 2001; Donohue 2002; Myeong *et al.* 2003; Tamminen 2003; Kulmatiski & Beard 2004; Whalen 2004; Bramley *et al.* 2005). However, all the previous studies about soil heterogeneity were only conducted in areas of intact soil mantles; no study was conducted in outcrop karst mountainous regions.

Attentions have been absorbed to outcrop karst soils (Drew 1983; Gillieson *et al.* 1994; Atalay 1997; Cerda 1998; Shang & Tiessen 2003; Parise & Pascali 2003; Wang *et al.* 2004a) for soil erosion (Cerda 1998), soil sediment recorded paleoenvironment (Sauro *et al.* 2009), environmental pollution (Parise & Pascali 2003) and soil carbon sequestration (Shang & Tiessen 2003). Because of susceptible karst environment, pressure of

sharp population growth accelerated soil erosion and karst desertification (or karst desert, rocky desert, bare karst). Karst desert area in Guizhou, China, one of the biggest karst desertification center in the world, is up to 11×10^4 km², and also accelerating in much higher speed (Wang *et al.* 2004a). With the high speed expanding of urbanization and industrialization, high quality arable land was invaded, ecosystems were interrupted, especially soils. Karst rocky desertification, one of the serious eco-environmental problems, happened in karst mountainous areas in southwestern China (Wang *et al.* 2004a, b), several studies have concerned on soil property in this region (Liu *et al.* 2005; Li *et al.* 2006; Zhang *et al.* 2006). Unfortunately, the distinct soil heterogeneity of karst mountainous regions was not taken into consideration in their soil sampling methods. How can we appropriately evaluate the change of outcrop karst soil quality?

The present study was undertaken with the following two objectives: (1) comprehending the pattern of soil heterogeneity of outcrop karst, and (2) proposing a valid way to sample in outcrop karst soil.

MATERIALS AND METHODS

STUDY AREA

Maolan karst forest (107°52'10" – 108°05'40"E, 25°09'20" – 25°20'50"N) is located in southern Guizhou Province, China. This area has preserved a primitive forest ecosystem with an area of 19,640 hectares, which is the only karst virgin forest survived at the same latitude on the Earth. This area topographically presents a typical, well-developed peak cluster karst (fengcong in Chinese). Peak clusters constitute a subtype of tower karst, characterized by many dolines scattered among groups of large stone peaks sharing the same base. The outcropped rocks are composed of limestone of Carboniferous period, covering around 70%~80% of the forest. The subtropical monsoon climate is warm and humid with an average annual temperature of 15.3°C and average annual precipitation of 1,752.5 mm. More than 81% precipitation (about 1,419.6 mm) occurs from April to September. Forest ecosystem has developed diverse plants, involving 154 families, 514 geneses, 120 species of vascular bundle plants, of which there are 143 families, 494 geneses, and 1,172 species of spermatophytes (Lan 1987; Chen 1997). Recently, Maolan karst forest ecosystem has attracted more attention worldwide to multidisciplinary researches pertaining to biology, ecology, geology, and environment.

The investigated area was specified on a typically plot of 20×30 m² (108°02'29"E, 25°16'21"N; Fig. 1) with a slope of 38°, with predominant yellow limestone soil. Soil layers are shallow from several cm to 60 cm, soil fragments distribute in patches among bare limestone and soil constructions generally show transition tendency from A-B-R to A-R.

FIELD INVESTIGATION

Prior to sampling soil, soil distribution was carefully investigated and classified. The identified 90 individual microhabitats could be roughly classified into 7 microhabitat types on the basis of micro-terrain, ratio of exposed rock, soil covering site and size, etc. (Fig. 1). (1) Rocky: almost all exposed rocks. (2) Soil: ratio of exposed rock was lower than 30%, soil covered uniformly, and the area was larger (length and width were more than 2 m). (3) Rock-soil: ratio of exposed rock was more than 30%; soil cover was less than (2). (4) Karren: Depth/width ratio was less than 2 with varied thickness of soil at the bottom of karren. (5) Grike: Depth/width ratio was more than 2 with thin soil at the bottom of the grike. (6) Stone pit: irregular deep-eroded depression with 0.1 ~ 1 m² soil cover. (7) Stone hole: various sizes of channels formed by water




Fig. 1: Location of test site and the distribution of microhabitats in the plot (arrow indicated the number of microhabitat, and the 3-53 indicated the microhabitat number in the plot).

corroded (Fig. 2, Tab. 1). Soil was covering 170 m², only 28.4% of the plot, but the outcropped carbonate rocks covering 71.6%. It was difficult to apply grids sampling method (Leo 1963; Shang & Tiessen 2003) in outcrop

karst for the reason that soil was isolated by limestone rocks, soil properties of each individual microhabitat was driven by different water, air, heat, organism, etc.

Tab. 1: Statistics of various microhabitat types in the investigated plot.

Microhabitat type	Description	Number	Size / m ²	Size proportion (Plot base) / %	Soil size / m ²	Soil proportion (Plot base) / %
Rock	Large size of carbonate exposure. Litter but less soil is available at small flat or hollow sites	46	291.1	48.5	0	0
Soil	Rock exposure rate < 30 %, well-proportioned soil layer, size > 2 m × 2 m.	14	49.7	8.4	26.5	15.6
Rock-soil	Rock exposure rate > 30 %, size of soil layer < 2 m × 2 m.	5	117.0	19.5	93.0	54.7
Karren	Rock erosion gully, depth/ width ratio < 2, disproportionate soil layer and litter accumulated at bottom.	6	120.6	20.1	41.5	24.4
Grike	Depth/width ratio > 2, small amounts of soil at bottom, litter accumulated.	3	5.1	0.8	2.3	1.4
Stone pit	Irregularly-shaped eroded depression, soil distributing size is around 0.1-1 m ² .	11	10.9	1.8	1.1	0.6
Stone hole	Tubular channels of various sizes from water erosion, 10-30 cm at depth.	5	5.6	0.9	5.6	3.3
Total		90	600	100	170	100




Fig. 2: Type of microhabitats (photo: a-rocky; b-soil; c-rock-soil; d-karren; e-grike; f-stone pit; g-stone hole. Photos: S. J. Wang).

SOIL SAMPLING AND ANALYSIS

Soil samples were collected from microhabitats to a depth of 20 cm (or less where soils were too shallow). Only one sample was collected at a microhabitat which is topographically flat with uniform litter layers, but three replicated samples were collected from 8 individual microhabitats existing significant differences in litter depth, slope, etc. At each sampling site, 3-5 points were collected and mixed to be a final sample. No soil sample was collected from the microhabitats of all rocky and grike,

some karrens and rock-soil for the reason of limited soil depth. 38 soil samples were collected from 22 individual microhabitats where sufficient soils are available in the plot. Each sample was 1 kg and air-dried, stones, visible litters and roots were removed before grinding to 2 mm, and plant micro piece was removed by using a plastic bar of static electricity produced by plastic bar friction. A sub-sample was milled to pass 0.25 mm sieve to test SOC by using potassium dichromate oxidation method of Walkley-Black procedure (Nelson & Sommers 1982).

RESULTS AND DISCUSSION

SOIL HETEROGENEITY IN TERMS OF SOIL DISTRIBUTION

Soil heterogeneity in the investigated plot was obvious in terms of soil distribution. Rocky, one of microhabitat types, predominated in size, accounts for almost 48.5% of the plot, soil and rock-soil for 28%, karren for 20%, and grike, stone pit and stone hole for 3.5% (Tab. 1). Soil covered in five microhabitat types (soil, rock-soil, karren, stone pit and stone hole) account for more than 99% of the total 170 m² soil area. Among them, three major microhabitat types involved of soil, rock-soil, karren, account for about 95% of the total soil area. Soil distributed in patches among bare limestone.

SOIL HETEROGENEITY IN ORGANIC CARBON

Soil organic carbon (SOC) was extremely heterogeneous in the plot, ranging from 40.1 g/kg to 203.5 g/kg (Tab. 2). Compared with red soil of 26 g/kg and yellow soil of 28 g/kg of forest and/or grasslands in China (Xiong & Li 1987), mass humus combined with Ca²⁺ and Mg²⁺ to form highly polymer for the reason of Ca-rich in karst soil, calcium humic acid was stabilized and accumulated in karst soil (He *et al.* 2001). The CV (coefficient of variation) value of SOC from 38 soil sampling points was 43% (Tab. 2), much higher than those in non-karst areas. For instance, the CV of organic carbon of 625 m² irrigation-silting soils in Aksu area of Xinjiang, China was not more than 17.84% (Yang *et al.* 2002).

With respect to 22 microhabitats, the CV of organic carbon was 41%, close to 43% from all the 38 soil-sampling points (Tab. 2).

SOC contents of the five microhabitat types, accounted for 99% of the total soil area in the plot, were 40.1~107.2 g/kg (Soil), 86.9~165.8 g/kg (rock-soil), 47.3~203.5 g/kg (karren), 41.0~72.0 g/kg (stone pit) and 102.5~154.7 g/kg (stone hole), respectively. Except for microhabitat type of karren, the variation coefficients of SOC in the other 4 of microhabitat types varied within the range of 22%~28%, much lower than 41% calculated on the basis of 22 individual microhabitats (Tab. 2). Meanwhile, it could be seen that average SOC content in microhabitat types showed significant variation from one another, the order was: stone pit (124.8 g/kg) > karren (121.1 g/kg) > rock-soil (112.4 g/kg) > Soil (81.5 g/kg) > stone hole (57.8 g/kg). Particularly, dominated 3 microhabitat types (soil, rock-soil and karren), more than 95%

ren had the greatest soil heterogeneity. Soil covering size for each of the 8 individual microhabitats ranged from 1.5 m² to 15.7 m² (Tab. 1). The CV values of SOC in the karst regions were relatively large when compared with that in non-karst regions in the same scale. For example, CV of surface soil organic carbon was 11.2% for a 50 m² plot in the Xilin River watershed in Inner Mongolia, but was only 6.25% for the 4 m² surface soil in the same plot (Wang *et al.* 1998). Therefore, it could be inferred that soil samples should be collected to represent individual microhabitat in the karst region by using a multi-point sampling protocol.

ESTIMATION OF SOIL ORGANIC CARBON IN OUTCROP KARST AREA AND SOIL SAMPLING PROTOCOL

It was evident from the above discussion that karst forest soil was significantly heterogeneous. This was manifested

Tab. 2: Concentrations of soil organic carbon in each investigated microhabitat types in the plot.

Microhabitat type	Number of samples	Minimum (g/kg)	Maximum (g/kg)	Average (g/kg)	SD	CV (%)
Karren	6	47.3	203.5	121.1	50.3	42
Soil	5	40.1	107.2	81.5	22.8	28
Rock-soil	3	86.9	165.8	112.4	26.9	24
Stone pit	3	102.5	154.7	124.8	26.9	22
Stone hole	5	41.0	72.0	57.8	13.0	23
All above microhabitat types	22	41.0	203.5	97.1	40.1	41
Total sampling points	38	40.1	203.5	98.5	42.4	43

of the total soil area of the plot, also showed significant variations in SOC one another. From above, highly varied SOC values clearly indicated that the microhabitat types were considerably different in soil properties one another. Therefore, it was unwise to collect soil sample only from one single microhabitat types for purpose of revealing soil nutrient levels.

HETEROGENEITY OF SOIL ORGANIC CARBON IN INDIVIDUAL MICROHABITAT

The variations of SOC in 8 individual microhabitats were listed in Tab. 3. CV values of soil organic carbon generally varied from 14% to 25%, slightly lower than the CV values (22%~28%) for the same microhabitat types (Tab. 2), the maximum CV value was up to 57.3% in karren. These results implied that karst microhabitats were significantly heterogeneous in terms of SOC not only in the same microhabitat types, but also among microhabitats and in an individual microhabitat. Moreover, the results also indicated that microhabitat type of kar-

ren not only by alternative distribution of soil and carbonate rock, but also by variations of SOC among sampling sites in an individual microhabitat. Because of high heterogeneity of outcrop karst forest soil, the results of the conventional sampling methods which disregarded soil heterogeneity may not be compared with one another. In the case of soil organic carbon, it was not representative based on the equivalent values from one single sample site or individual microhabitat, or from one microhabitat types.

There were two ways to calculate SOC content of the plot. One was to calculate arithmetic mean of 38 sampling points; another was to calculate weighted average of all soil sampling microhabitats. Because of the great heterogeneity of outcrop karst soil, arithmetic mean value was not more accurately indicated the content of SOC than weighted average value. Estimating soil organic carbon that might most approximately represent the soil property of a significantly heterogeneous outcrop karst region, the weighted value based on soil distributing size

Tab. 3: Variation of contents of soil organic carbon in selected samples.

Microhabitat type	Microhabitat No.	Number of sample	Minimum (g/kg)	Maximum (g/kg)	Average (g/kg)	SD	CV (%)
Soil	3-10	3	75.6	106.9	93.9	16.3	17
	3-25	3	51.5	70.7	62.4	9.8	16
	3-40	3	40.1	61.3	53.0	11.3	21
Rock-soil	3-17	3	107.5	165.8	140.6	29.9	21
	3-34	3	91.4	139.2	109.8	25.7	23
Karren	3-16	3	61.3	98.0	76.7	19.0	25
	3-21	3	134.2	176.5	160.5	22.9	14
	3-50	3	47.3	169.3	106.6	61.1	57

for all individual microhabitats in the plot must be taken into consideration. Weighted value based on soil distributing size could guarantee to obtain representative SOC contents in karst mountainous area.

In this study, a calculation method was proposed for estimating weighted SOC based on soil distributing size, and it was described by the following formula:

$$SOC = \sum_{j=1}^m \left[\sum_{i=1}^n \left(S_i \times \frac{A_i}{A_j} \right) \times \frac{A_j}{A} \right]$$

Where SOC was the weighted values of soil organic carbon based on soil distributing size in the plot (g/kg), *j* was microhabitat type *j*, *i* was each individual microhabitat *i*, *S_i* was the average soil organic carbon content of each individual microhabitat *i* (g/kg), *A_i* was the soil distributing size of individual microhabitat *i* (m²), *A_j* was the soil distributing size of microhabitat type *j* (m²), and *A* was the total soil distributing size in the plot (m²).

According to the above formula, SOC content of weighted average value was 95.6 g/kg from the total 38 sampling sites of the investigated plot (Approach 1), and 91.1 g/kg from the three major microhabitat types (soil, rock-soil and karren) of covering 95% soil distributing area of the plot (Approach 2; Tab. 4). Both SOC values were quite close (± 3%). If 95% soil distributing area of

these three major microhabitat types was used to represent the soil distributing area of the whole investigated plot (Approach 3), weighted SOC based on soil distributing size was 92.1 g/kg (Tab. 4), quite close to the values of Approach 1. Therefore, there were small differences among the above three estimations (Approaches 1-3). By the way, if randomly sampling (Liu *et al.* 2005; Li *et al.* 2006; Zhang *et al.* 2006) a few points in relatively large and intact soil fragments to represent the plot, SOC content was only 81.5 g/kg (Tab. 2), much lower than above values.

The weighted average SOC value of 92.1 g/kg from Approach 3 had minor discrepancy from the arithmetic mean SOC value 98.5 g/kg (Tab. 2) based on all sampling sites. This finding from Approach 3 was quite significant for study on nutrient status in heterogeneous soils of outcrop karst area. Firstly, in case of large numbers of soil microhabitats, only small numbers of soil samples were required to collect and analyze from dominating microhabitat types that accounted for around 95% soil distributing of the studied area. Secondly, it was significant to avoid large discrepancies of nutrient levels, such as SOC, among various research results in outcrop karst areas for purpose of comparison. Suggestion of soil sampling was to mix soil samples collected from Approach 3 in the plot were capable of representing the soil nutrient status of the whole plot.

Tab. 4: Calculated average content of topsoil organic carbon in the plot (unit in g/kg).

	100 % soil distributing area	95 % soil distributing area	95 % soil distributing area representing the whole plot
Estimated average value based on soil distributing size quotient	95.6 (Approach 1)	91.1 (Approach 2)	92.1 (Approach 3)

CONCLUSIONS

Soils in Maolan outcrop karst forest area were extremely heterogeneous in terms of soil distribution size, and SOC contents. It required an appropriate protocol to effectively study outcrop karst soil property. A representative sampling protocol based on soil distributing size weight proposed by this study successfully demonstrated advantage to effectively determine nutrient levels in heterogeneous soils of outcrop karst mountainous areas.

It is suggested that only small numbers of soil samples were required to collect and analyze from major mi-

crohabitats that account for around 95% soil distributing area of the studied karst. In this way, estimated average contents of nutrient matters such as SOC were capable of representing soil nutrient status of the whole studied karst areas, and of good comparisons among various research results as well. This soil sampling protocol could also contribute to study on heterogeneous soil property in non-karst areas.

ACKNOWLEDGEMENTS

This work was jointly funded by the National Key Basic Research Development Program (Grant No. 2006CB403200), Orientation Project of Knowledge Innovation Program of Chinese Academy of Sciences (No. kzcx2-yw-306), and the National Natural Science Foun-

dation of China (Grant No. 40721002 and 40672112). We sincerely thank to Dr. ZSENI Anikó and another anonymous reviewer for their valuable comments and suggestions for improving the manuscript.

REFERENCES

- Atalay, I., 1997: Red Mediterranean soils in karstic regions of Taurus Mountains, Turkey.- *Catena*, 28, 247–260.
- Barahona, E. & A. Iriarte, 2001: An overview of the present state of standardization of soil sampling in Spain.- *The Science of the Total Environment*, 264, 169–174.
- Boerner, R. E. J., Scherzer, A. J. & J. A. Brinkman, 1998: Spatial patterns of inorganic N, P availability, and organic C in relation to soil disturbance: a chronosequence analysis.- *Applied Soil Ecology*, 7, 159–177.
- Bramley, R. G. V. & L. J. Janik, 2005: Precision agriculture demands a new approach to soil and plant sampling and analysis – Examples from Australia.- *Communications in Soil Science and Plant Analysis*, 36, 9–22.
- Cerda, A., 1998: Relationships between climate and soil hydrological and erosional characteristics along climatic gradients in Mediterranean limestone areas.- *Geomorphology*, 25, 123–134.
- Chen, Z., R., 1997: Karst forest investigation in Maolan.- In: Zhu S. Q. (ed.) *Study of Karst forest ecosystem II*. Guizhou Science & Technology Press, pp.167–172, Guiyang [in Chinese].
- Donohue, S. J., 2002: Evaluation of soil nutrient variability for development of turfgrass soil test sampling methods.- *Communications in Soil Science and Plant Analysis*, 33, 3335–3345.
- Drew, D. P., 1983: Accelerated soil erosion in a karst area: the Burren, Western Ireland.- *Journal of Hydrology*, 61, 113–124.
- Fernando, A., Fernandes, J. P. A. & J. F. S. Oliveira, 2001: Comparative evaluation of European methods for sampling and sample preparation of soils – the Portuguese contribution.- *The Science of the Total Environment*, 264, 181–186.
- Gillieson, D. S., Cochrane, J. A. & A. Murray, 1994: Surface hydrology and soil movement in an arid karst: the Nullarbor Plain, Australia.- *Environmental Geology*, 23, 125–133.

- He, Z. P., Jiang, Z. C. & W. L. Lu, 2001: Effect of karst dynamic system on fertility of typical calcareous soils.- *Carsologica Sinica*, 20, 231–235 [in Chinese with English abstract].
- Jhorar, R. K., Dam, J. C. V. & W. G. M. Bastiaanssen, 2004: Calibration of effective soil hydraulic parameters of heterogeneous soil profiles.- *Journal of Hydrology*, 285, 233–247.
- Kulmatiski, A. & K. H. Beard, 2004: Reducing sampler error in soil research.- *Soil Biology & Biochemistry*, 36, 383–385.
- Lan, K. M., 1987: Karst forest preliminary investigation in Maolan.- In: Zhou, Z. X. (ed.) *Scientific investigation report of Maolan karst forest*. Guizhou Science & Technology Press, pp. 148–161, Guiyang [in Chinese].
- Leo, M. W. M., 1963: Heterogeneity of Soil of Agricultural Land in Relation to Soil Sampling.- *Agricultural Land Food Chemistry*, 11, 5, 432–434.
- Li, Y. B., Xie, D. T. & S. J. Wang, 2006: Impact of land cover types on the soil characteristics in karst area of Chongqing.- *Journal of Geographical Sciences*, 16, 143–154.
- Liu, F., Wang, S. J. & Y. S. Liu, 2005: The changes of soil quality in the process of karst rocky desertification and its evaluation of ecological environmental impact.- *Acta Ecologica Sinica*, 25, 639–644 [in Chinese, with English abstract].
- Myeong, H. Y., Seung, J. J. & S. Katsuyuki, 2003: Spatial variability of soil respiration in a larch plantation: estimation of the number of sampling points required.- *Forest Ecology and Management*, 175, 585–588.
- Nelson, D. W. & L. E. Sommers, 1982: Total carbon, organic carbon and organic matter.- In: Page, A.L., Miller, R.H. & D.R. Keeney (eds.) *Methods of Soil Analysis* (2nd edition). American Society of Agronomy, pp. 539–577, Madison, Wisconsin.
- Parise, M. & V. Pascali, 2003: Surface and subsurface environmental degradation in the karst of Apulia (southern Italy).- *Environmental Geology*, 44, 247–256.
- Park, S. J. & P. L. G. Vlek, 2002: Environmental correlation of three-dimensional soil spatial variability: a comparison of three adaptive techniques.- *Geoderma*, 109, 117–140.
- Sauro, U., Ferrarese, F., Francese, R., Miola, M., Mozzi, P., Rondo, G.Q., Trombino, L. & G. Valentini, 2009: Doline fills – case study of the Faverghe plateau (Venetian Pre-Alps, Italy).- *Acta carsologica*, 38, 1, 51–63.
- Shang, C. & H. Tiessen, 2003: Soil organic C sequestration and stabilization in karstic soils of Yucatan.- *Biogeochemistry*, 62, 177–196.
- Tamminen, P., 2003: Sampling and laboratory errors in forest soil analysis.- *Communications in Soil Science and Plant Analysis*, 34, 1193–1209.
- Theocharopoulos, S. P., Wagner, U. G. & J. Sprengart, 2001: European soil sampling guidelines for soil pollution studies.- *The Science of the Total Environment*, 264, 51–62.
- Wang, Q. B., Li, L. & H. Liu, 1998: Spatial heterogeneity of soil organic carbon and total nitrogen in a Xilin river basin grass land, Inner Mongolia.- *Acta Phytocologica Sinica*, 22, 409–414 [in Chinese with English abstract].
- Wang, S. J., Liu, Q. M. & D. F. Zhang, 2004a: Karst rocky desertification in southwestern China: Geomorphology, landuse, impact and rehabilitation.- *Land Degradation & Development*, 15, 115–121.
- Wang, S. J., Li, R. & L. Sun, 2004b: How types of carbonate rock assemblages constrain the distribution of karst rocky desertified land in Guizhou Province, P. R. China: Phenomena and mechanisms.- *Land Degradation & Development*, 15, 123–131.
- Whalen, J. K., 2004: Spatial and temporal distribution of earthworm patches in cornfield, hayfield and forest systems of southwestern Quebec, Canada.- *Applied Soil Ecology*, 27, 143–151.
- Yang, Y. L., Tiang, C. Y. & J. D. Sheng, 2002: Spatial variability of soil organic matter, total nitrogen phosphorus and potassium in cotton field.- *Agricultural Research in the Arid Areas*, 20, 3, 26–30 [in Chinese with English abstract].
- Xiong, Y. & Q. K. Li, 1987: *Chinese Soil* (2nd edition).- China Science Press, pp. 39–66. Beijing, [in Chinese].
- Zhang, P. J., Li, L. Q. & G. X. Pan, 2006: Soil quality changes in land degradation as indicated by soil chemical, biochemical and microbiological properties in a karst area of southwest Guizhou, China.- *Environmental Geology*, 51, 609–619.