


МИНИСТЕРСТВО НАУКИ И ВЫСШЕГО ОБРАЗОВАНИЯ
РОССИЙСКОЙ ФЕДЕРАЦИИ
НАЦИОНАЛЬНЫЙ ИССЛЕДОВАТЕЛЬСКИЙ
ТОМСКИЙ ГОСУДАРСТВЕННЫЙ УНИВЕРСИТЕТ
ПРАВИТЕЛЬСТВО РОССИЙСКОЙ ФЕДЕРАЦИИ
РОССИЙСКИЙ ФОНД ФУНДАМЕНТАЛЬНЫХ ИССЛЕДОВАНИЙ



Петрология магматических и метаморфических комплексов

Выпуск 10

Материалы X Всероссийской конференции
с международным участием

27 ноября – 30 ноября 2018 года



Томск 2018

THE LARGE IGNEOUS PROVINCE (LIP) RECORD OF RUSSIA THROUGH TIME: PRELIMINARY SUMMARY

ВРЕМЕННЫЕ РУБЕЖИ КРУПНЫХ ИЗВЕРЖЕННЫХ ПРОВИНЦИЙ РОССИИ: ПРЕДВАРИТЕЛЬНЫЙ ОБЗОР

R. E. Ernst^{1,2}, K.L. Buchan³, D.P. Gladkochub⁴, V.N. Puchkov^{5,6}, S.B. Botsyun⁷, I.F. Gertner²

¹ Department of Earth Sciences, Carleton University, Ottawa, Ontario, Canada

² Faculty of Geology and Geography, Tomsk State University, Russia

³ Geological Survey of Canada, Natural Resources Canada, Ottawa, Ontario, Canada

⁴ Institute of the Earth's Crust, Siberian Branch of the Russian Academy of Sciences, Irkutsk, Russia

⁵ Institute of Geology and Geochemistry, Uralian Branch of Russian Academy of Science, Ekaterinburg

⁶ Institute of Geology, Ufimian Federal Research Centre, Russian Academy of Science, Ufa, Russia

⁷ Department of Geosciences, University of Tübingen, Tübingen, Germany

This is an initial step toward a full-scale review paper and map of Large Igneous Provinces (LIPs) for Russia.

Данная работа представляет собой первый шаг в написании обширной обзорной статьи и создании карты крупных изверженных провинций России.

Introduction to Large Igneous Provinces (LIPs)

The precise definition of a LIP has varied with time, but generally emphasizes a huge volume of intraplate mafic magma emplaced in a short period (e.g. review in Ernst 2014). The following version is modified after Ernst (2014) and Ernst and Youbi (2017). LIPs represent large volume (>0.1 Mkm³; frequently above 1 Mkm³), mainly mafic (-ultramafic) magmatic events of intraplate affinity (based on tectonic setting and/or geochemistry) that occur in both continental and oceanic settings, and are typically either of short duration (<5 Ma) or consist of multiple short pulses over a maximum of a few 10s of Ma. LIPs consist of volcanic packages (flood basalts) and a plumbing system of regional dyke swarms (linear, radiating and a newly identified circumferential type), sill complexes, layered mafic-ultramafic (M-UM) intrusions, and crustal magmatic underplates. Continental LIPs can also be associated with major silicic magmatic events termed Silicic LIPs (SLIPs), as well as carbonatites and kimberlites. LIP events occur at variable rates through time. Since 2500 Ma, there has been an average of one LIP event every 20–30 Ma. The rate of LIP occurrences in the Archean, however, is less certain due to poorer preservation.

LIPs have been the focus of a significant amount of recent research, given their growing importance in constraining paleo-continental reconstructions (Ernst et al. 2013), as a tool in exploration targeting (e.g. Ernst and Jowitt 2013), as analogues for voluminous planetary intraplate magmatism (e.g. Head and Coffin 1997; Ernst 2014), and for their causal role in dramatic climate change throughout Earth history (Ernst and Youbi 2017; Bond and Grasby 2017).

Here we provide a brief summary of LIP events (and possible LIP events) of Russia. A number of more speculative LIP events have not been included. Note that in this compilation we only consider events that are dominantly mafic (cf. definition of LIPs above) and do not discuss the related Silicic LIP (SLIP) events that are also important in Russia (e.g. Kravhinsky 2012; Yarmolyuk et al. 2014). Pre-Mesozoic oceanic plateaus are likely present as fragments in orogenic belts (e.g. in central Asia, Safonova 2009), but have not been listed below. “Dykes” and “sills” mentioned below are mafic and “layered intrusions” are mafic-ultramafic unless otherwise noted.

Preliminary Survey of the Russian LIP Record

Cretaceous LIP [NE Russia]: >700 km long NW-trending Cretaceous dyke swarm (extracted from 1:1 million scale maps; Ernst et al. 2016c).

Jurassic LIP [NE Russia]: Possible >500 km long NW trending Jurassic dyke swarm (extracted from 1: 1 million scale maps; Ernst et al. 2016c).

120 Ma “Hawaii” LIP [NE Russia]: Tectonically fragmented magmatic event that is speculated to be associated with the start of the Hawaiian hot spot track (Portnyagin et al. 2008; Batanova et al. 2014).

125-90 Ma High Arctic LIP (HALIP) [northern Russia]: Part of a widespread LIP in the northern Arctic (Buchan and Ernst 2018a, Oakey and Saltus 2016; Petrov et al. 2016).

150 Ma Okhotsk oceanic plateau [far eastern Russia]: Underlies the Okhotsk sea (Bogdanov and Dobretsov 2002). Similar in age to the Shatsky oceanic plateau in the northern Pacific Ocean.

250 Ma Siberian Traps LIP [Siberian craton and West Siberia basin]: Covering more than 4 Mkm², and consists of extensive flood basalts, pyroclastic volcanism, sill complexes, giant radiating and circumferential dyke swarms, and magmatism associated with rifting in the west Siberian basin. Precise U-Pb dating suggests emplacement in less than 1 Ma (Burgess and Bowring 2015). There are also associated Ni-Cu-PGE ore deposits and associated hydrothermal ores (Ryabov et al. 2013; Pirajno et al. 2009), associated kimberlites (ca. 245-220 Ma (Sun et al. 2014) and carbonatites (e.g. Kiselev et al. 2012). Recognized as the cause of the end-Permian global mass extinction (Burgess and Bowring 2015).

370 Ma Yakutsk-Vilvui LIP [eastern Siberian craton]: Giant radiating swarm and rift system (Kiselev et al. 2012). U-Pb geochronology indicates emplacement in two pulses, at ca. 374 Ma, and ca. 363.4 Ma, that have been linked with mass extinctions. (Ricci et al. 2013; Polyansky et al. 2017, 2018). Associated with carbonatites and the diamondiferous kimberlites. Plume centre on the eastern margin of the craton defined by the radiating rift and dyke swarm systems (Kiselev et al. 2012).

370 Ma Kola-Dnieper LIP [Baltic/East European craton]: Includes coeval mafic magmatism of the Dnieper-Donets rift, a dyke swarm extending for 2000 km along the Urals, and magmatism in the Kola Peninsula and elsewhere in the Baltic craton

(Nikishin et al. 1996; Kravchinsky 2012; Puchkov et al. 2016). Two plume centres are recognized (Puchkov et al. 2016). Kimberlites (Archangelsk) and carbonatites (Kola Alkaline province) are also associated with this event. Correlated with the Yakutsk–Vilui event. Potential cause of a global mass extinction event.

440 Ma Suordakh LIP [Verkhovansk belt in eastern Siberia]: Sills and volcanics approximately dated as ca. 440 Ma (Khudoley et al. 2013). More precise dating will test whether it is the cause of the end-Ordovician mass extinction (Chamberlain et al. 2018 [this volume]).

440–450 Ushat complex [Urals]. Subalkaline volcanics within the older Igonino, Mashak and Aiskii volcanics of the western zones of the Urals. The early stages of development of the Vishnevogorsk/Ilmeny carbonatite complex (Puchkov, 2018 a,b). May be correlated with the Suordakh complex.

475–460 Ma Kidrvasovo complex [Urals]. Graben facies of the nascent passive continental margin of volcanic type along the whole Urals (coarse siliciclastic sediments and subalkaline volcanics (Puchkov 2018 a,b).

530–510 Ma Kharaulakh event [NE Siberian craton]: Widespread sills at the mouth of the Lena river and associated volcanics in the Kharaulakh mountains (Khudoley et al. 2013).

564–485 Ma Mankhambo event [Urals] Mankhambo A-granites, gabbro and contrast basalt–rhyolite volcanics in the western zones of the Urals (Puchkov, 2018b).

720 Ma Irkutsk LIP [southern Siberian craton]: Dykes, layered intrusions, and sills (Polyakov et al. 2013; Ernst et al. 2016a; Ariskin et al. 2018). Proposed to be the continuation of the Franklin LIP in a reconstruction of southern Siberia adjacent to northern Laurentia (Ernst et al. 2016a). May be correlated with the Igonino complex of the Urals (Puchkov 2018a).

706 and 735 Ma (two pulses) Igonino complex (Arshinskii - Barangulov- Serebrnaka event) [Urals]: (Krasnobaev et al. 2007, 2012; Puchkov 2012, 2018b; Maslov et al., 2018). Uncertain size and setting.

1005 Ma Sette-Daban [Verkhovansk belt in eastern Siberia]: Sill province (Rainbird et al. 1998; Khudoley et al. 2007; Gladkochub et al. 2010). Possibly linked with a carbonatite complex of similar age (Savelyeva et al. 2016) in Baikal region.

1260 Ma Srednecheremshanskii event [southern Siberia]: Wide dyke dated at 1260 Ma (U-Pb) and linked to the huge 1270 Ma Mackenzie LIP of northern Laurentia based on a continental reconstruction (Ernst et al. 2016a).

1350–1340 Ma event [southern Siberian craton]: A new event identified by U-Pb dating that may be linked to coeval magmatism in the formerly connected northern Laurentia (Ernst et al. 2016a; Gladkochub et al. 2018b).

1380 Ma Mashak LIP [eastern Baltica]: Widespread LIP recognized in the Ural mountains and in the subsurface further west (Ernst et al. 2008; Puchkov et al. 2013; Puchkov 2018a,b).

1380 Ma Chieress LIP [northern Siberia]: Dyke swarm in eastern Anabar shield (Ernst et al. 2000; Ernst et al. 2008). Likely present as sill in Taimyr (Priyatknia et al. 2017).

1470 (-1500 Ma) Tuna-Trond Gota - Ladoga event [northern Baltica]: Dykes and sills (Ernst et al. 2008; Lubnina et al. 2010).

1501 Ma Kuonamka LIP [northern Siberia]: Dykes and sills extending for 700 km across northern Siberia (Wingate et al. 2009; Ernst et al. 2000, 2016b) and proposed to have been reconstructed adjacent to the Sao Francisco–Congo craton (Ernst et al. 2016b).

1710–1730 Ma Bilvakhan-Ulkan event [SE Siberian craton]: Sills, dykes, rift systems, and associated layered intrusions (e.g., Kun Manye mineralized intrusion) (Larin 2014; Didenko et al. 2015; Guryanov and Peshkov 2017). Also associated with Ulkan–Dzhugdzhur ore-bearing AMCG (Larin 2014). Potential

link with Pelly Bay magmatic event in northern Laurentia (Ernst et al. 2016a).

1750 Ma Tampton LIP [Siberian craton]: Dykes and sills in the Aldan shield, Baikal region and Anabar shield, forming an overall radiating pattern marking a plume centre in eastern Siberia and likely later reused by the Vilyui rift arm of the 370 Ma Yakusk–Vilyui LIP (Gladkochub et al. 2010a,b). Linked with the Kivalliq Igneous Event of Northern Laurentia in the reconstruction of Ernst et al. (2016a).

1750 Ma Aiskii event [Urals, part of eastern margin of Baltica]: Aiskii volcanic suite dated by U-Pb (Krasnobaev et al. 2013). Of unknown scale and setting.

1800–1750 Ma Prutivka-Novosol event [Voronezh region, SW Russia]: AMCG (anorthosite-mangerite-charnockite-granite) magmatism, and mafic-ultramafic intrusions including dykes. Extends into the Ukrainian shield (Bodganova et al. 2013; Shumlyansky et al. 2016).

1870–1860 Ma Kalaro-Nimnvrskv- Malozadoiskv LIP [southern Siberian craton]: Dykes in the Aldan shield and in the Irkutsk promontory defining a swarm of potential 1500 km length (Ernst et al. 2016a). Can be linked with the coeval mineralized Ti–V Chiney gabbro (Gladkochub et al. 2018a). A reconstruction link is proposed with a coeval LIP of the Slave craton of northern Laurentia (Ernst et al. 2016a).

1900 Ma Angaul dyke [Irkutsk promontory, southern Siberia]: Dyke swarm dated at ca. 1920 Ma (Gladkochub et al. 2010a). A link is suggested with magmatism in northern Laurentia in the reconstruction of Ernst et al. (2016a).

1970 Ma Pechenga-Onega LIP [northern Karelia]: Sills in the Pechenga area (associated with Cu–Ni–PGE mineralization), dyke swarms and associated Onega sill province. (Ernst and Buchan 2001; Lubnina et al. 2016, 2017; Glushanin et al. 2011).

2115–2140 Ma events [Karelian craton]: Dyke swarms (Vuollo and Huhma 2005; Stepanova et al. 2014). Linked with coeval LIP events in Superior, Hearne and Wyoming cratons (Ernst and Bleeker 2010; Davey et al. 2016).

2330 Ma Taivalkoski-Kuito LIP [Karelian craton]: Regional dyke swarm (Salminen et al. 2014; Stepanova et al. 2015).

2210 Ma Koli (Karialitic) event [Karelian craton]: Widespread sills (Vuollo and Huhma 2005, Ernst 2014). Linked with the coeval Ungava-Nipissing LIP of Superior craton (Ernst and Bleeker 2010; Davey et al. 2018).

2400 Ma event [northern Karelia]: Dykes (Stepanova et al. 2017).

2500–2450 Ma Baltic (BLIP) [Karelia craton]: Layered intrusions (some mineralized), dyke swarms, and rifting (Smolkin 1996; Ernst and Buchan 2001; Lauri et al. 2012; Kulikov et al. 2010). In the formerly adjacent Superior craton there are distinct 2510 Ma Mistassini and 2480–2450 Ma Matachewan LIPs. On the basis of a reconstruction with Karelia the 2500–2450 Ma of Baltica should be parsed into two events (ca. 2510 and 2480–2450 Ma) (Kulikov et al. 2010).

Archean LIPs

The Russian LIP record continues into the Archean (e.g. Ernst and Buchan 2001), but is not well constrained and is not included in this overview.

Grant support

This study is partly supported by:

RNF grant 18-17-00240 (assisting in characterizing LIPs of eastern margin of the Siberian craton), RNF grant 16-17-10192 (assisting in characterizing LIPs of the Urals),

RNF grant 18-17-00101 (assisting in characterizing LIPs of southern part of the Siberian craton), and Mega-Grant 14.Y26.31.0012 (assisting in characterizing LIPs elsewhere in Russia).

References

- Ariskin, A., Danyushevsky, L., Nikolaev, G., et al. (2018). The Dovyren Intrusive Complex (Southern Siberia, Russia): Insights into dynamics of an open magma chamber with implications for parental magma origin, composition, and Cu-Ni-PGE fertility. *Lithos*, v. 302-308, p. 242-262.
- Batanova, V.G., Lyaskovskaya, Z.E., Savelieva, G.N., Sobolev, A.V. (2014). Peridotites from the Kamchatsky Mys: evidence of oceanic mantle melting near a hotspot. *Russian Geology and Geophysics*, v. 55, p. 1395-1403.
- Bogdanov, N.A., Dobretsov, N.L. (2002). The Okhotsk volcanic oceanic plateau. *Russian Geology and Geophysics*, v. 43, p. 87-99.
- Bogdanova, S.V., Gintov, O.B., Kurlovich, D.M., Lubnina, N.V., Nilsson, M.K.M., Orlyuk, M.I., Pashkevich, I.K., Shumlyansky, L.V., Starostenko, V.I. (2013) Late Palaeoproterozoic mafic dyking in the Ukrainian Shield of Volgo-Sarmatia caused by rotation during the assembly of supercontinent Columbia (Nuna). *Lithos*, v. 174, p. 196-216.
- Bond, D.P.G., Grasby, S.E. (2017). On the causes of mass extinctions. *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 478, p. 3-29.
- Buchan, K.L., Ernst, R.E. (2018a). A giant circumferential dyke swarm associated with the High Arctic Large Igneous Province (HALIP). *Gondwana Research*, v. 58, p. 39-57.
- Buchan, K.L., Ernst, R.E. (2018b). Giant Circumferential Dyke Swarms: Catalogue and Characteristics, in Srivastava, R.K., Ernst, R.E., Peng, P. (eds.) *Dyke Swarms of the World – A Modern Perspective*. Springer, p. 1-44.
- Burgess, S.D., Bowring, S.A. (2015). High-precision geochronology confirms voluminous magmatism before, during and after Earth's most severe extinction. *Sci. Adv.* 1 (7), e1500470.
- Chamberlain, K.R., Khudoley, A.K., Ernst, R.E. (2018). Improved U-Pb dating of the ca. 450 Ma Suordakh mafic event in eastern Siberia will test whether this is the missing LIP related to end-Ordovician mass extinction. (this volume).
- Davey, S.C., Bleeker, W., Kamo, S.L., Vuollo, J., Ernst, R.E., Cousens, B. (2016). Testing the Superia supercraton reconstruction: U-Pb age and geochemical comparison of Nipissing (Canada) and Karjalitic (Finland) sills and related dykes. GSA Annual Meeting, 25-28 September 2016, Denver, USA, <http://community.geosociety.org/gsa2016/home>.
- Davey, S.C., Bleeker, W., Kamo, S., Ernst, R., Cousens, B. (2018). Trace element geochemistry and Sm-Nd isotopes of 2.1 Ga mafic magmatism in the Karelia-Kola, Wyoming and Superior cratons. 33rd Nordic Geological Winter Meeting, Technical University of Denmark (DTU), Kgs. Lyngby, Copenhagen, Denmark – January 10-12, 2018, <https://2dggf.dk/foreningen/33rd-nordic-geological-winter-meeting/> p. 49 in Abstract volume.
- Didenko, A. N., Vodovozov, V. Y., Peskov, A. Y., Guryanov, V. A., Kosynkin, A. V. (2015). Paleomagnetism of the Ulkan massif (SE Siberian platform) and the apparent polar wander path for Siberia in late Paleoproterozoic-early Mesoproterozoic times. *Precambrian Research*, v. 259, p. 58-77.
- Ernst, R.E. (2014). *Large Igneous Provinces*. Cambridge University Press. 653 p.
- Ernst, R.E., Bleeker, W. (2010). Large igneous provinces (LIPs), giant dyke swarms, and mantle plumes: significance for breakup events within Canada and adjacent regions from 2.5 Ga. *Canadian Journal of Earth Sciences*, v. 47, p. 695-739.
- Ernst, R.E., Buchan, K.L. (2001). Large mafic magmatic events through time and links to mantle plume heads. In: R.E. Ernst, Buchan, K.L. (eds.) *Mantle Plumes: Their Identification Through Time*. Geological Society of America Special Paper 352, pp. 483-575.
- Ernst R.E., Jowitt S.M. (2013). Large igneous provinces (LIPs) and metallogeny. In: Colpron M. et al., eds. *Tectonics, metallogeny, and discovery: The North American Cordillera and similar accretionary settings*. *Society of Economic Geologists Special Publication*, v. 17, p. 17-51.
- Ernst, R.E., Jowitt, S.M. (2017). Multi-commodity, multi-scale exploration targeting using the Large Igneous Province record. In: Wyche, S. and Witt, W.K. (eds.) *TARGET 2017, Perth, Australia: Abstracts*. *Geological Survey of Western Australia, Record 2017/6*, p. 41-44.
- Ernst, R.E., Youbi, N. (2017). How Large Igneous Provinces affect global climate, sometimes cause mass extinctions, and represent natural markers in the geological record: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 478, p. 30-52.
- Ernst, R. E., Buchan, K. L., Hamilton, M. A., Okrugin, A. V., Tomshin, M. D. (2000) Integrated paleomagnetism and UPb geochronology of mafic dikes of the eastern Anabar Shield region, Siberia: implications for Mesoproterozoic paleolatitude of Siberia and comparison with Laurentia. *Journal of Geology*, v. 108, p. 381-401.
- Ernst, R.E., Wingate, M.T.D., Buchan, K.L., Li, Z.X. (2008). Global record of 1600–700 Ma Large Igneous Provinces (LIPs): implications for the reconstruction of the proposed Nuna (Columbia) and Rodinia supercontinents. *Precambrian Research*, 160: 159–178.
- Ernst, R.E., Bleeker, W., Söderlund, U., Kerr, A.C. (2013). Large Igneous Provinces and supercontinents: Toward completing the plate tectonic revolution. *Lithos*, v. 174, p. 1–14.
- Ernst, R.E., Hamilton, M.A., Söderlund, U., J.A. Hanes, J.A., Gladkochub, D.P., Okrugin, A.V., Kolotilina, T., Mekhonoshin, A.S., Bleeker, W., LeCheminant, A.N., Buchan, K.L., Chamberlain, K.R., Didenko, A.N. (2016a) Long-lived connection between southern Siberia and northern Laurentia in the Proterozoic. *Nature Geoscience*, v. 9, p. 464-469.
- Ernst, R.E., A.V. Okrugin, R.V. Veselovskiy, S.L. Kamo, M.A. Hamilton, V. Pavlov, U. Söderlund, K.R. Chamberlain, C. Rogers (2016b). The 1501 Ma Kuonamka Large Igneous Province of northern Siberia: U-Pb geochronology, geochemistry, and links with coeval magmatism on other crustal blocks. *Russian Geology and Geophysics*, v. 57, p. 653-671.
- Ernst, R.E., Buchan, K.L., Botsyun, S. (2016c). Map of Mafic Dyke Swarms and Related Units of Russia and Adjacent Regions. *Acta Geologica Sinica (English Edition)*, v. 90(supp. 1): p. 22-23.
- Gladkochub, D.P., Pisarevsky, S.A., Donskaya, T.V., et al. (2010a). Proterozoic mafic magmatism in Siberian craton: an overview and implications for paleocontinental reconstruction. *Precambrian Research*, v. 183, p. 660–668.
- Gladkochub, D.P., Pisarevsky, S.A., Ernst, R., Donskaya, T.V., Soderlund, U., Mazukabzov, A.M., Hanes, J. (2010b). Large Igneous Province of about 1750 Ma in the Siberian Craton. *Doklady Earth Sciences*, v. 430, p. 168-171.
- Gladkochub, D.P., Donskaya, T.V., Sklyarov, E.V. et al. (2017). The unique Katugin rare-metal deposit (southern Siberia): Constraints on age and genesis. *Ore Geology Reviews*, v. 91, p. 246-263.
- Gladkochub, D.P., Donskaya, T.V., Sklyarov, E.V., Kotov, A.B., Ernst, R.E. (2018a). Main stages of ore-formation processes within the Kalar-Udokan ore region (Aldan shield, Siberian craton) and testing links with Large Igneous

- Provinces In Special Session MIN11 at RFG 2018 conference, 17-22 June 2018 Vancouver, Canada.
29. Gladkochub, D.P., Donskaya, T.V., Ernst, R.E., Hamilton, M.A., Mazukabzov, A.M., Pisarevsky, S.A., Kamo, S. (2018b). New Ectasian event of basic magmatism in the Southern Siberian craton. *Dokl. Earth Sciences*, (in press)
 30. Glushanin, L.V., Sharov, N.V., Shchiptsov, V.V. (editors) (2011). PALAEOPROTEROZOIC ONEGA STRUCTURE (GEOLOGY, TECTONICS, DEEP STRUCTURE AND MINERALOGENY) [in Russian]. Karelian Science Centre, Russian Academy of Sciences, Petrozavodsk, Russia, 431 p.
 31. Guryanov, V.A. Peskov, A. Yu. (2017) Ulkan Paleorift Structure in the South-Eastern Environs of the Siberian Platform: Age, Conditions, Sources, and Geodynamic Setting. *Geosciences Research* v. 2, p. 59-71.
 32. Head, J.W. and Coffin, M.F. (1997). Large Igneous Provinces: a planetary perspective. In: Mahoney, J.J., and Coffin, M.F. (eds.) Large Igneous Provinces: Continental, Oceanic and Planetary Flood Volcanism. AGU Geophysical Monograph, v. 100, p. 411-438.
 33. Kiselev, A.I., Ernst, R.E., Yarmolyuk, V.V., Egorov, K.N. (2012). Radiating rifts and dyke swarms of the middle Paleozoic Yakutsk plume, of eastern Siberian craton: *Journal of Asian Earth Sciences*, v. 45, p. 1-16.
 34. Krasnobaev, A.A., Kozlov, V.I., Puchkov, V.N., Larionov, A.N., Nekhorosheva, A.G., Berezhnaya, N.G. (2007). The Polygenous-Polychronous Nature of Zircons and the Problem of the Age of the Barangulov Gabbro-Granite Complex. *Doklady Earth Sciences*, v. 416, p. 1070-1075.
 35. Krasnobaev, A.A., Koslov, V.I., Puchkov, V.N., Sergeeva, H.D., Busharina, S.V. (2012) New data by zircon geochronology of the Arshinski volcanics (Southern Urals) [in Russian]. *Lithosphaera*, No. 4, p. 127-139.
 36. Krasnobaev, A.A., Puchkov, V.N., Koslov, V.I., Sergeeva, N.D., Busharina, S.V., Lepekina, Ye. N. (2013). Цирконология навьшских вулканиитов айской свиты и проблема возраста нижней границы рифея на Южном Урале. [in Russian] *Dokladi Akamii Nauk*, v. 448, p. 437-442.
 37. Kravchinsky, V.A. (2012). Paleozoic large igneous provinces of Northern Eurasia: Correlation with mass extinction events. *Global and Planetary Change*, v. 86-87, p. 31-36.
 38. Khudoley, A.K., Kropachev, A.P., Tkachenko, V.I., Rublev, A.G., Sergeev, S.A., Matukov, D.I., Lyahnikskaya, O.Yu., 2007. Mesoproterozoic to Neoproterozoic evolution of the Siberian craton and adjacent microcontinents: an overview with constraints for a Laurentian connection. Spec. Publ.—SEPM 86, 209-226, Society for Sedimentary Geology Special Publication.
 39. Khudoley, A.K., Prokopiev, A.V., Chamberlain, K.R., Ernst, R.E., Jowitt, S.M., Malyshev, S.V., Zaitsev, A.I., Kropachev, A.P., Koroleva, O.V. (2013). Early Paleozoic mafic magmatic events on the eastern margin of the Siberian craton. *Lithos*, v. 174, p. 44-56.
 40. Kulikov, V.S., Bychkova, Y.V., Kulikova, V.V., Ernst, R.E. (2010). The Vetryny Poyas (Windy Belt) subprovince of southeastern Fennoscandia: an essential component of the ca. 2.5-2.4 Ga Sumian large igneous provinces. *Precambrian Research*, v. 183, p. 589-601.
 41. Larin, A.M. (2014). Ulkan-Dzhugdzhur Ore-Bearing Anorthosite-Rapakivi Granite-Peralkaline Granite Association, Siberian Craton: Age, Tectonic Setting, Sources, and Metallogeny. *Geology of Ore Deposits*, v. 56, p. 257-280.
 42. Lauri, L.S., Mikkola, P., Karinen, T. (2012). Early Paleoproterozoic felsic and mafic magmatism in the Karelian province of the Fennoscandian shield. *Lithos*, v. 151, p. 74-82.
 43. Lubnina, N.V., Mertanen, S., Söderlund, U., Bogdanova, S. Vasilieva, T.I., Frank-Kamenetsky, D. (2010). A new key pole for the East European Craton at 1452 Ma: Palaeomagnetic and geochronological constraints from mafic rocks in the Lake Ladoga region (Russian Karelia). *Precambrian Research*, v. 183, p. 442-462.
 44. Lubnina, N.V., Stepanova, A.V., Ernst, R.E., Nilsson, M., Söderlund, U. (2016). New U-Pb baddeleyite age, and AMS and paleomagnetic data for dolerites in the Lake Onega region belonging to the 1.98-1.95 Ga regional Pechenga-Onega Large Igneous Province. *GFF*, v. 138, p. 54-78.
 45. Lubnina N.V., Stepanova, A.V., Bogdanova, S.V. Sokolov, S.J. (2017). Fennoscandia before Nuna/Columbia: Paleomagnetism of 1.98-1.96 Ga mafic rocks of the Karelian craton and paleogeographic implications. *Precambrian Research*, v. 292, p. 1-12.
 46. Maslov A. V., Kovalev S. G., Puchkov V. N., Sergeeva N. D. (2018). The Riphean Arsha Group of the South Urals: A Problem of the Geodynamic Origin of Rock Associations. *Doklady Earth Sciences*, v. 480, part 1, p. 551-554.
 47. Nikishin, A.M., Ziegler, P.A., Stephenson, R.A. et al. (1996). Late Precambrian to Triassic history of the East European Craton: dynamics of sedimentary basin evolution. *Tectonophysics*, v. 268, p 23-63.
 48. Oakey, G.N., Saltus, R.W. (2016). Geophysical analysis of the Alpha-Mendeleev ridge complex: Characterization of the High Arctic Large Igneous Province. *Tectonophysics*, v. 691, p. 65-84.
 49. Peng, P., 2015. Precambrian mafic dyke swarms in the North China Craton and their geological implications. *Science China: Earth Sciences*, v. 58, p. 649-675.
 50. Petrov, O., Morozov, A., Shokalsky, S., Kashubin, S., Artemieva, I.M., Sobolev, N., Petrov, E., Ernst, R.E., Sergeev, S., Smelror, M. (2016) Crustal structure and tectonic model of the Arctic region. *Earth-Science Reviews*, v. 154, p. 29-71.
 51. Pirajno, F., Ernst, R.E., Borisenko, A.S., Fedoseev, G., Naumov, E.A. (2009). Intraplate magmatism in central Asia and China and associated metallogeny: *Ore Geology Reviews*, v. 35, p. 114-136.
 52. Portnyagin, M., Savelyev, D., Hoernle, K., Hauff, F., Gerbe-Schönberg, D. (2008) Mid-Cretaceous Hawaiian tholeiites preserved in Kamchatka. *Geology*, v. 36, p. 903-906.
 53. Polyakov, G.V., Tolstykh, N.D., Mekhonoshin, A.S., et al. (2013). Ultramafic-mafic igneous complexes of the Precambrian East Siberian metallogenic province (southern framing of the Siberian craton): age, composition, origin, and ore potential. *Russian Geology and Geophysics*, v. 54, p. 1319-1331.
 54. Polyansky, O.P., Prokopiev, A.V., Koroleva, O.V., Tomsin, M.D., Reverdatto, V.V., Selyatitsky, A.Yu., Travin, A.V., Vasiliev, D.A. (2017). Temporal correlation between dyke swarms and crustal extension in the middle Proterozoic Vilyui rift basin, Siberian platform. *Lithos*, v. 282-283, p. 45-64.
 55. Priyatkina, N., Collins, W.J., Khudoley, A. et al. (2017). The Proterozoic evolution of northern Siberian Craton margin: a comparison of U-Pb-Hf signatures from sedimentary units of the Taimyr orogenic belt and the Siberian platform. *International Geology Review*, v. 59, p. 1632-1656.
 56. Puchkov, V.N. (2012). Dyke swarms and associated magmatic complexes. *Geotectonics*, v. 1, p. 42-53.
 57. Puchkov, V.N., Bogdanova, S.V., Ernst, R.E., et al. (2013). The ca. 1380 Ma Mashak igneous event of the Southern Urals. *Lithos*, 174: 109-124.

58. Puchkov, V., Ernst, R.E., Hamilton, M.A., Söderlund, U., Sergeeva, N. (2016). A Devonian >2000-km long dolerite swarm-belt and associated basalts along the Urals-Novozemelian fold-belt: part of an East-European (Baltica) LIP tracing the Tuzo Superswell. *GFF*, v. 138, p. 6-16.
59. Puchkov, V., The Plumes – a new word in Geology of the Urals. *Lithosphere (Russia)*, 2018, v. 18, no. 4, p. 483–499 (in Russian)
60. Puchkov, V., The plume-dependent granite-rhyolite magmatism. *Lithosphere (Russia)*, 2018, v. 18, no. 5, p. 692–705 (in Russian)
61. Rainbird, R.H., Stern, R.A., Khudoley, A.K., et al. (1998). U–Pb geochronology of Riphean sandstone and gabbro from southeast Siberia and its bearing on the Laurentia-Siberia connection. *Earth and Planetary Science Letters*, v. 164, p. 409–420.
62. Ricci, J., Quidelleur, X., Pavlov, V., Orlov, S., Shatsillo, A., Courtillot, V. (2013). New $^{40}\text{Ar}/^{39}\text{Ar}$ and K–Ar ages of the Viluy traps (Eastern Siberia): Further evidence for a relationship with the Frasnian–Famennian mass extinction. *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 386, p. 531-540.
63. Ryabov, V.V., Shevko, A.Ya., Gora, M.P. (2013). Trap Magmatism and Ore Formation in the Siberian Noril'sk Region: Volume 1, Trap Petrology, Springer, 390 p.
64. Safonova, I. Yu. (2009). Intraplate magmatism and oceanic plate stratigraphy of the Paleo-Asian and Paleo-Pacific oceans from 600 to 140 Ma. *Ore Geology Reviews*, v. 35, p. 137–154.
65. Salminen, J., Halls, H.C., Mertanen, S., Pesonen, L.J., Vuollo, J., Söderlund, U. (2014). Paleomagnetic and geochronological studies on Paleoproterozoic diabase dykes of Karelia, East Finland—key for testing the Superia supercraton. *Precambrian Research*, v. 244, p. 87–99.
66. Samsonov, A.V., Stepanova, A.V., Salnikova, E.B., Larionova, Yu.O., Egorova, S.V., Larionov, A.N. (2016). Mafic Dyke Records of Paleoproterozoic Mantle Plume Activity in the Karelian Craton: U–Pb Baddeleyite/Zircon Geochronology and Sr–Nd Isotopic Data. *Acta Geologica Sinica (English Edition)*, 90(supp. 1): 118-119.
67. Savelyeva, V.B., Demonterova, E.I., Danilova, Yu.V., Bazarova, E.P. Ivanov, A.V., Kamenetsky, V.S. (2016). New carbonatite complex in the western Baikal area, southern Siberian craton: Mineralogy, age, geochemistry and petrogenesis. *Petrology*, v. 24, p. 271-302.
68. Smolkin, V.F. (1997). The Paleoproterozoic (2.5–1.7 Ga) Midcontinent rift system of the northeastern Fennoscandian Shield. *Canadian Journal of Earth Sciences*, 34: 426–443.
69. Stepanova, A.V., Samsonov, A.V., Salnikova, E.B., et al. (2014). Palaeoproterozoic Continental MORB-type Tholeiites in the Karelian Craton: Petrology, Geochronology, and Tectonic Setting. *Journal of Petrology*, v. 55, p. 1719-1751.
70. Stepanova, A.V., Salnikova, E.B., Samsonov, A.V., Egorova, S.V., Larionova, Y.O., Stepanov, V.S. (2015). The 2.31 Ga mafic dykes in the Karelian Craton, eastern Fennoscandian shield: U–Pb age, source characteristics and implications for continental break-up processes. *Precambrian Research*, v. 259, p. 43-57.
71. Stepanova A.V., Salnikova E.B., Samsonov A.V., Larionova Yu.O., Egorova S.V., Stepanov V.S. (2016). The 2405 Ma and 2310 Ma Mafic Dyke Swarms in the Karelian Craton: Age, Chemical and Sr–Nd Isotope Composition, and Tectonic Setting. *Acta Geologica Sinica (English Edition)*, 90(supp. 1): 123.
72. Stepanova, A.V., Salnikova, E.B., Samsonov, A.V., Larionova, Yu.O., Egorova, S.V., Savantekov, V.M. (2017). The 2405 Ma Doleritic Dykes in the Karelian Craton: A Fragment of a Paleoproterozoic Large Igneous Province. *Doklady Earth Sciences*, v. 472, p. 72-77.
73. Shumlyanskyy, L., Ernst, R.E., Billstrom, K., Wing, B.A., Bekker, A. (2016) Age and sulfur isotope composition of the Prutivka intrusion (the 1.78 Ga Prutivka-Novogol Large Igneous Province in Sarmatia). *Mineralogical Journal: Geochemistry (Ukraine)*, v. 38, no. 3, p. 91-101.
74. Sun, J., Liu, C-Z., Tappe, S., Kostrovitsky, S.I., Wu, F-Y., Yakovlev, D., Yang, Y-H., Yang, J-H., (2014). Repeated kimberlite magmatism beneath Yakutia and its relationship to Siberian flood volcanism: insights from in situ U–Pb and Sr–Nd perovskite isotope analysis. *Earth Planet. Sci. Lett.*, v. 404, p. 283–295.
75. Vuollo, J., Huhma, H. (2005). Paleoproterozoic mafic dikes in NE Finland. In: *Precambrian geology of Finland – Key to the evolution of the Fennoscandian Shield*. Elsevier, Amsterdam, pp. 195-236.
76. Wingate, M.T.D., Pisarevsky, S.A., Gladkochub, D.P., Donskaya, T.V., Konstantinov, K.M., Mazukabzov, A.M., Stanevich, A.M. (2009). Geochronology and paleomagnetism of mafic igneous rocks in the Olenek Uplift, northern Siberia: implications for Mesoproterozoic supercontinents and paleogeography. *Precambrian Research*, v. 170, p. 256–266.
77. Yarmolyuk, V.V., Kuzmin, M.I., Ernst, R.E. (2014). Intraplate geodynamics and magmatism in the evolution of the Central Asian Orogenic Belt: *Journal of Asian Earth Sciences*, v. 93, p. 158–179.