



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: June 4, 2013
REVISED: July 5, 2013
ACCEPTED: July 30, 2013
PUBLISHED: August 26, 2013

Weyl transformations and trace anomalies in $N=1$, $D=4$ supergravities

L. Bonora and S. Giaccari

*International School for Advanced Studies (SISSA),
Via Bonomea 265, 34136 Trieste, Italy
INFN, Sezione di Trieste,
Trieste, Italy*

E-mail: bonora@sissa.it, giaccari@sissa.it

ABSTRACT: We consider the supersymmetric extension of Weyl transformations in various types of supergravities, the minimal, $20+20$ and $16+16$ nonminimal $N=1$ SUGRA in $4D$, formulated in terms of superfields and study their local cohomology. Based also on previous results we conclude that there are only two types of trace anomalies in $20+20$ and $16+16$ nonminimal supergravities, which correspond to the two nontrivial cocycles of the minimal supergravity and, when reduced to component form, to the well-known squared Weyl density and Euler density.

KEYWORDS: Anomalies in Field and String Theories, Supergravity Models, Superspaces, Conformal and W Symmetry

Contents

1	Introduction	1
2	N=1 minimal supergravity in D=4 and its superfields	3
2.1	Superconformal symmetry and (super)anomalies	4
2.2	Meaning of superconformal transformations	5
2.3	Anomalies in components	7
2.3.1	The square Weyl cocycle	7
2.3.2	The Gauss-Bonnet cocycle	9
3	Non minimal supergravity	10
3.1	Superconformal transformations in the non minimal model	11
3.2	Cocycles in non minimal SUGRA	12
4	The 16+16 nonminimal model	12
4.1	Cocycles in 16+16 nonminimal SUGRA	13
5	Reduction to component form	14
6	Mapping formulas between different supergravity models	15
7	Cocycles from minimal supergravity	18
7.1	From minimal to nonminimal cocycles	18
7.2	From minimal to 16+16 nonminimal cocycles	19
8	Conclusions	20
A	Reduction formulae	22

1 Introduction

In this paper we want to analyze the supersymmetric extension of Weyl transformations in various types of supergravities, the minimal, 20+20 and 16+16 nonminimal N=1 SUGRA in 4D, and study the general structure of trace anomalies. To this end, rather than considering specific cases we carry out a cohomological analysis, whose validity is not limited to one-loop calculations.

The motivation for this research is twofold. On the one hand it has been pointed out recently that ‘old’ minimal supergravity in 4D (with 12+12 dofs) might be inconsistent due to the presence of an inherent global conserved current, [1]. It has also been suggested that a different type of SUGRA, referred to henceforth as *16+16 nonminimal*, characterized

by 16+16 dofs, may be exempt from this risk. This model has been identified with the supergravities studied in [2] and [3]. The study of conformal anomalies in these and other models is interesting not only in itself, but also because it allows us to identify what the ‘superWeyl group’ is, as will be seen below.

Another motivation arises from the proposal of [4] that a source of CP violation in a 4D theory coupled to gravity could come from the trace anomaly. The trace anomaly may contain, in principle, beside the Weyl density (square of the Weyl tensor)

$$\mathcal{R}_{nmkl}\mathcal{R}^{nmkl} - 2\mathcal{R}_{nm}\mathcal{R}^{nm} + \frac{1}{3}\mathcal{R}^2 \tag{1.1}$$

and the Gauss-Bonnet (or Euler) one

$$\mathcal{R}_{nmkl}\mathcal{R}^{nmkl} - 4\mathcal{R}_{nm}\mathcal{R}^{nm} + \mathcal{R}^2, \tag{1.2}$$

another nontrivial piece, the Pontryagin density

$$\epsilon^{nmkl}\mathcal{R}_{nmpq}\mathcal{R}_{lk}{}^{pq} \tag{1.3}$$

Each of these terms appears in the trace of the e.m. tensor with its own coefficient. The first two are denoted c and a , respectively. They are known at one-loop for any type of (Gaussian) matter [5, 6], and a is the protagonist of recent important developments, [7]. The coefficient of (1.3) is not sufficiently studied. The original purpose of this paper was to analyse whether the appearance of such a term in the trace anomaly is compatible with supersymmetry. Since it is hard to supersymmetrize these three pieces and relate them to one another in a supersymmetric context, the best course is to proceed in another way, that is to consider a conformal theory in 4D coupled to (external) supergravity formulated in terms of superfields and find all the potential superconformal anomalies. This will allow us to see whether (1.3) can be accommodated in an anomaly supermultiplet as a trace anomaly member.

This type of analysis was carried out long ago for minimal supergravity, see [8] and also [10]. Our purpose here is to extend it to other types of 4D supergravities,¹ in particular to the nonminimal SUGRA mentioned above, [2] and [3]. Unfortunately there is no unique choice of the torsion constraints for these theories and no unique superfield formalism, (see [11, 12] and [13, 14] for earlier ‘new minimal’ formulations and [15] for their equivalence; see [16, 17] for earlier non-minimal formulations; see moreover [19–21] and the textbooks [18, 22]). Thus we have chosen to follow the formalism of [23], further expanded in [2, 24–26]. Our original aim, the analysis of trace anomalies, has turned out to be anything but standard, contrary to the case of minimal supergravity. The reason is that in the latter case the cohomological analysis can be done on a differential space formed by polynomials of the superfields. In the other abovementioned versions of supergravities one has to admit in the differential space also nonpolynomial expressions of the superfields, due to the essential role of dimensionless prepotentials in these models. To solve in a satisfactory way

¹A general cohomological treatment of anomalies in 4D supergravities, which has some overlap with the present paper, is contained in [9].

the cohomology problem one has to start from minimal supergravity and map its cocycles to the other models with the superfield mappings of ref. [22, 24]. Once this is clarified the possible superconformal anomalies are rather easily identified. Based also on the analysis carried out long ago in [8], one can conclude that there are, not unexpectedly, two independent anomalies corresponding to the square Weyl and Gauss-Bonnet densities, much like in minimal supergravity. The anomaly corresponding to the Gauss-Bonnet density has a particularly complicated form in non-minimal supergravities, and could be identified only via the abovementioned mapping method.

The conclusion concerning the Pontryagin density (1.3) is negative: in all types of supergravities the Pontryagin density does not show up in the trace anomaly, but it appears in the chiral (Delbourgo-Salam) anomaly, which, as expected, belongs, together with the trace anomaly, to a unique supermultiplet.

The paper is organized as follows. In the next section we review the minimal supergravity case. This section does not contain new results, it has mainly pedagogical and reference purposes: the analysis is far clearer if we keep in mind the minimal case as a guide. In section 3 we introduce the (20+20) non minimal supergravity and the relevant superconformal transformations. We also easily identify a non-trivial cocycle corresponding to the square Weyl density. In section 4 we do the same for the 16+16 nonminimal supergravity. In section 5 we study the reduction to component form of the latter cocycle. In section 6 we briefly introduce the mapping method from one supergravity model to another and introduce the relevant formulas. In section 7 we move on to compute the remaining superWeyl cocycles. Section 8 is devoted to the conclusions.

2 N=1 minimal supergravity in D=4 and its superfields

For the notation we follow [23]. The superspace of $N = 1$ supergravity is spanned by the supercoordinates $Z^M = (x^m, \theta^\mu, \bar{\theta}_{\dot{\mu}})$. The minimal $N = 1$ supergravity in $D = 4$ can be formulated in terms of the superfields: $R(z), G_a(z)$ and $W_{\alpha\beta\gamma}(z)$. R and $W_{\alpha\beta\gamma}$ are chiral while G_a is real. We will also need the antichiral superfields $R^+(z)$ and $\bar{W}_{\dot{\alpha}\dot{\beta}\dot{\gamma}}(z)$, conjugate to R and $W_{\alpha\beta\gamma}$, respectively. $W_{\alpha\beta\gamma}$ is completely symmetric in the spinor indices α, β, \dots . These superfields are subject to the constraints:

$$\begin{aligned} \nabla^\alpha G_{\alpha\dot{\beta}} &= \bar{\nabla}_{\dot{\beta}} R^+, & \bar{\nabla}^{\dot{\beta}} G_{\alpha\dot{\beta}} &= \nabla_\alpha R \\ \nabla^\alpha W_{\alpha\beta\gamma} + \frac{i}{2}(\nabla_{\beta\dot{\beta}} G_{\delta}^{\dot{\beta}} + \nabla_{\beta\dot{\beta}} \dot{G}_{\dot{\beta}}^{\beta}) &= 0 \\ \bar{\nabla}^{\dot{\alpha}} W_{\dot{\alpha}\dot{\beta}\dot{\gamma}} + \frac{i}{2}(\nabla_{\beta\dot{\beta}} G^{\beta}_{\dot{\delta}} + \nabla_{\beta\dot{\beta}} \dot{G}^{\beta}_{\dot{\beta}}) &= 0 \end{aligned} \tag{2.1}$$

The latter are found by solving the (super)Bianchi identities for the supertorsion and the supercurvature

$$\begin{aligned} T^A &= dE^A + E^B \phi_B^A = \frac{1}{2} E^C E^B T_{BC}^A = \frac{1}{2} dz^M dz^N T_{NM}^A \\ R_A^B &= \frac{1}{2} E^C E^D R_{DCA}^B = dz^M dz^N \partial_N \phi_{MA}^B + dz^M \phi_{MA}^C dz^N \phi_{NC}^B \end{aligned} \tag{2.2}$$

where $\phi_{MA}{}^B$ is the superconnection and $E^A = dz^M E_M{}^A$ the supervierbein

$$E_M{}^A E_A{}^N = \delta_M{}^N, \quad E_A{}^M E_M{}^B = \delta_A{}^B,$$

after imposing by hand the restrictions

$$\begin{aligned} T_{\underline{\alpha}\underline{\beta}}{}^{\underline{\gamma}} &= 0, & T_{\alpha\beta}{}^c &= T_{\dot{\alpha}\dot{\beta}}{}^c = 0 \\ T_{\alpha\dot{\beta}}{}^c &= T_{\dot{\beta}\alpha}{}^c = 2i\sigma_{\alpha\dot{\beta}}{}^c \\ T_{\underline{a}\underline{b}}{}^c &= T_{b\underline{\alpha}}{}^c = 0, & T_{ab}{}^c &= 0 \end{aligned} \tag{2.3}$$

where $\underline{\alpha}$ denotes both α and $\dot{\alpha}$. The superdeterminant of the vierbein $E_M{}^A$ will be denoted by E .

The Bianchi identities are

$$\mathcal{D}\mathcal{D}E^A = E^B R_B{}^A, \quad \nabla T^A = E^B R_B{}^A \tag{2.4}$$

where $\mathcal{D} = dz^M \nabla_M$ and $\nabla_A = E_A{}^M \nabla_M$. Imposing (2.4) one gets all the components of T^A and $R_A{}^B$ in terms of $R, G_a, W^{\alpha\beta\gamma}$ and their conjugates. The other Bianchi identity

$$(\mathcal{D}R)_A{}^B = 0 \tag{2.5}$$

is automatically satisfied.

2.1 Superconformal symmetry and (super)anomalies

Superconformal transformations are defined by means of the chiral superfield parameter $\sigma = \sigma(z)$ and its conjugate $\bar{\sigma}$.

$$\begin{aligned} \delta E_M{}^a &= (\sigma + \bar{\sigma}) E_M{}^a & (2.6) \\ \delta E_M{}^\alpha &= (2\bar{\sigma} - \sigma) E_M{}^\alpha + \frac{i}{2} E_M{}^a \bar{\sigma}_a{}^{\dot{\alpha}\alpha} \nabla_{\dot{\alpha}} \bar{\sigma} \\ \delta E_M{}^{\dot{\alpha}} &= (2\sigma - \bar{\sigma}) E_M{}^{\dot{\alpha}} + \frac{i}{2} E_M{}^a \bar{\sigma}_a{}^{\dot{\alpha}\alpha} \nabla_{\alpha} \sigma \\ \delta \phi_{M\alpha\beta} &= E_{M\alpha} \nabla_{\beta} \sigma + E_{M\beta} \nabla_{\alpha} \sigma + (\sigma^{ab})_{\alpha\beta} E_{Ma} \nabla_b (\sigma + \bar{\sigma}) \end{aligned}$$

where

$$\phi_{M\alpha}{}^{\beta} = \frac{1}{2} \phi_{Mab} (\sigma^{ab})_{\alpha}{}^{\beta}, \quad \phi_{M\dot{\beta}}{}^{\dot{\alpha}} = \frac{1}{2} \phi_{Mab} (\bar{\sigma}^{ab})^{\dot{\alpha}}{}_{\dot{\beta}}$$

The transformations (2.6) entail

$$\begin{aligned} \delta E &= 2(\sigma + \bar{\sigma}) E & (2.7) \\ \delta R &= (2\bar{\sigma} - 4\sigma) R - \frac{1}{4} \nabla_{\dot{\alpha}} \nabla^{\dot{\alpha}} \bar{\sigma} \\ \delta R^+ &= (2\sigma - 4\bar{\sigma}) R^+ - \frac{1}{4} \nabla^{\alpha} \nabla_{\alpha} \sigma \\ \delta G_a &= -(\sigma + \bar{\sigma}) G_a + i \nabla_a (\bar{\sigma} - \sigma) \\ \delta W_{\alpha\beta\gamma} &= -3\sigma W_{\alpha\beta\gamma} \end{aligned}$$

If we promote the superfield σ to a superghost superfield, by inverting the spin-statistics connection, so that it becomes an anticommuting parameter, it is easy to prove that the above transformations are nilpotent.

Let us define the functional operator that implements these transformations, i.e.

$$\Sigma = \int_{x\theta} \delta\chi_i \frac{\delta}{\delta\chi_i}$$

where χ_i represent the various superfields in the game and $_{x\theta}$ denotes integration $d^4x d^4\theta$. This operator is nilpotent: $\Sigma^2 = 0$. As a consequence it defines a cohomology problem. The cochains are integrated local expressions of the superfields and their superderivatives, invariant under superdiffeomorphism and local superLorentz transformations. Candidates for superconformal anomalies are nontrivial cocycles of Σ which are not coboundaries, i.e. integrated local functionals Δ_σ , linear in σ , such that

$$\Sigma \Delta_\sigma = 0, \quad \text{and} \quad \Delta_\sigma \neq \Sigma \mathcal{C} \tag{2.8}$$

for any integrated local functional \mathcal{C} (not containing σ).

The complete analysis of all the possible nontrivial cocycles of the operator Σ was carried out long ago in [8]. It was shown there that the latter can be cast into the form

$$\Delta_\sigma = \int_{x\theta} \left[\frac{E(z)}{-8R(z)} \sigma(z) \mathcal{S}(z) + h.c. \right] \tag{2.9}$$

where $\mathcal{S}(z)$ is a suitable chiral superfield. In [8] all the possibilities for \mathcal{S} were classified. For pure supergravity (without matter) the only nontrivial possibilities turn out to be:

$$\mathcal{S}_1(z) = W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \quad \text{and} \quad \mathcal{S}_2(z) = (\bar{\nabla}_{\dot{\alpha}} \bar{\nabla}^{\dot{\alpha}} - 8R)(G_a G^a + 2RR^+) \tag{2.10}$$

(the operator $(\bar{\nabla}_{\dot{\alpha}} \bar{\nabla}^{\dot{\alpha}} - 8R)$ maps a real superfield into a chiral one).

It is well-known that the (2.9) cocycles contain not only the trace anomaly, but a full supermultiplet of anomalies. The local expressions of the latter are obtained by stripping off the corresponding parameters from the integrals in (2.9). Let us recall also that the conversion of σ to an anticommuting parameter is not strictly necessary: eq. (2.8) simply corresponds to the Wess-Zumino consistency conditions, i.e. to the invariance under reversing the order of two successive (Abelian) gauge transformations. But an anticommuting σ allows us to use the incomparably simpler formalism of cohomology.

2.2 Meaning of superconformal transformations

Eqs. (2.10) are rather implicit and it is opportune to see the corresponding expressions in component fields, at least as far as the dependence on the metric alone is concerned. This reduction has been done in [27]. We repeat it here for pedagogical reasons, but also because the formalism we use is different from the one of [27]. The method below will be used throughout the paper. In general the expressions of the above cocycles in components are extremely complicated and really unmanageable because of the presence of auxiliary fields. We are interested in recognizing the two cocycles (2.10) when only the metric is

taken into account while all the other fields are ignored, so that we can compare them with the usual Weyl cocycles (the squared Weyl tensor, the Gauss-Bonnet and the Pontryagin densities). Our task in the sequel is to extract such expressions from (2.10). We will refer to them as the *ordinary* parts of the cocycles.

We first introduce the relevant components fields and clarify the meaning of the components in the parameters superfield $\sigma(z)$. To start with let us define the lowest component fields of the supervierbein as in [23]

$$E_M^A(z)|_{\theta=\bar{\theta}=0} = \begin{pmatrix} e_m^a(x) & \frac{1}{2}\psi_m^\alpha(x) & \frac{1}{2}\bar{\psi}_{m\dot{\alpha}}(x) \\ 0 & \delta_\mu^\alpha & 0 \\ 0 & 0 & \delta^{\dot{\mu}}_{\dot{\alpha}} \end{pmatrix} \quad (2.11)$$

and

$$E_A^M(z)|_{\theta=\bar{\theta}=0} = \begin{pmatrix} e_a^m(x) & -\frac{1}{2}\psi_a^\mu(x) & -\frac{1}{2}\bar{\psi}_{a\dot{\mu}}(x) \\ 0 & \delta_\alpha^\mu & 0 \\ 0 & 0 & \delta^{\dot{\alpha}}_{\dot{\mu}} \end{pmatrix} \quad (2.12)$$

where e_m^a are the usual 4D vierbein and $\psi_m^\alpha(x), \bar{\psi}_{m\dot{\alpha}}(x)$ the gravitino field components. We have in addition

$$R(z)|_{\theta=\bar{\theta}=0} = -\frac{1}{6}M(x), \quad G_a(z)|_{\theta=\bar{\theta}=0} = -\frac{1}{3}b_a(x) \quad (2.13)$$

where M is a complex scalar field and b_a is a real vector field. As for the superconnection we have

$$\phi_{mA}^B|_{\theta=\bar{\theta}=0} = \omega_{mA}^B(x), \quad \phi_{\mu A}^B|_{\theta=\bar{\theta}=0} = 0, \quad \phi_{\dot{\mu} A}^B|_{\theta=\bar{\theta}=0} = 0, \quad (2.14)$$

and $\omega_{mA}^B(x)$ is of course of the Lorentz type. Its independent components turn out to be

$$\begin{aligned} \omega_{nml} &\equiv e_m^a e_{lb} \omega_{na}^b = \\ &= \frac{1}{2} \left[e_{na} (\partial_m e_l^a - \partial_l e_m^a) - e_{la} (\partial_n e_m^a - \partial_m e_n^a) - e_{ma} (\partial_l e_n^a - \partial_n e_l^a) \right] \\ &\quad + \frac{i}{4} \left[e_{na} (\psi_l \sigma^a \bar{\psi}_m - \psi_m \sigma^a \bar{\psi}_l) - e_{la} (\psi_m \sigma^a \bar{\psi}_n - \psi_n \sigma^a \bar{\psi}_m) - e_{ma} (\psi_n \sigma^a \bar{\psi}_l - \psi_l \sigma^a \bar{\psi}_n) \right] \end{aligned} \quad (2.15)$$

This has the same symmetry properties in the indices as the usual spin connection and reduces to it when the gravitino field is set to 0.

It is then easy to prove, using (2.3), that

$$R_{nma}^b|_{\theta=\bar{\theta}=0} = \partial_n \omega_{ma}^b - \partial_m \omega_{na}^b + \omega_{ma}^c \omega_{nc}^b - \omega_{na}^c \omega_{mc}^b \equiv \mathcal{R}_{nma}^b \quad (2.16)$$

This relation will be used later on. In conclusion the independent component fields are the vierbein, the gravitino and the two auxiliary fields M and b_a .

Let us come now to the interpretation of the superconformal transformations (2.6). To this end we expand the chiral superfield $\sigma(z)$ in the following way:

$$\sigma(z) = \omega(x) + i\alpha(x) + \sqrt{2}\Theta^\alpha \chi_\alpha(x) + \Theta^\alpha \Theta_\alpha (F(x) + iG(x)) \quad (2.17)$$

where we have introduced new anticommuting variables Θ^α , which, unlike θ^μ , carry Lorentz indices. This is always possible, see [23]: the first term on the r.h.s. corresponds to $\sigma|_{\theta=\bar{\theta}=0}$, χ_α to $\nabla_\alpha\sigma|_{\theta=\bar{\theta}=0}$, and $F(x) + iG(x)$ to $\nabla^\alpha\nabla_\alpha\sigma|_{\theta=\bar{\theta}=0}$. Comparing now with the first equation in (2.6), and taking into account (2.11), (2.12), we see that $\omega(x)$ is the parameter of the ordinary Weyl transformation, while comparing with the second and third equation in (2.6) one can see that ψ_α and $\bar{\psi}^{\dot{\alpha}}$ transform with opposite signs with respect to the parameter $\alpha(x)$. Thus $\alpha(x)$ is the parameter of an ordinary chiral transformation.

Therefore when (2.10) is inserted in (2.9) the term linear in $\omega(x)$ will represent a conformal anomaly, while the term linear in $\alpha(x)$ will represent a chiral (Delbourgo-Salam) anomaly. Similarly the term linear in χ_α is the supercurrent anomaly. For the meaning of the cocycles linear in $F(x)$ and $G(x)$ see for instance [27]. Not surprisingly all these anomalies form an $N = 1$ supermultiplet.

The next step is to derive the conformal and chiral anomalies in components.

2.3 Anomalies in components

To derive the anomalies in components we have to integrate out the anticommuting variables. To this end it is convenient to use, instead of the superdeterminant E , the chiral density \mathcal{E} (see [23]). The latter is defined by

$$\mathcal{E}(z) = a(x) + \sqrt{2}\Theta\rho(x) + \Theta\Theta f(x) \tag{2.18}$$

where $a(x) = \frac{1}{2}e(x) \equiv \frac{1}{2}\det e_m^a$. The ρ and f components contain, beside e the gravitino and/or the auxiliary field M , and they vanish when the latter are set to 0. We can rewrite our two integrated cocycles as follows

$$\Delta_\sigma^{(i)} = \int d^4x \left(\int d^2\Theta \mathcal{E}(z) \sigma(z) \mathcal{S}_i(z) + h.c. \right), \quad i = 1, 2 \tag{2.19}$$

This means that, given the interpretation of the lowest components of $\sigma(z)$ as the parameters of the conformal and chiral transformations, and due to (2.18), the *ordinary* part of the conformal and chiral anomaly terms (i.e. the terms linear in ω and α) will depend on $\nabla\nabla\mathcal{S}_i \equiv \nabla^\alpha\nabla_\alpha\mathcal{S}_i$, because this corresponds to the coefficient of $\Theta\Theta$ in the expansion of \mathcal{S}_i . So finally we can write

$$\Delta_\sigma^{(i)} \approx 4 \int d^4x \left(\frac{1}{2}e(\omega + i\alpha) \nabla\nabla\mathcal{S}_i|_{\theta=\bar{\theta}=0} + h.c. \right), \quad i = 1, 2 \tag{2.20}$$

where \approx means ‘up to terms that vanish when all the fields except the metric are set to 0’. The anomalous trace of the energy-momentum tensor and the divergence of the chiral current are obtained from the integral on the r.h.s. of (2.20) by stripping off it the parameters ω and α , respectively.

2.3.1 The square Weyl cocycle

Let us start from S_1 . The relevant terms to be considered are $\nabla^\alpha\nabla_\alpha W_{\beta\gamma\delta}W^{\beta\gamma\delta}$ and $\nabla^\alpha W^{\beta\gamma\delta}\nabla_\alpha W_{\beta\gamma\delta}$ at $\theta = \bar{\theta} = 0$. The term $W_{\beta\gamma\delta}|_{\theta=\bar{\theta}=0}$ is linear in the gravitino field

and in the field b_a . As a consequence this term does not affect the ordinary part of the anomaly. On the contrary the square derivative of W does affect the ordinary part of the anomaly. It is therefore necessary to compute it explicitly. The symmetric part of $\nabla_\alpha W_{\beta\gamma\delta}|_{\theta=\bar{\theta}=0}$ can be computed as follows. Let us consider the identity

$$R_{nma}{}^b = E_n{}^c E_m{}^d R_{cda}{}^b + E_n{}^\gamma E_m{}^d R_{\gamma da}{}^b + E_n{}^c E_m{}^{\bar{d}} R_{c\bar{d}a}{}^b - E_n{}^\gamma E_m{}^{\bar{d}} R_{\gamma\bar{d}a}{}^b \quad (2.21)$$

and evaluate it at $\theta = \bar{\theta} = 0$. We know the l.h.s. due to (2.16). The r.h.s. contains various expressions, and in particular the totally symmetrized derivative $\nabla_{(\alpha} W_{\beta\gamma\delta)}$. It is possible to project it out and get

$$\nabla_{(\alpha} W_{\beta\gamma\delta)}|_{\theta=\bar{\theta}=0} = -\frac{1}{16}(\sigma^a \bar{\sigma}^b \epsilon)_{(\alpha\beta}(\sigma^c \bar{\sigma}^d \epsilon)_{\gamma\delta)} \mathcal{R}_{abcd} \quad (2.22)$$

and similarly

$$\nabla_{(\dot{\alpha}} W_{\dot{\beta}\dot{\gamma}\dot{\delta}})|_{\theta=\bar{\theta}=0} = -\frac{1}{16}(\epsilon \bar{\sigma}^a \sigma^b)_{(\dot{\alpha}\dot{\beta}}(\epsilon \bar{\sigma}^c \sigma^d)_{\dot{\gamma}\dot{\delta}}) \mathcal{R}_{abcd} \quad (2.23)$$

where $\mathcal{R}_{abcd} = e_a{}^n e_b{}^m \mathcal{R}_{nmcd}$.

Using the second equation in (2.1) one can easily obtain

$$\begin{aligned} \nabla_\alpha W_{\beta\gamma\delta} &= \nabla_{(\alpha} W_{\beta\gamma\delta)} \\ &+ \frac{i}{4} \left(\epsilon_{\alpha\beta}(\sigma^{ab}\epsilon)_{\gamma\delta} + \epsilon_{\alpha\gamma}(\sigma^{ab}\epsilon)_{\beta\delta} + \epsilon_{\alpha\delta}(\sigma^{ab}\epsilon)_{\beta\gamma} \right) (\nabla_a G_b - \nabla_b G_a) \end{aligned} \quad (2.24)$$

and a similar equation for the conjugate derivative. Now let us see, as an example of arguments that will be repeatedly used in the sequel, that $\nabla_a G_b - \nabla_b G_a$ evaluated at $\theta = \bar{\theta} = 0$ does not contribute to the ordinary part of the anomaly. In fact $\nabla_a G_b$ cannot contribute to it, for we have

$$\nabla_a G_b = E_a{}^M \partial_M G_b + E_a{}^M \phi_{Mb}{}^c G_c$$

The last term, when evaluated at $\theta = \bar{\theta} = 0$ is linear in the field b_c . The second term in the r.h.s. can be written

$$E_a{}^M \partial_M G_b|_{\theta=\bar{\theta}=0} = -\frac{1}{3} e_a{}^m \partial_m b_a - \frac{1}{2} e_a{}^m \psi_m{}^\alpha \nabla_\alpha G_b| - \frac{1}{2} e_a{}^m \bar{\psi}_{m\dot{\alpha}} \nabla^{\dot{\alpha}} G_b|$$

where the vertical bar stands for $|_{\theta=\bar{\theta}=0}$. Since both $\nabla_\alpha G_b|$ and $\nabla^{\dot{\alpha}} G_b|$ vanish when the gravitino and the auxiliary fields are set to 0, it follows that also $\nabla_a G_b$ vanishes in the same circumstances. Therefore for our purposes only the completely symmetrized spinor derivative of W matters in (2.24). We will write

$$\nabla_\alpha W_{\beta\gamma\delta} \approx \nabla_{(\alpha} W_{\beta\gamma\delta)}, \quad \nabla_{\dot{\alpha}} W_{\dot{\beta}\dot{\gamma}\dot{\delta}} \approx \nabla_{(\dot{\alpha}} W_{\dot{\beta}\dot{\gamma}\dot{\delta})} \quad (2.25)$$

to signify that the l.h.s. is equal to the r.h.s. up to terms that vanish when the gravitino and the auxiliary fields are set to 0.

Now it is a lengthy but standard exercise to verify that

$$\nabla^\alpha W^{\beta\gamma\delta} \nabla_\alpha W_{\beta\gamma\delta} \approx \frac{1}{8} \left(\mathcal{R}_{nmkl} \mathcal{R}^{nmkl} - 2\mathcal{R}_{nm} \mathcal{R}^{nm} + \frac{1}{3} \mathcal{R}^2 + \frac{i}{2} \epsilon^{nmkl} \mathcal{R}_{nmcd} \mathcal{R}_{lk}{}^{cd} \right) \quad (2.26)$$

where $\mathcal{R}_{nmkl} = e_n^a e_m^b e_k^c e_l^d \mathcal{R}_{abcd}$, $\mathcal{R}_{nm} = e_k^a e^{kb} \mathcal{R}_{anbm}$ and $\mathcal{R} = e_n^a e^{nc} e_m^b e^{md} \mathcal{R}_{abcd}$. The first three terms in brackets in the r.h.s. are easily recognized to correspond to the ordinary Weyl density, while the fourth term is the Pontryagin density. We thus have

$$\begin{aligned} \Delta_\sigma^{(1)} &\approx 4 \int d^4x e \left[(\omega + i\alpha) \nabla^\alpha W^{\beta\gamma\delta} \nabla_\alpha W_{\beta\gamma\delta} + h.c. \right] \\ &\approx \frac{1}{2} \int d^4x e \left[(\omega + i\alpha) \left(\mathcal{R}_{nmkl} \mathcal{R}^{nmkl} - 2\mathcal{R}_{nm} \mathcal{R}^{nm} + \frac{1}{3} \mathcal{R}^2 + \frac{i}{2} \epsilon^{nmkl} \mathcal{R}_{nmcd} \mathcal{R}_{lk}{}^{cd} \right) + h.c. \right] \\ &= \int d^4x e \left\{ \omega \left(\mathcal{R}_{nmkl} \mathcal{R}^{nmkl} - 2\mathcal{R}_{nm} \mathcal{R}^{nm} + \frac{1}{3} \mathcal{R}^2 \right) - \frac{1}{2} \alpha \epsilon^{nmkl} \mathcal{R}_{nmpq} \mathcal{R}_{lk}{}^{pq} \right\} \end{aligned} \quad (2.27)$$

In the last line one recognizes the conformal Weyl anomaly linear in ω and the Delbourgo-Salam anomaly linear in α .

2.3.2 The Gauss-Bonnet cocycle

The second cocycle is determined by $\nabla \nabla \mathcal{S}_2|_{\theta=\bar{\theta}=0}$ and its hermitean conjugate. Since G_a, R, R^+ and their first order spinorial derivative evaluated at $\theta = \bar{\theta} = 0$ all vanish when the gravitino and auxiliary fields are set to 0, the *ordinary* part of the cocycle will be determined by

$$\nabla \nabla \mathcal{S}_2| \approx -4 \nabla^\beta \bar{\nabla}_{\dot{\alpha}} G_a \nabla_\beta \bar{\nabla}^{\dot{\alpha}} G^a + 2 \nabla^\alpha \nabla_\alpha R \bar{\nabla}_{\dot{\alpha}} \bar{\nabla}^{\dot{\alpha}} R^+ \quad (2.28)$$

The second term is well known, see [23]. We have $\nabla \nabla R| \approx -\frac{1}{3} \mathcal{R}$, so

$$\nabla^\alpha \nabla_\alpha R \bar{\nabla}_{\dot{\alpha}} \bar{\nabla}^{\dot{\alpha}} R^+| \approx \frac{1}{9} \mathcal{R}^2 \quad (2.29)$$

It remains for us to compute $\nabla_\beta \bar{\nabla}_{\dot{\alpha}} G_a|$. From (2.21) we can derive $\bar{\nabla}_{\dot{\alpha}} \nabla_\beta G_a|$. On the other hand we have

$$(\nabla_\beta \bar{\nabla}_{\dot{\alpha}} + \bar{\nabla}_{\dot{\alpha}} \nabla_\beta) G_a = R_{\dot{\alpha}\beta a}{}^b G_b - T_{\dot{\alpha}\beta}{}^B \nabla_B G_a = -2G_a \sigma^b{}_{\beta\dot{\alpha}} G_b - 2i\sigma^b{}_{\beta\dot{\alpha}} \nabla_b G_a \approx 0$$

Therefore

$$\nabla_\beta \bar{\nabla}_{\dot{\alpha}} G_a \approx -\bar{\nabla}_{\dot{\alpha}} \nabla_\beta G_a \quad (2.30)$$

Next, using the notation $\bar{\nabla}_{\dot{\alpha}} \nabla_\alpha G_{\beta\dot{\beta}} = \sigma^a{}_{\beta\dot{\beta}} \bar{\nabla}_{\dot{\alpha}} \nabla_\alpha G_a$, we introduce the following decomposition

$$\bar{\nabla}_{\dot{\alpha}} \nabla_\alpha G_{\beta\dot{\beta}} = A_{(\alpha\beta)(\dot{\alpha}\dot{\beta})} + \epsilon_{\alpha\beta} B_{\dot{\alpha}\dot{\beta}} + \epsilon_{\dot{\alpha}\dot{\beta}} C_{(\alpha\beta)} + \epsilon_{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\beta}} D \quad (2.31)$$

Now we remark that (2.21) contains the part of $\bar{\nabla}_{\dot{\alpha}} \nabla_\alpha G_{\beta\dot{\beta}}$ which is symmetric both in the couple α, β and $\dot{\alpha}, \dot{\beta}$. After some lengthy but straightforward calculation one can extract it and get

$$A_{(\alpha\beta)(\dot{\alpha}\dot{\beta})} = -\frac{1}{2} R_{abcd} (\sigma^{ab} \epsilon)_{\alpha\beta} (\bar{\epsilon} \bar{\sigma}^{cd})_{\dot{\gamma}\dot{\delta}} \quad (2.32)$$

Next, contracting the decomposition (2.31) with $\epsilon^{\beta\alpha}$ and using the first equation in (2.1) we get

$$\epsilon^{\beta\alpha}\bar{\nabla}_{\dot{\alpha}}\nabla_{\alpha}G_{\beta\dot{\beta}} = 2B_{\dot{\alpha}\dot{\beta}} + 2\epsilon_{\dot{\alpha}\dot{\beta}}D = \nabla_{\dot{\alpha}}\nabla_{\dot{\beta}}R^+ \approx -\frac{1}{2}\epsilon_{\dot{\alpha}\dot{\beta}}\bar{\nabla}\bar{\nabla}R^+ \quad (2.33)$$

A similar result one gets by contracting (2.31) with $\epsilon^{\dot{\alpha}\beta}$. We conclude that

$$\begin{aligned} B_{(\dot{\alpha}\dot{\beta})} &| \approx 0, & C_{(\alpha\beta)} &| \approx 0 \\ D &| \approx -\frac{1}{4}\nabla\nabla R & \approx -\frac{1}{4}\bar{\nabla}\bar{\nabla}R^+ & \approx \frac{1}{12}\mathcal{R} \end{aligned} \quad (2.34)$$

The remaining computation is straightforward. We get

$$\begin{aligned} \nabla\nabla S_2 &| \approx -4\nabla^{\beta}\bar{\nabla}_{\dot{\alpha}}G_a\nabla_{\beta}\bar{\nabla}^{\dot{\alpha}}G^a + 2\nabla^{\alpha}\nabla_{\alpha}R\bar{\nabla}_{\dot{\alpha}}\bar{\nabla}^{\dot{\alpha}}R^+ \\ &\approx \frac{4}{9}\mathcal{R}^2 - 2\mathcal{R}_{nm}\mathcal{R}^{nm} + \frac{2}{9}\mathcal{R}^2 = \frac{2}{3}\mathcal{R}^2 - 2\mathcal{R}_{nm}\mathcal{R}^{nm} \end{aligned} \quad (2.35)$$

that is

$$\Delta_{\sigma}^{(2)} = 4 \int d^4x e \omega \left(\frac{2}{3}\mathcal{R}^2 - 2\mathcal{R}_{nm}\mathcal{R}^{nm} \right) \quad (2.36)$$

This is not the Gauss-Bonnet density, as one could have expected. But it is easy to recover it by means of a linear combination of $\Delta_{\sigma}^{(1)}$ and $\Delta_{\sigma}^{(2)}$:

$$\Delta_{\sigma}^{(1)} + \frac{1}{2}\Delta_{\sigma}^{(2)} \approx \int d^4x e \left\{ \omega (\mathcal{R}_{nmkl}\mathcal{R}^{nmkl} - 4\mathcal{R}_{nm}\mathcal{R}^{nm} + \mathcal{R}^2) - \frac{1}{2}\alpha \epsilon^{nmlk}\mathcal{R}_{nmpq}\mathcal{R}_{lk}{}^{pq} \right\} \quad (2.37)$$

which contains precisely the Gauss-Bonnet density.²

In conclusion $\Delta_{\sigma}^{(1)}$ corresponds to a multiplet of anomalies, whose first component is the Weyl density multiplied by ω , accompanied by the Pontryagin density (the Delbourgo-Salam anomaly) multiplied by α . On the other hand $\Delta_{\sigma}^{(2)}$ does not contain the Pontryagin density and the part linear in ω is a combination of the Weyl and Gauss-Bonnet density.

3 Non minimal supergravity

In supergravity there is a freedom in imposing the torsion constraints. A convenient choice is in terms of the so-called ‘natural constraints’

$$\begin{aligned} T_{ab}{}^c &= 0, & T_{\alpha\beta}{}^a &= T_{\dot{\alpha}\dot{\beta}}{}^a = 0, & T_{\alpha\dot{\beta}}{}^a &= 2i\sigma_{\alpha\dot{\beta}}^a, \\ T_{\gamma}{}^{\dot{\beta}}{}_{\dot{\alpha}} &= (n-1)\delta_{\dot{\alpha}}^{\dot{\beta}}T_{\gamma}, & T^{\dot{\gamma}}{}_{\beta}{}^{\alpha} &= (n-1)\delta_{\beta}^{\alpha}\bar{T}^{\dot{\gamma}}, \\ T_{\gamma\beta}{}^{\alpha} &= (n+1)(\delta_{\gamma}^{\alpha}T_{\beta} + \delta_{\beta}^{\alpha}T_{\gamma}), & T^{\dot{\gamma}\dot{\beta}}{}_{\dot{\alpha}} &= (n+1)(\delta_{\dot{\alpha}}^{\dot{\beta}}\bar{T}^{\dot{\gamma}} + \delta_{\dot{\alpha}}^{\dot{\beta}}\bar{T}^{\dot{\gamma}}) \\ T_{\gamma b}{}^a &= 2n\delta_b^a T_{\gamma}, & T^{\dot{\gamma}}{}_{b}{}^a &= 2n\delta_b^a \bar{T}^{\dot{\gamma}} \end{aligned} \quad (3.1)$$

where n is a numerical parameter and $T_{\alpha}, \bar{T}_{\dot{\alpha}}$ are new (conjugate) superfields in addition to those of minimal supergravity. The latter is obtained by setting $T_{\alpha} = 0$. $T_{\alpha}, \bar{T}_{\dot{\alpha}}$ are U(1)

²For an early appearance of the Gauss-Bonnet and Weyl density anomalies in supergravity see [30, 31].

connections. The $U(1) \times U(1)$ gauge symmetry was added with the purpose of enlarging the minimal supergravity model. The solution for the Bianchi identities can be found in [24]. There are many significant changes with respect to the minimal model. For instance $W_{\alpha\beta\gamma}$ and R are not chiral anymore, but

$$(\bar{\mathcal{D}}_{\dot{\alpha}} + (3n + 1)\bar{T}_{\dot{\alpha}})W_{\alpha\beta\gamma} = 0 \tag{3.2}$$

$$(\bar{\mathcal{D}}_{\dot{\alpha}} + 2(n + 1)\bar{T}_{\dot{\alpha}})R = 0 \tag{3.3}$$

where \mathcal{D} replaces ∇ as covariant derivative.³

A distinguished superfield is S (and its conjugate \bar{S}), defined by

$$S = \mathcal{D}^{\alpha}T_{\alpha} - (n + 1)T^{\alpha}T_{\alpha}, \tag{3.4}$$

which satisfies

$$\mathcal{D}_{\alpha}S = 8T_{\alpha}R^{+} \tag{3.5}$$

The combination

$$Y = 8R + 2(n + 1)\bar{S} \tag{3.6}$$

is chiral, $\bar{\mathcal{D}}^{\dot{\alpha}}Y = 0$. The operator

$$\Delta = \mathcal{D}^{\alpha}\mathcal{D}_{\alpha} - 3(n + 1)T^{\alpha}\mathcal{D}_{\alpha} - Y \tag{3.7}$$

projects a superfield without Lorentz indices to an antichiral superfield and $\frac{\Delta}{Y}$ is a chiral projector.

The *non minimal model* for supergravity is obtained by further imposing the constraint

$$R = R^{+} = 0 \tag{3.8}$$

with nonvanishing T_{α} and $\bar{T}_{\dot{\alpha}}$.

The non minimal supergravity has 20+20 degrees of freedom. The bosonic dofs are those of the minimal model, excluding R and R^{+} , plus 10 additional ones which can be identified with the lowest components of the superfields $S, \bar{S}, \bar{\mathcal{D}}_{\dot{\alpha}}T_{\alpha} = c_{\alpha\dot{\alpha}} + id_{\alpha\dot{\alpha}}$ and $\bar{\mathcal{D}}_{\alpha}T_{\dot{\alpha}} = -c_{\alpha\dot{\alpha}} + id_{\alpha\dot{\alpha}}$. The additional fermionic dofs can be identified with the lowest components of $T_{\alpha}, \bar{T}_{\dot{\alpha}}$ and $\mathcal{D}_{\alpha}\bar{S}, \bar{\mathcal{D}}_{\dot{\alpha}}S$. We will refer to this model as 20+20 nonminimal, or nonminimal supergravity for short.

3.1 Superconformal transformations in the non minimal model

In the non minimal model there are transformations compatible with the constraints that correspond to local vierbein rescalings. We will refer to them generically as superconformal transformations. They are good candidates for superWeyl transformations (i.e, for supersymmetric extensions of the ordinary Weyl transformations) but, as we shall see, do not

³In principle there is no reason to use two different symbols for the covariant derivative, they denote the same covariant derivative in different settings. The use of two different symbols, however, will be instrumental in section 7.

automatically correspond to them. They are expressed in terms of an arbitrary (complex) superfield Σ

$$\begin{aligned}
 \delta E_\alpha{}^M &= -(2\bar{\Sigma} - \Sigma) E_\alpha{}^M \\
 \delta E_{\dot{\alpha}}{}^M &= -(2\Sigma - \bar{\Sigma}) E_{\dot{\alpha}}{}^M \\
 \delta E_a{}^M &= -(\Sigma + \bar{\Sigma}) E_a{}^M + \frac{i}{2} \bar{\sigma}_a^{\dot{\beta}\beta} \bar{\mathcal{D}}_{\dot{\beta}} \left(\bar{\Sigma} - \frac{3n-1}{3n+1} \Sigma \right) E_\beta{}^M \\
 &\quad + \frac{i}{2} \bar{\sigma}_a^{\beta\dot{\beta}} \mathcal{D}_\beta \left(\Sigma - \frac{3n-1}{3n+1} \bar{\Sigma} \right) E_{\dot{\beta}}{}^M \\
 \delta T_\alpha &= -(2\bar{\Sigma} - \Sigma) T_\alpha + \frac{3}{3n+1} \mathcal{D}_\alpha \bar{\Sigma} \\
 \delta W_{\alpha\beta\gamma} &= -3\Sigma W_{\alpha\beta\gamma} \\
 \delta G_a &= -(\Sigma + \bar{\Sigma}) G_a + i\mathcal{D}_a(\bar{\Sigma} - \Sigma) + \frac{1}{3} \bar{\sigma}_a^{\dot{\alpha}\alpha} (T_\alpha \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\Sigma} - \bar{T}_{\dot{\alpha}} \mathcal{D}_\alpha \Sigma) \\
 &\quad - \frac{3n-1}{3(3n+1)} \bar{\sigma}_a^{\dot{\alpha}\alpha} (T_\alpha \bar{\mathcal{D}}_{\dot{\alpha}} \Sigma - \bar{T}_{\dot{\alpha}} \mathcal{D}_\alpha \bar{\Sigma}) \\
 \delta R^+ &= -2(2\bar{\Sigma} - \Sigma) R^+ \\
 &\quad + \frac{1}{4(3n+1)} (\mathcal{D}^\alpha \mathcal{D}_\alpha + (n+1) T^\alpha \mathcal{D}_\alpha) [3n(\bar{\Sigma} - \Sigma) - (\bar{\Sigma} + \Sigma)]
 \end{aligned} \tag{3.9}$$

From (3.8) and (3.9) we see that the superfield Σ is constrained by the linear condition

$$(\mathcal{D}^\alpha \mathcal{D}_\alpha + (n+1) T^\alpha \mathcal{D}_\alpha) [3n(\bar{\Sigma} - \Sigma) - (\bar{\Sigma} + \Sigma)] = 0 \tag{3.10}$$

3.2 Cocycles in non minimal SUGRA

In the non minimal model it is easy to construct an invariant (0-cocycle)

$$I_{n.m.}^{(1)} = \int_{x,\theta} E W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \frac{\bar{T}_{\dot{\alpha}} \bar{T}^{\dot{\alpha}}}{\bar{S}^2} + h.c. \tag{3.11}$$

and a 1-cocycle

$$\Delta_{n.m.}^{(1)} = \int_{x,\theta} E \Sigma W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \frac{\bar{T}_{\dot{\alpha}} \bar{T}^{\dot{\alpha}}}{\bar{S}^2} + h.c. \tag{3.12}$$

It is easy to prove that $\delta I_{n.m.}^{(1)} = 0 = \delta \Delta_{n.m.}^{(1)}$ for any n . To this end the condition (3.10) is inessential. If $R \neq 0$ this is not true anymore.

The construction of a second cocycle corresponding to $\Delta_\sigma^{(2)}$ above, is not as straightforward and will be postponed to section 7, after the technique of mapping between different supergravity models has been introduced.

4 The 16+16 nonminimal model

One way to define the 16+16 nonminimal model is to introduce a 2-superform B_{AB} and impose natural constraints on its supercurvature. In this way we obtain a 16+16 model.

The independent bosonic dofs are the vierbein, the lowest component of S, \bar{S} , $c_{\alpha\dot{\alpha}}$ and $G_{\alpha\dot{\alpha}}$ (the components of $d_{\alpha\dot{\alpha}}$ are not independent in this model). The fermionic degrees of freedom are, beside the gravitino field, the lowest components of $T_\alpha, \bar{T}_{\dot{\alpha}}$ and $\mathcal{D}_\alpha \bar{S}, \mathcal{D}_{\dot{\alpha}} S$. The new dofs (with respect to the minimal model) are linked to the mode contained in B_{ab} . In 16+16 nonminimal supergravity the range of the parameter n is limited to $n > 0$ and $n < -\frac{1}{3}$.

In practice this means that

$$T_\alpha = \mathcal{D}_\alpha \psi, \quad T_{\dot{\alpha}} = \mathcal{D}_{\dot{\alpha}} \psi \quad (4.1)$$

where ψ is a (dimensionless) real superfield. The transformations corresponding to (3.9) on ψ are

$$\delta\psi = \frac{3}{3n+1}(\bar{\Sigma} - \bar{\Lambda}) = \frac{3}{3n+1}(\Sigma - \Lambda) \equiv \frac{3}{3n+1}L \quad (4.2)$$

where $\Lambda(\bar{\Lambda})$ is an arbitrary chiral (antichiral) superfield, and L is a real (vector) superfield. As a consequence the transformations (3.9), compatible with the constraints, for the surviving superfields take the form:

$$\begin{aligned} \delta E_\alpha{}^M &= -(L + 2\bar{\Lambda} - \Lambda) E_\alpha{}^M \\ \delta E_{\dot{\alpha}}{}^M &= -(L + 2\Lambda - \bar{\Lambda}) E_{\dot{\alpha}}{}^M \\ \delta T_\alpha &= -(L + 2\bar{\Lambda} - \Lambda) T_\alpha + \frac{3}{3n+1} \mathcal{D}_\alpha L \\ \delta W_{\alpha\beta\gamma} &= -3(L + \Lambda) W_{\alpha\beta\gamma} \\ \delta G_{\alpha\dot{\alpha}} &= -(2L + \Lambda + \bar{\Lambda}) G_{\alpha\dot{\alpha}} + i\mathcal{D}_{\alpha\dot{\alpha}}(\bar{\Lambda} - \Lambda) - \frac{2}{3} (T_\alpha \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\Lambda} - \bar{T}_{\dot{\alpha}} \mathcal{D}_\alpha \Lambda) \\ &\quad + \frac{2(3n-1)}{3(3n+1)} (T_\alpha \bar{\mathcal{D}}_{\dot{\alpha}} L - \bar{T}_{\dot{\alpha}} \mathcal{D}_\alpha L) \\ \delta S &= -2(L + 2\bar{\Lambda} - \Lambda) S + 4\mathcal{D}^\alpha(L + \Lambda) T_\alpha - \frac{21n+5}{3n+1} \mathcal{D}^\alpha L T_\alpha + \frac{3}{3n+1} \mathcal{D}^\alpha \mathcal{D}_\alpha L \end{aligned} \quad (4.3)$$

and (3.10) becomes

$$(\mathcal{D}^\alpha \mathcal{D}_\alpha + (n+1)T^\alpha \mathcal{D}_\alpha) (2L + (3n+1)\Lambda) = 0 \quad (4.4)$$

4.1 Cocycles in 16+16 nonminimal SUGRA

As in nonminimal SUGRA it is easy to construct an invariant

$$I_{new}^{(1)} = \int_{x,\theta} E W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \frac{\bar{T}_{\dot{\alpha}} \bar{T}^{\dot{\alpha}}}{\bar{S}^2} + h.c. \quad (4.5)$$

and a 1-cocycle

$$\Delta_{new}^{(1)} = \int_{x,\theta} E (L + \Lambda) W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \frac{\bar{T}_{\dot{\alpha}} \bar{T}^{\dot{\alpha}}}{\bar{S}^2} + h.c. \quad (4.6)$$

It is easy to prove that $\delta I_{new}^{(1)} = 0 = \delta \Delta_{new}^{(1)}$. Once again we don't need (4.4) to prove this. On the other hand it is not easy to construct a cocycle similar to $\Delta^{(2)}$, i.e. quadratic in the superfield G_a , which, after translation to component form, leads to the Gauss-Bonnet density.

5 Reduction to component form

In the following analysis the reduction of the cocycles to ordinary form will play a major role. Thus the purpose of this section is to outline the procedure to derive the component form of the cocycles in nonminimal and 16+16 nonminimal supergravities, as we have done in section 2 for the minimal supergravity anomalies. The operator, [24],

$$\bar{\Delta} = \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\mathcal{D}}^{\dot{\alpha}} - 3(n+1) \bar{T}_{\dot{\alpha}} \bar{\mathcal{D}}^{\dot{\alpha}} - 2(n+1) \bar{S} \quad (5.1)$$

projects a generic superfield to a chiral superfield. Let U be a superfield without Lorentz indices. It is not hard to see that (see [24])

$$\int_{x,\theta} E U = \frac{1}{4n} \int_{x,\theta} E \bar{\Delta} \left(e^{-\bar{\Omega}} \frac{U}{\bar{S}} \right) \quad (5.2)$$

where $\bar{\Omega} = 2(3n+1) \frac{\bar{T}\bar{T}}{\bar{S}}$. Therefore, introducing the appropriate chiral density \mathcal{E} , [25], we can write

$$\int_{x,\theta} E U = \frac{1}{4n} \int d^4x \int d^2\Theta \mathcal{E} \bar{\Delta} \left(U e^{\bar{\Omega}} \right) \quad (5.3)$$

For instance, when Σ is a chiral superfield the anomaly (3.12) can be written

$$\Delta_{\Sigma}^{(1)} = \frac{1}{4n} \int d^4x \int d^2\Theta \mathcal{E} \bar{\Delta} \left[\Sigma W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \frac{\bar{T}_{\dot{\alpha}} \bar{T}^{\dot{\alpha}}}{\bar{S}^2} \right] + h.c. \quad (5.4)$$

When Σ is not chiral there is in the r.h.s. an additional term which, however, is irrelevant for the following considerations and so will be dropped. In a similar way we can deal with (4.6). After some algebra we have in particular

$$\Delta_{\Sigma}^{(1)} = -\frac{1}{4n} \int d^4x \int d^2\Theta \mathcal{E} \left(\Sigma W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} + 2 W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \frac{\bar{T}_{\dot{\alpha}} \mathcal{D}^{\dot{\alpha}} \Sigma}{\bar{S}} \right) + h.c. \quad (5.5)$$

Therefore, proceeding as in section 5, $\Delta_{\Sigma}^{(1)}$ in components becomes (we disregard a multiplicative factor)

$$\Delta_{\Sigma}^{(1)} \approx \int d^4x e \left[\Sigma \mathcal{D}^{\alpha} W^{\beta\gamma\delta} \mathcal{D}_{\alpha} W_{\beta\gamma\delta} \right] + h.c. \quad (5.6)$$

Now (2.22), (2.23) remain valid in the non minimal SUGRA, but (2.24) is replaced by a far more complicated equation, so that (2.25) has to be re-demonstrated. This is not trivial, but can be done (see appendix A). Thus we can conclude that, up to a multiplicative factor,

$$\Delta_{\Sigma}^{(1)} \approx \frac{1}{4} \int d^4x e \left\{ \omega \left(\mathcal{R}_{nmkl} \mathcal{R}^{nmkl} - 2 \mathcal{R}_{nm} \mathcal{R}^{nm} + \frac{1}{3} \mathcal{R}^2 \right) - \frac{1}{2} \alpha \epsilon^{nmkl} \mathcal{R}_{nmpq} \mathcal{R}_{lk}{}^{pq} \right\} \quad (5.7)$$

where $\omega + i\alpha$ is the lowest component of the superfield Σ . In this case too ω corresponds to the ordinary Weyl rescaling, while α is the parameter of a chiral transformation.

The same reduction to ordinary form holds also for (4.6). In this case $\omega + i\alpha$ is the first component of $L + \Lambda$.

At this point it is worth making a comment on the (apparent) singularity of expressions such as (3.11), (3.12), (4.5), (4.6). For instance, the cocycle (3.12), written in terms of superfields has a nonlocal or singular aspect due to the presence of an inverse power of S (\bar{S}). It must be stressed that, in the spirit of this paper, gravity is an external (super)field source, all superfield configurations are generic (and regular) and, in particular, not constrained on shell. Therefore such a singularity problem simply does not exist. On a different level, however, one may need to evaluate these cocycles on particular superfield configurations, so the singularity problem may become relevant. Since (3.12), for instance, is nothing but the supersymmetrization of (5.7), which is local, we would naively expect that also (3.12), when expressed in terms of components fields be nonsingular (although it may be non-polynomial if dimensionless prepotentials have to be introduced). It would be nice if non-singularity were made manifest at the superfield level. In [25] it was noted that in some cases this is indeed possible by means of opportune field redefinitions. However when the cocycle is evaluated on particular field configurations there may arise the problems pointed out in ref. [28]. So the question in general remains open.

The scheme outlined in this section is general and will be applied to all the cocycles we will come across.

6 Mapping formulas between different supergravity models

A cocycle similar to $\Delta_\sigma^{(2)}$ (i.e. quadratic in G_a) is hard to construct with ordinary means (i.e. with a polynomial cohomological analysis). For this we have to resort to a mapping between different supergravity models. This mapping was outlined in [2, 25] and brought to a more explicit form in [22]. The latter reference is based on different torsion constraints with respect to (3.1). Therefore we have to rederive new appropriate mapping formulas.

Various different models of supergravity are defined by making a definite choice of the torsion constraints and, after such a choice, by identifying the dynamical degrees of freedom. This is the way minimal, nonminimal and 16+16 nonminimal models were introduced. However it is possible to transform the choices of constraints into one another by means of suitable linear transformation of the supervierbein and the superconnection, [22, 24]:

$$E'_M{}^A = E_M{}^B X_B{}^A, \quad E'_A{}^M = X^{-1}{}^B{}_A E_B{}^M, \quad \Phi'_{MA}{}^B = \Phi_{MA}{}^B + \chi_{MA}{}^B \quad (6.1)$$

For instance, if we want to pass from a set of unprimed constraints to primed ones the required transformations are as follows

$$E'_\alpha = U E_\alpha, \quad E'^{\dot{\alpha}} = \bar{U} E^{\dot{\alpha}}, \quad E' = U^{-2} \bar{U}^{-2} E \quad (6.2)$$

$$E'_{\alpha\dot{\alpha}} = U \bar{U} E_{\alpha\dot{\alpha}} + i \frac{U \bar{U}}{3n+1} \left(E_{\dot{\alpha}}{}^M \partial_M \ln \left(\frac{U^{n+1}}{\bar{U}^{n-1}} \right) E_\alpha + E_\alpha{}^M \partial_M \ln \left(\frac{\bar{U}^{n+1}}{U^{n-1}} \right) E_{\dot{\alpha}} \right) \quad (6.3)$$

where U is a suitable expression of the superfields. Moreover

$$T'_\alpha = UT_\alpha - \frac{1}{6n+2} \mathcal{D}'_\alpha \ln(\bar{U}^2 U^4) \quad (6.4)$$

$$\Phi'_{\alpha\beta\gamma} = U \Phi_{\alpha\beta\gamma} - \frac{1}{3n+1} (\epsilon_{\alpha\gamma} \mathcal{D}'_\beta + \epsilon_{\alpha\beta} \mathcal{D}'_\gamma) \ln\left(\frac{U^{n-1}}{\bar{U}^{n+1}}\right) \quad (6.5)$$

$$W'_{\alpha\beta\gamma} = U \bar{U}^2 W_{\alpha\beta\gamma} \quad (6.6)$$

$$8R' + 2(n+1)\bar{S}' = -\left(\bar{\mathcal{D}}'_\alpha \bar{\mathcal{D}}'^{\dot{\alpha}} - 3(n+1)\bar{T}'_\alpha \bar{\mathcal{D}}'^{\dot{\alpha}} - 8R - 2(n+1)\bar{S}\right) \bar{U}^2 \quad (6.7)$$

where \mathcal{D}' denotes the covariant derivative in the primed system, together with the conjugate relations. The analogous transformation for the G_a superfield is more complicated:

$$\begin{aligned} G'_{\alpha\dot{\alpha}} = U\bar{U} & \left(G_{\alpha\dot{\alpha}} - \frac{i}{3} \mathcal{D}'_{\alpha\dot{\alpha}} \ln \frac{U}{\bar{U}} + \frac{1}{(3n+1)^2} \bar{\mathcal{D}}'_\alpha \ln \frac{U^{n+1}}{\bar{U}^{n-1}} \mathcal{D}'_\alpha \ln \frac{\bar{U}^{n+1}}{U^{n-1}} \right. \\ & + \frac{1}{3(3n+1)} \bar{\mathcal{D}}'_\alpha \ln \frac{U^{n+1}}{\bar{U}^{n-1}} \mathcal{D}'_\alpha \frac{U}{\bar{U}} + \frac{1}{3(3n+1)} \mathcal{D}'_\alpha \ln \frac{\bar{U}^{n+1}}{U^{n-1}} \bar{\mathcal{D}}'_\alpha \frac{U}{\bar{U}} \\ & \left. + \frac{2}{3(3n+1)} \mathcal{D}'_\alpha \ln \frac{\bar{U}^{n+1}}{U^{n-1}} \bar{T}'_\alpha - \frac{2}{3(3n+1)} \bar{\mathcal{D}}'_\alpha \ln \frac{U^{n+1}}{\bar{U}^{n-1}} T_\alpha \right) \end{aligned} \quad (6.8)$$

These formulas can be inverted. To this end we have to replace U with U^{-1} everywhere, replace the primed quantities with unprimed ones in the l.h.s., and the unprimed with the primed ones in r.h.s.; in this case the covariant derivatives on the r.h.s. are the primed ones.⁴

For instance, if we want to pass from the minimal to the nonminimal constraints we have to choose

$$U = \exp \left[2(3n+1) \left(\frac{\bar{\psi}}{6} - \frac{\psi}{3} \right) \right] \quad (6.9)$$

ψ is a ‘prepotential’ such that $T_\alpha = \mathcal{D}_\alpha \psi$ and $\bar{T}'_{\dot{\alpha}} = \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\psi}$. Of course if we wish to pass from the nonminimal to the minimal constraints we have simply to use the same formulas with inverted U .

One can verify that

$$(\bar{\nabla}'_{\dot{\alpha}} \bar{\nabla}'^{\dot{\alpha}} - 8R) \bar{U}^2 = -2(n+1) \bar{S} \quad (6.10)$$

We recall that ∇ denotes specifically the covariant derivative in minimal supergravity.

Let us consider next the superconformal transformations. We wish to compare the transformations (2.6), (2.7) with (3.9). Given the transformation of T_α and $T_\alpha = \mathcal{D}_\alpha \psi$, we can assume that $\psi, \bar{\psi}$ transform as follows

$$\delta\psi = \frac{3}{3n+1} (\bar{\Sigma} - \bar{\sigma}), \quad \delta\bar{\psi} = \frac{3}{3n+1} (\Sigma - \sigma), \quad (6.11)$$

where σ is an arbitrary chiral superfield. Taking the variation of both sides of (6.2) and applying (6.11) we can easily see that we can identify the σ superfield in (6.11) with the

⁴For more details on these transformations, see [29].

σ in (2.6). The same is easily done also for (6.6). The transformation of (6.7) is more complicated. We first derive, using (6.5),

$$\nabla^\alpha \nabla_\alpha \Phi = U^{-2} \left(\mathcal{D} \mathcal{D} \Phi - \frac{4}{3} (3n+1) \mathcal{D}^\alpha \bar{T} \mathcal{D}_\alpha \Phi + \frac{15n-1}{3} T^\alpha \mathcal{D}_\alpha \Phi \right) \quad (6.12)$$

for any scalar superfield Φ . Inverting (6.7) we can write

$$-8R^+ = (\mathcal{D}^\alpha \mathcal{D}_\alpha - 3(n+1)T^\alpha \mathcal{D}_\alpha - 2(n+1)S) U^{-2} \quad (6.13)$$

The l.h.s. represents R in the minimal model, while the r.h.s. refers to the nonminimal one. Taking the variation of both sides and using (6.11), (6.12), one can show that

$$\delta R^+ = -2(2\bar{\sigma} - \sigma)R^+ - \frac{1}{4} \nabla \nabla \sigma \quad (6.14)$$

This is identical to the transformation of R^+ in the minimal model, (2.7).

We can do the same with G_a . Taking the variation of l.h.s. and r.h.s. of the inverted eq. (6.8), and using

$$i\nabla_{\alpha\dot{\alpha}}(\bar{\sigma} - \sigma) = U^{-1} \bar{U}^{-1} \left(i\mathcal{D}_{\alpha\dot{\alpha}}(\bar{\sigma} - \sigma) - \frac{1}{3n+1} \bar{\mathcal{D}}_{\dot{\alpha}} \ln \frac{U^{n+1}}{\bar{U}^{n-1}} \mathcal{D}_\alpha \sigma + \frac{1}{3n+1} \mathcal{D}_\alpha \ln \frac{\bar{U}^{n+1}}{U^{n-1}} \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\sigma} \right)$$

one finds

$$\delta G_{\alpha\dot{\alpha}} = -(\sigma + \bar{\sigma})G_{\alpha\dot{\alpha}} + i\nabla_{\alpha\dot{\alpha}}(\bar{\sigma} - \sigma) \quad (6.15)$$

as expected.

Therefore (6.11) connects the superconformal transformations of the minimal and non-minimal models. It is however useful to consider this passage in two steps. Let us split U in (6.9) as follows:

$$U = U_c U_n, \quad U_c = e^{X-2\bar{X}}, \quad U_n = e^{\frac{\Omega}{3} - \frac{\bar{\Omega}}{6}} \quad (6.16)$$

where

$$X = \frac{1}{3}(3n+1)\bar{\psi} + \frac{\bar{\Omega}}{6}, \quad \bar{X} = \frac{1}{3}(3n+1)\psi + \frac{\Omega}{6} \quad (6.17)$$

Recall that $\Omega = 2(3n+1)\frac{T^\alpha T_\alpha}{S}$ and $T_\alpha = \mathcal{D}_\alpha \psi$, etc. It follows that X is a chiral and \bar{X} an antichiral superfield. Moreover $U_c \bar{U}_c^2 = e^{-3X}$ is chiral and $\bar{U}_c U_c^2 = e^{-3\bar{X}}$ is antichiral. Operating on the superfields according to (6.2), (6.4), (6.6) we see that, for instance $T_\alpha = 0$ is mapped to $T_\alpha = 0$ by the transformation induced by U_c , i.e. after such transformation the model is still minimal supergravity.

For later use we remark that (see also [25])

$$\delta \bar{\Omega} = \Gamma_\Sigma - 6\Sigma, \quad \Gamma_\Sigma = -\frac{3}{3n+1} \bar{\Delta} \left(\frac{\bar{\Omega}\Sigma}{\bar{S}} \right) \quad (6.18)$$

where $\bar{\Delta}$ is the chiral projector. By repeating the previous verifications one can see that Γ_Σ is an intermediate step between σ and Σ . The important property of Γ_Σ is that it is chiral,

but expressed in terms of the nonminimal superfields. Moreover it is consistent with the nonminimal transformation properties and, in particular, $\delta\Gamma_\Sigma = 0$. In parallel with (6.18) we have of course the conjugate formulas.

Analogous things hold if we replace the non minimal with the 16+16 nonminimal model. In this case of course we have to set $\psi = \bar{\psi}$ and the appropriate transformations are (4.2), (4.3). It is easy to see that the above superfield redefinitions connect the minimal supergravity transformations with (4.3). Also in this case we have an intermediate step which will turn out instrumental later on. In this case we have

$$\delta\bar{\Omega} = \Gamma_{L+\Lambda} - 6(L + \Lambda), \quad \Gamma_{L+\Lambda} = -\frac{3}{3n+1}\bar{\Delta}\left(\frac{\bar{\Omega}(L+\Lambda)}{\bar{S}}\right) = \Gamma_L + 6\Lambda \quad (6.19)$$

where Γ_L is chiral.

All this means one important thing: the possibility to construct invariants and cocycles of any supergravity model starting from the invariants and cocycles of a fixed one, for instance the minimal supergravity (such an idea is present in [9]).

7 Cocycles from minimal supergravity

We are now ready to construct the cocycles from those of minimal supergravity. The idea is very simple. We start from the cocycles of minimal supergravity and replace the superfields of the latter with the formulas of the previous subsection expressing them in terms of the superfields of other models. Since all the symmetry operations are coherent, the resulting expressions must also be cocycles. The invariants are a subcase of the discussion for 1-cocycles, thus in the sequel we explicitly deal only with the latter. We will consider first the 16+16 nonminimal case.

7.1 From minimal to nonminimal cocycles

Let us start from $\Delta_\sigma^{(1)}$. All the superfields therein must be expressed in terms of the new superfields. It is convenient to proceed in two steps, as just outlined. In the first step it is mapped to

$$\int_{x,\theta} \frac{E}{-8R} \sigma WW + h.c. = \int_{x,\theta} E' \Gamma_\Sigma \frac{W'W'}{\bar{U}_c^2 (\nabla'_\alpha \nabla'^{\dot{\alpha}} - 8R')} \bar{U}_c^{-2} + h.c. \quad (7.1)$$

where WW is a compact notation for $W^{\alpha\beta\gamma}W_{\alpha\beta\gamma}$ and primes denote the superfields in the new representation (which still corresponds to minimal supergravity). We recall that $\nabla'_\alpha \nabla'^{\dot{\alpha}} - 8R'$ projects to a chiral superfield. Therefore we can write

$$\int_{x,\theta} \frac{E \sigma}{-8R} WW = \int_{x,\theta} \frac{E'}{-8R'} \Gamma_\Sigma (\nabla'_\alpha \nabla'^{\dot{\alpha}} - 8R') \left(\frac{W'W' \bar{U}_c^{-2}}{(\nabla'_\alpha \nabla'^{\dot{\alpha}} - 8R') \bar{U}_c^{-2}} \right) \quad (7.2)$$

$$= \int_{x,\theta} \frac{E'}{-8R'} \Gamma_\Sigma W'W' \quad (7.3)$$

Now we complete the passage to the nonminimal model by performing the U_n transformation. This means

$$\int_{x,\theta} \frac{E'}{-8R'} \Gamma_\Sigma W' W' = \int_{x,\theta} E'' \Gamma_\Sigma \frac{W'' W''}{\bar{U}_n^2 \bar{\Delta}'' \bar{U}_n^{-2}} \quad (7.4)$$

where $\Delta'' = \mathcal{D}'' \mathcal{D}'' - 3(n+1) T''^\alpha \mathcal{D}''_\alpha - 2(n+1) S''$ is the antichiral projector in the nonminimal model (endpoint of the overall transformation). For simplicity, from now on, we drop primes, understanding that we are operating in the nonminimal model.

Next we use the identity, demonstrated in [24] by partial integration,

$$4n \int_{x,\theta} E e^{\bar{\Omega}} U = \int_{x,\theta} E \frac{\Phi}{\bar{S}} \quad (7.5)$$

where U is any superfield expression without Lorentz indices and $\Phi = \bar{\Delta} U$. Applying this identity with $U = \Sigma \frac{e^{-\bar{\Omega}} W W \bar{U}^{-2}}{\bar{\Delta} \bar{U}^{-2}}$ we get

$$\int_{x,\theta} E \Gamma_\Sigma \frac{W W}{\bar{U}^2 \bar{\Delta} \bar{U}^{-2}} = \frac{1}{4n} \int_{x,\theta} \frac{E}{\bar{S}} \Gamma_\Sigma e^{-\bar{\Omega}} W W \quad (7.6)$$

Applying (7.5) again with $U = \Sigma e^{-\bar{\Omega}} \frac{W W}{\bar{S}}$, so that $\Phi = -2(n+1) \Gamma_\Sigma e^{-\bar{\Omega}} W W$, we obtain

$$\frac{1}{4n} \int_{x,\theta} \frac{E}{\bar{S}} \Gamma_\Sigma e^{-\bar{\Omega}} W W = -\frac{1}{2(n+1)} \int_{x,\theta} \frac{E}{\bar{S}} \Gamma_\Sigma W W \quad (7.7)$$

Replacing now the explicit expression of Γ_Σ , (6.18), and integrating by parts, we find that $\Delta_\sigma^{(1)}$ is mapped to

$$3 \frac{5n+1}{n+1} \int_{x,\theta} \frac{E}{\bar{S}^2} \Sigma \bar{T}_\alpha \bar{T}^\alpha W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} + h.c. \quad (7.8)$$

which is proportional to the already obtained cocycle $\Delta_{n.m.}^{(1)}$, (3.12). The second cocycle is readily constructed in the same way:

$$\begin{aligned} \Delta_\Sigma^{(2)} &= \int_{x,\theta} E' (\Gamma_\Sigma + \bar{\Gamma}_\Sigma) U^2 \bar{U}^2 (G_a(G', T', U) G^a(G', T', U) + 2R(\bar{S}', \bar{T}', U) R^+(S', T', U)) \\ &= c \int_{x,\theta} E' (\Sigma + \bar{\Sigma}) (G'_a G'^a + \dots) \end{aligned} \quad (7.9)$$

after repeated partial integrations. $G_a(G', T', U)$ is given by the inverted (6.8), while $R^+(S, T, U)$ is given by (6.13). c is a suitable number. By construction $\Delta_\Sigma^{(2)}$ satisfies the consistency conditions with generic Σ . Its ordinary form is the same as $\Delta_\sigma^{(2)}$ in section 2.

7.2 From minimal to 16+16 nonminimal cocycles

Let us start again with $\Delta_\sigma^{(1)}$. Proceeding as above with the relevant new formulas outlined at the end of the previous section we get

$$\begin{aligned} \Delta_\sigma^{(1)} &= \int_{x,\theta} \frac{E}{-8R} \sigma W W + h.c. \\ &= \int_{x,\theta} \frac{E' \Gamma_{L+\Lambda}}{\bar{U}^2 (\bar{\mathcal{D}} \bar{\mathcal{D}} - 3(n+1) \bar{T}'_\alpha \bar{\mathcal{D}} - 2(n+1) \bar{S}') \bar{U}^{-2}} \frac{W' W'}{\bar{U}^{-2}} + h.c. \equiv \tilde{\Delta}_{L+\Lambda}^{(1)} \end{aligned} \quad (7.10)$$

where primed superfields refers to 16+16 nonminimal supergravity. From now on we drop primes, understanding that all the superfields are in the 16+16 nonminimal supergravity. Working out the derivatives in (7.10) we get

$$\tilde{\Delta}_{L+\Lambda}^{(1)} = -\frac{3}{4} \int_{x,\theta} \frac{E}{\bar{S}} \Gamma_{L+\Lambda} WW \left(1 - \frac{2}{3}(3n+1) \frac{\bar{T}_{\dot{\alpha}} \bar{T}^{\dot{\alpha}}}{\bar{S}} \right) + h.c. \quad (7.11)$$

This is not (4.6) yet, as we would have expected. However, using (5.2) and integrating by parts the spinor derivatives contained in $\Gamma_{L+\Lambda}$, as we have done above for the nonminimal case, one easily finds that $\tilde{\Delta}_{L+\Lambda}^{(1)}$ is proportional to (4.6).

Let us come now to the second cocycle. As above we start from the minimal cocycle $\Delta_{\sigma}^{(2)}$ and transform the superfields according to (6.13) and (6.8). We get

$$\begin{aligned} \Delta_{\sigma}^{(2)} &= \int_{x,\theta} E(\sigma + \bar{\sigma})(G_a G^a + 2RR^+) \\ &= \int_{x,\theta} E'(\Gamma_{L+\Lambda} + h.c.) \left(-\frac{1}{2} \left(G'_{\alpha\dot{\alpha}} + \frac{4}{9} T_{\alpha} \bar{T}_{\dot{\alpha}} \right) \left(G'^{\alpha\dot{\alpha}} + \frac{4}{9} T^{\alpha} \bar{T}^{\dot{\alpha}} \right) \right. \\ &\quad \left. + 2 \left(\frac{1}{6} S - \frac{n}{3}(3n+1) T^{\alpha} T_{\alpha} \right) \left(\frac{1}{6} \bar{S} - \frac{n}{3}(3n+1) \bar{T}_{\dot{\alpha}} \bar{T}^{\dot{\alpha}} \right) \right) \equiv \Delta_{L+\Lambda}^{(2)} \end{aligned} \quad (7.12)$$

where superfields and covariant derivatives in the r.h.s. are 16+16 nonminimal superfields. Of course since nothing has changed concerning the metric, the ordinary form of $\Delta_{\Lambda}^{(2)}$ is the same as the ordinary form of $\Delta_{\sigma}^{(2)}$, computed in section 2.

8 Conclusions

In this paper we have determined the possible trace anomalies in the 16+16 nonminimal supergravity as well as in the non minimal one. There are in all cases two independent nontrivial cocycles whose densities are given by the square Weyl tensor and by the Gauss-Bonnet density, respectively.

Concerning the Pontryagin density, it appears in the anomaly supermultiplets only in the form of chiral anomaly (Delbourgo-Salam anomaly), but never in the form of trace anomaly.

At this point we must clarify the question of whether the cocycles we have found in nonminimal and 16+16 nonminimal supergravities are the only ones. In this paper we have not done a systematic search of such nontrivial cocycles in the nonminimal and 16+16 nonminimal case, the reason being that when a dimensionless field like ψ and $\bar{\psi}$ are present in a theory a polynomial analysis is not sufficient (and a non-polynomial one is of course very complicated). But we can argue as follows: consider a nontrivial cocycle in nonminimal or 16+16 nonminimal supergravity; it can be mapped to a minimal cocycle which either vanishes or coincides with the ones classified in [8]. There is no other possibility because in minimal supergravity there are no dimensionless superfields (apart from the vielbein) and the polynomial analysis carried out in [8] is sufficient to identify all cocycles. We conclude that the nonminimal and 16+16 nonminimal nontrivial cocycles, which reduce in the ordinary form to a nonvanishing expression, correspond to $\Delta_{\sigma}^{(1)}$ and $\Delta_{\sigma}^{(2)}$ in minimal supergravity.

Finally we would like to make a comment on an aspect of our results that could raise at first sight some perplexity. Although one cannot claim the previous results to be a theorem, they nevertheless point in the direction of the non-existence of a supersymmetric anomaly multiplet that has, as its e.m. tensor trace component, the Pontryagin density. On the other hand we know systems with chiral fermions that at first sight can be supersymmetrized and coupled to supergravity. In such system we expect the trace of the e.m. tensor at one loop to contain the Pontryagin density, [33]; thus why couldn't we have an anomaly multiplet that contains as trace component the Pontryagin density? The point is that in such a chiral case there can exist an obstruction to that, as we try to explain next. Suppose that the e.m. tensor of a system like the one just mentioned, has, at one loop, an integrated nonvanishing trace $\Delta_\omega^{(P)}$, containing a term given by ω multiplied by the Pontryagin density. We cannot expect, in general this term to be supersymmetric. On the contrary, denoting by ϵ the supersymmetric local parameter we expect there to exist a partner cocycle $\Delta_\epsilon^{(P)}$ such that

$$\delta_\omega \Delta_\omega^{(P)} = 0, \quad \delta_\epsilon \Delta_\omega^{(P)} + \delta_\omega \Delta_\epsilon^{(P)} = 0, \quad \delta_\epsilon \Delta_\epsilon^{(P)} = 0 \quad (8.1)$$

The cocycle $\Delta_\epsilon^{(P)}$ to our best knowledge has not yet been computed in supergravity. So we have to rely on plausibility arguments. There are two possibilities: it might happen that $\Delta_\epsilon^{(P)}$ is trivial, i.e. $\Delta_\epsilon^{(P)} = \delta_\epsilon \mathcal{C}^{(P)}$, so that (8.1) implies that $\delta_\epsilon (\Delta_\omega^{(P)} - \delta_\omega \mathcal{C}^{(P)}) = 0$. The end result would be a supersymmetric Weyl cocycle. This is, for instance, what happens for the chiral ABJ anomaly in rigid supersymmetry, where the supersymmetric partner of the usual chiral anomaly must be trivial, [32], and, precisely as above, the chiral anomaly can be cast in supersymmetric form, see [34–36].

The second possibility is that no such counterterm $\mathcal{C}^{(P)}$ exists, in which case the cocycle $\Delta_\epsilon^{(P)}$ is nontrivial and there is no possibility to supersymmetrize $\Delta_\omega^{(P)}$. This seems to be the case for the chiral ABJ anomaly in the presence of local supersymmetry, [37]. And this may be the case also for $\Delta_\omega^{(P)}$, which would explain the origin of our inability to find a Weyl cocycle containing the Pontryagin form in the first position (trace anomaly) in terms of superfield.⁵ In both cases the origin of the obstruction is the same, i.e. the non-trivial breaking of local supersymmetry. In turn this would explain why a supersymmetry preserving regularization has never been found in such types of systems. Such converging arguments seem to nicely fit together and close the circle.

Acknowledgments

L.B. would like to thank Taichiro Kugo, Jorge Russo and Adam Schwimmer for discussions and suggestions. We would like to thank Friedemann Brandt and Yu Nakayama for a useful exchange of messages. We would like to thank the referee of this paper for valuable criticisms and remarks. The work of L.B. and S.G. was supported in part by the MIUR-PRIN contract 2009-KHZKRX.

⁵A plausible explanation for the difference between local and global supersymmetry is that the nontrivial part of $\Delta_\epsilon^{(P)}$ may be an integral of ϵ multiplied by (as it often happens) a total derivative; if ϵ is a (generic) local parameter $\Delta_\epsilon^{(P)}$ is a nonvanishing nontrivial cocycle, but it vanishes if ϵ is constant.

A Reduction formulae

In this appendix we collect from ([24]) the formulas that are needed to reduce superfield expressions to component form. The equations below are not complete, they contain only the terms essential to recover the ordinary parts of the expressions (that is only the parts that survive once all the fields except the metric are disregarded). The complete form can be found in ([24]), or in [23] for the minimal model. The first formula when evaluated at $\theta = \bar{\theta} = 0$, connects the Riemann curvature to specific superfield components and it is basic for reducing cocycles to ordinary form

$$\begin{aligned}
 & \sigma_{\alpha\dot{\alpha}}^a \sigma_{\beta\dot{\beta}}^b \sigma_{\gamma\dot{\gamma}}^c \sigma_{\delta\dot{\delta}}^d R_{cdba} \tag{A.1} \\
 & \approx 4\epsilon_{\gamma\delta}\epsilon_{\beta\alpha} \left[\frac{1}{4} \left(\bar{\mathcal{D}}_{\dot{\gamma}} \bar{W}_{\delta\dot{\beta}\dot{\alpha}} + \bar{\mathcal{D}}_{\dot{\delta}} \bar{W}_{\beta\dot{\alpha}\dot{\gamma}} + \bar{\mathcal{D}}_{\dot{\beta}} \bar{W}_{\dot{\alpha}\dot{\gamma}\dot{\delta}} + \bar{\mathcal{D}}_{\dot{\alpha}} \bar{W}_{\dot{\gamma}\dot{\delta}\dot{\beta}} \right) \right. \\
 & \quad - \frac{1}{8} \sum_{\dot{\gamma}\dot{\delta}} \sum_{\dot{\beta}\dot{\alpha}} \epsilon_{\dot{\gamma}\dot{\beta}} \sum_{\dot{\delta}\dot{\alpha}} \left[\bar{\mathcal{D}}_{\dot{\delta}} \mathcal{D}^\epsilon G_{\epsilon\dot{\alpha}} + \frac{1}{4} \bar{\mathcal{D}}_{\dot{\delta}} \mathcal{D}^\epsilon \left(\frac{1}{3} c_{\epsilon\dot{\alpha}} - i n d_{\epsilon\dot{\alpha}} \right) + \frac{i}{2} (n-1) \bar{\mathcal{D}}_{\dot{\delta}} \mathcal{D}_{\epsilon\dot{\alpha}} T^\epsilon \right. \\
 & \quad \left. \left. + n \mathcal{D}_{\epsilon\dot{\delta}} d^\epsilon_{\dot{\alpha}} + \frac{1}{6} \mathcal{D}_{\epsilon\dot{\delta}} c^\epsilon_{\dot{\alpha}} \right] + (\epsilon_{\dot{\gamma}\dot{\alpha}} \epsilon_{\dot{\beta}\dot{\delta}} + \epsilon_{\dot{\delta}\dot{\alpha}} \epsilon_{\dot{\beta}\dot{\gamma}}) \Lambda \right] \\
 & \quad - \frac{1}{2} \epsilon_{\gamma\delta} \epsilon_{\beta\dot{\alpha}} \sum_{\alpha\beta} \sum_{\dot{\gamma}\dot{\delta}} \left[i \mathcal{D}_{\beta\dot{\gamma}} \left(G_{\alpha\dot{\delta}} + \frac{1}{3} c_{\alpha\dot{\delta}} - i n d_{\alpha\dot{\delta}} \right) + \bar{\mathcal{D}}_{\dot{\gamma}} \mathcal{D}_\beta \left(G_{\alpha\dot{\delta}} + \frac{1}{3} c_{\alpha\dot{\delta}} \right) + \frac{1}{3} \bar{\mathcal{D}}_{\dot{\gamma}} \mathcal{D}_\beta c_{\alpha\dot{\delta}} \right] \\
 & \quad - \frac{1}{2} \epsilon_{\dot{\gamma}\dot{\delta}} \epsilon_{\beta\alpha} \sum_{\dot{\alpha}\dot{\beta}} \sum_{\gamma\delta} \left[i \mathcal{D}_{\gamma\dot{\beta}} \left(G_{\delta\dot{\alpha}} + \frac{1}{3} c_{\delta\dot{\alpha}} - i n d_{\delta\dot{\alpha}} \right) + \bar{\mathcal{D}}_{\dot{\beta}} \mathcal{D}_\gamma \left(G_{\delta\dot{\alpha}} + \frac{1}{3} c_{\delta\dot{\alpha}} \right) + \frac{1}{3} \bar{\mathcal{D}}_{\dot{\beta}} \mathcal{D}_\gamma c_{\delta\dot{\alpha}} \right] \\
 & \quad + 4\epsilon_{\dot{\gamma}\dot{\delta}} \epsilon_{\beta\dot{\alpha}} \left[-\frac{1}{4} \left(\mathcal{D}_\gamma W_{\delta\beta\alpha} + \mathcal{D}_\delta W_{\beta\alpha\gamma} + \mathcal{D}_\beta W_{\alpha\gamma\delta} + \mathcal{D}_\alpha W_{\gamma\delta\beta} \right) \right. \\
 & \quad \left. + \frac{1}{8} \sum_{\gamma\delta} \sum_{\beta\alpha} \epsilon_{\gamma\beta} \sum_{\dot{\delta}\dot{\alpha}} \left[\mathcal{D}_\delta \bar{\mathcal{D}}^{\dot{\epsilon}} G_{\alpha\dot{\epsilon}} + \frac{1}{4} \mathcal{D}_\delta \mathcal{D}^{\dot{\epsilon}} \left(\frac{1}{3} c_{\alpha\dot{\epsilon}} - i n d_{\alpha\dot{\epsilon}} \right) + \frac{i}{2} (n-1) \mathcal{D}_\delta \mathcal{D}_{\dot{\epsilon}\dot{\alpha}} \bar{T}^{\dot{\epsilon}} \right. \right. \\
 & \quad \left. \left. + n \mathcal{D}_{\delta\dot{\epsilon}} d^\epsilon_{\dot{\alpha}} + \frac{1}{6} \mathcal{D}_{\delta\dot{\epsilon}} c^\epsilon_{\dot{\alpha}} \right] + (\epsilon_{\gamma\alpha} \epsilon_{\beta\dot{\delta}} + \epsilon_{\delta\alpha} \epsilon_{\beta\dot{\gamma}}) \Lambda \right]
 \end{aligned}$$

where

$$\begin{aligned}
 \Lambda & \approx \frac{1}{24} \left(\mathcal{D}^\alpha \mathcal{D}_\alpha R + \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\mathcal{D}}^{\dot{\alpha}} R^+ \right) + \frac{1}{48} \left(\mathcal{D}^\alpha \bar{\mathcal{D}}^{\dot{\alpha}} - \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}^\alpha \right) G_{\alpha\dot{\alpha}} \\
 & \quad + \frac{1}{24} \left(-\frac{1}{12} \mathcal{D}^\alpha \bar{\mathcal{D}}^{\dot{\alpha}} c_{\alpha\dot{\alpha}} - \frac{i}{4} n \mathcal{D}^\alpha \bar{\mathcal{D}}^{\dot{\alpha}} d_{\alpha\dot{\alpha}} - \frac{i}{2} (n-1) \mathcal{D}^\alpha \mathcal{D}_{\alpha\dot{\alpha}} \bar{T}^{\dot{\alpha}} \right) \\
 & \quad + \frac{1}{24} \left(\frac{1}{12} \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}^\alpha c_{\alpha\dot{\alpha}} - \frac{i}{4} n \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}^\alpha d_{\alpha\dot{\alpha}} + \frac{i}{2} (n-1) \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}_{\alpha\dot{\alpha}} T^\alpha \right) - \frac{n}{16} \mathcal{D}^{\alpha\dot{\alpha}} d_{\alpha\dot{\alpha}} \tag{A.2}
 \end{aligned}$$

Other relations come from constraints among the various superfields

$$\begin{aligned}
 \mathcal{D}^\alpha \mathcal{D}_\alpha R - \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\mathcal{D}}^{\dot{\alpha}} R^+ & \approx -2i \mathcal{D}^{\alpha\dot{\alpha}} G_{\alpha\dot{\alpha}} \tag{A.3} \\
 & \quad + \left(\frac{1}{12} \mathcal{D}^\alpha \bar{\mathcal{D}}^{\dot{\alpha}} c_{\alpha\dot{\alpha}} + \frac{i}{4} n \mathcal{D}^\alpha \bar{\mathcal{D}}^{\dot{\alpha}} d_{\alpha\dot{\alpha}} + \frac{i}{2} (n-1) \mathcal{D}^\alpha \mathcal{D}_{\alpha\dot{\alpha}} \bar{T}^{\dot{\alpha}} \right) \\
 & \quad + \left(\frac{1}{12} \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}^\alpha c_{\alpha\dot{\alpha}} - \frac{i}{4} n \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}^\alpha d_{\alpha\dot{\alpha}} + \frac{i}{2} (n-1) \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}_{\alpha\dot{\alpha}} T^\alpha \right) + i \mathcal{D}^{\alpha\dot{\alpha}} c_{\alpha\dot{\alpha}}
 \end{aligned}$$

together with

$$\mathcal{D}_\alpha \bar{\mathcal{D}}_{\dot{\alpha}} R \approx \frac{i}{8} (n+1) \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\mathcal{D}}_{\dot{\beta}} d_{\alpha\dot{\beta}} - \frac{i}{4} (n-1) \bar{\mathcal{D}}_{\dot{\alpha}} \mathcal{D}_{\alpha\dot{\beta}} \bar{T}^{\dot{\beta}} \quad (\text{A.4})$$

$$\bar{\mathcal{D}}_{\dot{\alpha}} \mathcal{D}_\alpha R \approx \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\mathcal{D}}^{\dot{\beta}} G_{\alpha\dot{\beta}} - \frac{i}{8} n \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\mathcal{D}}^{\dot{\beta}} d_{\alpha\dot{\beta}} + \frac{5}{24} \bar{\mathcal{D}}_{\dot{\alpha}} \bar{\mathcal{D}}^{\dot{\beta}} c_{\alpha\dot{\beta}} - \frac{i}{4} (n-1) \bar{\mathcal{D}}_{\dot{\alpha}} \mathcal{D}_{\alpha\dot{\beta}} \bar{T}^{\dot{\beta}} \quad (\text{A.5})$$

$$\begin{aligned} \mathcal{D}^\gamma W_{\alpha\beta\gamma} \approx & \frac{1}{16} (\mathcal{D}_\alpha \bar{\mathcal{D}}^{\dot{\gamma}} G_{\beta\dot{\gamma}} + \mathcal{D}_\beta \bar{\mathcal{D}}^{\dot{\gamma}} G_{\alpha\dot{\gamma}}) - \frac{7}{144} (\mathcal{D}_\alpha \bar{\mathcal{D}}^{\dot{\gamma}} c_{\beta\dot{\gamma}} + \mathcal{D}_\beta \bar{\mathcal{D}}^{\dot{\gamma}} c_{\alpha\dot{\gamma}}) \\ & + \frac{in}{48} (\mathcal{D}_\alpha \bar{\mathcal{D}}^{\dot{\gamma}} d_{\beta\dot{\gamma}} + \mathcal{D}_\beta \bar{\mathcal{D}}^{\dot{\gamma}} d_{\alpha\dot{\gamma}}) \end{aligned} \quad (\text{A.6})$$

with the respective conjugate relations.

The last equation above, together with

$$\mathcal{D}_\alpha W_{\beta\gamma\delta} = \mathcal{D}_{(\alpha} W_{\beta\gamma\delta)} + \frac{1}{4} (\epsilon_{\alpha\beta} \mathcal{D}^\zeta W_{\gamma\delta\zeta} + \epsilon_{\alpha\gamma} \mathcal{D}^\zeta W_{\delta\beta\zeta} + \epsilon_{\alpha\delta} \mathcal{D}^\zeta W_{\beta\gamma\zeta}), \quad (\text{A.7})$$

allows us to conclude that

$$\mathcal{D}_\alpha W_{\beta\gamma\delta} \approx \mathcal{D}_{(\alpha} W_{\beta\gamma\delta)} \quad (\text{A.8})$$

Finally we quote in its exact form a constraint equation

$$\begin{aligned} \bar{\mathcal{D}}^\alpha G_{\alpha\dot{\alpha}} - \mathcal{D}_\alpha R = & \frac{1}{4} \left(n + \frac{1}{3} \right) \bar{\mathcal{D}} \bar{\mathcal{D}} T_\alpha - \frac{1}{4} \left(n - \frac{1}{3} \right) \bar{\mathcal{D}}^{\dot{\alpha}} \mathcal{D}_\alpha \bar{T}_{\dot{\alpha}} - \frac{1}{3} (n+1)(3n-1) \bar{T}^{\dot{\alpha}} \mathcal{D}_{\dot{\alpha}} T_\alpha \\ & - \frac{1}{4} (3n-5) \bar{T}^{\dot{\alpha}} G_{\alpha\dot{\alpha}} - \frac{1}{6} (6n^2 + 3n + 1) \bar{T}^{\dot{\alpha}} \mathcal{D}_\alpha \bar{T}_{\dot{\alpha}} + \frac{1}{6} (n-1) T_\alpha \bar{\mathcal{D}} \bar{T} \\ & - \frac{1}{3} (n-1) T_\alpha \bar{T} \bar{T} - \frac{i}{2} (n+1) \mathcal{D}_{\alpha\dot{\alpha}} \bar{T}^{\dot{\alpha}} \end{aligned} \quad (\text{A.9})$$

which, together with its conjugate, is needed for the cohomological analysis of cocycles.

References

- [1] Z. Komargodski and N. Seiberg, *Comments on Supercurrent Multiplets, Supersymmetric Field Theories and Supergravity*, *JHEP* **07** (2010) 017 [[arXiv:1002.2228](#)] [[INSPIRE](#)].
- [2] G. Girardi, R. Grimm, M. Muller and J. Wess, *Antisymmetric tensor gauge potential in curved superspace and a 16+16 supergravity multiplet*, *Phys. Lett. B* **147** (1984) 81 [[INSPIRE](#)].
- [3] W. Lang, J. Louis and B.A. Ovrut, *16+16 supergravity coupled to matter: the low-energy limit of the superstring*, *Phys. Lett. B* **158** (1985) 40 [[INSPIRE](#)].
- [4] Y. Nakayama, *CP-violating CFT and trace anomaly*, *Nucl. Phys. B* **859** (2012) 288 [[arXiv:1201.3428](#)] [[INSPIRE](#)].
- [5] J. Dowker and R. Critchley, *Stress-tensor conformal anomaly for scalar and spinor fields*, *Phys. Rev. D* **16** (1977) 3390 [[INSPIRE](#)].
- [6] S. Christensen and M. Duff, *Axial and conformal anomalies for arbitrary spin in gravity and supergravity*, *Phys. Lett. B* **76** (1978) 571 [[INSPIRE](#)].
- [7] Z. Komargodski and A. Schwimmer, *On Renormalization Group Flows in Four Dimensions*, *JHEP* **12** (2011) 099 [[arXiv:1107.3987](#)] [[INSPIRE](#)].

- [8] L. Bonora, P. Pasti and M. Tonin, *Cohomologies and anomalies in supersymmetric theories*, *Nucl. Phys. B* **252** (1985) 458 [[INSPIRE](#)].
- [9] F. Brandt, *Anomaly candidates and invariants of $D = 4$, $N = 1$ supergravity theories*, *Class. Quant. Grav.* **11** (1994) 849 [[hep-th/9306054](#)] [[INSPIRE](#)].
- [10] F. Brandt, *Local BRST cohomology in minimal $D = 4$, $N = 1$ supergravity*, *Annals Phys.* **259** (1997) 253 [[hep-th/9609192](#)] [[INSPIRE](#)].
- [11] K. Stelle and P.C. West, *Minimal Auxiliary Fields for Supergravity*, *Phys. Lett. B* **74** (1978) 330 [[INSPIRE](#)].
- [12] S. Ferrara and P. van Nieuwenhuizen, *The Auxiliary Fields of Supergravity*, *Phys. Lett. B* **74** (1978) 333 [[INSPIRE](#)].
- [13] V. Akulov, D. Volkov and V. Soroka, *On the General Covariant Theory of Calibrating Poles in Superspace*, *Theor. Math. Phys.* **31** (1977) 285 [[INSPIRE](#)].
- [14] M.F. Sohnius and P.C. West, *An Alternative Minimal Off-Shell Version of $N = 1$ Supergravity*, *Phys. Lett. B* **105** (1981) 353 [[INSPIRE](#)].
- [15] S. Ferrara, L. Girardello, T. Kugo and A. Van Proeyen, *Relation between different auxiliary field formulations of $N = 1$ supergravity coupled to matter*, *Nucl. Phys. B* **223** (1983) 191 [[INSPIRE](#)].
- [16] P. Breitenlohner, *Some invariant lagrangians for local supersymmetry*, *Nucl. Phys. B* **124** (1977) 500 [[INSPIRE](#)].
- [17] W. Siegel and S.J. Gates Jr., *Superfield supergravity*, *Nucl. Phys. B* **147** (1979) 77 [[INSPIRE](#)].
- [18] S.J. Gates Jr., M.T. Grisaru, M. Roček and W. Siegel, *Superspace or one thousand and one lessons in supersymmetry*, *Front. Phys.* **58** (1983) 1 [[hep-th/0108200](#)] [[INSPIRE](#)].
- [19] I. Buchbinder, S. Kuzenko and O. Solovev, *One loop counterterms of Wess-Zumino model in the $N = 1$ nonminimal supergravity background*, *Nucl. Phys. B* **322** (1989) 277 [[INSPIRE](#)].
- [20] D. Butter, *$N = 1$ Conformal Superspace in Four Dimensions*, *Annals Phys.* **325** (2010) 1026 [[arXiv:0906.4399](#)] [[INSPIRE](#)].
- [21] D. Butter and S.M. Kuzenko, *A dual formulation of supergravity-matter theories*, *Nucl. Phys. B* **854** (2012) 1 [[arXiv:1106.3038](#)] [[INSPIRE](#)].
- [22] I.L. Buchbinder and S.M. Kuzenko, *Ideas and Methods of Supersymmetry and Supergravity*, Taylor & Francis, New York U.S.A., London U.K. (1998).
- [23] J. Wess and J. Bagger, *Supersymmetry and supergravity*, Princeton University Press, Princeton U.S.A. (1992).
- [24] G. Girardi, R. Grimm, M. Müller and J. Wess, *Superspace geometry and the minimal, nonminimal, and new minimal supergravity multiplets*, *Z. Phys. C* **26** (1984) 123 [[INSPIRE](#)].
- [25] R. Grimm, M. Müller and J. Wess, *Supersymmetric gauge theories coupled to nonminimal and new minimal supergravity*, *Z. Phys. C* **26** (1984) 427 [[INSPIRE](#)].
- [26] M. Müller, *Supergravity in $U(1)$ superspace with a two form gauge potential*, *Nucl. Phys. B* **264** (1986) 292 [[INSPIRE](#)].
- [27] D. Anselmi, D. Freedman, M.T. Grisaru and A. Johansen, *Nonperturbative formulas for central functions of supersymmetric gauge theories*, *Nucl. Phys. B* **526** (1998) 543 [[hep-th/9708042](#)] [[INSPIRE](#)].

- [28] J. Gates, S.J., M.T. Grisaru and W. Siegel, *Auxiliary-field anomalies*, *Nucl. Phys. B* **203** (1982) 189 [INSPIRE].
- [29] S. Giaccari, Ph.D. Thesis, to appear.
- [30] S. Christensen and M. Duff, *New gravitational index theorems and supertheorems*, *Nucl. Phys. B* **154** (1979) 301 [INSPIRE].
- [31] P. Townsend and P. van Nieuwenhuizen, *Anomalies, topological invariants and the Gauss-Bonnet theorem in supergravity*, *Phys. Rev. D* **19** (1979) 3592 [INSPIRE].
- [32] O. Piguet and K. Sibold, *The anomaly in the slavnov identity for $N = 1$ supersymmetric Yang-Mills theories*, *Nucl. Phys. B* **247** (1984) 484 [INSPIRE].
- [33] L. Bonora and S. Giaccari, to appear.
- [34] G. Girardi, R. Grimm and R. Stora, *Chiral anomalies in $N = 1$ supersymmetric Yang-Mills theories*, *Phys. Lett. B* **156** (1985) 203 [INSPIRE].
- [35] L. Bonora, P. Pasti and M. Tonin, *ABJ anomalies in supersymmetric Yang-Mills theories*, *Phys. Lett. B* **156** (1985) 341 [INSPIRE].
- [36] L. Bonora, P. Pasti and M. Tonin, *The consistent chiral anomaly in supersymmetric Yang-Mills theories*, *Nucl. Phys. B* **261** (1985) 249 [Erratum *ibid.* **B 269** (1986) 745] [INSPIRE].
- [37] L. Bonora, P. Pasti and M. Tonin, *The chiral anomaly in supersymmetric gauge theories coupled to supergravity*, *Phys. Lett. B* **167** (1986) 191 [INSPIRE].