

SOIL PHYSICS

Relationship of Methane Consumption with the Respiration of Soil and Grass–Moss Layers in Forest Ecosystems of the Southern Taiga in Western Siberia

A. F. Sabrekov^{a–d}, M. V. Glagolev^{a–d}, I. A. Fastovets^a, B. A. Smolentsev^e,
D. V. Il'yasov^d, and Sh. Sh. Maksyutov^{b,f}

^a Faculty of Soil Science, Moscow State University, Moscow, 119991 Russia
e-mail: sabrekovaf@gmail.com, m_glagolev@mail.ru

^b Tomsk State University, ul. Lenina 36, Tomsk, 643050 Russia

^c Yugra State University, ul. Chekhova 16, Khanty-Mansiisk, 628012 Russia

^d Institute of Forest Science, Russian Academy of Sciences, ul. Sovetskaya 2, Uspenskoe, Moscow oblast, 143030 Russia,

^e Institute of Soil Science and Agrochemistry, Siberian Branch, Russian Academy of Sciences, pr. Ak. Lavrent'eva 8/2, Novosibirsk, 630099 Russia

^f National Institute for Environmental Studies, Nishi Odori str., Tsukuba, 305-8506 Japan

Received May 13, 2014

Abstract—The consumption of methane by some soils in the southern taiga of Western Siberia was studied by the static chamber method in the summer of 2013. The median of the specific CH₄ flux through the soil was -0.05 mg C/(m² h) for the entire set of measurements (the negative flux indicates the consumption of methane by the soil). A statistically significant ($R^2 = 0.81$) linear relationship has been found between the specific CH₄ flux to the soil and the total respiration of the soil and the grass–moss layers in the studied forest ecosystems. The quantitative theoretical explanation of this relationship is based on the plant-associated and free methanotrophy.

Keywords: rhizospheric methanotrophs, total soil and plant respiration, greenhouse gases, Fluvisol, Luvisol, Gleysol

DOI: 10.1134/S1064229315080062

INTRODUCTION

The accumulation of greenhouse gases in the atmosphere is one of the current global environmental problems [31]. The major contributions to the greenhouse effect are made by CO₂ and CH₄: 60 and 15%, respectively [43]. For understanding the natural changes, the inventory of sources and sinks, the assessment of their intensities and dynamics, and the investigation of their functioning mechanisms are necessary [31].

As for the study of specific CH₄ fluxes, the most numerous measurements were performed in Western Siberia, but the studies were somewhat unbalanced: only the emission of methane from natural bogs and lakes was studied [11, 29, 30, 42]. Single measurements of methane consumption, if any, were occasional: the emission of CH₄ was expected to be revealed in the methane-generating ecosystems, but its consumption was rarely observed [9, 12, 26]. In European Russia, both the emission of methane from the soils of forest, bogs, and areas under agricultural use, as well as their consumption, were studied [13, 15, 16]. The comparison of specific fluxes (in mass units of

the gas released from the soil surface in unit time) shows that the bogs as methane sources significantly exceed any soil sinks of this gas. However, the total bog area is significantly smaller than the area of CH₄ sinks (i.e., soils of fields, forests, etc.), so the latter cannot be ignored; on the contrary, they should be thoroughly studied to obtain an accurate balance. Earlier theoretical calculations using five different procedures showed that the soils of Russia consume 3.6 Mt of methane annually [2], while its emission is estimated by different authors at 7.5 to 23.5 Mt/year [7, 19, 46].

Regrettably, the assessment of methane consumption by the soil is a more difficult technical problem, because the CH₄ concentration should be measured in the range from the atmospheric to the null level in this case, while the measurements of methane emission include the analysis of concentrations exceeding the atmospheric level (frequently by several times). Therefore, along with the direct measurements of methane consumption, of special importance is the search of correlations between the CH₄ consumption rate and some parameters easier to measure.

Studies of oligotrophic bogs in Western Siberia revealed only a loose reliable correlation between the specific CH_4 flux and the emission of CO_2 due to the total respiration of the soil and the grass–moss layers [4]. In theoretical terms, a stronger correlation between these parameters could be expected in less wet soils. In addition, the significant role of CO_2 in the greenhouse effect attracts the attention of researchers to measurements of its specific fluxes [10, 14, 28, 39].

The general goal of recent studies is to close these gaps in studying the role of biogeocenoses in the cycle of methane as a greenhouse gas; therefore, this work was aimed at measuring the rate of methane consumption by forest soils in the southern taiga of Western Siberia and revealing (if possible) the quantitative relationship between the consumption of methane and the respiration of the soil and the grass–moss layers.

OBJECTS AND METHODS

The measurements were conducted on several key plots near the settlement of Plotnikovo in the Bakchar district of Tomsk oblast, the southern taiga subzone of Western Siberia, in the summer of 2013. The sampling sites were selected in order to reflect the natural diversity of forest communities in the area studied. The detailed description of the sampling sites is given in Table 1.

The specific CH_4 fluxes were measured by the static chamber method [13, 15, 32] described earlier [32]. A stainless steel support (37×37 or 40×40 cm² in cross-section and 15 cm in height) was inserted into the soil to a depth of 10–15 cm no earlier than 15 min before the measurements, in order for the disturbances of the methane concentration profile in the soil caused by the insertion of the support to be leveled. Then, a Plexiglas chamber (parallelepiped $30 \times 30 \times 40$ cm³ in size, without the lower face) was installed on the support, and the contact between the chamber and the support was sealed with a hydraulic lock (water was filled in the support grooves so that the contact between the lower part of the chamber and the support was immersed in water to a depth of at least 1 cm). A rubber cork, through which a metallic tube with a fitted rubber tube was passed, was then inserted into the hole in the upper face of the chamber. An SFM syringe for gas sampling was connected to the rubber tube. The sampling was performed in time moments $t_0 = 0$, t_1 , t_2 , and t_3 with equal intervals. The time of exposure ($t_3 - t_0$) was 60 min.

The syringes with the samples were kept in soil solution to prevent leakage of methane, which is poorly soluble in salt water. The gas concentration in the samples was measured on a modified KhPM-4 chromatograph (Khromatograf, Moscow, USSR) equipped with a gas line and a flame ionization detector from an LKhM-80 chromatograph (Khromatograf, Moscow, USSR). A steel column (1 m long, 2.5 mm in diameter) packed with Sovpol (80–

100 mesh) was used. The measurements were performed at 40°C; hydrogen was used as a carrier gas (flow rate 5 mL/min); the injector loop volume was 0.5 mL. The sample volume was 3 mL. Standard methane–air mixtures with CH_4 concentrations of 1.99, 5.00, and 9.84 ppmv (National Institute for Environmental Studies, Japan) were used for calibration; the standard concentration accuracy was ± 0.01 ppmv.

Air samples for the analysis of carbon dioxide concentrations were taken from light-proof chambers in an analogous way, simultaneously with the samples for methane determination, but the time intervals between each of the four sampling events were 3 min to prevent the excess increase in CO_2 concentration. Plants were not removed during the measurements. Carbon dioxide–air mixtures with CO_2 concentrations of 357 ± 5 and 708 ± 10 ppmv (VNIIMEM, St. Petersburg, Russia) were used for calibration.

The concentration of carbon dioxide was analyzed no later than several hours after sampling on a DX-6100 infrared gas analyzer (RMT Ltd., Moscow, Russia).

Soil and air temperatures were measured with Thermochron iButton DS1921G temperature loggers (Dallas Semiconductor, USA). Vegetation descriptions were performed according to [3]. Soil names are given in accordance with the current classification of Russian soils [6] and the international WRB soil classification [45].

Specific fluxes were calculated by the regression method in the time–concentration coordinates using linear regression for the emission of CO_2 and nonlinear regression for the consumption of CH_4 as described earlier [1]. The statistical processing of data (including the comparison of the means using the Wilcoxon test, as well as linear and nonlinear one-parameter regressions) was performed using interactive MATLAB environment (v. 7.0, MathWorks, USA).

RESULTS AND DISCUSSION

The results of individual measurements of specific CO_2 fluxes, as well as air and soil temperatures at the measurement moments, are given in Table 2. Analogous data for specific methane fluxes are given in Table 3. The positive fluxes denote the emission of methane and carbon dioxide to the atmosphere, and the negative fluxes correspond to their consumption. Some statistical characteristics of the distributions of the measured specific CH_4 and CO_2 fluxes and other integral parameters are given in Table 4. The median of specific CH_4 fluxes was -0.05 mg C/(m² h) for the entire set of measurements, which indicated the consumption of methane by the soil. These results well agree with the data obtained for other forest ecosystems in the zones of broad-leaved forests and southern taiga (Table 5).

A statistically significant correlation ($n = 8$, $R^2 = 0.81$) was found between the median of CH_4 specific

Table 1. Characterization of the objects of study

Measurement point (data)	Coordinates	Forest characterization	Dominant plant species*	Soil type (WRB)	Organic horizon depth, cm	Some soil properties (at 0–10 cm, without forest litter or moss)		
						pH _{water}	C _{org} , %	density, g/cm ³
T.Plo.For.1.25 (24.06.2013) T.Plo.For.1.26 (27.06.2013)	56.8617° N, 83.077° E, 56.8616° N, 83.0723° E	Overmoistened birch forest in the Iksa River floodplain	<i>Bet, Al, Cal</i>	alluvial clay-mucky gley soils (Gleyic Histic Fluvisols (Eutric, Siltic))	17	6.2	7.7	1.1
Plo.For.Iksa.R (18.07.2013 ÷ 19.08.2013)	56.9621° N, 83.0700° E	Strongly moistened southern-taiga spruce forest	<i>Pic, Car, Bet, Sor, Mat, Ox, Sp</i>	Gleyic soddy-podzolic soils with a second humus horizon (Albic Stagnic Luvisol (Cutanic, Siltic))	12	5.3	2.5	1.2
Plo.For.Razr (10.08.2013)	56.9116° N, 83.0527° E	Thick grass-green moss cedar-fir forest	<i>Ab, Pin, Pr, Car, Ox, Mai, Eq, Gal</i> various mosses	Gleyic soddy-podzolic soils with a second humus horizon (Albic Luvisol (Cutanic, Siltic))	7	5.5	2.9	1.3
Plo.For.Razr2 (12.08.2013) Plo.For.Razr2.1 (12.08.2013)	56.9091° N, 83.0547° E, 56.9090° N, 83.0550° E	Polydominant small grass-sedge-sphagnum-green moss fir-spruce-cedar forest	<i>Pin, Pic, Ab, Bet, Sor, Ox, Mai, Sp</i>	Mucky-humus gley soils (Eutric Histic Gleysol (Clayic, Humic))	24	5.7	6.7	1.2
Birch.For (16.08.2013)	56.8766° N, 82.9042° E	Secondary young (40-year-old) forest	<i>Bet, Cal, Eq</i>	Soddy-mucky gley soils (Eutric Gleysol (Clayic, Humic))	29	5.8	8.3	1.1

* (*Bet*) *Betula pendula*; (*Cal*) *Calamagrostis purpurea*; (*Ab*) *Abies sibirica*; (*Al*) *Alnus glutinosa*; (*Ox*) *Oxalis acetosella*; (*Mai*) *Maianthemum bifolium*; (*Eq*) *Equisetum sylvaticum*; (*Gal*) *Galium palustre*; (*Sp*) *Sphagnum*; (*Pic*) *Picea obovata*; (*Pin*) *Pinus sibirica*; (*Pr*) *Prunus padus*; (*Sor*) *Sorbus sibirica*; (*Car*) *Caragana arborescens*; (*Mat*) *Matteuccia struthiopteris*.

Table 2. Emission of CO₂ from taiga soils of Western Siberia, the Plotnikovo key plot

Temperature, °C					Specific CO ₂ flux, mg C-CO ₂ / (m ² h)		Temperature, °C					Specific CO ₂ flux, mg C-CO ₂ / (m ² h)	
air	soil, depth, cm				mean	S*	air	soil, depth, cm				mean	S*
	0	5	10	15				0	5	10	15		
T. Plo.For. 1.25, birch forest							22.0	21.5	16.1	15.0	14.0	489	34
18.7	16.6	13.4	11.8	11.8	50	8	22.0	21.5	16.2	15.0	14.0	417	21
18.7	16.6	13.4	11.8	11.8	29	7	21.5	21.2	16.5	15.0	14.0	447	28
12.2	13.2	12.7	11.5	11.9	77	14	21.3	21.0	16.4	15.0	14.0	641	62
12.2	13.2	12.7	11.5	11.9	104	6	21.5	21.0	16.5	15.0	14.0	448	12
12.2	13.2	12.7	11.5	11.9	90	10	21.5	21.4	16.5	15.0	14.0	419	29
T. Plo.For. 1.26 (measurement date 27.06.2013) birch forest, southern taiga							21.5	21.0	16.5	15.0	14.0	571	24
23.5	21.7	15.5	14.5	12.9	235	41	21.3	21.0	16.5	15.0	14.0	861	99
23.5	21.7	15.5	14.5	12.9	155	5	T. Birch.For birch forest						
T. Plo.For. Razr cedar–fir forest							18.5	18.7	15.5	14.5	14.0	196	24
24.1	20.2	18.5	18.0	18.0	1242	168	18.5	18.7	15.5	14.5	14.0	310	14
23.8	20.2	18.3	17.8	17.8	821	59	18.5	18.7	15.5	14.5	14.0	452	30
23.5	20.1	18.3	17.7	17.7	636	33	18.5	18.7	15.5	14.5	14.0	278	23
23.2	20.0	18.0	17.3	17.3	464	21	18.1	18.0	15.5	14.5	14.0	672	51
23.0	19.9	17.9	17.2	17.2	422	18	18.0	17.9	15.5	14.5	14.0	755	37
22.8	19.9	17.8	17.0	17.0	474	24	18.0	17.9	15.5	14.5	14.0	583	26
20.8	19.3	16.6	15.5	15.3	1145	37	17.1	16.9	15.5	14.5	14.0	384	69
20.5	19.2	16.5	15.3	15.0	1599	64	16.6	16.4	15.5	14.5	14.0	745	32
20.1	20.1	16.9	15.2	15.1	1311	83	16.5	16.1	15.5	14.5	14.0	546	26
22.4	19.7	17.6	16.7	16.7	448	12	14.9	15.3	15.5	14.5	14.0	1512	182
22.2	19.7	17.4	16.5	16.5	538	47	14.3	14.9	15.0	14.5	14.0	514	17
21.9	19.6	17.3	16.4	16.4	534	30	13.8	14.5	15.0	14.5	14.0	1409	174
21.6	19.5	17.1	16.1	16.1	1235	221	13.4	14.1	15.0	14.5	14.0	1200	115
21.4	19.4	17.0	15.9	15.9	968	44	12.5	13.5	15.0	14.5	14.0	569	13
21.1	19.4	16.9	15.8	15.7	961	255	12.1	13.3	15.0	14.5	14.0	906	10
20.0	19.1	16.0	15.0	14.5	751	67	12.1	13.3	15.0	14.5	14.0	833	50
19.7	19.0	16.0	14.7	14.2	739	31	11.5	13.0	15.0	14.5	14.0	675	146
19.5	18.9	16.0	14.5	14.0	681	58	18.1	18.0	15.5	14.5	14.0	877	48
T. Plo.For.Razr2 fir–spruce–cedar forest							18.1	18.0	15.5	14.5	14.0	681	47
21.0	21.5	16.6	14.5	14.3	525	41	18.1	18.0	15.5	14.5	14.0	457	17
21.0	21.5	16.1	14.5	14.0	585	72	18.0	17.9	15.5	14.5	14.0	678	35
21.0	21.1	16.0	14.5	14.0	292	40	18.0	17.9	15.5	14.5	14.0	576	57
22.3	21.9	16.0	14.5	14.0	411	8	16.5	16.5	15.5	14.5	14.0	455	15
23.4	22.4	16.0	14.5	14.0	384	20	16.5	16.3	15.5	14.5	14.0	366	23
22.6	22.5	16.0	14.8	14.0	383	27	16.1	16.0	15.5	14.5	14.0	735	75
22.5	22.5	16.0	15.0	14.0	344	9	15.6	15.9	15.5	14.5	14.0	920	19
23.1	22.9	16.0	15.0	14.0	639	81	15.2	15.5	15.5	14.5	14.0	1550	179
22.8	22.5	16.0	15.0	14.0	583	29	12.5	13.5	15.0	14.5	14.0	296	32
T. Plo.For.Razr2.1 ir–spruce–cedar forest							12.5	13.5	15.0	14.5	14.0	326	18
22.0	21.9	16.0	15.0	14.0	471	22	12.5	13.5	15.0	14.5	14.0	272	40

Table 2. (Contd.)

Temperature, °C					Specific CO ₂ flux, mg C–CO ₂ / (m ² h)		Temperature, °C					Specific CO ₂ flux, mg C–CO ₂ / (m ² h)	
air	soil, depth, cm				mean	S*	air	soil, depth, cm				mean	S*
	0	5	10	15				0	5	10	15		
12.1	13.3	15.0	14.5	14.0	649	31	18.3	18.8	17.6	15.6	14.4	579	35
12.1	13.3	15.0	14.5	14.0	681	25	18.0	18.3	17.0	15.0	14.0	343	24
11.5	13.0	15.0	14.5	14.0	789	70	18.0	18.0	17.0	15.0	14.0	475	8
11.5	13.0	15.0	14.5	14.0	896	62	18.0	18.0	17.0	15.0	14.0	229	17
11.5	13.0	15.0	14.5	14.0	658	78	19.0	18.5	16.5	14.7	14.0	416	17
T. Plo.For.Iksa.R (measurement date 18.07.2013) spruce forest							19.0	18.5	16.5	14.7	14.0	335	9
23.6	21.6	16.0	15.6	15.2	291	7	18.8	18.5	16.5	14.8	14.0	422	133
23.6	21.6	16.0	15.6	15.2	265	8	18.5	18.1	16.5	14.7	14.0	465	29
23.6	21.6	16.0	15.6	15.2	266	9	18.5	18.0	16.5	14.8	14.0	517	68
23.6	21.6	16.0	15.6	15.2	452	15	18.6	18.5	16.5	14.8	14.0	390	23
23.2	21.2	15.9	15.5	15.0	346	20	19.0	18.5	16.5	14.7	14.0	364	37
23.2	21.2	15.9	15.5	15.0	367	79	19.0	18.5	16.5	14.8	14.0	387	37
21.7	20.1	15.5	15.0	14.9	352	28	18.8	18.5	16.5	14.8	14.0	300	56
21.7	20.1	15.5	15.0	14.9	400	8	18.5	18.3	16.5	14.9	14.0	307	15
21.4	20.0	15.5	15.0	14.8	451	87	18.0	18.0	16.6	15.0	14.0	420	8
21.4	20.0	15.5	15.0	14.8	422	12	18.4	18.0	16.5	15.0	14.0	391	14
23.0	21.0	15.5	15.0	15.0	345	12	19.0	18.5	16.5	14.8	14.0	356	20
23.0	21.0	15.5	15.0	15.0	191	41	18.5	18.5	16.5	14.9	14.0	303	50
23.2	21.0	15.5	15.0	15.0	376	88	18.8	18.5	16.5	14.9	14.0	344	10
23.2	21.0	15.5	15.0	15.0	288	53	18.5	18.0	16.5	14.9	14.0	367	11
22.0	20.5	15.5	15.0	15.0	426	56	18.0	18.0	16.5	14.8	14.0	331	29
22.0	20.5	15.5	15.0	15.0	408	18	18.0	18.0	16.5	14.8	14.0	367	6
21.0	19.7	15.5	15.0	15.0	518	33	18.8	18.4	16.5	14.8	14.0	421	94
21.0	19.7	15.5	15.0	15.0	495	24	18.5	18.0	16.5	14.9	14.0	357	20
20.5	19.5	15.5	15.0	15.0	481	121	18.1	18.0	16.5	14.8	14.0	292	11
20.5	19.5	15.5	15.0	15.0	254	49	17.9	17.8	16.5	14.9	14.0	295	100
T. Plo.For.Iksa.R (measurement date 19.08.2013) spruce forest							17.5	17.5	16.5	14.7	14.0	213	40
18.4	18.9	17.8	15.8	14.6	335	106	17.4	17.5	16.5	14.5	14.0	281	54

* (S) standard deviation.

flux (SF, mg C–CH₄/(m² h)) from the soil in each of the studied points and the median of the total respiration of soil and grass–moss layers (TR, mg C–CO₂/(m² h)):

$$SF \text{ CH}_4 = a \text{ TR} + b, \quad (1)$$

where $a = -(0.000157 \pm 0.000031)$ mg C–CH₄/mg C–CO₂; $b = 0.015 \pm 0.014$ mg C–CH₄/(m² h); and the standard error of the regression is 0.018 mg C–CH₄/(m² h) (figure). It can be seen that when the

respiration rate of plants increases, the specific methane consumption rate also increases.

This relationship can be explained from the positions of two types of methanotrophy related to the activity of free-living methanotrophs [24, 34, 35] and plant-associated methanotrophs [23, 33, 38]. It should be noted that 33% of the total methane consumed by microorganisms is assimilated, and 60% is oxidized to carbon dioxide [8].

Methanotrophic bacteria are known to be permanently present in the phyllosphere and the rhizo-

Table 3. Methane emission from taiga forests of Western Siberia, the Plotnikovo key plot

		Temperature, °C				CH ₄ flux, mg C-CH ₄ /(m ² h)	
air	soil, depth, cm				mean	S	
	0	5	10	15			
T. Plo.For.1.25, birch forest							
18.7	16.6	13.4	11.8	11.8	-0.020	0.057	
18.7	16.6	13.4	11.8	11.8	0.013	0.032	
12.2	13.2	12.7	11.5	11.9	0.050	0.021	
12.2	13.2	12.7	11.5	11.9	0.006	0.078	
12.2	13.2	12.7	11.5	11.9	-0.012	0.469	
12.2	13.2	12.7	11.5	11.9	-0.029	0.154	
T. T.Plo.For.1.26 birch forest, southern taiga							
23.8	21.3	14.5	13.7	12.3	-0.040	0.113	
23.8	21.3	14.5	13.7	12.3	-0.016	0.139	
T. Plo.For.Razr cedar-fir forest							
23.5	20.1	18.2	17.6	17.6	-0.064	0.277	
23.5	20.1	18.2	17.6	17.6	-0.151	0.045	
21.9	19.6	17.3	16.4	16.4	-0.098	0.362	
21.9	19.6	17.3	16.4	16.4	-0.085	0.086	
20.4	19.2	16.4	15.2	14.9	-0.010	0.149	
20.4	19.2	16.4	15.2	14.9	-0.149	0.101	
Plo.For.Razr2 (on the right) and Razr2.1(on the left), fir-spruce-cedar forest							
21.4	21.5	16.1	14.5	14.0	-0.007	0.136	
21.4	21.5	16.1	14.5	14.0	-0.055	0.094	
22.7	22.5	16.0	14.9	14.0	-0.045	0.268	
22.7	22.5	16.0	14.9	14.0	-0.027	0.058	
21.8	21.5	16.2	15.0	14.0	-0.052	0.290	
21.8	21.5	16.2	15.0	14.0	-0.090	0.225	
T. Birch.For birch forest							
16.8	16.6	15.5	14.5	14.0	-0.113	1.362	
16.8	16.6	15.5	14.5	14.0	-0.129	0.228	
16.8	16.6	15.5	14.5	14.0	-0.122	0.035	
15.6	15.8	15.5	14.5	14.0	-0.120	0.211	
15.6	15.8	15.5	14.5	14.0	-0.095	0.097	
15.6	15.8	15.5	14.5	14.0	-0.065	0.060	
T. Plo.For.Iksa.R (measurement date 18.07.2013) spruce forest							
28.2	19.4	16.9	16.6	—	-0.54	0.30	
28.7	20.8	19.0	17.5	—	0.07	0.09	
31.7	23.6	20.4	19.7	—	0.16	0.07	
30.7	22.5	20.0	19.5	—	-0.01	0.32	
27.3	22.0	20.0	19.5	—	-0.16	1.01	
—	22.0	21.0	19.5	—	-0.02	0.05	
T. Plo.For.Iksa.R (measurement date 19.08.2013) spruce forest							
18.1	18.3	17.0	15.2	14.1	-0.037	0.261	
18.3	18.2	16.6	14.9	14.0	0.000	0.200	
18.7	18.3	16.5	14.8	14.0	-0.022	0.103	
17.3	17.4	16.5	14.6	14.0	-0.264	0.338	
17.3	17.4	16.5	14.6	14.0	-0.021	0.076	
18.8	18.4	16.5	14.8	14.0	-0.070	0.307	
18.0	17.8	16.5	14.8	14.0	-0.078	0.073	

(Dash) Parameter was not measured.

Table 4. Some statistical parameters of the distributions of the measured specific CH₄ and CO₂ fluxes (mg C/(m² h)) and other integral parameters

Measurement point	Median of specific fluxes (number of measurements)		Variability (for the set of specific CO ₂ fluxes)*	Mean temperature, °C				
	CO ₂	CH ₄		air	soil, depth, cm			
					0	5	10	15
T.Plo.For.1.25	76 (5)	−0.00 (6)	0.26	14.8	14.5	13.0	11.6	11.9
T.Plo.For.1.26	195 (2)	−0.03 (2)	0.10	23.5	21.7	15.5	14.5	12.9
Plo.For.Iksa.R								
18.07.2013	371 (20)	−0.02 (6)	0.19	22.3	20.6	15.6	15.2	15.0
19.08.2013	361 (30)	−0.04 (7)	0.15	18.4	18.2	16.6	14.9	14.0
Plo.For.Razr	745 (18)	−0.09 (6)	0.38	21.8	19.6	17.2	16.3	16.3
Plo.For.Razr2	410 (9)	−0.05 (3)	0.24	22.2	22.1	16.1	14.7	14.0
Plo.For.Razr2.1	470 (9)	−0.05 (3)	0.13	21.6	21.3	16.4	15.0	14.0
Birch.For	664 (36)	−0.12 (6)	0.26	15.2	15.7	15.3	14.5	14.0

* Calculated as the ratio of the difference between the third and first quartiles divided by 2 to the median of the set of specific CO₂ fluxes.


Table 5. Consumption of methane by soils in different forest ecosystems

Ecosystem	Measurement point coordinates	Specific methane fluxes, mg C–CH ₄ /(m ² h)	Source
Mixed forest on gray forest soils, broad-leaved forest zone	54.81° N, 37.59° E	−0.05...−0.06 (May–October, mean monthly values)	[15]
Mixed forest on gray forest soils with different degrees of erosion, broad-leaved forest zone	54.82° N, 37.58° E	−0.004...−0.04 (June–September, individual values)	[14]
Birch forest on tundra soils, forest-tundra zone	54.72° N, 66.70° E	−0.01...−0.07 (June, mean monthly values)	[18]
Mixed pine–oak forest on gray forest soils, broad-leaved forest zone	43.94° N, 69.57° E	−0.11 (June, mean monthly value)	[18]
Pine forest on podzolic soils, taiga zone	42.50° N, 72.17° E	−0.10 (April–December, mean for 6 years)	[21]
Pine forest on podzolic soils, taiga zone	42.50° N, 72.17° E	−0.12 (April–December, mean for 6 years)	[21]
Beech–oak forest on mountain-meadow soils, broad-leaved forest zone	51.00° N, 9.85° E	−0.12 (April), −0.11 (June)	[36]
Beech forest on podzolic soils, broad-leaved forest zone	51.57° N, 10.17° E	−0.05 (April), −0.04 (June)	[36]

sphere. The neutrophilic mesophilic methanotrophs are associated with not only aquatic plants (hydrophytes), but also mesophilic terrestrial woody and herbaceous plants (in particular, the formation of stable associations of methanotrophs with some grasses and other plants was demonstrated in vitro) [5]. The consumption of methane by rhizospheric bacteria may be very significant [23, 33, 38].

The known relationship between the specific consumption (and not specific flux) of methane by plant roots (V , mg C–CH₄/g dry root biomass per hour), or, more precisely, by rhizospheric bacteria, and the concentration of methane in soil pores (C , mg C–CH₄/m³) is described by the Michaelis–Menten equation:

$$V_r = V_{r, \max} C / (K_{r, m} + C), \quad (2)$$



Relationship between the consumption of methane and the total respiration (TR) of the soil and the grass–moss layers: (1) medians of experimental data for each point; (2) linear regression on these data for specific fluxes (SFs) of $\text{CH}_4 = -(0.000157 \pm 0.000031) \text{TR} + (0.015 \pm 0.014)$, $R^2 = 0.81$; (3) F_{ma} values calculated from the medians of the measured values of total soil respiration for each point; (4) sum of the F_{ma} values calculated from the medians of the measured values of total soil respiration for each point and the calculated F_s value.

where $V_{r, \text{max}}$ is the maximum specific consumption rate by plant roots ($\text{mg C-CH}_4/\text{g dry root biomass per hour}$), and $K_{r, m}$ is the Michaelis constant equal to the methane concentration at which the methane consumption rate is half of the maximum value ($\text{mg C-CH}_4/\text{m}^3$). Under the assumption that the oxidation of methane in the soil mainly occurs in the upper 15-cm thick layer [17, 36, 37], the mean C value in this layer of analogous soils is $0.65 \pm 0.07 \text{ mg C-CH}_4/\text{m}^3$ [36, 37]. Highly variable values of $V_{r, \text{max}}$ and $K_{r, m}$ are reported by different authors; therefore, they should be averaged to avoid gross errors. The mean value of $K_{r, m}$ is $59.2 \pm 12.6 \text{ mg C-CH}_4/\text{m}^3$ [27, 33], and the mean value of $V_{r, \text{max}}$ is $0.023 \pm 0.012 \text{ mg C-CH}_4/\text{g dry root biomass per hour}$.

The respiration rate of living plant biomass I_{lb} varies among the different species growing under temperate climatic conditions in the range from 0.5 to 2 $\text{mg C-CH}_4/\text{g dry root biomass per hour}$ at 20°C [22]. From other sources, for umbraticolous sciophytes under forest canopy, this value varies from 0.4 to 1.1 mg C at the same temperature [41]. Therefore, we take $I_{\text{lb}} = 1 \pm 0.3 \text{ mg C-CO}_2/\text{g dry root biomass per hour}$ (our measurements were also performed at temperatures close to 20°C). Thus, the living aboveground biomass of the grass–moss layer B_1 ($\text{g dry organic matter}/\text{m}^2$) can be estimated as follows:

$$B_1 = (\text{TR} - \text{RSR})/I_{\text{lb}}, \quad (3)$$

where RSR is the respiration of soil and roots, $\text{mg C-CO}_2/(\text{m}^2 \text{ h})$. The mean RSR value is about $160 \pm$

$40 \text{ mg C-CO}_2/(\text{m}^2 \text{ h})$ for mixed forests on podzolic soils in July–August [25, 39, 40]. The living aboveground/root biomass ratio for the grass–moss layer (p) varies in the range of 0.50–0.65 [26, 44]; hence, its mean value is 0.57 ± 0.07 . Thus, the root biomass B_r ($\text{g dry organic matter}/\text{m}^2$) can be expressed as follows:

$$B_r = p(\text{TR} - \text{RSR})/I_{\text{lb}}. \quad (4)$$

Then, the simulated value of CH_4 consumption due to the root-associated methanotrophy (T_m , $\text{mg C-CH}_4/(\text{m}^2 \text{ h})$) can be expressed as follows:

$$F_{\text{ma}} = p(\text{TR} - \text{RSR})V_{r, \text{max}}C/((K_{r, m} + C)I_{\text{lb}}). \quad (5)$$

The calculated F_{ma} values (average error was $0.025 \text{ mg C-CH}_4/(\text{m}^2 \text{ h})$) were found to be close to the measured values of specific CH_4 fluxes, and the slope of the linear relationship between the F_{ma} values and the total soil respiration was -0.000202 , which well agreed with the slope of the relationship between the specific CH_4 fluxes and the total soil respiration calculated from experimental data using linear regression: $-(0.000157 \pm 0.000031)$.

The activity of free-living methanotrophs can be estimated from the data on the consumption of methane by the soil free from plants. In this case, the Michaelis–Menten equation will be as follows:

$$V_s = V_{s, \text{max}}C/(K_{s, m} + C), \quad (6)$$

where $V_{s, \text{max}}$ is the maximum rate of specific consumption by the soil ($\mu\text{g C-CH}_4/\text{kg dry soil per hour}$), and $K_{s, m}$ is the Michaelis semisaturation constant for soils ($\text{mg C-CH}_4/\text{m}^3$). According to literature data

[20, 35], $V_{s, \max} = 1.62 \pm 4.26$ and $K_{s, m} = 8.64 \pm 27.24$ (median \pm standard deviation for the soils analogous to those under study). Then, the consumption of methane by the free-living methanotrophs can be calculated from its consumption by the soil (F_s , mg C–CH₄/(m² h)) using the following equation:

$$F_s = dhV_s, \quad (7)$$

where d is the soil density, kg/m³ (for the studied soils, the typical value is 1200), and h is the thickness of the soil layer in which an intensive oxidation of methane by free-living methanotrophs occurs, m. As was noted above, h can be taken equal to 0.15 ± 0.03 m. Then, F_s is 0.022 ± 0.080 mg C–CH₄/(m² h).

Thus, the theoretically calculated sum of plant-associated and free methanotrophies well agrees with the experimentally measured values, although it slightly exceeds them (figure). However, the performed calculations have some vulnerabilities. At the calculation of plant-related methanotrophy, the parameters $V_{r, \max}$ and $K_{r, m}$ for the Michaelis–Menten equation were found in the literature only for the bog and waterlogged ecosystems; therefore, their use for forest ecosystems bears some risks. The calculation of free methanotrophy involves the high variability of parameters $V_{s, \max}$ and $K_{s, m}$, which can also make the consumption prediction unreliable. Thus, the presented theoretical concept needs further support by data.

The effect of other factors like moisture, soil pH and temperature [21], and mineral nitrogen inhibiting the oxidation of methane [8] must also not be ignored; however, no statistically significant correlation between the soil temperature at any depth and the specific CH₄ fluxes was found in the current work.

CONCLUSIONS

(1) The median of specific CH₄ fluxes to the soil was -0.05 mg C/(m² h) for the entire set of measurements (the negative flux indicates the consumption of methane by the soil), which almost coincided with the values recorded in other forest ecosystems of the boreal zone.

(2) A statistically significant ($R^2 = 0.81$) linear relationship was revealed between the specific CH₄ fluxes to the soil and the total respiration of the grass–moss layer in forest ecosystems.

(3) The methane consumption by the soil with consideration for root-related methanotrophy was calculated from available literature data; from the obtained results, the experimental values of specific CH₄ fluxes were predicted, and the relationship between the consumption of methane by the soil and the total respiration of the grass–moss layer was determined. The consideration of free methanotrophy not related to plants resulted in an even better agreement with the experimental data, which confirmed the importance of both sinks for atmospheric methane.

ACKNOWLEDGMENTS

We thank E.V. Shein (Faculty of Soil Science, Moscow State University) for valuable remarks on the paper. Sabrekov thanks M.V. Semenov (Faculty of Soil Science, Moscow State University) for valuable advices and assistance in paper writing.

This work was performed within the Program for increasing the competitiveness of the Tomsk State University

REFERENCES

1. M. V. Glagolev, A. F. Sabrekov, and V. S. Kazantsev, *Physical Chemistry and Biology of Peat. Methods to Measure Gas Exchange at the Soil–Atmosphere Interface* (Tomsk State Pedagogical University, Tomsk, 2010) [in Russian].
2. M. V. Glagolev and I. V. Filippov, Inventory of methane consumption by soil,” *Dyn. Okruzh. Sredy Global’nye Izmen. Klim.* **2** (2/4), EDCCrev0002 (2011). <http://elibrary.ru/item.asp?id=18248134>. Cited May 5, 2014.
3. E. A. Golovatskaya and E. V. Porokhina, *Botany with the Principles of Phytocenology. Biological Productivity of Biogeocenoses* (Tomsk State Pedagogical University, Tomsk, 2005) [in Russian].
4. T. T. Efremova, N. M. Bazhin, I. M. Gadzhiev, S. P. Efremov, and G. A. Makhov, “Specific features of methanogenesis on oligotrophic bogs of Western Siberia and analysis of environmental factors related to correct extrapolation of CH₄ fluxes over larger territories,” *Sib. Ekol. Zh.*, No. 6, 563–570 (1998).
5. E. G. Ivanova, N. V. Doronina, and Yu. A. Trotsenko, “Aerobic methanotrophs as the plant symbionts,” *Tr. Inst. Mikrobiol. im. S.N. Vinogradskogo*, No. 13, 263–284 (2006).
6. *Classification and Diagnostic System of Russian Soils* (Oikumena, Smolensk, 2004) [in Russian].
7. K. Ya. Kondrat’ev and V. F. Krapivin, *Modeling of Global Carbon Cycle* (Fizmatlit, Moscow, 2004) [in Russian].
8. I. K. Kravchenko, S. A. Bykova, L. E. Dulov, V. F. Gal’chenko, V. M. Semenov, T. V. Kuznetsova, D. Pardini, M. Gispert, P. Boeckx, and O. van Cleemput, “Physicochemical and biological factors affecting atmospheric methane oxidation in gray forest soils,” *Microbiology (Moscow)* **74** (2), 216–220 (2005).
9. L. A. Krivenko, M. V. Glagolev, I. A. Fastovets, B. A. Smolentsov, and Sh. Sh. Maksyukov, “Specific methane flux from southern tundra in Western Siberia,” *Din. Okruzh. Sredy Global’nye Izmen. Klim.* **4** (2/8), (2013).
10. V. N. Kudayarov, “The role of soils in the carbon cycle,” *Eurasian Soil Sci.* **38** (8), 808–815 (2005).
11. A. V. Naumov, “Mires as a source of greenhouse gases in Western Siberia,” in *Second International Conference “Emission and Flow of Greenhouse Gases in Northern Eurasia”* (Pushchino, 2003), pp. 86–87.

12. A. V. Naumov, "Carbon dioxide and methane in soils and atmosphere of the mire systems of Western Siberia," *Sib. Ekol. Zh.*, No. 3, 313–318 (2002).
13. V. V. Novikov and A. V. Rusakov, "Release and absorption of greenhouse gases in ameliorated peat soils of the Rostov Lowland," *Eurasian Soil Sci.* **38** (7), 745–751 (2005).
14. V. V. Novikov, A. L. Stepanov, A. I. Pozdnyakov, and E. V. Lebedeva, "Seasonal dynamics of CO₂, CH₄, N₂O, and NO emissions from peat soils of the Yakhroma River floodplain," *Eurasian Soil Sci.* **37** (7), 755–761 (2004).
15. M. V. Semenov, I. K. Kravchenko, V. M. Semenov, T. V. Kuznetsova, L. E. Dulov, S. N. Udaltsov and A. L. Stepanov, "Carbon dioxide, methane, and nitrous oxide fluxes in soil catena across the right bank of the Oka River (Moscow oblast)," *Eurasian Soil Sci.* **43** (5), 541–549 (2010).
16. V. M. Semenov, I. K. Kravchenko, T. V. Kuznetsova, N. A. Semenova, S. A. Bykova, L. E. Dulov, V. F. Gal'chenko, G. Pardini, M. Gispert, P. Boeckx, and O. van Cleemput, "Seasonal dynamics of atmospheric methane oxidation in gray forest soils," *Microbiology (Moscow)* **73** (3), 356–362 (2004).
17. A. A. Sirin, G. G. Suvorov, M. V. Chistotin, and M. V. Glagolev, "Values of methane emission from drainage ditches," *Dyn. Okruzh. Sredy Global'nye Izmen. Klim.* **3** (2/6) (2012). http://www.ugrasu.ru/uploads/files/EDCC_3_2_Sirin.pdf. Cited November 23, 2013.
18. A. P. S. Adamsen and G. M. King, "Methane consumption in temperate and subarctic forest soils: rates, vertical zonation, and responses to water and nitrogen," *Appl. Environ. Microbiol.* **59** (2), 485–490 (1993).
19. N. G. Andronova and I. L. Karol, "The contribution of USSR sources to global methane emission," *Chemosphere* **26**, 111–126 (1993). doi: 10.1016/0045-6535(93)90416-3
20. M. Bender and R. Conrad, "Kinetics of CH₄ oxidation in toxic soils exposed to ambient air or high CH₄ mixing ratios," *FEMS Microbiol. Lett.* **101** (4), 261–269 (1992). doi: 10.1111/j.1574-6968.1992.tb05783.x
21. M. S. Castro, P. A. Steudler, J. M. Melillo, J. D. Aber, and R. D. Bowden, "Factors controlling atmospheric methane consumption by temperate forest soils," *Global Biogeochem. Cycles* **9** (1), 1–10 (1995). doi: 10.1029/94GB02651
22. D. E. Collier, "No difference in leaf respiration rates among temperate, subarctic, and arctic species grown under controlled conditions," *Can. J. Bot.* **74** (2), 317–320 (1996). doi: 10.1139/b96-039
23. R. Conrad, "Microbial ecology of methanogens and methanotrophs," *Adv. Agron.* **96**, 1–63 (2007). doi: 10.1016/S0065-2113(07)96005-8
24. R. Conrad, "Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS, N₂O, and NO)," *Microbiol. Rev.* **60** (4), 609–640 (1996).
25. P. M. Crill, "Seasonal patterns of methane uptake and carbon dioxide release by a temperate woodland soil," *Global Biogeochem. Cycles* **5** (4), 319–334 (1991). doi: 10.1029/91GB02466.
26. E. Garnier, "Growth analysis of congeneric annual and perennial grass species," *J. Ecol.* **80** (4), 665–675 (1992).
27. G. Gerard and J. Chanton, "Quantification of methane oxidation in the rhizosphere of emergent aquatic macrophytes: defining upper limits," *Biogeochemistry* **23** (2), 79–97 (1993). doi: 10.1007/BF00000444
28. M. V. Glagolev, E. A. Golovatskaya, and N. A. Shnyrev, "Greenhouse gas emission in West Siberia," *Contemp. Probl. Ecol.* **1** (1), 136–146 (2008). doi: 10.1134/S0001433811020113
29. M. Glagolev, I. Kleptsova, I. Filippov, S. Maksyutov, and T. Machida, "Regional methane emission from West Siberia mire landscapes," *Environ. Res. Lett.* **6**, 045214 (2011). doi: 10.1088/1748-9326/6/4/045214
30. M. V. Glagolev, A. F. Sabrekov, I. E. Kleptsova, I. V. Filippov, E. D. Lapshina, and T. Machida, and Sh. Sh. Maksyutov, "Methane emission from bogs in the subtaiga of Western Siberia: the development of standard model," *Eurasian Soil Sci.* **45** (10), 947–957 (2012). doi: 10.1134/S106422931210002X
31. M. Heimann, "Atmospheric science: enigma of the recent methane budget," *Nature* **476** (7359), 157–158 (2011). doi: 10.1038/476157a
32. G. L. Hutchinson and A. R. Mosier, "Improved soil cover method for field measurement of nitrous-oxide fluxes," *Soil Sci. Soc. Am. J.* **45**, 311–316 (1981).
33. G. M. King, "Associations of methanotrophs with the roots and rhizomes of aquatic vegetation," *Appl. Environ. Microbiol.* **60** (9), 3220–3227 (1994).
34. C. Knief and P. F. Dunfield, "Response and adaptation of different methanotrophic bacteria to low methane mixing ratios," *Environ. Microbiol.* **7** (9), 1307–1317 (2005). doi: 10.1111/j.1462-2920.2005.00814.x
35. C. Knief, A. Lipski, and P. F. Dunfield, "Diversity and activity of methanotrophic bacteria in different upland soils," *Appl. Environ. Microbiol.* **69** (11), 6703–6714 (2003). doi: 10.1128/AEM.69.11.6703-6714.2003
36. S. Kolb, C. Knief, P. F. Dunfield, and R. Conrad, "Abundance and activity of uncultured methanotrophic bacteria involved in the consumption of atmospheric methane in two forest soils," *Environ. Microbiol.* **7** (8), 1150–1161 (2005). doi: 10.1111/j.1462-2920.2005.00791.x
37. M. Koschorreck and R. Conrad, "Oxidation of atmospheric methane in soil: measurements in the field, in soil cores and in soil samples," *Global Biogeochem. Cycles* **7** (1), 109–121 (1993). doi: 10.1029/92GB02814
38. M. Krüger, P. Frenzel, and R. Conrad, "Microbial processes influencing methane emission from rice fields," *Global Change Biol.* **7** (1), 49–63 (2001). doi: 10.1046/j.1365-2486.2001.00395.x
39. I. Kurganova, V. Lopes de Gerenyu, L. Rozanova, D. Sapronov, T. Myakshina, and V. Kudeyarov, "Annual and seasonal CO₂ fluxes from Russian southern taiga soils," *Tellus B* **55** (2), 338–344 (2003). doi: 10.1034/j.1600-0889.2003.00047.x
40. M. B. Rayment and P. G. Jarvis, "An improved open chamber system for measuring soil CO₂ effluxes in the field," *J. Geophys. Res.: Atmos.* **102** (24), 28779–28784 (1997).

41. P. B. Reich, M. B. Walters, M. G. Tjoelker, D. Vanderklein, and C. Buschena, "Photosynthesis and respiration rates depend on leaf and root morphology and nitrogen concentration in nine boreal tree species differing in relative growth rate," *Funct. Ecol.* **12** (3), 395–405 (1998). doi: 10.1046/j.1365-2435.1998.00209.x
42. M. E. Repo, J. T. Huttunen, A. V. Naumov, A. V. Chichulin, E. D. Lapshina, W. Bleuten, and P. J. Martikainen, "Release of CO₂ and CH₄ from small wetland lakes in western Siberia," *Tellus B* **59**, 788–796 (2007). doi: 10.1111/j.1600-0889.2007.00301
43. H. Rodhe, "A comparison of the contribution of various gases to the greenhouse effect," *Science* **248**, 1217–1219 (1990). doi: 10.1126/science.248.4960.1217
44. B. Shipley and D. Meziane, "The balanced growth hypothesis and the allometry of leaf and root biomass allocation," *Funct. Ecol.* **16** (3), 326–331 (2002).
45. World Reference Base for Soil Resources 2014, International soil classification system for naming soils and creating legends for soil maps," in *IUSS Working Group WRB. World Soil Resources Reports No. 106* (Food and Agriculture Organisation, Rome, 2014).
46. X. Zhu, Q. Zhuang, Z. Qin, M. Glagolev, and L. Song, "Estimating wetland methane emissions from the northern high latitudes from 1990 to 2009 using artificial neural networks," *Gl. Biogeochem. Cycles* **27** (2), 592–604 (2013). doi: 10.1002/gbc.20052

Translated by K. Pankratova