

# Reconstructing metabolic pathways of hydrocarbon-degrading bacteria from the Deepwater Horizon oil spill

Nina Dombrowski<sup>1</sup>, John A. Donaho<sup>1</sup>, Tony Gutierrez<sup>2</sup>, Kiley W. Seitz<sup>1</sup>, Andreas P. Teske<sup>3</sup> and Brett J. Baker<sup>1\*</sup>

**1 The Deepwater Horizon blowout in the Gulf of Mexico in 2010, one of the largest marine oil spills<sup>1</sup>, changed bacterial communities in the water column and sediment as they responded to complex hydrocarbon mixtures<sup>2-4</sup>. Shifts in community composition have been correlated to the microbial degradation and use of hydrocarbons<sup>2,5,6</sup>, but the full genetic potential and taxon-specific metabolisms of bacterial hydrocarbon degraders remain unresolved. Here, we have reconstructed draft genomes of marine bacteria enriched from sea surface and deep plume waters of the spill that assimilate alkane and polycyclic aromatic hydrocarbons during stable-isotope probing experiments, and we identify genes of hydrocarbon degradation pathways. Alkane degradation genes were ubiquitous in the assembled genomes. *Marinobacter* was enriched with *n*-hexadecane, and uncultured *Alpha*- and *Gammaproteobacteria* populations were enriched in the polycyclic aromatic hydrocarbon-degrading communities and contained a broad gene set for degrading phenanthrene and naphthalene. The repertoire of polycyclic aromatic hydrocarbon use varied among different bacterial taxa and the combined capabilities of the microbial community exceeded those of its individual components, indicating that the degradation of complex hydrocarbon mixtures requires the non-redundant capabilities of a complex oil-degrading community.**

Q1


Marine microorganisms derive energy and carbon from the degradation of petroleum hydrocarbons and drive the bioremediation process during anthropogenic oil spills, such as the Deepwater Horizon (DWH) spill<sup>2,4,7-9</sup>. Various uncultured bacteria, primarily belonging to *Oceanospirillales* and other *Gammaproteobacteria*, were enriched during the DWH spill and are believed to have played a major role in oil degradation<sup>2,3,10-12</sup>. However, this assumption relies on the dominance of these organisms in environmental sequencing surveys, whereas the metabolic potential and pathways for degrading hydrocarbons in bacterial populations, which demonstrably assimilate these compounds, remain to be determined. To address this question, we obtained ~113 Gb of shotgun metagenomic data from hydrocarbon-degrading enrichments using stable-isotope probing (SIP) experiments<sup>13</sup>. Enrichments were obtained from weathered hydrocarbons floating on the sea surface and from the subsurface hydrocarbon plume, collected during the DWH oil spill in 2010. The plume sample was incubated with <sup>13</sup>C-labelled *n*-hexadecane (HEX), and the sea surface sample was separately incubated with <sup>13</sup>C-labelled naphthalene (NAP) or phenanthrene (PHE) in SIP experiments<sup>13</sup>. *De novo* assembly and (tetranucleotide signature) binning of metagenomic sequences from the <sup>13</sup>C-labelled fractions allowed the reconstruction of 7 high-quality (completeness of 52–95%) and 17

fragmentary draft genomes (completeness <50%), as well as ~13% of unbinned scaffolds (see Methods, Supplementary Table 1 and Supplementary Fig. 1).

We first annotated genes within all assembled genomes. Taxonomic assignment of annotated genes in the HEX, NAP and PHE assemblies revealed distinct bacterial communities (Supplementary Fig. 2a,b). *Marinobacter* and *Alcanivorax* dominated the HEX-degrading community, and *Alteromonas* and *Cycloclasticus* were abundant in NAP and PHE samples, respectively. This community spectrum coincided with results from the previous SIP-based analysis of the same samples using 16S rRNA clone sequencing<sup>13</sup>. To identify taxa representing the seven high-quality draft genomes, we constructed a phylogenetic tree using up to 15 ribosomal proteins (Fig. 1 and Supplementary Table 1). We recovered a 95% complete *Marinobacter* genome from the HEX enrichment, designated H-Mar (99% inferred protein similarity to *Marinobacter salarius* R9SW1, Fig. 1 and Supplementary Table 1). From the NAP-degrading community we reconstructed two gammaproteobacterial genomes belonging to *Oceanospirillales* (N-Alc, 73% similar to *Alcanivorax* sp. 43B\_GOM-46m, 84% complete) and *Alteromonadales* (N-Alt, 86% similar to *Alteromonas macleodii* English Channel 673, 80% complete). An additional 93% complete alphaproteobacterial genome belonged to the order *Rhodospirillales* (N-Tha, 99% similar to *Thalassospira profundimaris* WP0211). Three genomes from the PHE-degrading communities included a bacterium 83% similar to *Cycloclasticus pugetii* PS-134H (P-Cyc, 52% complete), an *Oceanospirillales* member with 73% similarity to *Neptuniibacter caesariensis* MED92 (P-Nep, 86% complete) and a member of the *Alteromonadales* with 64% similarity to *Colwellia psychrerythraea* 34H (P-Col, 70% complete).

The NAP-enriched taxa *Rhodospirillales*, *Oceanospirillales* and *Alteromonadales* (N-Tha, N-Alc and N-Alt) were previously associated with hydrocarbon degradation. For example, isolates of the order *Alteromonadales* showed alkane and polycyclic aromatic hydrocarbon (PAH)-degrading activity in previous SIP experiments<sup>13</sup>. The order *Rhodospirillales* accumulated during the DWH spill in sea surface samples, and isolates of this order, such as *Thalassospira tepidiphila*, degrade PAHs<sup>3,14</sup>. We also assembled one genome belonging to *Alcanivorax* from the NAP-degrading community (N-Alc), a genus that was barely detected in the spill itself but was present in the previous SIP experiments<sup>13</sup>. Additionally, *Alcanivorax* was abundant in HEX enrichments (Supplementary Fig. 2b). Although *Alcanivorax* isolates degrade alkane hydrocarbons and none have been described to metabolize PAHs<sup>13,15</sup>, *Alcanivorax* was previously detected in bacterial consortia growing

<sup>1</sup>Department of Marine Science, University of Texas Austin, Marine Science Institute, Port Aransas, Texas 78373, USA. <sup>2</sup>School of Life Sciences, Heriot-Watt University, Edinburgh EH14 4AS, UK. <sup>3</sup>Department of Marine Sciences, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599 USA. \*e-mail: acidophile@gmail.com



**Figure 1 | Phylogenetic characterization of genomic bins reconstructed from three SIP enrichments.** Maximum-likelihood-based phylogenetic tree of up to 15 concatenated ribosomal proteins (rpl2, 3, 4, 5, 6, 14, 15, 18, 22, 24 and rpS3, 8, 10, 17, 19) from seven high-quality genomes assembled from the *n*-hexadecane (H), naphthalene (N) and phenanthrene (P) SIP enrichments (bold). Two genomes from previously published DWH plume-derived metagenomes are indicated in italics<sup>10,46</sup>. Bootstrap values were generated using MrBayes. Two *Planctomycetes* strains were used as outgroup. Black, grey and white circles: nodes with bootstrap values of 100, 75-100 and <75% (1,000 replicates).


1 on pyrene as the sole carbon source<sup>16,17</sup>. Thus, the NAP enrichment  
 2 of *Alcanivorax* supports a broader versatility in *Alcanivorax* for  
 3 degrading not only alkanes but also PAHs, potentially by participating  
 4 in bacterial consortia that fully metabolize PAHs.

5 Focusing on the PHE-degrading community, the genera  
 6 *Cycloclasticus*, *Colwellia* and *Neptuniibacter* (corresponding to  
 7 P-Cyc, P-Col and P-Nep) represented 66, 13 and 3% of clones in  
 8 previous PHE-SIP experiments, respectively<sup>13</sup>. Notably, the  
 9 *Cycloclasticus* 16S rRNA phylotype recovered from the previous SIP  
 10 study was >99% similar to the most dominant *Cycloclasticus* phylotype  
 11 that was enriched at the sea surface during the spill<sup>13</sup>. Concordantly,  
 12 *Colwellia* was enriched in laboratory experiments in dispersant-oil  
 13 microcosms<sup>18</sup>. Thus, the assembled genomes represent abundant, as  
 14 well as rare, hydrocarbon-degrading DWH spill community members.

15 16S rRNA gene sequences recovered from the SIP assemblies co-  
 16 clusted with sequences derived from the previous SIP experiments  
 17 and partially with sequences enriched in surface and plume samples  
 18 during the spill (Supplementary Fig. 3)<sup>2,3,13</sup>. These clustering patterns  
 19 therefore indicate that the bacterial lineages present in our metagenomic  
 20 assemblies were present during the spill and do not represent artefacts of  
 21 the SIP experiment. To confirm this inference, we mapped our SIP  
 22 assemblies to published plume-derived metagenomic and metatranscriptomic  
 23 data sets<sup>10</sup>. The HEX-degrading assembly recruited 2.7  
 24 and 0.8% of metagenomic reads from plume and unpolluted samples,  
 25 respectively (Supplementary Table 2). Annotated genes from the HEX  
 26 assembly matched 6.3 and 1.3% of metatranscriptomic reads from  
 27 samples collected proximal and distal, respectively, from the plume  
 28 (Supplementary Table 2 and Supplementary Figs 4 and 5). This suggests  
 29 that genes from our assembled genomes were enriched and more highly

expressed with increasing hydrocarbon exposure. However, despite the  
 percentage of mapped reads, we were unable to locate specific alkane-  
 degradation genes, possibly due to insufficient sequencing depth that  
 would allow covering non-abundant taxa (Supplementary Fig. 5).  
 No metagenomic data set from the sea surface of the spill is available;  
 but based on previous 16S rRNA gene studies, *Alphaproteobacteria*  
 (that is, order *Rhodospirillales*) as well as *Gammaproteobacteria*  
 (orders *Alteromonadales* and *Oceanospirillales*) were enriched in  
 oil-contaminated sea surface samples<sup>3</sup>. Thus, the lineages recon-  
 structured in our assemblies reflect (or resemble) those enriched in  
 hydrocarbon-contaminated sea surface communities.

To identify genes that enabled bacterial community members to  
 be enriched during the DWH spill, we compared the gene content  
 between the different hydrocarbon-degrading communities obtained  
 in the SIP assemblies. Of the detected 4,756 unique gene functions,  
 ~40% were shared and 4, 13 and 22% were unique among the  
 HEX, NAP and PHE assemblies, respectively (Supplementary Fig. 6  
 and Supplementary Data 1). The PHE enrichment showed selection  
 for serine/threonine kinases, while the HEX samples encoded for  
 abundant short-chain dehydrogenases and the NAP assembly was  
 enriched for tripartite ATP-independent periplasmic (TRAP) trans-  
 porters. The serine/threonine kinases phosphorylate serine, threonine  
 and tyrosine are proposed mediators of bacteria-bacteria interactions,  
 while TRAP transporters function in osmoregulated solute transport<sup>19,20</sup>.  
 Additionally, several functional categories were present in individ-  
 ual genomic bins (Fig. 2 and Supplementary Data 2). Five of seven  
 genomes carried genes for bacterial chemotaxis and six of seven  
 for flagella biosynthesis, hinting at active motility and therefore the  
 ability to move towards oil substrates. Additionally, P-Col encoded




**Figure 2 | Abundance of functional categories detected within the reconstructed genomes.** Functional categories detected in the seven draft genomes assembled from the *n*-hexadecane (H), naphthalene (N) and phenanthrene (P) SIP enrichments. Functional categories are based on the SEED subsystems<sup>36</sup> and were compared against genes annotated within each metagenomic bin using blastp (e-value cutoff of 1e-20). Numbers in parentheses: total number of genes within each functional category. Percentage: per cent of total number of genes per subsystem found within each bin.

1 for several denitrification genes (13 of 21 tested genes), including  
 2 *norB/C*, *nosR* and *nosD*, while P-Nep appears able to oxidize  
 3 sulfur (that is, including *soxA-D* and sulfite oxidase), implying rel-  
 4 evance for degrading sulfur-containing components of oil and the  
 5 applied dispersant Corexit. Suggesting physiological adaptations to  
 6 phosphorous and iron limitation, most genomic bins contained  
 7 genes relevant for phosphate starvation or siderophore biosynthesis,  
 8 such as staphylobactin (that is, *phoH* and *sbnA*; Supplementary Data  
 9 2). The presence of major genes for motility and for use of scarce  
 10 nutrients suggests that the enriched organisms are well adapted  
 11 for chemotactic motility towards their hydrocarbon substrates and  
 12 for physiological responses to nutrient-limiting conditions that  
 13 characterize oil-induced bacterial blooms<sup>21</sup>. Alternatively, these  
 14 genes might be of general importance for survival and growth in the  
 15 Gulf of Mexico.

16 To resolve the hydrocarbon degradation pathways in the assembled  
 17 bacterial genomes, we searched for homologues to known degradation  
 18 genes. An entire pathway for alkane degradation was found in H-Mar

(*Marinobacter*), the dominant genus in the HEX-degrading commu- 19  
 nity (Fig. 3a and Supplementary Figs 1 and 2). This result is consistent 20  
 with isolates of *Marinobacter* being able to degrade HEX, increasing 21  
*Marinobacter* abundance in DWH plume samples as validated by 22  
 fluorescence *in situ* hybridization (FISH), and enrichment of 23  
 genes for alkane degradation in metatranscriptomes from the 24  
 DWH spill and the Gulf coast<sup>2,10,22,23</sup>. Additionally, we recon- 25  
 structed the alkane-degradation pathway in most of the genomes 26  
 from the NAP- and PHE-degrading bins (Fig. 3a). However, 27  
 based on the SIP approach, only a subset of bacteria appears to 28  
 actively employ these pathways—*Marinobacter* (H-Mar)—and 29  
 potentially outcompete other alkane-degrading bacteria, such as 30  
*Alcanivorax* (N-Alc). To validate this finding, metatranscriptomic 31  
 sequencing could be used to map active gene expression to our 32  
 assembled draft genomes. One unique feature of the active alkane- 33  
 degrader H-Mar is the presence of genes encoding for the type VI 34  
 secretion system<sup>24,25</sup>, which were mostly absent from all other 35  
 genomic bins (Fig. 2). The type VI secretion system is involved in 36



**Figure 3 | Distribution of key alkane and polycyclic hydrocarbon degradation pathways in reconstructed genomes. a**, Metabolic reconstruction of the *n*-hexadecane (HEX) degradation pathway within seven high-quality metagenomic bins. **b**, Metabolic reconstruction of naphthalene (NAP) and phenanthrene (PHE) degradation pathways within seven high-quality genomes. Purple, green, blue: HEX, NAP and PHE-degrading bins, respectively.

1 inter-bacterial competition and has been discussed to be relevant  
2 for alkane assimilation<sup>24,25</sup>, and thus could provide H-Mar with a  
3 competitive advantage over other non-active alkane degraders.

To characterize PAH degradation pathways in SIP-enriched bac- 4  
teria, we identified genes encoding for enzymes in both NAP- and 5  
PHE-degradation pathways in our assembled genomes (Fig. 3b). 6

1 Of 41 described NAP- and PHE-degradation genes, 19 were present  
2 in >50% of genomes, while a subset of genes, such as *nahC*, salicy-  
3 late hydroxylase or *phdF*, were only detectable in a few genomes. For  
4 example, within N-Tha, N-Alc and N-Alt, we were unable to anno-  
5 tate the full known NAP-degradation pathway. N-Tha  
6 (*Thalassospira profundimaris*) is equipped to degrade NAP to sali-  
7 cylate, but lacks the salicylate hydroxylase, which converts salicylate  
8 to catechol. Only the fragmentary genome NAP-25  
9 (*Oceanospirillales*) encoded the gene for this step among the  
10 assembled genomes from the NAP enrichments. Similarly, the  
11 extradiol dioxygenase (*PhdF*), which converts 3,4-dihydroxyphen-  
12 nanthrene to 2-hydroxybenzochromene-2-carboxylic acid early on  
13 in the PHE-degradation pathway, was only detected in P-Cyc  
14 (*Cycloclasticus*). Unless all pathway gaps are without exception  
15 ascribed to sequencing gaps within the genomic bins, a more parsimo-  
16 nious interpretation of these results suggests that alternative  
17 genes and enzymes for these individual steps exist, which are not  
18 represented in gene databases. Alternatively, coordinated commu-  
19 nity activity might be required to completely degrade polycyclic  
20 hydrocarbons during the DWH spill.

21 Our study combines metagenomics and SIP, allowing the  
22 assembly of genomes from active hydrocarbon-degrading bacteria  
23 present during the DWH spill, and enriched from deep plume and  
24 surface samples based on alkane and PAH assimilation. All  
25 assembled genomes possessed pathways for alkane and PAH  
26 degradation, suggesting that several taxa actively degraded  
27 hydrocarbons during the spill (Fig. 3). For example, H-Mar  
28 (*Oceanospirillales*) corresponds to *Marinobacter salarius*, which  
29 degrades alkanes in the plume, and the bacterium that constitutes  
30 N-Tha (*Rhodospirillales*) was responsible for degradation of PAHs  
31 at the sea surface. These taxa were enriched during the spill and  
32 their genes for hydrocarbon degradation were detected in the  
33 plume and sea surface<sup>2,3,10,22,26</sup>. However, the plume-derived  
34 genome (H-Mar) appears to be enriched only at a low relative  
35 abundance compared to previous data sets, where an uncharacter-  
36 ized *Oceanospirillales* represented >60% of sequence reads<sup>10</sup>. This  
37 disparity suggests that low-abundant community members were  
38 active during the spill, as observed in deep-sea hydrothermal  
39 plumes<sup>27</sup>. Additionally, differences between bacterial taxa detected  
40 during the DWH spill compared to SIP enrichments could be a  
41 reflection of the experimental set-up. For example, no dispersant  
42 was added during the experiment, but Corexit was added during  
43 the spill<sup>13,28</sup> and found to select against *Marinobacter* in micro-  
44 cosm experiments<sup>18</sup>. Additionally, the SIP procedure required  
45 incubation steps at room temperature instead of 4 °C to obtain  
46 sufficient incorporation of the <sup>13</sup>C label into DNA for SIP<sup>13</sup>.  
47 Such temperature shifts have been demonstrated to induce  
48 changes in plume bacterial community composition<sup>3</sup>. Whole-  
49 community gene expression studies will be required to evaluate  
50 the activity of our assembled genomes in their native  
51 community context.

52 Within the constraints inherent to the SIP experimental  
53 approach, combining SIP with metagenomics enabled us to identify  
54 and analyse hydrocarbon-degrading pathways within several uncultured  
55 bacteria. Bacteria that are usually not associated with PAH  
56 degradation (that is, members of *Alcanivorax* in the NAP enrich-  
57 ments) persisted at sufficiently high levels for metagenomic recon-  
58 struction, indicating that complete pathways for the degradation  
59 of PAHs may not be obligatory for individual community  
60 members that thrive in PAH-degrading enrichments. Although we  
61 cannot exclude that genes for hydrocarbon degradation were  
62 missed in our assemblies, the apparent partitioning of key  
63 pathway steps to individual community members suggests an alter-  
64 nate hypothesis, in that complete degradation of a complex mixture  
65 of PAHs requires the coordinated response of a complex bacterial  
66 community and the coordination of its metabolic pathways.

## Methods

**Sample collection and preparation.** DNA samples for metagenomic reconstruction  
were obtained from a previously published stable-isotope probing (SIP)  
experiment<sup>13</sup>. Briefly, an oil-contaminated sea surface sample (original ID PE5) was  
collected near a site of the DWH spill (28°44.175' N, 88°22.335' W) in the Gulf of  
Mexico on 5 May 2010. Two deep hydrocarbon plume samples (original ID B3  
and B6, biological replicates) were collected near the Macondo wellhead  
(28°41.686' N, 88°26.081' W) at depths of 1,170 and 1,210 m on 31 May 2010 and  
combined for the SIP experiment. Sea surface and plume samples were collected at  
~0.86 and 3.5 m from the wellhead. SIP experiments for the plume sample (B3+B6)  
were performed with <sup>13</sup>C-labelled HEX and for the sea surface sample (PE5) with  
<sup>13</sup>C-labelled NAP or PHE (yielding a total of three samples). Incubations containing  
solely <sup>12</sup>C-unlabelled substrate were run in parallel to act as unlabelled controls.  
Total DNA from labelled and unlabelled incubations from each of the three SIP  
experiments was extracted as described previously<sup>29</sup>. Extracted labelled and  
unlabelled DNA (0.7 to 1.0 µg for each sample) was separated using CsCl gradient  
ultracentrifugation<sup>13</sup>. To prepare these previously processed SIP samples for  
sequencing, DNA concentrations were measured using a Qubit 3.0 Fluorometer and  
a final concentration of 10 ng µl<sup>-1</sup> for each sample (using a total amount of 100 ng)  
from the heavy-labelled fraction was used to prepare libraries for paired-end  
Illumina (HiSeq 2500) sequencing.

**Metagenomic assembly and binning.** Illumina library preparation and sequencing  
was performed by the Genome Sequencing and Analysis Facility (GSAF) at the  
University of Texas at Austin. For each of the three samples, two Illumina libraries  
were prepared and sequenced (technical replicates). Sequencing was performed on  
an Illumina HiSeq 2500 with the following specifications: high-throughput run  
mode, run type paired end 2 × 125 bp, 6 × 4.0E8 target reads (millions), insert size of  
~360–420 bp and ~5% PhiX control spike-in. This sequencing approach provided  
~113 Gb of sequencing data (373,279,006, 289,782,708 and 246,243,202 reads from  
the HEX, NAP and PHE samples, respectively). Raw Illumina shotgun genomic  
reads were separated from Illumina artefacts by removing the adaptors and DNA  
spike-ins from the forward and reverse reads. Therefore, a sliding window approach  
using a kmer size of 28 and a step size of 1 was used. Reads with ≥3 Ns, an average  
score of <Q20 and a read length of <50 bps were removed using cutadapt (yielding  
886,823,725 total reads)<sup>30</sup>. Afterwards, reads were interleaved using interleave\_fasta.  
py and the interleaved sequences were trimmed using Sickle and a minimum quality  
cutoff of 5 (yielding 877,414,119 total reads)<sup>31</sup>. The script for interleave\_fasta.py can  
be found at [https://github.com/jorvis/biocode/blob/master/fasta/interleave\\_fasta.py](https://github.com/jorvis/biocode/blob/master/fasta/interleave_fasta.py).  
Metagenomic reads from all SIP samples were individually assembled using  
IDBA-UD and the following parameters: -pre\_correction, -mink 75, -maxk 105,  
-step 10, -seed\_kmer 55 (ref. 32). This yielded a total of 2,739,076 scaffolds from all  
three SIP enrichments. The minimum and maximum scaffold length ranged from  
200 to 130,027 bp.

Metagenomic binning was performed on assembled SIP enrichments using  
tetranucleotide frequencies on scaffolds with a length ≥4,000 bp (including a total of  
4,874 scaffolds)<sup>33</sup>. The resulting emerging self-organizing maps (ESOMs) were  
manually sorted and curated (Supplementary Fig. 1)<sup>33</sup>. Metagenomic binning was  
enhanced by incorporating reference genomes as genetic signatures for the  
assembled contigs into ESOMs<sup>33,34</sup>. In this way we assembled 7 high-quality and 17  
fragmentary metagenomic bins (completeness threshold of 50%). The seven high-  
quality genomes showed a completeness ranging from 52–95%, included 1,492/4,874  
scaffolds with a scaffold length between 4,006 and 257,386 bp. The 17 fragmentary  
genomes displayed a completeness below 50% and comprised 2,737/4,874 scaffolds  
that ranged from 4,000 to 44,889 bp. Additionally, 645/4,874 scaffolds remained  
unbinned (4,002–77,080 bp length). After binning, draft genomes were linked to the  
SIP enrichments based on their unique scaffold ID. CheckM was used to evaluate the  
accuracy of the binning approach by determining the percentage of completeness  
and contamination (Supplementary Table 1)<sup>35</sup>. Contaminants that were identified  
based on their phylogenetic placement (wrong taxonomic assignment compared to  
the average taxonomic assignment of the genes assigned to each bin; see section on  
taxonomic assignment), GC content (>25% difference compared to the mean of all  
scaffolds assigned to each bin) or confidence level (below 0.5) were manually  
removed from the bins.

**Gene calling, taxonomic assignment and functional characterization.** Gene  
calling and taxonomic assignment of the three SIP enrichments and seven  
high-quality draft genomes were performed using the Joint Genome  
Institute–Production Genomics Facility (JGI-PGF) Integrated Microbial Genomes  
with Microbiome (IMG/M) system. To achieve this, the complete assemblies  
from the three SIP enrichments were uploaded onto the JGI server. The retrieved JGI  
output was linked to the seven draft genomes using their unique scaffold IDs.  
The JGI output summarizing the taxonomic assignment of all gene annotations was  
used to depict the taxonomic composition as shown in Supplementary Fig. 2.  
The JGI-based amino-acid sequences were used as a database for blastp searches  
(as indicated in the text).

For the functional characterization of our draft genomes, a reference gene  
database was assembled by downloading SEED subsystems using the network-based  
SEED API (function svr\_all\_subsystems)<sup>36</sup>. The SEED-based reference database was

1 manually curated to include only subsystems of interest by downloading the gene  
2 functions and gene IDs of individual subsystems (functions svr\_subsystem\_roles  
3 and svr\_subsystem\_spreadsheet), which were then used to retrieve their respective  
4 fasta sequences (function svr\_fasta). The SEED subsystems included are shown in  
5 Fig. 2, and the numbers in parentheses include the total number of genes screened  
6 per subsystem (Fig. 2 and Supplementary Data 2). Additionally, we manually  
7 curated a reference database including only hydrocarbon degradation genes by  
8 searching the KEGG and NCBI databases for hydrocarbon degradation pathways  
9 and corresponding genes (Fig. 3). Both of these reference databases were screened  
10 against annotated genes from the seven draft genomes using their JGI-derived  
11 amino-acid sequences as input for blastp (e-value threshold of  $1e-20$ )<sup>37</sup>.

12 **Phylogenetic analyses.** Phylosift was used to extract marker genes for the  
13 phylogenetic placement of the assembled metagenomic bins<sup>38</sup>. These marker genes  
14 consist of up to 15 syntenic ribosomal protein genes that have been demonstrated to  
15 undergo limited lateral gene transfer (rpl2, 3, 4, 5, 6, 14, 15, 18, 22, 24 and rpS3, 8,  
16 10, 17, 19)<sup>39</sup>. This gene set was derived from a reference database as detailed in  
17 ref. 40. To search for ribosomal protein sequences, all metagenomic bins (fasta files)  
18 were used as an input in Phylosift, which was used with default parameters.  
19 Moreover, we searched NCBI to include amino-acid sequences from bacterial  
20 reference strains for phylogenetic analyses. Amino-acid alignments of the individual  
21 ribosomal protein genes were generated using MUSCLE and curated manually<sup>41</sup>.  
22 Afterwards, the curated alignments of the ribosomal proteins were concatenated for  
23 further phylogenetic analyses. In addition to analysing ribosomal proteins, we  
24 employed EMIRGE to retrieve 16S rRNA genes from the three SIP enrichments<sup>42</sup>.  
25 We therefore ran an EMIRGE analysis comparing the SILVA database against the  
26 raw, short reads from the three SIP enrichments using default parameters. Retrieved  
27 16S rRNA genes were aligned to the SILVA database using the ARB alignment tool  
28 and were curated manually<sup>43</sup>. For both the ribosomal protein and 16S rRNA gene  
29 alignments, phylogenetic trees were generated using a maximum likelihood-based  
30 analysis (RAxML; rate distribution models: PROTGAMMA for ribosomal proteins  
31 and GTRGAMMA for 16S rRNA gene sequences)<sup>44</sup>. Bootstrap values were  
32 calculated using MrBayes with 100,000 generations of Markov chain Monte Carlo  
33 (MCMC) analyses with 100 sample and print frequencies<sup>45</sup>. As a point of  
34 comparison, we included published 16S rRNA gene sequences from the plume and  
35 sea surface as well as from the previous SIP experiment (16S rRNA genes from clone  
36 libraries and isolated bacterial strains) for the 16S rRNA phylogenetic tree<sup>2,3,10,13,46</sup>.

37 **Meta analysis.** To determine whether genes from the assembled metagenomic bins  
38 in this study were enriched and active in the plume during the spill, we compared  
39 our three SIP enrichments with previously published plume-derived metagenomic  
40 and metatranscriptomic data sets ([http://mason.eoas.fsu.edu/DWH\\_plume/](http://mason.eoas.fsu.edu/DWH_plume/))<sup>10</sup>. Two  
41 reference databases were generated, which were compared against the published  
42 metagenomic and metatranscriptomic data sets. Reference database 1 consisted of  
43 DNA sequences comprising scaffolds of >2,500 bp that were derived from the three  
44 SIP assemblies. Reference database 2 included amino-acid sequences from the seven  
45 high-quality draft genomes. Reference database 1 (DNA sequences) was mapped  
46 against the published metagenomic and metatranscriptomic data set using BWA.  
47 Database 2 (amino-acid sequences) was used to search against the published  
48 metatranscriptomic data using Rapsearch2 to retrieve the exact genes that mapped  
49 against this data set<sup>47,48</sup>. For BWA the default parameters were used and an e-value  
50 cutoff of 0.001 was used for RapSearch2.

51 **Accession codes.** The genomes are available in NCBI Genbank under BioProjectID  
52 PRJNA301966. The whole genome shotgun projects have been deposited under  
53 accession nos. LSM000000000 (H-Mar), LSMN000000000 (N-Tha),  
54 LSMO000000000 (N-Alc), LSMP000000000 (N-Alt), LSMQ000000000 (P-Cyc),  
55 LSMR000000000 (P-Nep) and LMS000000000 (P-Col). The reference numbers for  
56 the original raw sequencing data are SRX1562986 (*n*-hexadecane), SRX1585241  
57 (naphthalene) and SRX1586894 (phenanthrene).

58 Received 11 December 2015; accepted 29 March 2016;  
59 published XX XX 2016

## 60 References

- 61 1. Crone, T. J. & Tolstoy, M. Magnitude of the 2010 Gulf of Mexico oil leak. *Science*  
62 **330**, 634–634 (2010).
- 63 2. Hazen, T. C. *et al.* Deep-sea oil plume enriches indigenous oil-degrading  
64 bacteria. *Science* **330**, 204–208 (2010).
- 65 3. Redmond, M. C. & Valentine, D. L. Natural gas and temperature structured a  
66 microbial community response to the Deepwater Horizon oil spill. *Proc. Natl*  
67 *Acad. Sci. USA* **109**, 20292–20297 (2012).
- 68 4. Yang, T. *et al.* Pulsed blooms and persistent oil-degrading bacterial populations  
69 in the water column during and after the Deepwater Horizon blowout. *Deep Sea*  
70 *Res. II* doi:10.1016/j.dsr2.2014.01.014 (2014).
- 71 5. Crespo-Medina, M. *et al.* The rise and fall of methanotrophy following a  
72 deepwater oil-well blowout. *Nature Geosci.* **7**, 423–427 (2014).
- 73 6. Valentine, D. L. *et al.* Propane respiration jump-starts microbial response to a  
74 deep oil spill. *Science* **330**, 208–211 (2010).

7. Chauhan, A., Fazlurrahman, Oakeshott, J. G. & Jain, R. K. Bacterial metabolism  
of polycyclic aromatic hydrocarbons: strategies for bioremediation. *Indian J.*  
*Microbiol.* **48**, 95–113 (2008). Q7
8. Kimes, N. E., Callaghan, A. V., Suflita, J. M. & Morris, P. J. Microbial  
transformation of the Deepwater Horizon oil spill—past, present, and future  
perspectives. *Front. Microbiol.* **5** (2014). Q8
9. Lea-Smith, D. J. *et al.* Contribution of cyanobacterial alkane production to the  
ocean hydrocarbon cycle. *Proc. Natl Acad. Sci. USA* **112**, 13591–13596 (2015). 81
10. Mason, O. U. *et al.* Metagenome, metatranscriptome and single-cell  
sequencing reveal microbial response to Deepwater Horizon oil spill.  
*ISME J.* **6**, 1715–1727 (2012). 85
11. Bælum, J. *et al.* Deep-sea bacteria enriched by oil and dispersant from the  
Deepwater Horizon spill. *Environ. Microbiol.* **14**, 2405–2416 (2012). 86
12. Kleindienst, S. *et al.* Diverse, rare microbial taxa responded to the Deepwater  
Horizon deep-sea hydrocarbon plume. *ISME J.* **10**, 400–415 (2015). Q9
13. Gutierrez, T. *et al.* Hydrocarbon-degrading bacteria enriched by the  
Deepwater Horizon oil spill identified by cultivation and DNA-SIP. *ISME J.* **7**,  
2091–2104 (2013). 90
14. Kodama, Y., Stiknowati, L. I., Ueki, A., Ueki, K. & Watanabe, K. *Thalassospira*  
*tepidiphila* sp. nov., a polycyclic aromatic hydrocarbon-degrading bacterium  
isolated from seawater. *Int. J. Syst. Evol. Microbiol.* **58**, 711–715 (2008). 94
15. Schneiker, S. *et al.* Genome sequence of the ubiquitous hydrocarbon-  
degrading marine bacterium *Alcanivorax borkumensis*. *Nature Biotechnol.* **24**,  
997–1004 (2006). 96
16. Lai, Q. *et al.* *Alcanivorax pacificus* sp. nov., isolated from a deep-sea pyrene-  
degrading consortium. *Int. J. Syst. Evol. Microbiol.* **61**, 1370–1374 (2011). 98
17. Wang, B., Lai, Q., Cui, Z., Tan, T. & Shao, Z. A pyrene-degrading consortium  
from deep-sea sediment of the West Pacific and its key member *Cycloclasticus* sp.  
P1. *Environ. Microbiol.* **10**, 1948–1963 (2008). 100
18. Kleindienst, S. *et al.* Chemical dispersants can suppress the activity of natural  
oil-degrading microorganisms. *Proc. Natl Acad. Sci. USA* **112**,  
14900–14905 (2015). Q10
19. Cozzone, A. J. Role of protein phosphorylation on serine/threonine and  
tyrosine in the virulence of bacterial pathogens. *J. Mol. Microbiol. Biotechnol.* **9**,  
198–213 (2005). 107
20. Grammann, K., Volke, A. & Kunte, H. J. New type of osmoregulated solute  
transporter identified in halophilic members of the bacteria domain: TRAP  
transporter TeaABC mediates uptake of ectoine and hydroxyectoine in  
*Halomonas elongata* DSM 2581T. *J. Bacteriol.* **184**, 3078–3085 (2002). 110
21. Edwards, B. R. *et al.* Rapid microbial respiration of oil from the Deepwater  
Horizon spill in offshore surface waters of the Gulf of Mexico. *Environ. Res. Lett.*  
**6**, 035301 (2011). 111
22. McKay, L. J., Gutierrez, T. & Teske, A. P. Development of a group-specific 16S  
rRNA-targeted probe set for the identification of *Marinobacter* by fluorescence  
in situ hybridization. *Deep Sea Res. II* doi:10.1016/j.dsr2.2013.10.009 (2014). Q11
23. Lamendella, R. *et al.* Assessment of the Deepwater Horizon oil spill impact on  
Gulf coast microbial communities. *Aquat. Microbiol.* **5**, 130 (2014). 112
24. Salomon, D. *et al.* Type VI secretion system toxins horizontally shared between  
marine bacteria. *PLoS Pathogens* **11** e1005128 (2015). Q12
25. Vaysse, P.-J. *et al.* Proteomic analysis of *Marinobacter hydrocarbonoclasticus*  
SP17 biofilm formation at the alkane–water interface reveals novel proteins and  
cellular processes involved in hexadecane assimilation. *Res. Microbiol.* **160**,  
829–837 (2009). 123
26. Arnosti, C., Ziervogel, K., Yang, T. & Teske, A. Oil-derived marine aggregates—  
hot spots of polysaccharide degradation by specialized bacterial communities.  
*Deep Sea Res. II* doi:10.1016/j.dsr2.2014.12.008. Q13
27. Baker, B. J., Lesniewski, R. A. & Dick, G. J. Genome-enabled transcriptomics  
reveals archaeal populations that drive nitrification in a deep-sea hydrothermal  
plume. *ISME J.* **6**, 2269–2279 (2012). 129
28. Kujawinski, E. B. *et al.* Fate of dispersants associated with the Deepwater  
Horizon oil spill. *Environ. Sci. Technol.* **45**, 1298–1306 (2011). 130
29. Tillett, D. & Neilan, B. A. Xanthogenate nucleic acid isolation from cultured and  
environmental cyanobacteria. *J. Phycol.* **36**, 251–258 (2000). 131
30. Martin, M. Cutadapt removes adapter sequences from high-throughput  
sequencing reads. *EMBnet.journal* **17**, 10–12 (2011). 132
31. Joshi, N. & Fass, J. Sickle: a sliding-window, adaptive, quality-based  
trimming tool for FastQ files (Version 1.33) (Software) (2011);  
<https://github.com/najoshi/sickle> 133
32. Peng, Y., Leung, H. C. M., Yiu, S. M. & Chin, F. Y. L. IDBA-UD: a *de novo*  
assembler for single-cell and metagenomic sequencing data with highly uneven  
depth. *Bioinformatics* **28**, 1420–1428 (2012). Q14
33. Dick, G. J. *et al.* Community-wide analysis of microbial genome sequence  
signatures. *Genome Biol.* **10**, R85 (2009). 143
34. Baker, B. J. Omic approaches in microbial ecology: charting the unknown.  
*Microbe* **8** (2013). 144
35. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W.  
CheckM: Assessing the quality of microbial genomes recovered from isolates,  
single cells, and metagenomes. *Genome Res.* **25**, 1043–1055 (2015). 145

- 1 36. Overbeek, R. *et al.* The SEED and the Rapid Annotation of microbial  
2 genomes using Subsystems Technology (RAST). *Nucleic Acids Res.* **42**,  
3 D206–D214 (2014).
- 4 37. Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein  
5 database search programs. *Nucleic Acids Res.* **25**, 3389–3402 (1997).
- 6 38. Darling, A. E. *et al.* PhyloSift: phylogenetic analysis of genomes and  
7 metagenomes. *PeerJ* **2**, e243 (2014).
- 8 39. Sorek, R. *et al.* Genome-wide experimental determination of barriers to  
9 horizontal gene transfer. *Science* **318**, 1449–1452 (2007).
- 10 40. Castelle, C. J. *et al.* Genomic expansion of domain archaea highlights roles  
11 for organisms from new phyla in anaerobic carbon cycling. *Curr. Biol.* **25**,  
12 690–701 (2015).
- 13 41. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and  
14 high throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).
- 15 42. Miller, C. S., Baker, B. J., Thomas, B. C., Singer, S. W. & Banfield, J. F. EMIRGE  
16 reconstruction of full-length ribosomal genes from microbial community short  
17 read sequencing data. *Genome Biol.* **12**, R44 (2011).
- 18 43. Ludwig, W. *et al.* ARB: a software environment for sequence data. *Nucleic Acids*  
19 *Res.* **32**, 1363–1371 (2004).
- 20 44. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-  
21 analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).
- 22 45. Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference  
23 under mixed models. *Bioinformatics* **19**, 1572–1574 (2003).
- 24 46. Eren, A. M. *et al.* Anvi'o: an advanced analysis and visualization platform for  
25 'omics data. *PeerJ* **3**, e1319 (2015).
- 26 47. Li, H. Aligning sequence reads, clone sequences and assembly contigs with  
Q1632 BWA-MEM. arXiv:1303.3997 (2013).

48. Zhao, Y., Tang, H. & Ye, Y. RAPSearch2: a fast and memory-efficient protein  
28 similarity search tool for next-generation sequencing data. *Bioinformatics* **28**,  
29 125–126 (2012).  
30

### Acknowledgements

The metagenomic DNA originated from work that was supported by a Marie Curie  
International Outgoing Fellowship (PIOF-GA-2008-220129) to T.G. within the 7th  
European Community Framework Programme. Sampling in the Gulf of Mexico and SIP  
experiments underlying this study were made possible in part by a grant from The Gulf of  
Mexico Research Initiative and in part by a Marie Curie Fellowship to T.G. A.T. also  
acknowledges funding from the National Science Foundation (RAPID Response: the  
microbial response to the Deepwater Horizon Oil Spill; NSF-OCE 1045115). Data are  
publicly available through the Gulf of Mexico Research Initiative Information & Data  
Cooperative (GRIIDC) at <https://data.gulfresearchinitiative.org> (doi:10.7266/N7GH9FZ8).  
This is ECOGIG contribution 431.

### Author contributions

N.D., T.G. and B.J.B. conceived this study. N.D. and B.J.B. supervised experiments and  
analyses. N.D., J.A.D., K.W.S. and B.J.B. performed analyses. N.D., T.G., A.P.T. and B.J.B.  
wrote the paper with contributions from all authors.

### Additional information

Supplementary information is available online. Reprints and permissions information is  
available online at [www.nature.com/reprints](http://www.nature.com/reprints). Correspondence and requests for materials should  
be addressed to B.J.B.

### Competing interests

The authors declare no competing financial interests.

Journal: Nature Microbiology  
Article ID: nmicrobiol.2016.57  
Article title: Reconstructing metabolic pathways of hydrocarbon-degrading bacteria from the Deepwater Horizon oil spill  
Author(s): Nina Dombrowski *et al.*

Query no.	Queries	Response
Q1	Please check that the bold paragraph is OK as amended	
Q2	Please check that the sentence beginning “The serine” is OK as amended	
Q3	Please check that the sentence beginning “However” is OK as amended	
Q4	In the sentence beginning “Reads” – please ensure that the different parameters are clearly identified.	
Q5	Please clarify where you mean by “in the text” here	
Q6	Ref 4 – please provide updated information	
Q7	Ref 7 – the author names appear to be incomplete – please check and provide full names.	
Q8	Ref 8 – please check volume number and provide page range	
Q9	Ref 12 – please check inserted volume number and page range	
Q10	Ref 18 – please check inserted volume number and page range	
Q11	Ref 22 – please provide updated information	
Q12	Ref 24 – please check inserted article number	
Q13	Ref 26 – please provide updated information	
Q14	Ref 31 – please check added web link	
Q15	Ref 34 – please check volume number and provide page range	
Q16	Ref 47 – please provide updated information	
Q17	Nearly all the text is in italics in Figure 1 – is this statement about italics in the caption valid?	