



Enhanced Working Memory Binding by Direct Electrical Stimulation of the Parietal Cortex

Agustina Birba^{1,2}, Eugenia Hesse^{1,2,3}, Lucas Sedeño^{1,2}, Ezequiel P. Mikulan^{1,2}, María del C. García⁴, Juan Ávalos⁴, Federico Adolffi^{1,2}, Agustina Legaz¹, Tristán A. Bekinschtein⁵, Máximo Zimmerman¹, Mario Parra^{6,7,8,9}, Adolfo M. García^{1,2,10} and Agustín Ibáñez^{1,2,9,11,12*}

¹Laboratory of Experimental Psychology and Neuroscience (LPEN), Institute of Cognitive and Translational Neuroscience (INCYT), INECO Foundation, Favaloro University, Buenos Aires, Argentina, ²National Scientific and Technical Research Council (CONICET), Buenos Aires, Argentina, ³Instituto de Ingeniería Biomédica, Facultad de Ingeniería, Universidad de Buenos Aires, Buenos Aires, Argentina, ⁴Hospital Italiano de Buenos Aires, Buenos Aires, Argentina, ⁵Consciousness and Cognition Laboratory, Department of Psychology, University of Cambridge, Cambridge, United Kingdom, ⁶Department of Psychology, School of Social Sciences, Heriot-Watt University, Edinburgh, United Kingdom, ⁷Human Cognitive Neuroscience, Centre for Cognitive Ageing and Cognitive Epidemiology, Alzheimer Scotland Dementia Research Centre, Department of Psychology, University of Edinburgh, Edinburgh, United Kingdom, ⁸Neuroprogressive and Dementia Network, NHS Research Scotland, Edinburgh, United Kingdom, ⁹Facultad de Psicología, Universidad Autónoma del Caribe, Barranquilla, Colombia, ¹⁰Faculty of Education, National University of Cuyo (UNCuyo), Mendoza, Argentina, ¹¹Center for Social and Cognitive Neuroscience (CSCN), School of Psychology, Universidad Adolfo Ibáñez, Santiago, Chile, ¹²Centre of Excellence in Cognition and its Disorders, Australian Research Council (ARC), Sydney, NSW, Australia

OPEN ACCESS

Edited by:

Alessio Avenanti,
Università di Bologna, Italy

Reviewed by:

Lucia Serenella De Federicis,
National Institute of Social Welfare,
Italy
John Fredy Ochoa,
University of Antioquia, Colombia

*Correspondence:

Agustín Ibáñez
aibanez@ineco.org.ar

Received: 27 March 2017

Accepted: 19 May 2017

Published: 08 June 2017

Citation:

Birba A, Hesse E, Sedeño L, Mikulan EP, García MdC, Ávalos J, Adolffi F, Legaz A, Bekinschtein TA, Zimmerman M, Parra M, García AM and Ibáñez A (2017) Enhanced Working Memory Binding by Direct Electrical Stimulation of the Parietal Cortex. *Front. Aging Neurosci.* 9:178. doi: 10.3389/fnagi.2017.00178

Recent works evince the critical role of visual short-term memory (STM) binding deficits as a clinical and preclinical marker of Alzheimer's disease (AD). These studies suggest a potential role of posterior brain regions in both the neurocognitive deficits of Alzheimer's patients and STM binding in general. Thereupon, we surmised that stimulation of the posterior parietal cortex (PPC) might be a successful approach to tackle working memory deficits in this condition, especially at early stages. To date, no causal evidence exists of the role of the parietal cortex in STM binding. A unique approach to assess this issue is afforded by single-subject direct intracranial electrical stimulation of specific brain regions during a relevant cognitive task. Electrical stimulation has been used both for clinical purposes and to causally probe brain mechanisms. Previous evidence of electrical currents spreading through white matter along well defined functional circuits indicates that visual working memory mechanisms are subserved by a specific widely distributed network. Here, we stimulated the parietal cortex of a subject with intracranial electrodes as he performed the visual STM task. We compared the ensuing results to those from a non-stimulated condition and to the performance of a matched control group. In brief, direct stimulation of the parietal cortex induced a selective improvement in STM. These results, together with previous studies, provide very preliminary but promising ground to examine behavioral changes upon parietal stimulation in AD. We discuss our results regarding: (a) the usefulness of the task to target prodromal stages of AD; (b) the role of a posterior network in STM binding and in AD; and (c) the potential opportunity to improve STM binding through brain stimulation.

Keywords: working memory binding, Alzheimer's disease, direct electrical stimulation, short term memory, single case study

WORKING MEMORY BINDING: ANTECEDENTS AND CASE REPORT

Recent works (Parra et al., 2009, 2010, 2015) have evidenced the critical role of visual short-term memory (STM) binding deficits as a clinical and preclinical marker of Alzheimer's disease (AD). These studies (and other related reports, see below) show that, in AD, working memory is selectively impaired for tasks requiring binding of multiple elements, but preserved for processing isolated features. Moreover, they suggest a potential role of parieto-temporo-posterior regions in both the neurocognitive deficits of AD patients and STM binding at large. Here we show that direct electrical stimulation of the parietal cortex through invasive electrodes selectively enhances working memory binding, with no effects on feature-level working memory processes.

Memory binding is the function that allows integrating multiple elements of complex events into unified wholes (von der Malsburg, 1999; Baddeley, 2000; Tulving, 2002; Zimmer et al., 2006). STM or working memory binding underpins the temporary retention of arrays of features (e.g., shapes with colors) as integrated complex objects (Treisman and Zhang, 2006). Parra et al. (2010) visual working memory task discriminates between patients with familial AD and preclinical carriers of the causative E280A mutation in the presenilin-1 gene (Lemere et al., 1996) from non-carriers of such a mutation. Notably, despite the impairment in STM binding, asymptomatic carriers and healthy controls did not differ in tasks assessing general memory, attention and executive functions. Impairments in this function are associated with lower white matter integrity in familial Alzheimer disease (Parra et al., 2015). Furthermore, this STM binding deficit is absent in non-AD dementias, such as frontotemporal, vascular, or Lewy body dementias (Della Sala et al., 2012). STM binding seems to place minimal demands on executive functions and appears to be subserved by components of the memory network impaired in AD, but not in other dementias (Della Sala et al., 2012). On the other hand, working memory deficits have been reported in epilepsy patients. For example, temporal lobe epilepsy typically has disabling effects on verbal memory functions. Furthermore, juvenile myoclonic epilepsy and benign epilepsy in children are associated with impaired performance on visual working memory tasks (Elger et al., 2004).


STM binding yields specific activation increases across neural generators that collectively support temporary visual memory for isolated and integrated features. Within the left hemisphere, binding of object features mainly engages the superior and inferior parietal cortex, the fusiform gyrus and the dorsal premotor cortex (Parra et al., 2014). Moreover, research on the temporal dynamics of STM-relevant networks suggest an early role of posterior regions in binding processes (Smith et al., 2017).

Following this evidence, we inferred that stimulation of the posterior parietal cortex (PPC) might offer new opportunities to approach working memory deficits in AD, especially at early stages. However, no study has yet demonstrated a causal

role of such a region in STM binding. A unique approach to define necessary hubs in brain networks and infer reliable mechanisms in cognitive neuroscience consists in applying direct intracranial electrical stimulation to single subjects (epilepsy patients implanted with depth electrodes) in specific brain regions to causally modulated cognitive task performance. Here, as part of an ongoing program of research (Chennu et al., 2013; Canales-Johnson et al., 2015; Hesse et al., 2016), we profited from the unique opportunity to produce such causal evidence via single-subject direct intracranial electrical stimulation (Parvizi et al., 2012; Mégevand et al., 2014). Specifically, we stimulated the left parietal cortex (precuneus) during the visual STM task in a patient implanted with intracranial electrodes, and compared the results with a non-stimulated condition and to the performance of a matched control group (for details about the participant and the controls, see "Participants" Section; intracranial recording specifications are offered in "Subject's Intracranial Recording" Section). The patient presented a specific pattern of working memory deficits (with impairments in the binding condition but not in the shape condition) resembling the typical performance of AD patients on this task (Parra et al., 2009, 2010). This allowed us to test the selective and specific effect observed in the binding condition alone, as a relevant model mirroring the pattern which characterizes AD.

We used the visual working memory task developed by Parra and colleagues (Parra et al., 2010, 2015; Pietto et al., 2016). This task involves two sets of stimuli, one with black shapes, used to evaluate memory for shapes; and a second one with colored shapes, assessing shape-color binding in working memory (see **Figure 1**). The subject performed the task with and without stimulation. During task performance we directly stimulated the left precuneus with 1 mA at 50 Hz in 2-s intervals (stimulation block). In the control condition (sham block) the patient continued the task under a simulated stimulation (for details see "Experimental Design and Stimuli" and "Stimulation Protocol" Sections, respectively). The subject was not able to distinguish between stimulation and sham blocks.

Direct stimulation of the parietal cortex induced a selective improvement in STM binding. Relative to controls ("Behavioral Analysis" Section), the subject only reached normal STM binding performance upon stimulation (Crawford's $t = 0.20$, $p = 0.84$, $Z_{ccc} = 0.21$; **Figure 2A**). In addition, compared with the SHAM condition, electrical stimulation during the binding condition significantly enhanced performance (Crawford's $t = -2.93$, $p = 0.02$, $Z_{ccc} = -3.31$; **Figure 2A**). Moreover, the subject's performance in the shapes condition did not significantly differ from that of controls, in any of the treatments (PPC: Crawford's $t = 1.33$, $p = 0.23$, $Z_{ccc} = -1.42$; SHAM: Crawford's $t = 2$, $p = 0.08$, $Z_{ccc} = 2$; **Figure 2A**). To assess the impact and spread of the electrical stimulation, we compared intracranial event-related activity in all the recording electrodes between stimulated and SHAM trials (**Figure 2B**; for data pre-processing and processing see "Signal Preprocessing and Data Quality" Section, respectively). Propagation



however, important limitations. While we have accounted for the known caveats of intracranial EEG recordings by adopting several measures (see “Signal Preprocessing and Data Quality” Section), future studies could further test our conclusions while circumventing method-specific limitations.

As reported in the pioneering work (Penfield and Boldrey, 1937), electrical stimulation has been used both for clinical purposes and to causally probe brain mechanisms. The evidence for electrical currents to spread through white matter along functionally well-defined brain networks (Duffau et al., 2005; Tolia et al., 2005) supports our result that a wide network

underlies specific visual working memory mechanisms. Direct cortical stimulation becomes much more than a blunt tool to modulate a simple area, and turns into a unique causal instrument to probe into the process of the investigated function.

Finally, given the usefulness of the STM binding task as a marker of prodromal AD, and alongside the potential to improve STM binding by stimulation, our results open a new area of research centered in the non-invasive brain stimulation of the PPC in clinical and preclinical populations. Future studies with this approach may shed light on functional restoration options for AD patients and

subjects at risk for the disease (mild cognitive impairment, MCI), paving the way for new treatments to delay the development of neurocognitive deficits associated with this pathology.

MATERIALS AND METHODS

Data Acquisition and Protocol Design

Participants

As part of an ongoing protocol (Chennu et al., 2013; Canales-Johnson et al., 2015; Hesse et al., 2016), we recruited one patient with intractable epilepsy who was offered surgical intervention to alleviate his condition. The subject was an 18-year-old, right-handed male who had completed high school and suffered from drug-resistant epilepsy since the age of six. He was attentive and cooperative throughout the task. In addition, we recruited a control group comprised of seven healthy male participants matched with the patient for age ($M = 18.57$, $SD = 1.27$; $t = -0.35$, $p = 0.70$) and years of education ($M = 12.71$, $SD = 1.25$; $t = -0.53$, $p = 0.61$). None of these subjects reported a history of psychiatric or neurological disease. This study was carried out in accordance with the recommendations of the Declaration of Helsinki, as well as the Guidelines of the Ethics Committee of INECO (approved protocol number FONCyT-PICT 2012-0412 and 2012-1309). All subjects gave written informed consent in accordance with the Declaration of Helsinki.

Subject's Intracranial Recording

Direct cortical recordings were obtained from semi-rigid, multi-lead electrodes implanted in the patient's brain. The electrodes were 0.8 mm in diameter and consisted of 5, 10 or 15 2-mm wide contact leads placed 1.5 mm apart from each other (DIXI Medical Instruments). We used a Micromed video-SEEG monitoring system which records as many as 128 depth-EEG electrode sites simultaneously. Recordings were obtained from 127 sites and sampled at 512 Hz. The data were collected from the precuneus, the cuneus, the hippocampus, the posterior cingulate and the postcentral gyrus. The recordings obtained were distal to the epileptogenic foci, and no single recording site presented epileptogenic activity (see below). We also obtained post-implantation MRI and CT scans were obtained from each patient. Both volumetric images were affine registered and normalized on the SPM8 MATLAB toolbox. Using MRIcron, we established the coordinates of each contact site and their respective Brodmann areas.

Experimental Design and Stimuli

The task assessed memory for shapes and combinations of shapes and colors. Stimuli were randomly selected from a set of eight shapes and eight colors and presented as individual features or as features combined into integrated objects. Each type of stimulus was presented in a separate condition. Two experimental conditions were used, each consisting of 32 test trials, leading to a total of 64 test trials. Trials were fully

randomized and, for the control participants, conditions were delivered in a counter-balanced order. In the "Shapes" condition, arrays of shapes were presented in the study display. In the test display for the "different" trials, two new shapes from the study array were replaced with two new shapes (Figure 1). Hence, in these conditions, participants were required to detect changes in individual features. In the "Shape-color binding" condition, the study display showed combinations of shapes and colors. In the test display for "different" trials, two shapes swapped the colors in which they had been shown in the study display (Figure 1). Hence, detection of this change relied on shape-color bindings. No shape or color was repeated within a given array. Fifty percent of the test trials were "same" trials (the study and test displays presented identical items) and 50% were "different" trials.

Stimulation Protocol

Electrical stimulation was delivered in bipolar square waves between two adjacent electrode contacts in the left precuneus, 54 ($-4 -60 56$, MNI coordinates) 55 ($-2 -60 56$, MNI coordinates). Stimulation occurred at 1 mA for the real stimulation condition and at 0 mA for the control SHAM condition using a 200 ms pulse width at a frequency of 50 Hz, during 2 s. Each condition involved 10 trials under real stimulation and another 10 under SHAM stimulation. The stimulation began at the onset of the study display and continued through the test display. First we performed the real and SHAM stimulation (in that order) for the shape-color binding condition and then for the shape-only condition. EEG signals were simultaneously monitored before and after discharges. Electrodes and trials compromised by seizures or leading to epileptic activity were excluded. The subject was asked to describe any perceptual or physical changes he experienced during or after each stimulation trial.

Data Analysis

Behavioral Analysis

As in previous studies (Parra et al., 2010, 2015; Pietto et al., 2016), recognition during the visual working memory task was calculated by subtracting the proportion of false alarms from the hits. The subject's indexes were compared to those of a control sample through a modified two-tailed t test (Crawford and Garthwaite, 2002; Crawford et al., 2009, 2011). This methodology allows the assessment of significance by comparing test scores of one or several individuals with norms derived from small samples. This modified test is robust for non-normal distributions, presents low values of type I error, and has already been reported in single case studies (Straube et al., 2010). The alpha level was set at $p < 0.05$.

Signal Preprocessing and Data Quality

Several measures were adopted to circumvent the caveats of data obtained from an epileptic patient. We excluded channels in epileptogenic foci, used stringent inclusion criteria for the remaining channels, and ensured the

absence of neuroanatomical abnormalities or major cognitive deficits in the subject. We discarded the contact sites that presented pathological waveforms. Electrodes with epileptic activity were discarded upon visual identification by two professional neurologists (MCG and JA). Moreover, we discarded channels whose values exceeded five times the signal's mean and/or consecutive signal samples exceeding five standard deviations (SD) from the gradient's mean (Chennu et al., 2013; Hesse et al., 2016). A total of 83 contact sites remained after applying these criteria, and all of them were processed. Then, the data were referenced to the mean value of the non-stimulated sites (averages of such sites were subtracted from each recording). Finally, the data were segmented into 2000 ms epochs, including a -200 to 0 ms pre-stimulus baseline period. The epochs were baseline corrected.

REFERENCES

- Baddeley, A. (2000). The episodic buffer: a new component of working memory? *Trends Cogn. Sci.* 4, 417–423. doi: 10.1016/s1364-6613(00)01538-2
- Canales-Johnson, A., Silva, C., Huepe, D., Rivera-Rei, Á., Noreika, V., Garcia Mdel, C., et al. (2015). Auditory feedback differentially modulates behavioral and neural markers of objective and subjective performance when tapping to your heartbeat. *Cereb. Cortex* 25, 4490–4503. doi: 10.1093/cercor/bhv076
- Chennu, S., Noreika, V., Gueorguiev, D., Blenkman, A., Kochen, S., Ibáñez, A., et al. (2013). Expectation and attention in hierarchical auditory prediction. *J. Neurosci.* 33, 11194–11205. doi: 10.1523/JNEUROSCI.0114-13.2013
- Crawford, J. R., and Garthwaite, P. H. (2002). Investigation of the single case in neuropsychology: confidence limits on the abnormality of test scores and test score differences. *Neuropsychologia* 40, 1196–1208. doi: 10.1016/s0028-3932(01)00224-x
- Crawford, J. R., Garthwaite, P. H., and Howell, D. C. (2009). On comparing a single case with a control sample: an alternative perspective. *Neuropsychologia* 47, 2690–2695. doi: 10.1016/j.neuropsychologia.2009.04.011
- Crawford, J. R., Garthwaite, P. H., and Ryan, K. (2011). Comparing a single case to a control sample: testing for neuropsychological deficits and dissociations in the presence of covariates. *Cortex* 47, 1166–1178. doi: 10.1016/j.cortex.2011.02.017
- Della Sala, S., Parra, M. A., Fabi, K., Luzzi, S., and Abrahams, S. (2012). Short-term memory binding is impaired in AD but not in non-AD dementias. *Neuropsychologia* 50, 833–840. doi: 10.1016/j.neuropsychologia.2012.01.018
- Duffau, H., Gatignol, P., Mandonnet, E., Peruzzi, P., Tzourio-Mazoyer, N., and Capelle, L. (2005). New insights into the anatomo-functional connectivity of the semantic system: a study using cortico-subcortical electrostimulations. *Brain* 128, 797–810. doi: 10.1093/brain/awh423
- Elger, C. E., Helmstaedter, C., and Kurthen, M. (2004). Chronic epilepsy and cognition. *Lancet Neurol.* 3, 663–672. doi: 10.1016/s1474-4422(04)00906-8
- Gazzaley, A., and Nobre, A. C. (2012). Top-down modulation: bridging selective attention and working memory. *Trends Cogn. Sci.* 16, 129–135. doi: 10.1016/j.tics.2011.11.014
- Hesse, E., Mikulan, E., Decety, J., Sigman, M., Garcia Mdel, C., Silva, W., et al. (2016). Early detection of intentional harm in the human amygdala. *Brain* 139, 54–61. doi: 10.1093/brain/awv336
- Jacobs, H. I., Radua, J., Lückmann, H. C., and Sack, A. T. (2013). Meta-analysis of functional network alterations in Alzheimer's disease: toward a network biomarker. *Neurosci. Biobehav. Rev.* 37, 753–765. doi: 10.1016/j.neubiorev.2013.03.009
- Karvigh, S. A., Motamedi, M., Arzani, M., and Roshan, J. H. N. (2017). HD-tDCS in refractory lateral frontal lobe epilepsy patients. *Seizure* 47, 74–80. doi: 10.1016/j.seizure.2017.03.005
- Lemere, C. A., Lopera, F., Kosik, K., Lendon, C., Ossa, J., Saido, T., et al. (1996). The E280A presenilin 1 Alzheimer mutation produces increased *Ab42* deposition and severe cerebellar pathology. *Nat. Med.* 2, 1146–1150.
- Mégevand, P., Groppe, D. M., Goldfinger, M. S., Hwang, S. T., Kingsley, P. B., Davidesco, I., et al. (2014). Seeing scenes: topographic visual hallucinations evoked by direct electrical stimulation of the parahippocampal place area. *J. Neurosci.* 34, 5399–5405. doi: 10.1523/JNEUROSCI.5202-13.2014
- Parra, M. A., Abrahams, S., Fabi, K., Logie, R., Luzzi, S., and Della Sala, S. (2009). Short-term memory binding deficits in Alzheimer's disease. *Brain* 132, 1057–1066. doi: 10.1093/brain/awp036
- Parra, M. A., Abrahams, S., Logie, R. H., Méndez, L. G., Lopera, F., and Della Sala, S. (2010). Visual short-term memory binding deficits in familial Alzheimer's disease. *Brain* 133, 2702–2713. doi: 10.1093/brain/awQ158
- Parra, M. A., Della Sala, S., Logie, R. H., and Morcom, A. M. (2014). Neural correlates of shape-color binding in visual working memory. *Neuropsychologia* 52, 27–36. doi: 10.1016/j.neuropsychologia.2013.09.036
- Parra, M. A., Saarimäki, H., Bastin, M. E., Londoño, A. C., Pettit, L., Lopera, F., et al. (2015). Memory binding and white matter integrity in familial Alzheimer's disease. *Brain* 138, 1355–1369. doi: 10.1093/brain/awv048
- Parvizi, J., Jacques, C., Foster, B. L., Withoft, N., Rangarajan, V., Weiner, K. S., et al. (2012). Electrical stimulation of human fusiform face-selective regions distorts face perception. *J. Neurosci.* 32, 14915–14920. doi: 10.1523/JNEUROSCI.2609-12.2012
- Penfield, W., and Boldrey, E. (1937). Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain* 60, 389–443. doi: 10.1093/brain/60.4.389
- Pietto, M., Parra, M. A., Trujillo, N., Flores, F., García, A. M., Bustin, J., et al. (2016). Behavioral and electrophysiological correlates of memory binding deficits in patients at different risk levels for Alzheimer's disease. *J. Alzheimers Dis.* 53, 1325–1340. doi: 10.3233/JAD-160056
- Smith, K., Ricaud, B., Shahid, N., Rhodes, S., Starr, J. M., Ibáñez, A., et al. (2017). Locating temporal functional dynamics of visual short-term memory binding using graph modular dirichlet energy. *Sci. Rep.* 7:42013. doi: 10.1038/srep42013
- Straube, T., Weisbrod, A., Schmidt, S., Raschdorf, C., Preul, C., Mentzel, H.-J., et al. (2010). No impairment of recognition and experience of disgust in a patient with a right-hemispheric lesion of the insula and basal ganglia. *Neuropsychologia* 48, 1735–1741. doi: 10.1016/j.neuropsychologia.2010.02.022

AUTHOR CONTRIBUTIONS

AB and AI designed the study. AB, EH, LS, EPM, MdCG, JA, FA, AL, TAB, MZ, MP, AMG and AI carried out and analyzed the experiments. AB and AI wrote the article. AI and AB conceived the study and wrote the final article, together with the other authors. All authors have approved the manuscript.

ACKNOWLEDGMENTS

Partial support was obtained from grants by CONICET; CONICYT/FONDECYT Regular (1170010); FONCYT-PICT (2012-0412 and 2012-1309), FONDAP (15150012); and the INECO Foundation. MAP work is supported by Alzheimer's Society, Grant ASSF-14-008.

- Tekturk, P., Erdogan, E. T., Kurt, A., Vanli-yavuz, E. N., Ekizoglu, E., Kocagoncu, E., et al. (2016). The effect of transcranial direct current stimulation on seizure frequency of patients with mesial temporal lobe epilepsy with hippocampal sclerosis. *Clin. Neurol. Neurosurg.* 149, 27–32. doi: 10.1016/j.clineuro.2016.07.014
- Tolias, A. S., Sultan, F., Augath, M., Oeltermann, A., Tehovnik, E. J., Schiller, P. H., et al. (2005). Mapping cortical activity elicited with electrical microstimulation using fMRI in the macaque. *Neuron* 48, 901–911. doi: 10.1016/j.neuron.2005.11.034
- Treisman, A., and Zhang, W. (2006). Location and binding in visual working memory. *Mem. Cognit.* 34, 1704–1719. doi: 10.3758/bf03195932
- Tulving, E. (2002). Episodic memory: from mind to brain. *Annu. Rev. Psychol.* 53, 1–25. doi: 10.1146/annurev.psych.53.100901.135114
- von der Malsburg, C. (1999). The what and why of binding: the modeler's perspective. *Neuron* 24, 95–104. doi: 10.1016/S0896-6273(00)80825-9
- Zimmer, H. D., Mecklinger, A., and Lindenberger, U. (Eds) (2006). “Levels of binding: types, mechanisms, and functions of binding in remembering,” in *Handbook of Binding and Memory: Perspectives from Cognitive Neuroscience*, (Oxford: Oxford University Press), 3–22.
- Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Birba, Hesse, Sedeño, Mikulan, García, Ávalos, Adolphi, Legaz, Bekinschtein, Zimmerman, Parra, García and Ibáñez. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.